

1 **Full title**

2 The association between limiting longstanding illness and serious psychological distress in
3 adolescents: A secondary analysis of the UK Millennium Cohort Study

4

5 **Short title**

6 Limiting longstanding illness and serious psychological distress in UK adolescents

7

8 Gareth M. Palliser^{1*}, Lorna K. Fraser², Kate E. Mooney^{1,3}, and Stuart W. Jarvis¹

9

10 ¹ Department of Health Sciences, University of York, York, UK

11 ² Cicely Saunders Institute, King's College London, London, UK

12 ³ Better Start Bradford Innovation Hub, Bradford Institute for Health Research

13

14 *Corresponding author

15 E-mail: gareth.palliser@york.ac.uk

16

17

18

19

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

20 **Abstract**

21 **Background**

22 Previous studies have shown associations between specific limiting longstanding illnesses and
23 mental health difficulties using cross-sectional studies in the UK. This study explored the
24 association between having any limiting longstanding illness and serious psychological
25 distress or of currently receiving treatment for depression or serious anxiety at age 17 years.

26 **Methods**

27 A secondary analysis of the UK Millennium Cohort Study was conducted. Outcome measures
28 were dichotomised responses from the self-administered Kessler-6 questionnaire for
29 nonspecific psychological distress, and self-reported currently receiving treatment for
30 depression or serious anxiety. Limiting longstanding illness data and covariates were taken
31 from questionnaires with parents and adolescents, from birth up to age 17 years. Data were
32 analysed using multiple binary logistic regression, first using complete-case analysis, and then
33 using multiple imputation using chained equations.

34 **Results**

35 Adolescents with a history of limiting longstanding illness were at an increased odds of both
36 serious psychological distress (odds ratio = 1.53, 95%CI = 1.27-1.86) and self-reporting
37 currently receiving treatment for depression or serious anxiety at age 17 years (odds ratio =
38 3.02, 95%CI 2.24-4.07).

39 **Conclusion**

40 Children and young people with a limiting longstanding illness are at increased risk of having
41 serious psychological distress, depression, and serious anxiety. Practitioners should be aware
42 of this and routine screening for psychological distress, plus additional preventative support,
43 may be beneficial.

44

45 Introduction

46 Adolescence marks the period between childhood and adulthood, representing a key period
47 in both physical and psychological development. National UK survey data found a prevalence
48 of probable mental illness of 17.6% in those aged 6-16 years old in England^[1], compared with
49 16.7% in adults in similar surveys ^[2]. Recently, mental health disorders for children and
50 adolescents have become a public health crisis in England. Child referrals to mental health
51 services more than doubled between 2020 and 2021^[3], and one in every five children aged 8-
52 16 years old now experiences a mental health disorder^[4]. A previous analysis of US national
53 survey data ^[5] found that half of lifetime mental health disorders in the Diagnostic and
54 Statistical Manual of Mental Disorders (DSM-IV)^[6] start by 14 years of age, and three-quarters
55 occur by 24 years. Adolescent mental illness is therefore a critical period for research. Mental
56 illness experienced in adolescence is associated with significant potential harms including
57 lower educational achievement ^[7], alcohol and substance misuse ^[8], social isolation ^[9], and
58 suicide and self-inflicted injury ^[10].

59 Longstanding illness is defined by The UK Department of Health ^[11] as any illness or condition
60 which cannot be cured, but can instead be treated with pharmacological or other treatments;
61 it is limiting if it reduces the ability of someone to take part in normal activities. As medical
62 care improves, childhood mortality rates in those aged 1 to 15 years in England and Wales
63 have decreased from 33 deaths per 100,000 in 1981 to 8 deaths per 100,000 in 2021^[12]. This,
64 amongst other factors including a rise in sedentary lifestyles, has resulted in a global rise in
65 the proportion of those surviving to live with a limiting longstanding illness (LLSI) in all age
66 groups, including children and adolescents^[13]. Adolescents with a limiting longstanding illness
67 (LLSI) may face difficulties such as hospital appointments, disruptions to their routine and

68 education, and reduced capability to take part in regular activities; these challenges could
69 lead to mental health difficulties^[14]. It is important to understand any association between
70 physical and mental illnesses to inform prevention strategies and paediatric service delivery,
71 and to understand whether the early provision of psychological support may be necessary for
72 all children and adolescents with a chronic illness.

73 Previous cross-sectional and longitudinal studies have identified associations between LLSIs
74 and mental health difficulties in children outside of the UK^[15–17]. Systematic reviews ^[18–20]
75 have identified that in adolescents, studies are largely constrained to identifying associations
76 between diabetes or asthma and mental illness, with some evidence suggesting higher
77 prevalence of mental health difficulties in those with inflammatory bowel disease, cystic
78 fibrosis, and sickle cell disease. Further, few studies exploring longitudinal relationships
79 between longstanding illnesses and mental health difficulties were at a low risk of
80 methodological bias.

81 An analysis of a UK-based survey (n=9,834) found that children and adolescents aged 5-15
82 years old with asthma were at a slightly increased odds of anxiety (OR=1.38, 95%CI 1.05-1.82)
83 and hyperkinetic disorders (OR=1.64, 95%CI 1.00-2.39) when compared with those without
84 asthma, measured using the Development and Well-Being Assessment^[21]. Children with
85 asthma who had parent-rated poor health were at significantly increased odds of scoring as
86 borderline or abnormal for the parent (OR 3.1, 95%CI 2.4-4.1), teacher (OR 1.7, 95%CI 1.2-
87 2.4), and child- (OR 1.9, 95%CI 1.2-3.0) rated total difficulties score on the Strengths and
88 Difficulties Questionnaire, when compared with children without asthma who were in good
89 health. Those with asthma but were in good health were not at a significantly increased odds

90 of increased score on the Strengths and Difficulties Questionnaire, indicating that increased
91 symptom severity in children with asthma may lead to increased risk of mental health
92 difficulties.

93 One UK study (n=14,775) using a secondary analysis of the Avon Longitudinal Study of Parents
94 and Children found that those with a chronic illness were at an (OR 1.6, 95%CI 1.14-2.25)
95 increased odds of having mental illness as measured by the Development and Well-Being
96 Assessment at aged 15 years, compared with those without a chronic illness^[22]. However,
97 participants in this study were recruited from a single region in southwestern England, with
98 those from disadvantaged or ethnic minority backgrounds being under-represented, and
99 those from white ethnicity and advantaged backgrounds were over-represented.

100 As studies in adults^[23,24] have previously identified a bi-directional relationship between LLSI
101 and mental health difficulties seen in other studies, this must be considered when
102 interpreting estimates from cross-sectional studies. This study therefore explored the
103 association between having any LLSI before aged 17 years with serious psychological distress
104 at age 17 years in a large representative UK birth cohort- the Millennium Cohort Study. This
105 study also investigated the relationship between LLSI and receiving treatment for depression
106 or serious anxiety.

107

108 **Methods**

109 **Data source**

110 This study uses a retrospective secondary analysis of data from the Millennium Cohort Study,
111 a nationally representative UK birth cohort consisting of 19,517 cohort members and their
112 families. Cohort members were recruited from the Child Benefit Register at age 9 months in
113 data collection sweep 1 between the years 2000-2002, with some participants whose
114 geographical data was unavailable recruited in sweep 2 at age 3 years ^[25]. A stratified sampling
115 design was used, recruiting additional participants from areas with higher rates of minority
116 ethnic groups and deprivation than the UK average. There are currently 7 sweeps of data;
117 most recently when cohort members were aged 17 years ^[25]. Parents provided consent to
118 take part in the study on behalf of cohort members up to age 11 years; from 14 years onwards,
119 cohort members provided consent themselves.

120 **Outcome measures**

121 Cohort members completed the Kessler-self-administered 6-item questionnaire (K6) at age
122 17 years. The scale is designed to measure non-specific psychological distress, and is scored
123 from 0-24, with a cut-point of ≥ 13 indicating serious distress used frequently throughout the
124 literature ^[26-28]. All participants with scores between 0-12 were coded as not having
125 psychological distress, and participants with scores between 12-24 were considered as having
126 serious psychological distress^[28]. The K6 has an area under the receiver operating
127 characteristic curve (AUROC) of 86.5%, compared with 85.4% in the 10-item version of the
128 Kessler scale (K10), indicating the K6 performs slightly better than the K10 at distinguishing

129 between those with and without serious mental illness^[29]. The scale demonstrates high
130 internal consistency, with Cronbach's alpha estimates across varying languages, countries,
131 and settings ranging from 0.81 to 0.89 in a single factor structure^[29–32].

132 At age 17, cohort members were asked 'Has a doctor ever told you that you suffer from
133 depression or serious anxiety?'. Where cohort members responded with 'yes' they were
134 further asked 'are you currently being treated for depression or serious anxiety?'. Those who
135 answered 'yes' to both questions were considered to have an LLSI for the purposes of this
136 study. This enabled some exploration of contact with healthcare services and treatment.

137 **Exposure**

138 LLSI status was gathered on behalf of cohort members from parent/main carer interviews,
139 between the ages of 3 and 14 years. The parent or main carer of each cohort member was
140 asked 'Does [cohort member] have a longstanding illness or condition?' at each sweep; for
141 those who answered 'yes' they were further asked 'Does this illness limit [cohort member]'s
142 activity in any way?'. For the purposes of this study, respondents whose parent or main carer
143 answered 'yes' to both questions at any sweep were considered to have an LLSI, and
144 respondents whose parents answered no to both, or no to the second question were
145 considered to not have an LLSI, and were used as the reference group. At age 11 and 14 years,
146 participants who answered 'yes' were also asked which health conditions affect the cohort
147 member; those who reported being affected exclusively by a mental health condition, and
148 not by any other condition, were not considered to have an LLSI.

149

150 Covariates

151 All covariates were taken from face-to-face interviews with each cohort member's parent or
152 main carer, at age 9 months for those in the original sample, and age 3 years for those in the
153 booster sample. Covariates selection for this study was informed by previous studies; all
154 covariates, are theorised to confound the relationship between LLSI and serious psychological
155 distress^[33–36] [see Figure 1]. Covariates were cohort member sex (coded as 0=male,
156 1=female), ethnic group (coded as 0=White British (reference group), 1=Pakistani and
157 Bangladeshi, 2= Black or Black British, 3=Mixed or Other Ethnic group), family structure
158 (coded as 0=dual-parent family, 1=single-parent family), being above or below the 60%
159 median on the Organisation for Economic Co-operation and Development equivalised income
160 scale (coded as 0=above median, 1=below median), highest parental National Vocational
161 Qualification (NVQ) or equivalent (coded as 0=none or overseas qualification, 1=NVQ level 1,
162 2=NVQ level 2, 3=NVQ level 3, 4=NVQ level 4, 5=NVQ level 5) parental self-reported LLSI
163 (coded as 0=no LLSI, 1=LLSI present), and parental K6 score (continuous)^[33–36].

164 [INSERT FIGURE 1 HERE].

165 Figure 1: Directed acyclic graph of relationships between measured variables in this study

166

167 Note: arrows between confounders have been removed to simplify the presentation of the
168 DAG

169

170 **Statistical analyses**

171 For the primary outcome of psychological distress measured by K6 score, complete-case
172 analyses were used. Complete case analyses were also used for the secondary outcome of
173 whether a cohort member reported currently receiving treatment for depression or serious
174 anxiety. To minimise attrition, participants with LLSI data for at least one sweep were
175 included, providing that all covariate data were available. Where there was more than one
176 cohort member in a family, the second cohort member was excluded to prevent clustering by
177 household. The MCS used a stratified sampling design, over-sampling from electoral wards
178 with higher levels of ethnic diversity and socioeconomic deprivation in the 1991 UK Census^[37].
179 Sampling and response weights were used to account for both sampling design and attrition
180 in all analyses.

181 Binary logistic regression was used to produce crude and adjusted odds ratios (ORs) for the
182 association between LLSI and dichotomised K6 score, and between LLSI and receiving
183 treatment for depression or serious anxiety. Two-sided statistical tests using a p-value
184 threshold of <0.05 were used throughout. All covariates listed above were included in
185 adjusted models for both outcomes, using multiple logistic regression, which were compared
186 with unadjusted ORs from models using simple binary logistic regression. Additionally, a
187 sensitivity analysis for the model used for the primary outcome of K6 score was also
188 conducted using multiple imputation using chained equations (MICE) to impute missing data
189 for all variables included in the logistic regression model investigating the relationship
190 between LLSI and K6 score (see Figure 1 for numbers of missing values). All variables used in
191 the analysis model for the primary outcome were included in the imputation model. The

192 Stata command ‘mi impute chained’ was used to impute 20 datasets. This was used to assess
193 how the use of complete-case analyses influenced model estimates. All analyses were
194 conducted in Stata Version 17.

195 **Results**

196 Of the 19,482 cohort members in end-user license agreement dataset, 7,866 cohort members
197 provide complete exposure, outcome, and covariate data, representing 40.4 % of the original
198 sample.

199 Figure 2: Attrition diagram showing those included in complete-case analyses

200 [INSERT FIGURE 2 HERE]

201 Unweighted cohort member characteristics for the complete-case dataset are presented in
202 Table 1; weighted figures are available in Appendix S1. 15.8% of cohort members scored
203 above the threshold for serious distress on the K6, with 21.9% of participants who reported
204 LLSI before age 17 years scoring above the cutoff, compared with 14.5% in the unexposed
205 group. Greater proportional differences were seen in receiving treatment for depression or
206 serious anxiety; 9.7% in those with reported LLSI, compared to 3.5% in those without. Males
207 had higher prevalence of LLSI, with 53.6% of those reporting an LLSI being male. Those with a
208 reported LLSI also saw slightly higher prevalence of LLSI in proportions of those from single-
209 parent families, those below the 60% median equivalised household income threshold, and
210 those from the White ethnic group, although the study population is also overwhelmingly
211 white at 88.1%.

212

213 **Table 1: Unweighted summary statistics of covariates by LLSI status in complete case dataset**

Variable	No LLSI (%)	LLSI (%)	Total (%)
Categorical variables			
Sex			
Female	3,369 (51.9)	640 (46.4)	4,009 (51.0)
Male	3,119 (48.1)	738 (53.6)	3,857 (49.0)
Ethnic group			
White	5,696 (87.8)	1,227 (89.0)	6,923 (88.0)
Indian	135 (2.1)	28 (2.0)	163 (2.0)
Pakistani and Bangladeshi	267 (4.1)	50 (3.6)	317 (4.0)
Black or Black British	159 (2.5)	29 (2.1)	188 (2.4)
Mixed or Other Ethnic group	231 (3.6)	44 (3.2)	275 (3.5)
Family structure			
Two parents/carers	5,769 (88.9)	1,160 (84.2)	6,929 (88.1)
Single parent/carer	719 (11.1)	218 (15.8)	937 (11.9)
Parental education			
None or overseas qualification	642 (9.9)	157 (11.4)	799 (10.2)
NVQ level 1	459 (7.1)	114 (8.3)	573 (7.3)
NVQ level 2	1,825 (28.1)	417 (30.3)	2,242 (28.5)
NVQ level 3	990 (15.3)	212 (15.4)	1,202 (15.3)
NVQ level 4	2,258 (34.8)	423 (30.7)	2,681 (34.1)
NVQ level 5	314 (4.8)	55 (4.0)	369 (4.7)
Parental limiting longstanding illness			
No	5,217 (80.4)	967 (70.2)	6,184 (78.6)
Yes	1,271 (19.6)	411 (29.8)	1,682 (21.4)
OECD equivalised income			
Above 60% median	4,879 (75.2)	909 (66.0)	5,788 (73.6)
Below 60% median	1,609 (24.8)	469 (34.0)	2,078 (26.4)
Continuous variables			
	No LLSI	LLSI	Total
	mean (SD)	mean (SD)	mean (SD)
Parental Kessler-6 score	2.96 (3.40)	3.93 (4.20)	3.13 (3.57)

214

215 **LLSI and serious psychological distress**

216 21.9% (302) of those with an LLSI scored above the threshold for serious psychological distress
 217 on the Kessler-6 compared with 14.5% (938) of those without an LLSI. The crude OR for serious
 218 psychological distress at age 17 years in those with LLSI compared to those without was 1.53
 219 (95%CI 1.29-1.83). The weighted adjusted OR was also 1.53 (95%CI 1.27-1.86). Unweighted
 220 complete-case analyses can be seen in Appendix S2. Models using MICE to account for missing
 221 data retained 19,229 cohort members and produced an adjusted OR of 1.58 (95%CI 1.36-
 222 1.84), which was similar to the estimates from complete-case analyses (Appendix S3).

223 **Table 2: Weighted complete-case multiple logistic regression of dichotomised Kessler-6 score**

Covariate	Weighted adjusted OR	95% CI	P-value
Limiting longstanding illness			
No limiting longstanding illness	1.00		(ref)
Limiting longstanding illness	1.53	1.27-1.86	<0.001
Sex			
Female	1.00		
Male	0.37	0.31-0.45	<0.001
Ethnic group			
White	1.00		(ref)
Indian	0.43	0.22-0.84	0.013
Pakistani/ Bangladeshi	0.59	0.37-0.94	0.025
Black/ Black British	1.07	0.51-2.24	0.854
Mixed/ Other	0.89	0.52-1.52	0.670
Family structure			
Dual parents/carers	1.00		(ref)
Single parent/carer	1.10	0.61-1.49	0.561
Parental education			
None or overseas only	1.00		(ref)
NVQ level 1	1.02	0.61-1.70	0.942
NVQ level 2	0.99	0.71-1.40	0.968
NVQ level 3	1.00	0.69-1.47	0.983
NVQ level 4	0.94	0.66-1.33	0.733
NVQ level 5	0.94	0.60-1.34	0.779
OECD equivalised income			
Above 60% median	1.00		(ref)
Below 60% median	1.17	0.97-1.41	0.108
Parental limiting longstanding illness			
No	1.00		(ref)
Yes	1.08	0.89-1.30	0.423
Parental Kessler-6 score	1.04	1.02-1.07	<0.001
Constant	0.22	0.15-0.31	<0.001

224 *Cutoff Kessler-6 score of ≥ 13 used to indicate serious psychological distress*

225 **LLSI and treatment for depression or serious anxiety**

226 9.7% (134) of those with an LLSI reported currently receiving treatment for depression or
 227 serious anxiety, compared with just 3.5% (225) in those without an LLSI. The crude OR for
 228 children and adolescents who reported currently receiving treatment for depression or
 229 serious anxiety by age 17 years in those with an LLSI compared to those without was 3.18

230 (95%CI 2.40-4.22), and the adjusted value was 3.02 (95%CI 2.24-4.07). Unweighted complete-
 231 case analyses can be seen in Appendix S4.

232 Table 3: Weighted complete-case logistic regression of currently receiving treatment for
 233 depression or serious anxiety

Covariate			
	Weighted adjusted OR	95% CI	P-value
Limiting longstanding illness			
No limiting longstanding illness	1.00		(ref)
Limiting longstanding illness	3.02	2.24-4.07	<0.001
Sex			
Female	1.00		
Male	0.29	0.21-0.39	<0.001
Ethnic group			
White	1.00		(ref)
Indian	0.67	0.16-2.71	0.572
Pakistani/ Bangladeshi	0.32	0.12-0.86	0.025
Black/ Black British	0.48	0.14-1.64	0.241
Mixed/ Other	0.22	0.07-0.70	0.010
Family structure			
Dual parents/carers	1.00		(ref)
Single parent/carer	0.79	0.46-1.36	0.398
Parental education			
None or overseas only	1.00		(ref)
NVQ level 1	1.28	0.69-2.38	0.435
NVQ level 2	0.80	0.47-1.33	0.384
NVQ level 3	0.61	0.35-1.08	0.087
NVQ level 4	0.71	0.43-1.17	0.173
NVQ level 5	0.97	0.49-1.90	0.921
OECD equivalised income			
Above 60% median	1.00		(ref)
Below 60% median	1.09	0.76-1.55	0.646
Parental limiting longstanding illness			
No	1.00		(ref)
Yes	1.35	0.98-1.86	0.069
Parental Kessler-6 score			
	1.06	1.02-1.09	0.003
Constant	0.06	0.04-0.09	<0.001

234

235 Discussion

236 This was the first UK-wide longitudinal study investigating the relationship between LLSI and
 237 psychological distress. This study found that children and adolescents with a known history

238 of LLSI had an odds of 1.53 having serious psychological distress at age 17, when compared
239 with those without an LLSI. Those with a known history of LLSI had an odds of 3.02 for
240 reporting currently receiving treatment for depression or serious anxiety by a healthcare
241 professional at age 17 years, when compared with those without an LLSI.

242 The weighted prevalence of LLSI in this study of 15.7% (Appendix S1) is greater than that seen
243 in national survey data for children aged 0-15 years in a comparable time period such as The
244 Health Survey for England at 8% [38]. The use of a longitudinal measure of LLSI in this study,
245 which placed those with any history of self-reporting an LLSI in the exposure group may be
246 responsible for some of this difference. Varying age ranges, areas, and definitions of LLSI
247 across studies also make between-studies comparison of LLSI prevalence difficult, with other
248 England surveys showing higher rates at 23% [39].

249 All LLSI data were taken before both outcome measurements at age 17 years. Those who
250 reported having an LLSI at age 11-14 years, but self-categorised this as only having a mental
251 health condition with no other type of LLSI were excluded from the exposure group for the
252 purposes of this study. However, some participants were missing data at these sweeps, so it
253 is not possible to assess the mental health status of some participants between the ages of
254 11-14 years. The longitudinal design of the current study therefore does facilitate some causal
255 interpretation to indicate that LLSI may increase the risk of serious psychological distress, and
256 of receiving treatment for depression or serious anxiety at age 17 years. Mediation analyses
257 and assessment of dose-response relationships, such as severity or duration of LLSI in other
258 populations could further explore this.

259

260 This study identified a greater proportion of adolescents with recent serious nonspecific
261 psychological distress (15.8%) than those who were undergoing treatment for depression or
262 serious anxiety (4.6%). This could indicate underdiagnosis of depression and serious anxiety,
263 although some cohort members could have experienced psychological distress without
264 meeting the threshold for treatment for depression or serious anxiety. It is important for
265 healthcare practitioners to be aware of the increased psychological distress in adolescents
266 with an LLSI. A separate pilot study implemented routine screening and an early stepped
267 intervention pathway into a paediatric epilepsy clinic and was able to identify children with
268 previously unidentified mental health difficulties and enable them to appropriately access
269 care ^[46]. In another study, routine depression screening in adolescents with type 1 diabetes
270 mellitus at quarterly appointments in a Cincinnati Children's Hospital was able to detect
271 depression symptoms early and create opportunities for early intervention ^[47]. As the current
272 study identified increased odds of serious psychological distress or of receiving treatment for
273 depression or serious anxiety, it may be beneficial to screen for psychological distress and
274 provide early interventions more broadly in adolescents with any LLSI, to facilitate early
275 detection of mental health difficulties. Future studies could investigate the relationship
276 between contact with health professionals and diagnosis and treatment of mental health
277 difficulties to understand this further, and to improve access to mental health support in
278 young people.

279 The OR for receiving treatment for serious depression or anxiety (3.02) was greater than when
280 measuring serious psychological distress using the K6 (1.53). As LLSIs often require treatment
281 or regular monitoring, adolescents with an LLSI may have increased contact with healthcare
282 professionals when compared with their peers. This may lead to increased opportunity for

283 healthcare professionals to identify mental health difficulties in children and young people
284 with LLSIs. It is possible that those without LLSIs may constitute a larger proportion of those
285 with undiagnosed mental health difficulties. Further exploration of the potential benefits of
286 screening for psychological distress outside of those with an LLSI may also be useful for the
287 early identification of mental health difficulties.

288 This study adjusted for the effects of ethnicity in all adjusted models, however, the
289 representativeness of some ethnic minority groups in the Millennium Cohort Study is limited.
290 Previous literature has shown that young people from ethnic minority and socioeconomically
291 deprived backgrounds face substantial barriers to accessing quality support to manage their
292 physical and mental health^[48]. Future research could examine whether the association found
293 here varies across different ethnic groups, to establish whether the association may be
294 stronger for groups who have more difficulties with managing their mental health.

295 The increased odds of serious psychological distress in those with an LLSI in this study is
296 comparable to odds ratios from the previously mentioned UK studies investigating
297 associations between chronic illness and mental illness (1.6)^[22] and asthma and mental illness
298 (1.38)^[21]. The consistency of results suggests that the risk of mental illness may be higher for
299 children with any LLSI. However, the added benefit of this study was the use of a longitudinal
300 design in a more representative UK sample, covering a broader range of LLSIs than those seen
301 in previous studies.

302 Although this study has found increased odds of mental health difficulties in adolescents with
303 an LLSI, it is important to consider the moderating and potentially neutralising effects of
304 family support, education, and management of the physical condition itself before

305 considering mental health interventions. Adolescence often marks the transition from parent-
306 to self-management of LLSI, with differences in the age this occurs seen across individuals and
307 conditions ^[40]. As a result of this and other developmental and environmental changes, non-
308 adherence to treatments across illnesses including asthma ^[41] and diabetes ^[42] is particularly
309 high in adolescents. This is concerning, as non-adherence with treatment of LLSI may
310 exacerbate symptoms or illness progression; it is possible that this could also have adverse
311 effects on mental health outcomes. The reverse may also be true – a systematic review
312 identified that adolescents self-reported that poor mental health made it more difficult to
313 adhere to treatment for physical illnesses ^[43]. Few studies appear to have examined the
314 mental health implications of improving LLSI treatment adherence in adolescents. As
315 smartphone use becomes increasingly popular, one study found that a Mobile-health app
316 which included symptom monitoring questionnaires, medication reminders, information
317 videos, and capacity to talk to pharmacists or peers improved treatment adherence in
318 adolescents with a low adherence at baseline ^[44]. A trial found that adolescents with type 1
319 diabetes who had received motivational interviewing had improved glycaemic control and life
320 satisfaction, and had reduced worry ^[45]. Future studies could further explore the mental
321 health impacts of comparable interventions and alternative approaches such as coaching
322 across other chronic conditions.

323

324 **Strengths and limitations**

325 This study has several strengths. The Millennium Cohort Study provided a large nationally
326 representative UK sample, with a prospective study design which reduced recall bias

327 compared to a single cross-sectional analysis, and sufficient demographic and socioeconomic
328 information for use as confounding variables.

329 This study also has some limitations. The high attrition rate and use of complete case analysis
330 in this study meant that 41% of singleton cohort members recruited to the Millennium Cohort
331 Study were included in the complete-case analyses. This may reduce the representativeness
332 of results and lead to attrition bias. Although sensitivity analyses using multiple imputation
333 by chained equations to account for missingness did not lead to substantial changes in model
334 estimates, if the missing at random assumption does not hold, then estimates using multiple
335 imputation are also likely to be biased^[49].

336 Prior to age 11 years, parents and carers were not asked what type of health condition cohort
337 members were affected by. This may have led to some cohort members with a limiting
338 mental health condition prior and no LLSI prior to age 11 years being misclassified as having
339 an LLSI, which may have inflated the relationship between LLSI and serious psychological
340 distress in this study.

341 **Conclusion**

342 This was the first large-scale longitudinal study to investigate the relationship between LLSI
343 and psychological distress in a representative sample of UK adolescents. This study found that
344 children and adolescents with a known LLSI are at an increased risk of serious psychological
345 distress and of currently receiving treatment for depression or serious anxiety at age 17 years
346 when compared with those without an LLSI. Working alongside young people to provide
347 support, improve treatment adherence, and routinely screen for and identify psychological
348 distress in young people with any LLSI may be beneficial.

349 **Acknowledgements**

350 **Author contributions**

351 Substantial contributions to conception and design, acquisition of data, or analysis and
352 interpretation of data: Palliser, G.M., Fraser, L.K., Mooney, K.L., and Jarvis, S.W.

353 Drafting the article or revising it critically for important intellectual content: Palliser, G.M.,
354 Fraser, L.K., Mooney, K.L., and Jarvis, S.W.

355 Final approval of the version to published: Palliser, G.M., Fraser, L.K., Mooney, K.L., and
356 Jarvis, S.W.

357 **Competing interests**

358 The authors have no competing interests to declare.

359 **Consent statement**

360 This study only included data from the Millennium Cohort Study, which have previously
361 obtained consent from all participants. No additional patient consent was required to
362 conduct the secondary analyses in this study.

363 **Data availability**

364 The study data are freely available from the UK Data Service, under an End User Licence
365 Agreement. Data can be obtained from

366 <https://beta.ukdataservice.ac.uk/datacatalogue/series/series?id=2000031>

367

368 References

- 369 **1** Newlove-Delgado T, Williams T, Robertson K, McManus S, Sadler K, Vizard T, Cartwright
370 C, Mathews F, Norman S, Marcheselli F, Ford T. Mental health of children and young
371 people in England. Leeds: NHS Digital Available from:
372 https://files.digital.nhs.uk/97/B09EF8/mhcyp_2021_rep.pdf#
- 373 **2** Bebbington PE, McManus S. Revisiting the one in four: the prevalence of psychiatric
374 disorder in the population of England 2000–2014. *Br J Psychiatry* 2020; **216**: 55–57.
375 [DOI: 10.1192/bjp.2019.196]
- 376 **3** Mental Health of Children and Young People in England 2021 - wave 2 follow up to the
377 2017 survey. NHS England Digital. Available from: [https://digital.nhs.uk/data-and-](https://digital.nhs.uk/data-and-information/publications/statistical/mental-health-of-children-and-young-people-in-england/2021-follow-up-to-the-2017-survey)
378 [information/publications/statistical/mental-health-of-children-and-young-people-in-](https://digital.nhs.uk/data-and-information/publications/statistical/mental-health-of-children-and-young-people-in-england/2021-follow-up-to-the-2017-survey)
379 [england/2021-follow-up-to-the-2017-survey](https://digital.nhs.uk/data-and-information/publications/statistical/mental-health-of-children-and-young-people-in-england/2021-follow-up-to-the-2017-survey)
- 380 **4** Mental Health of Children and Young People in England, 2023 - wave 4 follow up to the
381 2017 survey. NHS England Digital. Available from: [https://digital.nhs.uk/data-and-](https://digital.nhs.uk/data-and-information/publications/statistical/mental-health-of-children-and-young-people-in-england/2023-wave-4-follow-up)
382 [information/publications/statistical/mental-health-of-children-and-young-people-in-](https://digital.nhs.uk/data-and-information/publications/statistical/mental-health-of-children-and-young-people-in-england/2023-wave-4-follow-up)
383 [england/2023-wave-4-follow-up](https://digital.nhs.uk/data-and-information/publications/statistical/mental-health-of-children-and-young-people-in-england/2023-wave-4-follow-up)
- 384 **5** Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE. Lifetime Prevalence
385 and Age-of-Onset Distributions of DSM-IV Disorders in the National Comorbidity Survey
386 Replication. *Archives of General Psychiatry* 2005; **62**: 593–602. [DOI:
387 10.1001/archpsyc.62.6.593]
- 388 **6** American Psychiatric Association. Diagnostic and statistical manual of mental disorders,
389 4th edition. 4th ed. Washington, D.C.: American Psychiatric Association
- 390 **7** Agnafors S, Barmark M, Sydsjö G. Mental health and academic performance: a study on
391 selection and causation effects from childhood to early adulthood. *Soc Psychiatry*
392 *Psychiatr Epidemiol* 2021; **56**: 857–866. [DOI: 10.1007/s00127-020-01934-5]
- 393 **8** Meier MH, Hall W, Caspi A, Belsky DW, Cerdá M, Harrington H, Houts R, Poulton R,
394 Moffitt TE. Which adolescents develop persistent substance dependence in adulthood?
395 Using population-representative longitudinal data to inform universal risk assessment.
396 *Psychol Med* 2016; **46**: 877–889. [PMID: 26620720 DOI: 10.1017/S0033291715002482]
- 397 **9** Hards E, Loades ME, Higson-Sweeney N, Shafran R, Serafimova T, Brigden A, Reynolds S,
398 Crawley E, Chatburn E, Linney C, McManus M, Borwick C. Loneliness and mental health
399 in children and adolescents with pre-existing mental health problems: A rapid
400 systematic review. *British Journal of Clinical Psychology* 2022; **61**: 313–334. [DOI:
401 10.1111/bjc.12331]
- 402 **10** Manor I, Gutnik I, Ben-Dor DH, Apter A, Sever J, Tyano S, Weizman A, Zalsman G.
403 Possible association between attention deficit hyperactivity disorder and attempted
404 suicide in adolescents - a pilot study. *Eur Psychiatry* 2010; **25**: 146–150. [PMID:
405 19699060 DOI: 10.1016/j.eurpsy.2009.06.001]

- 406 **11** Department of Health. Long term conditions compendium of information. Leeds:
407 Department of Health Available from:
408 [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachm](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/216528/dh_134486.pdf)
409 [ent_data/file/216528/dh_134486.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/216528/dh_134486.pdf)
- 410 **12** Child and infant mortality in England and Wales - Office for National Statistics. 2021.
411 Available from:
412 [https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/d](https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/bulletins/childhoodinfantandperinatalmortalityinenglandandwales/2021#trends-in-child-and-infant-mortality)
413 [eaths/bulletins/childhoodinfantandperinatalmortalityinenglandandwales/2021#trends-](https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/bulletins/childhoodinfantandperinatalmortalityinenglandandwales/2021#trends-in-child-and-infant-mortality)
414 [in-child-and-infant-mortality](https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/bulletins/childhoodinfantandperinatalmortalityinenglandandwales/2021#trends-in-child-and-infant-mortality)
- 415 **13** Anderson E, Durstine JL. Physical activity, exercise, and chronic diseases: A brief review.
416 *Sports Medicine and Health Science* 2019; **1**: 3–10. [DOI: 10.1016/j.smhs.2019.08.006]
- 417 **14** Allen T, Reda S, Martin S, Long P, Franklin A, Bedoya SZ, Wiener L, Wolters PL. The Needs
418 of Adolescents and Young Adults with Chronic Illness: Results of a Quality Improvement
419 Survey. *Children (Basel)* 2022; **9**: 500. [PMID: 35455544 DOI: 10.3390/children9040500]
- 420 **15** Delamater AM, Guzman A, Aparicio K. Mental health issues in children and adolescents
421 with chronic illness. *International Journal of Human Rights in Healthcare* 2017; **10**: 163–
422 173. [DOI: 10.1108/IJHRH-05-2017-0020]
- 423 **16** Ferro MA, Gorter JW, Boyle MH. Trajectories of depressive symptoms during the
424 transition to young adulthood: The role of chronic illness. *Journal of Affective Disorders*
425 2015; **174**: 594–601. [DOI: 10.1016/j.jad.2014.12.014]
- 426 **17** Speechley KN, Ferro MA, Camfield CS, Huang W, Levin SD, Smith ML, Wiebe S, Zou G.
427 Quality of life in children with new-onset epilepsy. *Neurology* 2012; **79**: 1548–1555.
428 [DOI: 10.1212/WNL.0b013e31826e25aa]
- 429 **18** Zheng K, Abraham C, Bruzzese J-M, Smaldone A. Longitudinal Relationships between
430 Depression and Chronic Illness in Adolescents: An Integrative Review. *J Pediatr Health*
431 *Care* 2020; **34**: 333–345. [PMID: 32171610 DOI: 10.1016/j.pedhc.2020.01.008]
- 432 **19** Cobham VE, Hickling A, Kimball H, Thomas HJ, Scott JG, Middeldorp CM. Systematic
433 Review: Anxiety in Children and Adolescents With Chronic Medical Conditions. *Journal*
434 *of the American Academy of Child & Adolescent Psychiatry* 2020; **59**: 595–618. [DOI:
435 10.1016/j.jaac.2019.10.010]
- 436 **20** Brady AM, Deighton J, Stansfeld S. Psychiatric outcomes associated with chronic illness
437 in adolescence: A systematic review. *Journal of Adolescence* 2017; **59**: 112–123. [DOI:
438 10.1016/j.adolescence.2017.05.014]
- 439 **21** Calam R, Gregg L, Goodman R. Psychological Adjustment and Asthma in Children and
440 Adolescents: The UK Nationwide Mental Health Survey. *Psychosomatic Medicine* 2005;
441 **67**: 105–110. [DOI: 10.1097/01.psy.0000151490.77622.37]

- 442 **22** Brady AM, Deighton J, Stansfeld S. Chronic illness in childhood and early adolescence: A
443 longitudinal exploration of co-occurring mental illness. *Development and*
444 *Psychopathology* 2021; **33**: 885–898. [DOI: 10.1017/S0954579420000206]
- 445 **23** Mersha AG, Tollosa DN, Bagade T, Eftekhari P. A bidirectional relationship between
446 diabetes mellitus and anxiety: A systematic review and meta-analysis. *Journal of*
447 *Psychosomatic Research* 2022; **162**: 110991. [DOI: 10.1016/j.jpsychores.2022.110991]
- 448 **24** Grassi L, Riba M. Cancer and severe mental illness: Bi-directional problems and potential
449 solutions. *Psycho-Oncology* 2020; **29**: 1445–1451. [DOI: 10.1002/pon.5534]
- 450 **25** Centre for Longitudinal Studies. Millennium Cohort Study User Guide: Surveys 1-5, 9th
451 edn. London: Centre for Longitudinal Studies Available from:
452 [http://doc.ukdataservice.ac.uk/doc/5795/mrdoc/pdf/mcs1-5_user_guide_ed9_2020-](http://doc.ukdataservice.ac.uk/doc/5795/mrdoc/pdf/mcs1-5_user_guide_ed9_2020-08-07.pdf)
453 [08-07.pdf](http://doc.ukdataservice.ac.uk/doc/5795/mrdoc/pdf/mcs1-5_user_guide_ed9_2020-08-07.pdf)
- 454 **26** Hope S, Deighton J, Micali N, Law C. Maternal mental health and childhood injury:
455 evidence from the UK Millennium Cohort Study. *Arch Dis Child* 2019; **104**: 268–274.
456 [DOI: 10.1136/archdischild-2017-313809]
- 457 **27** Rosenberg AR, Bradford MC, McCauley E, Curtis JR, Wolfe J, Baker KS, Yi-Frazier JP.
458 Promoting resilience in adolescents and young adults with cancer: Results from the
459 PRISM randomized controlled trial. *Cancer* 2018; **124**: 3909–3917. [DOI:
460 10.1002/cncr.31666]
- 461 **28** Peiper N, Clayton R, Wilson R, Illback R. The performance of the K6 Scale in a large
462 school sample. *Psychological Assessment* 2015; **27**: 228. [DOI: 10.1037/pas0000025]
- 463 **29** Kessler R C, Barker P R, Colpe L J, Epstein J F, Gfroerer J C, Hiripi E, Howes M J, Normand
464 S-L T, Manderscheid R W, Walters E E, Zaslavsky A M. Screening for serious mental
465 illness in the general population. *Arch Gen Psychiatry* 2003; **60**: 184–189. [DOI:
466 10.1001/archpsyc.60.2.184]
- 467 **30** Hajebi A, Motevalian A, Amin-Esmaeili M, Rahimi-Movaghar A, Sharifi V, Hoseini L,
468 Shadloo B, Mojtabei R. Adaptation and validation of short scales for assessment of
469 psychological distress in Iran: The Persian K10 and K6. *Int J Methods Psychiatr Res* 2018;
470 **27**: e1726. [PMID: 29888523 DOI: 10.1002/mpr.1726]
- 471 **31** Easton SD, Safadi NS, Wang Y, Hasson RG. The Kessler psychological distress scale:
472 translation and validation of an Arabic version. *Health Qual Life Outcomes* 2017; **15**:
473 215. [PMID: 29078774 DOI: 10.1186/s12955-017-0783-9]
- 474 **32** Sakurai K, Nishi A, Kondo K, Yanagida K, Kawakami N. Screening performance of K6/K10
475 and other screening instruments for mood and anxiety disorders in Japan. *Psychiatry*
476 *and Clinical Neurosciences* 2011; **65**: 434–441. [DOI: 10.1111/j.1440-1819.2011.02236.x]
- 477 **33** Yang M, Carson C, Creswell C, Violato M. Child mental health and income gradient from
478 early childhood to adolescence: Evidence from the UK. *SSM Popul Health* 2023; **24**:
479 101534. [PMID: 37954013 DOI: 10.1016/j.ssmph.2023.101534]

- 480 **34** Straatmann VS, Lai E, Lange T, Campbell MC, Wickham S, Andersen A-MN, Strandberg-
481 Larsen K, Taylor-Robinson D. How do early-life factors explain social inequalities in
482 adolescent mental health? Findings from the UK Millennium Cohort Study. *J Epidemiol*
483 *Community Health* 2019; **73**: 1049–1060. [PMID: 31492761 DOI: 10.1136/jech-2019-
484 212367]
- 485 **35** Patalay P, Fitzsimons E. Development and predictors of mental ill-health and wellbeing
486 from childhood to adolescence. *Soc Psychiatry Psychiatr Epidemiol* 2018; **53**: 1311–
487 1323. [DOI: 10.1007/s00127-018-1604-0]
- 488 **36** Ferro MA, Boyle MH. The Impact of Chronic Physical Illness, Maternal Depressive
489 Symptoms, Family Functioning, and Self-esteem on Symptoms of Anxiety and
490 Depression in Children. *J Abnorm Child Psychol* 2015; **43**: 177–187. [DOI:
491 10.1007/s10802-014-9893-6]
- 492 **37** Plewis I. Technical Report on Sampling 4th Edition. London: Centre for Longitudinal
493 Studies
- 494 **38** NHS Digital. Health Survey for England, 2019: Data tables. NHS Digital. 2020. Available
495 from: [https://digital.nhs.uk/data-and-information/publications/statistical/health-](https://digital.nhs.uk/data-and-information/publications/statistical/health-survey-for-england/2019/health-survey-for-england-2019-data-tables)
496 [survey-for-england/2019/health-survey-for-england-2019-data-tables](https://digital.nhs.uk/data-and-information/publications/statistical/health-survey-for-england/2019/health-survey-for-england-2019-data-tables)
- 497 **39** Brooks F, Klemere E, Chester K, Magnusson J, Spencer N. Health behaviour in school
498 aged children. Hertfordshire: Centre for Research in Public Health and Community Care
499 Available from: [http://hbscengland.org/wp-content/uploads/2020/01/HBSC-England-](http://hbscengland.org/wp-content/uploads/2020/01/HBSC-England-National-Report-2020.pdf)
500 [National-Report-2020.pdf](http://hbscengland.org/wp-content/uploads/2020/01/HBSC-England-National-Report-2020.pdf)
- 501 **40** Lerch MF, Thrane SE. Adolescents with chronic illness and the transition to self-
502 management: A systematic review. *Journal of Adolescence* 2019; **72**: 152–161. [DOI:
503 10.1016/j.adolescence.2019.02.010]
- 504 **41** Morton RW, Everard ML, Elphick HE. Adherence in childhood asthma: the elephant in
505 the room. *Archives of Disease in Childhood* 2014; **99**: 949–953. [PMID: 24876303 DOI:
506 10.1136/archdischild-2014-306243]
- 507 **42** Shah R, McKay SV, Levitt Katz LE, ghormli LE, Anderson BJ, Casey TL, Higgins L, Izquierdo
508 R, Wauters AD, Chang N. Adherence to multiple medications in the TODAY (Treatment
509 Options for type 2 Diabetes in Adolescents and Youth) cohort: effect of additional
510 medications on adherence to primary diabetes medication. *J Pediatr Endocrinol Metab*
511 2020; **33**: 191–198. [PMID: 31809265 DOI: 10.1515/jpem-2019-0315]
- 512 **43** Hanghøj S, Boisen KA. Self-Reported Barriers to Medication Adherence Among
513 Chronically Ill Adolescents: A Systematic Review. *Journal of Adolescent Health* 2014; **54**:
514 121–138. [DOI: 10.1016/j.jadohealth.2013.08.009]
- 515 **44** Kosse RC, Bouvy ML, Belitser SV, Vries TW de, Wal PS van der, Koster ES. Effective
516 Engagement of Adolescent Asthma Patients With Mobile Health–Supporting Medication
517 Adherence. *JMIR mHealth and uHealth* 2019; **7**: e12411. [DOI: 10.2196/12411]

- 518 **45** Channon SJ, Huws-Thomas MV, Rollnick S, Hood K, Cannings-John RL, Rogers C, Gregory
519 JW. A Multicenter Randomized Controlled Trial of Motivational Interviewing in
520 Teenagers With Diabetes. *Diabetes Care* 2007; **30**: 1390–1395. [DOI: 10.2337/dc06-
521 2260]
- 522 **46** George C, Felix SA, McLellan A, Shetty J, Middleton J, Chin RF, Poveda B, Brand C, Small
523 M, Verity K. Pilot project of psychological services integrated into a pediatric epilepsy
524 clinic: Psychology Adding Value – Epilepsy Screening (PAVES). *Epilepsy Behav* 2021; **120**.
525 [PMID: 33971391 DOI: 10.1016/j.yebeh.2021.107968]
- 526 **47** Corathers S, Mara CA, Chundi PK, Kichler JC. Depression Screening of Adolescents With
527 Diabetes: 5-Years of Implementation and Outcomes. *Journal of the American Academy*
528 *of Child & Adolescent Psychiatry* 2019; **58**: 628–632. [PMID: 30802493 DOI:
529 10.1016/j.jaac.2019.01.013]
- 530 **48** Chauhan A, Walton M, Manias E, Walpola RL, Seale H, Latanik M, Leone D, Mears S,
531 Harrison R. The safety of health care for ethnic minority patients: a systematic review.
532 *International Journal for Equity in Health* 2020; **19**: 118. [DOI: 10.1186/s12939-020-
533 01223-2]
- 534 **49** Mera-Gaona M, Neumann U, Vargas-Canas R, López DM. Evaluating the impact of
535 multivariate imputation by MICE in feature selection. *PLoS One* 2021; **16**: e0254720.
536 [PMID: 34320016 DOI: 10.1371/journal.pone.0254720]
- 537




Figure 1- Simplified DAG




Figure 2- Attrition Diagram