1	On the cause and consequences of coinfection:
2	A general mechanistic framework of within-host parasite competition
3	
4	Ashwini Ramesh ^{1,2*} and Spencer R Hall ²
5	¹ Department of Integrative Biology; Department of Molecular & Microbial Genetics, Michigan State
6	University, East Lansing, USA – 48823
7	² Department of Biology, Indiana University, Bloomington, Indiana, USA – 47405
8	
9	*Corresponding author email: <u>rameshas@msu.edu</u>
10	Co-author email: <u>sprhall@iu.edu</u>
11	
12	Competing Interest: Authors declare no competing interests
13	Short title: On the cause and consequences of coinfection
1/	Koy words, competition, coinfaction, priority effects, founder control, putrient supply, best bealth
17	nutritional immunity infactious disease
15	nutrional minumity, miectious disease
16	Type: Synthesis; review; framework; mathematical modelling; infectious disease; public health
17	Number of words in abstract: 200
18	Number of words in main text: 7154
19	Number of references: 54
20	Number of figures, and tables: 9 figures and 1 table
21	
22	
23	
24	
25	
26	NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.
	· · · ·

27

28 ABSTRACT

29 Coinfections pose serious threats to health and exacerbate parasite burden. If coinfection is 30 detrimental, then what within-host factors facilitate it? Equally importantly, what hinders it, say via 31 exclusion or priority effects? Such interactions ought to stem from their within-host environment ('niche'), 32 i.e., resources that parasites steal from hosts and immune cells that kill them. Yet, despite two decades of 33 empirical focus on within-host infection dynamics, we lack a mechanistic framework to understand why 34 coinfection arises and the diverse range of its' consequences. Hence, we construct a trait-based niche 35 framework, one that illustrates general principles that govern parasite competition for a resource and apparent competition for immune cells. We show that coinfection requires a competition-resistance tradeoff 36 37 and that each parasite most impacts the niche factor to which it is most sensitive. These predictions then 38 provide mechanistic interpretation for infection outcomes seen in a variety of extant experiments: Why does 39 nutrient supplementation shift relative frequencies of coinfecting parasites? When and how does sequence 40 of parasite invasion allow only early invading parasites to win? How does intrinsic variation in immune 41 response shape coinfection burden? Together, this mechanistic framework of parasite competition offers 42 new perspectives to better predict within-host infection dynamics and improve individual health.

43

45 INTRODUCTION

46 Coinfections pose a serious threat to health both at the individual and population scale for hosts. 47 Broadly, coinfection is the successful concurrent infestation of a host with two or more parasite species. 48 Coinfection worsens human health (76% of 2000 publications in a meta-analysis) and exacerbates infection 49 burden (57%; Griffiths et al 2011). For instance, bacterial co-infections cause two-fold increase in mortality 50 in COVID patients (Shah et al 2023). Similarly, coinfecting helminth species increased the odds of anaemia 51 five-to-eight-fold among children (Ezeamama et al 2005). Coinfection at the individual scale can also 52 influence population-scale disease dynamics (Ezenwa & Jolles 2011). Populations having higher frequency 53 of coinfection experience larger epidemics than those with epidemics of one parasite (Susi et al 2015). Yet, 54 despite virulent costs, coinfections are pervasive amongst hosts, including humans, wildlife, and livestock 55 / agriculture (Vogels et al 2018; Ezenwa et al 2010; Halliday, Penczykowski et al 2020). Therefore, we 56 need to better understand why and how parasites coinfect their hosts.

57 Nonetheless, coinfection represents just one of numerous outcomes of concurrent pathogenesis. That 58 fact poses fundamental questions about coinfection. First, if coinfection is indeed detrimental, then what 59 within-host factors facilitate it? Equally importantly, what prevents it? Within-host parasite competition 60 can lead to coinfection (*i.e.*, within-host parasite coexistence), singly infection (through exclusion or 61 priority effects), or clearance (no infection) from hosts (Vogel et al 2018). Yet, we lack theoretical 62 predictions to elucidate those outcomes of competition. For instance, irrespective of timing, sequential 63 exposure to some parasites always lead to coinfection, yet to others, only early infecting parasites prevail 64 (Fig. 1A; Clay et al 2019; Devevey et al 2015). Second, when they coinfect why do some parasites become 65 more abundant than others in relative and absolute senses? For instance, nutrient supply or resources can 66 shift this 'community structure' of parasites within hosts favouring one parasite over the other (Fig. 1B; Fellous & Koella 2009, Budischak et al 2015). Finally, not all coinfected hosts are equal. Some hosts 67 68 present higher coinfection burden than others, hinting at variation in immunological resistance (Fig. 1C; 69 Fuess et al 2021, Halliday et al 2018). How does such variation govern coinfection burden? Presently, these 70 disparate bits about coinfection remain unconnected, lacking a synthetic glue. Here, we seek to catalyse 71 creation of a framework linking genesis of coinfection to its consequences (Lively et al 2014; Restif & 72 Graham 2015).

73 A mechanistic, within-host framework of parasite competition provides a start. To illustrate, we 74 develop and evaluate within-host models of coinfection, synthesizing insights through an ecological lens 75 (following Ramesh & Hall 2023). Broadly, coinfection (*i.e.*, within-host coexistence of parasites) occurs 76 when each parasite species competes more strongly with themselves (intraspecifically) than with each other 77 (interspecifically; Fig. 1D). Those competitive differences could be measured phenomenologically. Even 78 better, they can be quantified from trait-based niche models for parasites competing within hosts (Graham 79 et al 2008, Cressler et al 2014). In such models, parasites interact within host 'ecosystems' for shared 80 resources while facing attack by energetically costly immune cells. Simple ecological rules, then, govern within-host parasite competition. To illustrate, we first borrow from old and new niche models to outline 81 82 how shared energy (resources) and/or immune cells could govern divergent coinfection outcomes (Fig. 2). 83 Second, using a case study of a two parasite – immune cells – energy model (2PIE) niche model, we link 84 outcomes of infection to competitive abilities, nullclines (niches), feedback, and key traits (Figs. 3-7; Table 85 S1,2). Third, our approach scripts how to link a priori predictions to interpretation of experimental 86 outcomes using these types of models (Fig. 8-9). Finally, we offer suggestions for future theory and 87 experiments (Fig. 10, Table 1). Taken together, we lay a general mechanistic framework to understand 88 coinfection using within-host niche-based competition model.

89

90 Section I: A phenomenological approach to coinfection

91 To visualize these interactions, we label the direct effect of species j on growth rate of species i, 92 yielding (hereafter) interspecific *positive effects* (black arrow) and *negative effects* (red arrow) and 93 intraspecific *self-limitation* (red curve; Fig. 1,2). Then, we can write the two-species Lotka–Volterra 94 competition model in terms of intra (a_{ii}) - and inter (a_{ij}) -specific per capita competition coefficients:

95
$$\underbrace{\frac{1}{P_i} \cdot \frac{dP_i}{dt}}_{\text{per captita}} = \underbrace{r_i}_{\text{intrinsic}} \left(1 - \underbrace{a_{ii}P_i}_{\text{effect of intrasp}} - \underbrace{a_{ij}P_j}_{\text{effect of intersp}}\right)$$
(1)

where P_i is the density of parasite species *i*, and r_i is its intrinsic (maximum per capita) growth rate. This model assumes that per capita growth rates (fitness) decrease linearly with density of each species with slopes a_{ii} and a_{ij} (Table 1A). Successful coinfection satisfies two conditions. First, each single-species

99 ('boundary') carrying capacity $(1/a_{ii})$ must be invasible by the other parasite when rare. That condition is 100 met when each species reduces its own growth rate more than that of its competitor (i.e., $a_{11} > a_{21}$ and $a_{22} >$ 101 a_{12}). Second, joint intraspecific ($a_{11} a_{22}$) must exceed joint interspecific competition ($a_{12} a_{21}$). When both 102 are met, the feasible coexistence ('interior') equilibrium is stable with negative feedback (Table 1A).

103 Coinfection is hindered in one of two ways. First, strong positive feedback at the interior generates 104 priority effects when joint inter- exceeds intra-specific competition $(a_{12} a_{21} > a_{11} a_{22})$. Priority effects also 105 require mutually uninvasible boundary equilibria and a feasible interior (which are both met when $a_{11} < a_{21}$ 106 and $a_{22} < a_{12}$). Here, high enough initial density of parasites determines the winner. Second, competitive 107 exclusion arises with asymmetric competition. Specifically, if P_2 cannot invade P_1 's boundary $(a_{11} < a_{21})$ 108 but P_1 can invade P_2 's boundary $(a_{22} > a_{12})$, then P_1 excludes P_2 . Conversely, P_2 excludes P_1 when $a_{11} > a_{21}$ 109 but $a_{22} < a_{12}$. Hence, mutually or asymmetrically strong interspecific competition can hinder coinfection.

110 Empirically, estimates of competition coefficients can come from densities and growth rates of 111 parasites. Various methods exist, including a response surface design. In the context of macroparasites, this 112 design involves growing species either alone or at the lowest densities that lead to infection. It also involves 113 growing species with increasing density of intra- and inter-specific competitors. Through this design, a_{ii} 114 and a_{ii} can be estimated, then used to predict if species can successfully infect when rare (Freckleton & 115 Watkinson 2000; reviewed in Hart et al 2018; see Table 1A). However, this approach remains 116 phenomenological, measuring coefficients only in a specific environment. That context-dependence limits 117 predictions of infection outcomes across gradients of traits, nutrient supply, or in allocation to immunity. 118 Instead, predictions in these other contexts require a mechanistic framework of competition between 119 parasites involving host immunity and/or host resources.

120

121 Section II: Within-host feedbacks driving divergent infection outcomes

To move toward creation of such a framework, we outline six models of within-host competition between parasites involving immunity, resources, and/or direct interference (Fig. 2). All potentially contain mechanisms for coinfection, single infection due to exclusion or priority effects (aka, alternative states or founder control), and no infection. Although details can vary, such divergent outcomes in these models follow general principles of species competition as outlined above. In essence, coinfection occurs when

strong intraspecific competition (mediated directly or looped indirectly via 'niche' factors such as resources or immune cells) outweighs strong interspecific competition. Stronger intraspecific competition leads to net negative feedback on the feasible ('interior') stable equilibrium, facilitating coinfection.

130

131 Two parasite – shared energy (2PE) or – shared immune cells (2PI)

132 Parasite species that share a resource (energy) or suffer attack from a shared immune system engage 133 in exploitative or apparent competition, respectively (Fig. 2A,B). In exploitative competition (Fig. 2A), a 134 parasite that can survive at the lower equilibrium energy (resource) level (E^*) can outcompete the other. For 135 instance, among two clones of a rodent malaria (P. chabaudi), the superior competitor for red blood cells 136 (RBC) always excludes the inferior clone (De Roode et al 2005). Competition for a single, shared resource 137 can lead to a competitive hierarchy among multiple parasite species. For instance, in a malarial-hookworm 138 system, one Plasmodium species (P. falciparum) outcompetes hookworms which themselves outcompete 139 another Plasmodium (P. vivax) for red blood cells (Budischak et al 2018). Analogous rules can characterize 140 host systems with a shared immune system (Fig. 2B). In models, when two exponentially growing parasites 141 share immune cells, the species that can withstand highest density of immune cells (*i.e.*, highest I^*) wins 142 via apparent competition (Fenton & Perkins 2010). In a possibly analogous experiment, immune-mediated 143 interactions led to competitive suppression of an avirulent malarial clone, allowing the virulent clone to 144 dominate (Råberg et al 2006). Therefore, exploitative competition for a single resource or apparent 145 competition involving shared immune cells can lead to competitive exclusion alone.

146 Three aspects of within-host parasite biology can introduce coinfection or priority effects to pure 147 resource or apparent competition. First, if some mechanism imposes self-limitation on both parasites, then 148 they can coinfect even if they share only one immune niche. For instance, donor-controlled reproduction can create strong self-limitation on macroparasite density (Fenton & Perkins 2010). Second, non-linearity 149 150 in immune clearance rates can generate self-limitation, particularly when clearance accelerates at higher 151 parasite density (Fenton & Perkins 2010). Third, interference competition between parasites can facilitate 152 priority effects or coinfection (Amarasekare 2002; Fig. 2A-B with mutually inhibitory effects between P_1 153 and P_2 [grey shading]). If the inferior energy competitor is superior at interference, priority effects ensue. 154 Then, at high enough initial densities, the inferior energy competitor can win instead of being excluded.

However, if the interference also confers a benefit to the interacting species (e.g., if killing competitor larvae
and consuming it increases per capita growth rate), then coinfection ensues (following Amarasekare 2002).
These general principles for joint interference and exploitative competition (Fig. 2A) likely also apply to
immune-mediated apparent competition, a possibility for future exploration (Fig. 2B).

159

160 *Two parasite – two energy (2P2E; Resource ratio) or Two parasite – two immune cells (2P2I; Immune*

161 *ratio*)

162 When parasites (P_i) simultaneously compete for two resource or energy sources (E_i) via exploitative 163 competition, infection outcomes can include coinfection and priority effects (Tilman 1982). Consider two 164 parasite species competing for two substitutable resources (like in the resource ratio model: Fig. 2C). Here, 165 coinfection minimally requires each species to trade off their requirements for each energy resource. Then 166 sufficiently intermediate supply ratios must permit each single-species ('boundary') equilibria to fall within 167 each competitors' niches (enabling mutual invasibility). With both conditions met, coinfection arises if each 168 parasite has larger impact on the resource to which its fitness is most sensitive (see Appendix Section 2). 169 In testing the resource ratio model, Lacroix *et al.* observed a distinct competitive hierarchy, where one plant 170 virus altered success of a competing cereal dwarf virus under nitrogen: phosphorus (N:P) supply. In coinoculated plants, the cereal dwarf virus infection rates decreased with elevated P supply rate, while the 171 172 addition of N significantly increased its' infection rate (Lacroix et al 2014; Smith 2014). This suggests that, 173 at the very least, N:P shapes how each parasite species differentially impacts their resources, setting the 174 stage for coinfection.

175 Analogous rules likely apply to hosts with two types of immune response (2P2I, the immune ratio 176 model; Fig. 2D). If parasites act as substitutable resources for two generalist immune responses, coinfection arises if each parasite also has larger impact on the immune response to which its' fitness is most sensitive 177 178 (Appendix Section 2; derived for case with parasite self-limitation). Modification of this 2P2I framework 179 captures well-studied within-host interactions. First, the host can mount an independent, specialist immune 180 response to each parasite (with no links between $P_i - I_j$, Fenton & Perkins 2010). Therefore, the separate 181 dynamics of each immune-parasite pair determines coinfection. Similar results emerge for parasites 182 separated spatially (e.g., in different host tissues: Cervi et al 2004; Karvonen et al 2006). Second, two

parasites can each be attacked by specialist immune responses that inhibit each other. For instance, in mice, competing parasites interact *via* T-helper cells, where Th1 attacks intra-cellular malaria, Th2 attacks intestinal nematodes, but Th1 and Th2 inhibit each other (Griffiths *et al* 2015). Such interference creates negative feedback that can further enable coinfection (*e.g.*, links between I_1 and I_2 [grey shading]; Fenton & Perkins 2010).

188

189 Two parasite – immune cell – energy (2PIE)

190 Parasites can also engage in simultaneous exploitative and immune-mediated apparent competition 191 (Cressler *et al* 2014; Ramesh & Hall 2023). Consider common variations of two competing parasites (P_i) 192 that steal energy (E) from hosts and immune cells (I) that kill them (2PIE; Fig. 2E-F; Table S1). In one, 193 only parasites induce production of immune cells (2PIEi, induced immunity). In the other, energy is 194 continuously allocated to maintain baseline immune function, even without parasites (2PIEc, constitutive 195 immunity). This structure resembles a food web in which two prey species share a resource while attacked 196 by a generalist predator (Holt et al 1994; Leibold 1996). This keystone predation (diamond) model 197 anticipates the tradeoffs and niche dimensions that would govern exploitative and apparent competition 198 parasites (Ramesh & Hall 2023). When combined, these forms of competition enable coinfection, priority 199 effects, or competitive exclusion (see below Fig. 3; Ramesh & Hall 2023). Constitutive immunity (Fig. 2F) 200 reduces opportunities (parameter space) for coinfection, lowering parasite burden while maintaining higher 201 energy for other metabolic work (see below; Fig. 9).

The remainder of this review will focus on joint immune and energy (resource) competition in the 2PIE model. Most organisms have some type of (costly) immune defences to fight parasites that steal host energy (*reviewed in* Lochmiller & Deerenberg 2000; Zuk & Stoehr 2002). Furthermore, principles that govern strength of intra- *v* inter-specific competition in 2PIE models should apply to others. With the 2PIE model, then, we ask three questions: (a) What within-host feedback drives divergent outcomes of infection? (b) How does this framework offer new ways to resolve results in previous coinfection experiments (Fig. 1A-C)? (c) How can this framework guide future theory and experiments?

- 209
- 210

211 Section III: A mechanistic framework linking feedbacks, trait ratios, and infection outcomes

In this section, we construct a framework for within-host competition of parasites that connects trade-offs, nullclines (niches), feedbacks, and traits. As we will learn, the 2PIE model follows the familiar tale of intra- vs inter-specific competition governing coexistence vs priority effects (given a feasible interior equilibrium; Fig. 1D). Although specifics between 2PIEi and the others (Fig. 2A-F) surely differ, this approach lays a general script to generate mechanistic *a priori* predictions and glean better understanding of coinfection dynamics from within-host niche models. Many of the mathematical details are found elsewhere (Appendix Section 1).

219

220 *The Model* (see also S1, Table S1):

Growth rate of immune cells I (Eq. 2A): In 2PIEi, immune cells (I) increase after attack on parasite j (P_j) at rate f_{IP_j} , inducing consumption of e_{IP_j} of energy, E, with conversion efficiency e_I of energy into an immune cell after debiting loss rate m_I . In 2PIEc, I also increases with allocation of E at baseline rate a_b .

Growth rate of parasite P_j (Eq. 2B): The two parasites consume host energy with feeding rate f_{P_j} and energy per parasite conversion, $e_{P_jE}^{-1}$. The parasites are lost due to attack by immune cells (f_{IP_j}) and die at (shared) background rate m_P .

228 Growth rate of energy, E (Eq. 2C): The host consumes a resource S via a Monod function with 229 maximal assimilation rate f_E and half-saturation constant h. (This function merely pays homage to non-230 linear feeding behaviour of hosts following Cressler et al 2014). That resource, converted to energy within 231 the host (E), is lost at fixed rate r for use by hosts (for metabolic needs). Its net production, then, is f(S) - r232 E. Additionally, host energy is consumed via induction from immune attack on parasites (proportional to 233 the triple product $P_i E I$, and from consumption (theft) by parasites themselves. Hence, parasite j 234 'consumes' energy both indirect and directly described, a sum grouped below as $f_i(I)$. In 2PIEc, energy is 235 also lost via allocation to baseline immune function at rate a_b . The 2PIEi, c models are thus:

$$\frac{dI}{dt} = \frac{a_b E + \Sigma e_{IP_j} f_{IP_j} P_j E I}{e_I} - m_I I,$$
 Eq. 2A

$$\frac{dP_j}{dt} = \left(\frac{f_{P_j}E}{e_{P_jE}} - f_{IP_j}I - m_P\right)P_j,$$
 Eq. 2B

$$\frac{dE}{dt} = \frac{f_E S}{h+S} - rE - (\Sigma e_{IP_j} f_{IP_j} I + \Sigma f_{P_j}) P_j E - a_b E.$$
 Eq. 2C

236

237 Outcomes of competition

238 Depending on their traits, competing parasites can successfully coinfect, singly infect (via 239 competitive exclusion or priority effects), or produce no infection in a host. Such outcomes can be captured 240 along gradients of, say, supply of resources to hosts, S, and feeding rate of parasites on energy within hosts, 241 f_{P_1} (Fig. 3A, Table S1). The lines within the 2D bifurcation diagram represent qualitative shifts of within-242 host composition determined by minimal requirements. We highlight two such shifts or 'transcriticals' $(T_1$ 243 and T_2) governing coinfection, but we build to it. Very low S prevents persistence of either parasite, resulting 244 in no infection (a nutritional clearance; Fig. 3A, yellow). Increasing S, now supports only single infection by parasite 1 (P_1) because it excludes the other via resource competition (P_2 , at $E-P_1$ [lighter] or $E-P_1-I$ 245 246 [darker] 'boundary equilibria'). Now, intermediate S, such that $T_2 < S < T_1$, allows parasite 2 (P_2) to invade 247 that P_1 boundary and P_1 invasion into the P_2 's boundary equilibrium (E-P₂-I). Hence, that region represents 248 mutual invasibility and coinfection (Fig. 3A, orange), situated between regions of competitive exclusion 249 ('single infection'). At higher S, P_1 is excluded allowing P_2 to win (E-P₂-I; Fig. 3A, dark purple). Notice, 250 increasing both f_{P_1} and S flips the geometry of those transcriticals. Now, higher S, such that $T_1 < S < T_2$, 251 does not allow P_1 invasion into the P_2 's boundary equilibrium and vice-versa. Hence, this region represents 252 mutual uninvasibility and priority effects aka founder control (Fig. 3A, grey; see Appendix S1 for details). 253 The offsets show sample dynamics when hosts are exposed to concurrent infection, started at one or two 254 initial densities, in each of these regions (Fig. 3B - F). In the no infection region, insufficient energy caused 255 failed infection, even at high initial densities of parasites (Fig. 3D). In the single infection regions, the 256 winner always excludes the other, even with initially low densities (Fig. 3A, E). In the coinfection region, 257 all trajectories head toward a stable (interior) equilibrium (Fig. 3C). With priority effects, the parasite 258 starting more densely wins and excludes the other (i.e., top: P_1 wins; bottom: P_2 wins; Fig. 3F).

260 Mechanism of competition: A within-host "assembly rules" and nullcline approach

These predictions hinge on intra- vs inter-specific competition. Intraspecific competition, mediated indirectly via host energy and immune cells, produces negative feedback loops while interspecific competition yields positive ones (Fig. 1D). When strength of intra- exceeds inter-specific competition, net negative feedback on the feasible interior equilibrium can enable coinfection (coexistence). Conversely, greater inter- than intra-specific competition generates net positive feedback that can produce priority effects. This section actualizes these concepts for 2PIEi using niches and nullclines.

267 Assembly rules: Broadly, coinfection requires a competition-resistance trade-off: the species that 268 competes superiorly for energy (without immune cell) must remain more vulnerable to immune attack. It 269 also requires intermediate resource supply, S, as consumed by hosts, f(S), falling in between two thresholds 270 (described below: $T_2 \le f(S) \le T_1$). The assembly of a within-host 'community' of parasites can be understood 271 via hosts exposed to (i) single species (Fig. 4A, 5A dotted and dashed lines) and (ii) both species (Fig. 4B, 272 5B solid lines). The single species 'boundary' equilibria provide insights into their minimum resource requirements of parasites for energy $(E_j^*, akin to R^* sensu$ Tilman 1982) and of immune cells for the product 273 274 of energy and parasites (EP_i^*) . They also highlight maximal immune cell densities supported by each parasite $(I_i^*, akin to P^* sensu$ Holt 1977). These quantities provide a start for assembly of coinfection. 275

276 First, the route to single infection follows resource- and immune-based assembly rules along a 277 nutrient supply gradient, S (Figs. 4B, 5B). Very low nutrient supply (S) meets neither parasite's minimum 278 energy requirements ($E^* < E_i^*$). Hence, very low S prevents infection (i.e., clearance via starvation; yellow). However, with high enough S, E^* increases, eventually meeting that minimum (E_j^*) . Then, P_j can invade 279 280 and increase with S (pinning E at E_i^* ; E-P_i regions in lighter). Eventually, P_i's density crosses a minimal 281 threshold (EP_i^*) that induces immune activation (darker colors). With higher S still, immune attack reduces density of P_j freeing up more energy (where energy with immune cells is denoted by $E_{j,I}^{*}$). While both 282 283 parasite species follow similar assembly rules, they differ in a key trade-off: P_1 is the superior competitor without immune cells $(E_I^* < E_2^*)$ but is more vulnerable to immune attack $(f_{IP_1} > f_{IP_2})$. That competition-284 285 resistance trade-off provides a first prerequisite for coinfection.

286 Comparison of those key energy and immune quantities yields more conditions for coinfection (Fig.
287 4B). Along the nutrient gradient, *S*, the more vulnerable parasite, *P*₁ invades (*E*-*P*₁). It outcompetes

(excludes) the more resistant parasite, P_2 , via resource competition (since $E_1^* < E_2^*$). At some S following immune activation (EP_1^*), P_1 becomes the superior apparent competitor ($I_1^* > I_2^*$) but inferior energy competitor ($E_{2,I}^* < E_{I,I}^*$) – a shift in competitive dominance. Then, with further S still, the *E-I* niche environment set by P_1 enables P_2 to invade (at T_2 : orange).

292 In a window of coinfection, the within-host niche environment remains constant while parasite densities shift (Fig. 4B). Here, energy and immune cells stay at E_{int}^* and I_{int}^* , respectively. However, with 293 294 higher S, the structure of this community shifts, with P_2 increasing while P_1 declines. With this shift, the 295 energy and parasite density needed to keep the immune system activated, $E_{int}^{*}(P_{1}^{*}+P_{2}^{*})$, increases until P_{2} 296 could support immune activation alone (at EP_2^* ; Fig. 4D). Broadly, in this region, the more vulnerable P_1 remains the superior apparent competitor (Fig. 4D, $I_1^* > I_2^*$) but weaker competitor for energy (Fig. 4C, E_2^* 297 $< E_{I,I}^*$ shifts to $E_{2,I}^* < E_{I,I}^*$). Conversely, the more resistant P_2 is the superior energy competitor but weaker 298 299 apparent competitor. This biology means that the more vulnerable parasite species also produces more 300 immune cells, the niche factor to which it is most sensitive (see below). That combination imposes a brake on its own growth rate (negative feedback), thus promoting coinfection. With S past this threshold (T_1) , P_2 301 excludes P_1 (via resource competition: $E_{2,I}^* < E_{I,I}^*$) but more immune cells lower P_2 's density (purple, Figs. 302 303 5B-E).

Priority effects can arise with different trait combinations for the two parasites (here: at higher f_{P_1} ; 304 305 Fig. 5B). Along a gradient of nutrient supply to hosts, S, the region of priority effects is approached similarly 306 to the coinfection case: first the more vulnerable parasite invades (when its E_l^* is met), then the immune system is activated (when its EP_1^* is reached). However, P_1 always remains the superior energy competitor, 307 308 unlike the flip seen before the coinfection window for coinfection (Fig. 5C). Past T_1 , more resistant P_2 309 cannot invade until T_2 is crossed; with higher S, more resistant P_2 excludes P_1 via apparent competition 310 (since $I_2^* > I_1^*$; Fig. 5D). Biologically, this means that the more resistant species produces more immune 311 cells, the niche factor to which their competitors are most sensitive (see below), creating positive feedback 312 and thus priority effects. With decreasing S, P_1 could not invade until T_1 was crossed. Hence, the interior 313 equilibrium in the priority effects window is unstable (a saddle), separating two stable states, dominance 314 by P_1 (at its *E*- P_1 -*I* boundary equilibrium) or by P_2 (at *E*- P_2 -*I*).

315

A nullcline (niche) interpretation: Nullclines help to characterize the niche environment that

316 enables it coinfection. At a given nutrient supply S, nullclines denote conditions in which a parasite (P_i) or 317 niche component (energy, immune cells) neither grows nor declines. Combinations of energy (E) and 318 immune (I) cells that fall to the right of a parasite's nullcline (higher energy, lower immune cells) sit within 319 its fundamental niche (Fig. 6A; P1, blue shading; P2, purple). Then, the impact of each parasite on their 320 within-host niche determines I and E nullclines. The nullcline for E is the combination of P_1 and P_2 that 321 'consumes' all production of energy at the interior (E_{int}^*) . Below this line, energy grows (dE/dt > 0, yellow322 in Fig. 6B). The P_i -axis intercepts (\tilde{P}_i) note the density of each parasite, when alone, that consumes all net energy production. The nullcline for I is the combination of P_1 and P_2 that meets the minimal EP^* needed 323 to maintain its activation. All points above the line increase production of immune cells (dI/dt > 0; green 324 in Fig. 6B). At each intercept (\hat{P}_i), the parasite alone supports immune activation (at EP_i^*) given E_{int}^* . 325

Coinfection (Fig. 6C - E; black, closed circle): Co-infection requires that parasite nullclines cross, 326 327 niche nullclines cross, and nutrient supply is intermediate. The nullclines of the parasites cross in E-I space 328 because of their competition-resistance trade-off (Appendix section S1). The shallower slope of less 329 resistant P_1 's nullcline means it is more sensitive to I while more resistant P_2 is more sensitive to E. When 330 they coinfect, both parasites set their relevant single-species (boundary) equilibria within the fundamental 331 niche of the other. Hence, each single-parasite environment can be invaded by the other (Fig. 6C; P_I 332 equilibrium [blue dot] falls within P_2 's niche [purple shaded] and vice-versa). Now P_1 is the superior apparent competitor $(I_1^* > I_2^*)$ while P_2 is the superior energy competitor (i.e., has lower $E^*: E_{I,I}^* > E_{2,I}^*$ 333 [as shown] or E_2^* ; Fig. 4C). Meanwhile, the *I* nullcline is steeper in P_1 - P_2 space (Fig. 6E). Hence P_1 has 334 335 larger impact on immune cells while P_2 has larger impact on energy. Therefore, each parasite exerts largest 336 impact on the niche factor to which it is most sensitive. Based on the nullcline intercepts, coinfection requires that the interior niche would support surplus $P_1(\hat{P}_1 < \tilde{P}_1)$ but deficient P_2 (so $\hat{P}_2 > \tilde{P}_2$) than needed 337 to meet immune system's EP^* requirement (Fig. 6E; when $f(S) = T_2$, $\hat{P}_1 = \tilde{P}_1$; at T_1 , $\hat{P}_2 = \tilde{P}_2$). The coinfection 338 339 equilibrium, then, combines densities of the superior apparent (P_1) and energy competitor (P_2) that 340 consumes all net energy production while meeting minimal requirements of immune cells - eliminating 341 those surpluses and deficits.



Priority effects (Fig. 6F-H; white, open circle): During priority effects, those three elements change.

343 First, while parasite nullclines still cross (due to the competition-resistance trade-off), now both parasites 344 (P_i) alone create uninvasible immune (I) – energy (E) niche environments (Fig. 6F; P_i blue dot outside P_2 345 purple shaded niche and vice-versa). Furthermore, in this region P_1 is always the superior energy competitor, without $(E_1^* \le E_2^*)$ and with immune cells $(E_{I,I}^* \le E_{2,I}^*)$, while P_2 becomes the superior 346 apparent competitor ($I_2^* > I_1^*$; Fig. 6F). Second, the *E* nullcline is now steeper than the *I* nullcline (Fig. 6J). 347 348 Hence, impacts flip: less resistant P_1 has larger impact on E while more resistant P_2 has larger impact on I. 349 Therefore, each parasite has a larger impact on the niche factor to which its competitor is most sensitive. Thus, the system would provide excess of the more resistant parasite $(\hat{P}_2 < \tilde{P}_2)$ and a deficiency of the 350 351 vulnerable one $(\hat{P}_1 > \tilde{P}_1)$ enabling priority effects.

352

353 Factors enabling coinfection: Feedback loops and trade-offs of traits

354 Outcomes of coinfection v priority effects depend on feedback and traits that govern them. At the 355 most fundamental scale, co-infection must be 'feasible' (set by an environment supporting both species) at 356 an equilibrium having net negative feedback. Such net negative feedback emerges when intraspecific 357 competition exceeds interspecific competition. That competition depends on loops connecting parasites to 358 their niche (Fig. 7A, S1). Loops are chains of interactions linking increased density of a species to growth rates of others, eventually returning to that species (Puccia & Levins 1991). For instance, interspecific 359 360 competition can be traced starting with an increase in P_1 . (i) An increase in P_1 's density can 'fuel immune 361 cells' that suppress its competitor, freeing up energy resources, thereby increasing its own growth rate via 362 higher 'births'. (ii) P_1 can also 'starve immune cells' by reducing P_2 's density via consumption of E. 363 Through both chains of interactions, P_l indirectly benefits from an increase in its own density via gains in 364 energy, hence birth rate, or lower death rate, respectively. Those two positive (destabilizing) loops then 365 push against two negative loops, (iii and iv) "P_i is attacked, P_i eats". These latter loops add the stabilizing, 366 consumer-resource-like, intraspecific competitive interactions within which each parasite is enmeshed. For instance, (iii) a small increase in P_1 increases immune attack, thereby reducing P_1 (P_1 -I loop) while P_2 is 367 slowed by resource consumption (P_2 -E loop). Then, (iv) those roles reverse, *i.e.*, resources brake P_1 while 368 369 the immune system slows parasite P_2 . Summed, those two loops (iii and iv) jointly determine the amount of intraspecific competition (negative feedback). Then, the 2PIE model follows the recognizable script of intra- vs inter-specific competition governing coexistence vs priority effects. If the strength of intraspecific competition loops (iii + iv: negative) exceeds that of interspecific loops (i + ii: positive), net negative feedback leads to coinfection. If, instead, interspecific exceed intraspecific loop strength, net positive feedback generates priority effects.

375 With some rearrangement, these intra- and inter-specific loops governing feedback can more 376 intimately connect to the niche environment and assembly rules revealed above (Fig. 7B, S1). The first 377 involves the difference in ratios of sensitivity of each parasite to immune cells, I, vs. energy, E (denoted by 378 α_i ; following Pásztor et al 2016). The second involves the ratio of impacts of each parasite on I vs. E (ε_i). 379 Both ratios can be written as proportional to ratios of traits or traits and minimal requirements (Fig. 7D and 380 E, respectively; see details in Appendix Section S1). Coinfection requires that each parasite has a larger 381 impact on the niche factor to which it is most sensitive. If the more vulnerable P_1 is most sensitive to and 382 has highest impact on I, then $\alpha_1 > \alpha_2$ and $\varepsilon_1 < \varepsilon_2$. Such symmetry in ratios generates net negative feedback. 383 For priority effects, the competition-resistance trade-off still yields $\alpha_l > \alpha_2$. However, because P_l has highest 384 impact on E (to which P_2 is most sensitive), the impact ratios flip, $\varepsilon_1 < \varepsilon_2$. This asymmetry in sensitivity vs. 385 impact ratios produces net positive feedback.

386

387 Section IV: A focus on the traits and quantities to measure in the future

Presently, most within-host competition experiments observe patterns of coinfection, then infer mechanisms. The next phase of experimentation should shift toward feedback, minimal requirements, and traits of mechanistic niche models (like 2PIEi). Above, we learned that coinfection requires that each parasite has greater impact on the niche factor to which it is most sensitive (generating net negative feedback); with priority effects, each parasite has greatest impact on the niche factor to which its competitor is most sensitive (yielding net positive feedback). Future research on coinfection dynamics could aim to quantify the traits and minima's underlying these sensitivity and impact frameworks.

(1) *Quantifying traits:* The (a)symmetry of the *sensitivity-impacts* ratios governing coinfection and
 priority effects can be better understood by delineating the combination of traits (Fig. 7D, Appendix Section

S1). For instance, the *sensitivity* ratios are governed by killing rate of immune cells on parasites (f_{IP_j}) , feeding rate of parasites (f_{P_j}) , and the cost of producing a new parasite (e_{P_jE}) . On the other hand, the *impact* ratios depend on how attacked parasites induce energy allocation to immune cells $(e_{IP_j}f_{IP_j})$ and the per capita 'consumption' of energy by parasite *j* directly and indirectly $(f_{P_j}(I))$.

401 (3) *Quantifying key traits and minimal requirements:* Experimental tests of *sensitivity-impact* ratios 402 could also centre on measurement of key quantities such as the minimal requirement of each parasite (E^*) 403 for energy and of the immune system for activation (EP^* ; Fig. 7E, Table 1C). For instance, the difference 404 in sensitivity ratios boils down to killing rate of immune cells on parasites, f_{P_j} , and each parasite's minimal 405 energy requirement without immune cells, E_j^* Similarly, the difference in impact ratios depend on per capita 406 'consumption' of energy by parasites, $f_{P_j}(I)$, and the immune cells' minimal energy-parasite requirement

407 EP_j^* (Appendix S1). These traits and quantities could be estimated via experiments (in principle).

408 Despite this call for a focus on traits and minimal requirements, we acknowledge inherent challenges 409 in measuring them. To move forward, experimentalists can leverage interdisciplinary methodologies. For 410 instance, measurement of state variables like parasite (P_j^*) , energy (E_j^*) , or immune densities (I_j^*) over time 411 can allow estimation of parameters of the model. Alternatively, traits could be measured with radioisotope 412 labelling of resources to track parasite feeding rate (Gomez-Amaro et al 2015), live imaging to track 413 immune killing rate (Galli et al 2021), ICP-MS to measure energetic content of infected host (Cassat et al 414 2018), etc. As a pay-off, the combination of trait measurements and models generates a priori predictions 415 of infection outcomes and the feedback mechanisms that govern them (notably demonstrated in Budischak 416 et al 2015; Griffiths et al 2015). Ultimately, such approaches may catalyse a new wave of theory-grounded, 417 niche-mechanistic combinations of modelling and experimentation.

418

419 Section V: Explanations for divergent infection outcomes using 2PIE

The model of exploitative and apparent competition between parasites, 2PIEi,c, makes predictions (Fig. 8A, 9) that can help to contextualize and interpret previous experiments (Fig. 8B-F). In this next section, we envision how 2PIEi could produce these various outcomes, backing-out mechanism *post hoc*. Thus, 2PIEi offers a way to potentially resolve otherwise seemingly inconsistent experimental outcomes.

- 424 Then, it can guide creation of new coinfection models tested in future experiments. Ideally, such efforts
 425 would start with parameterized predictions that test divergence of infection outcomes *a priori*.
- 426

427 Coinfected or not? What within-host factors facilitate or inhibit coinfection? (Fig. 8B, C)

428 Priority effects

429 *Theory*: Stronger inter- than intra-specific competition ensures priority effects (via net positive 430 feedback) where the parasite with sufficiently high initial dose wins (Fig. 8A, B; Table 1B).

Empirical evidence: In a rare, unequivocal demonstration of priority effects, the order of arrival of
bacterial (*B. burgdorferi*) strains determined which won and excluded the other within a host mouse (Fig.
8B). Immune response likely did not explain the priority effects (Devevey *et al* 2015), so future work can
pinpoint mechanisms (e.g., resource competition, interference, etc.) facilitating them (Fig. 2B; Table 1B).

Guiding future experiments: Varying initial parasite dose (dose-dependent assays) provide on possible way to delineate coinfection vs priority effects. When parasites coinfect, dose does not change outcome, despite varying initial P_2 and constant initial P_1 . With priority effects, sufficiently high initial density determines winner (Table 1B). Such an experiment would reveal whether competitive exclusion, or priority effects led to single infection. Even better, measurement of key quantities involved in *sensitivityimpact* ratios can delineate mechanism (e.g., superior resource v resistant competitor as in Fig. 5).

441

442 *Competitive exclusion*

Theory: Differences in parasite traits can separate exclusion from coinfection. For example, for a
given nutrient supply to hosts, a parasite with higher feeding rate (all else equal), can exclude its competitor
resulting in single infection (Fig. 8A, C). In contrast, one with lower feeding rate may successfully coinfect. *Empirical evidence:* Mice infected with malarial strain P₁ or P₃ (blue) were allowed to compete
pairwise against a common malarial strain P₂ (purple; Fig. 8C; De Roode *et al* 2005). In the P₃ - P₂ pairing,
regardless of order or arrival or delay between infections, P₃ competitively excluded P₂ (Fig. 8C).

449 *Guiding future experiments:* Higher competitive ability of P_3 may reflect higher feeding rate on 450 host energy. In contrast, if strain P_1 had lower feeding rate (hence less competitive), it could stably coinfect 451 with P_2 (Fig. 8C). Such hypotheses about resource acquisition traits could be tested in the future (Table 1B).

452

453 *Coinfection hierarchy and community structure* (Fig. 8D-F)

454 *Theory*: When parasites coinfect, nutrient supply to hosts (*S*) can mediate densities of the within-455 host energy and immunity, and hence relative and absolute densities of parasites (Fig. 8D). Thus, nutrient 456 supply can shift this 'community structure' of parasites within hosts. For instance, increasing nutrient 457 supply favours the more resistant P_2 over the less resistant P_1 .

Empirical evidence: This prediction could explain shifts in community structure of parasites in two different systems (Fig. 8E). The first arose with mice infected by two species of gastrointestinal worms (case I; Budischak *et al* 2015); the second used mosquito larvae infected by a microsporidian and protozoan parasite (case II; Fellous & Koella 2009). Both demonstrate how changing nutrient supply (*S*) favours one species (purple; positive slope) over the other (blue; negative slope; Fig. 8D). In both cases, relative abundance of coinfecting species can shift (or not) depending on *S*, too. However, future trait measurements would need to establish that higher *S* favoured the more resistant parasite as 2PIE predicts (Table 1C).

465 Guiding future experiments: Much may be learned about coinfection and nutrient supply with focus on 466 individual hosts (reviewed in Ezenwa 2021). Hosts can exhibit high intra- and inter-individual variation in 467 infection outcomes (Merill & Cáceres 2018). To illustrate, parasite densities in the mosquito case ranged 468 among all individuals from coinfection to single infection via exclusion / priority effects to no infection 469 (see reference to shaded regions in Fig. 8D v 8F; replotting of Fellous & Koella 2009). Such variation 470 among hosts could arise from individual differences in resource acquisition. If so, individuals nominally 471 fed the same amount (in a treatment) may fall functionally along different supply points (like in Fig. 8D). 472 Individuals consuming less resources would favour the less resistant parasite (P_1) while those eating much 473 more would favour the more resistant one (P_2) - yielding exclusion in both cases (Fig. 8D, F). The remaining 474 hosts became coinfected, with parasites reaching different densities among hosts. Future, experimental tests 475 can test for the full range of possible infection outcomes (and hence 'assembly rules' leading to coinfection 476 outcomes) at the individual scale, say, along broader nutrient gradients (Table 1C).

Furthermore, niche-based insights can guide more predictive experiments at the within-host scale which can then be scaled to the population linking within- to between-host dynamics. For instance, fluctuation in nutrient supply to host could shift competitive outcomes within hosts that then alters multi-parasite

480 outbreaks at the population scale (Hite & Cressler 2018; Ezenwa & Jolles 2011) or ecosystem processes
481 (Kendig *et al* 2020).

482

483 *Coinfection burden* (Fig. 9)

484 Finally, some hosts present higher coinfection burden (total density) than others, hinting at intrinsic 485 host resistance via immune clearance. How does variation in immune response govern coinfection burden? 486 Theory: In comparison to the model with only immune induction (2PIEi), baseline energy 487 allocation to immunity squeezes parameter space for coinfection and reduces parasite burden (2PIEc, 488 constitutive immunity; Fig. 9A-B). Despite the higher allocation (via a_b), reduced burden allows the infected host to maintain more energy for other metabolic work (Fig. 9A-B). At lower nutrient supply (S), 489 490 'no infection' by parasite shifts from nutritional clearance (E, yellow; Fig, 9A) to combination of immune 491 and nutritional form of clearance (E-I, green; Fig, 9B). The 2PIEc model also predicts that increasing 492 baseline allocation favours the more resistant P_2 over P_1 , eventually excluding it (Fig. 9E).

493 *Empirical evidence*: Consistent with these predictions, plant hosts treated with immune-signalling 494 hormone experienced a lower prevalence of a less aggressive parasite, increased burden of infection by a 495 more aggressive parasite, and experienced fewer co-infections (Halliday *et al* 2018). Thus, hosts with higher 496 allocation to immunity can resist infection more than those with lower allocation, leading to variation 497 among hosts in infection burden (all else equal).

498 *Guiding future experiments:* First, one could test such predictions using strains with immune 499 suppression of specific genes or metabolites involved in constitutive immunity (like in Chen *et al* 2005). 500 That could shift competition from a 2PIEc framework (with $a_b > 0$) to a more 2PIEi like one (with $a_b = 0$ at 501 the extreme). Another test could involve measuring competitive outcomes in host genotypes differing 502 allocation to baseline immunity (Fuess *et al* 2021; Table 1D).

503

504 Section VI: Conclusion

505 Why do divergent infection dynamics arise within a host? Competing parasites within a host can 506 coinfect (coexist), singly infect (through exclusion, or priority effects), or become cleared (no infection; 507 Fig. 1, 3). Those differing outcomes significantly affect health of individual hosts and even alter population-

level disease outbreaks (Mideo *et al* 2008). Such divergent infection outcomes ought to stem from niche interactions, *i.e.*, with the resources that parasites steal from hosts and the immune cells that kill them (Cressler *et al* 2014, Graham 2008). Yet, despite two decades of empirical focus on infection dynamics, many of these studies present a collection of disparate results without a synthetic glue. Clearly then, we need a comprehensive framework that explains why parasites coinfect and why they might not.

513 Here we build a within-host framework of parasite competition based on ecological theory (Fig. 2). Using a two parasite - immune cells - energy model (2PIE), we illustrate general principles governing 514 515 parasite competition via join exploitative and immune-mediated apparent competition (Ramesh & Hall 516 2023). Those forms of competition underlie a within-host framework that connects infection outcomes to 517 competitive abilities, nullclines (niches), feedbacks, and traits (Figs. 3 - 7; see details in Appendix Section S1). Notably, we delineate how the interplay of three quantities – minimum resource requirements of (1) 518 519 parasites for energy $(E_i^*, \text{akin to } R^*)$, and of (2) the product of energy and parasites (EP_i^*) for immune cells, 520 and (3) maximal immune cells supported by each parasite (I_i^*) – provide a start for assembly of coinfection. 521 We show that coinfection requires that each parasite exerts greater impact on its more sensitive niche factor; 522 that arrangement introduces net stabilizing, negative feedback. With priority effects, each parasite more 523 strongly impacts the factor to which its competitor is most sensitive, leading to net destablizing, positive 524 feedback (Fig. 3, 4).

525 That framework also provides mechanistic insights into and explanations for some experimental 526 results (Fig. 8-9). First, it explains two ways to hinder coinfection: priority effects can favour early invaders, 527 while competitive exclusion always inhibits one parasite (arising via *e.g.*, fast feeders at low nutrient supply; 528 Fig. 5, 8B-C). Second, increasing nutrient supply in the coinfection region can favour the more resistant 529 over the less resistant parasite, shifting community structure along nutrient gradients (Fig. 8D). Finally, 530 greater investment in constitutive immunity squeezes opportunity (parameter space) for coinfection and 531 reduces parasite burden, freeing up energy for metabolic needs (Fig. 9). Together, the 2PIE models provide 532 interpretation for otherwise puzzling outcomes from various experiments. Although specifics will vary, 533 these principles should apply other niche models involving, say, two energy or immune sources (Fig. 2, 534 2P2E, 2P2I; see Appendix Section S2). This synthesis of experiments with models underscores need for 535 parametrized trait-based experiments that better predict within-host infection outcomes (Table 1).

- 536 Unpacking such within-host mechanisms can improve our understanding of individual health. For 537 instance, deworming trials of hosts coinfected with malaria and gastrointestinal worms show that increased 538 availability of RBCs allow malaria to proliferate within the host, ultimately making them sicker (Budischak 539 et al 2018). Disentangling the mechanisms can then allow for correct course of treatment plan to improve 540 individual health (e.g., malarial drugs or vaccines followed by deworming). Together, our work provides a 541 synthesis of within-host niche-based frameworks, laying a theoretical groundwork for a mechanistic 542 understanding of competition outcomes. We then used model predictions to contextualize key findings from 543 the past two decades of experimentation. With this synthesis, we aim to catalyse a new wave of theory-544 grounded, niche-mechanistic combinations of modelling and experimentation. Ultimately, such a new wave 545 could help to mitigate the severity and comorbidities of coinfections but also advance the development of 546 preventive drug therapeutics and vaccines, ultimately enhancing individual health.
- 547

548 Acknowledgements:

549 AR was supported by the IU Center for the Integrative Study of Animal Behavior, NSF DEB (1655656),

and a MSU Presidential Postdoctoral Fellowship. We thank J Lau and the Klausmeier-Litchman lab for

helpful comments on the manuscript; D Brisson, J de Roode, S Budischak, S Fellous for generously sharing

their data. AR credits J Coltrane and A Ali Khan for artistic inspiration.

553 FIGURE CAPTIONS

554 Figure 1: Unpacking causes and consequences of coinfection: (A-C) Examples from experiments probing why coinfection arises and the diverse range of its' consequences, including: (A) Coinfected or 555 556 not? What within-host factors facilitate coinfection and what prevents it? Sequential exposure to parasite 557 species can lead to coinfection irrespective of timing (Case I); in other scenarios, early-infecting parasites 558 prevailed (impeding coinfection; Case II). (B) Coinfection community structure: When they coinfect, why 559 do some parasites become more abundant than others? When can nutrient supply or resources increase a 560 parasite's relative abundance (Case I) and both relative and absolute abundance (Case II)? (C) Coinfection 561 burden: Finally, some hosts or host genotypes present higher (co)infection burden than others hinting at 562 differences in immunological resistance. (D) Parasite competition hinges on two main interactions: how (a) 563 species compete with themselves or intra-specific competition (negative intraspecific direct effect, red 564 curve), and how (b) species compete interspecifically (negative interspecific direct effect, red arrows). 565 Generally, coinfection (or within-host coexistence) occurs when intra- exceeds inter-specific competition, 566 and vice-versa for priority effects, at a feasible 'interior' equilibrium (Table S2). Here, those outcomes are 567 illustrated with direct effects (in the Lotka-Volterra model). Subsequently, those competitive effects work 568 indirectly through niche factors (like energy or immune cells).

569 Figure 2: Models of within-host competition of parasites, with potential infection outcomes. Top 570 row, single host energy or immune cells model: Two parasites can compete indirectly for (A) host energy 571 (exploitative: 2PE) or (B) via immune cells (apparent competition: 2PI). Parasites can also engage directly in interference competition (denoted by $P_1 - P_2$ interactions: grey shading). Middle row, two energy or 572 573 immune models: Two parasites can compete for (C) two sources of energy or other resources within hosts 574 (2P2E; i.e., like a resource ratio model) or for (D) two different forms of immune cells or classes (2P2I; 575 'immune ratio'; mutual inhibition between $I_1 - I_2$ is possible [grey shading]). Bottom row, energy and 576 *immune models*: Competing parasites can be attacked by shared immune cells and compete for host energy 577 with (E) induced immunity (2PIEi), and (F) with constitutive immunity (2PIEc). Immune cells require host 578 energy for proliferation resulting in additional *E-I* loops beyond those above (B, D). Red (black) arrows: 579 negative (positive) interspecific direct effects, evaluated at positive densities (a feasible interior 580 equilibrium); red curved arrow: self-limitation (negative intraspecific effect).

Figure 3: *Outcomes of within-host competition in the 2PIEi model* (A) Outcomes of competition are characterized here in a 2D-bifurcation diagram over gradients of nutrient supply point (S) and feeding rate of a parasite, $P_1(f_{P_1})$. The (a)symmetry in inter- and intra-specific competition involved governs single infection (via competitive exclusion: P_1 [blue] or P_2 [purple] only), coinfection (orange), priority effects (gray), or no infection (yellow). (B – F) Sample dynamics for one or two starting densities of P_1 and P_2 .

586 Figure 4: Elements of coinfection (coexistence) in the 2PIEi model. (A-B) Equilibrial densities 587 along a gradient of nutrient supply (S) of (A) one single parasite, P_1 (blue region, tiny dash, left panel) or 588 the other, P_2 (purple, large dash, right) or (B) by both parasites (coinfection in orange) for a fixed feeding 589 rate of P_I ($f_{P_1} = 7.2$). (C-D) A closer comparison of energy (E) and immune (I) quantities of each parasite 590 provides insight into outcomes of joint resource (energy) and apparent competition. (C) Energy 591 *competition*: Without immune attack, P_1 is the superior energy competitor $(E_1^* < E_2^*)$ allowing P_1 to win 592 (sensu the R^* rule). Mortality from immune activation weaken P_1 's competitive ability, allowing P_2 to 593 become the superior energy competitor $(E_{I,I}^* > E_{2,I}^*)$. (D) Immune- mediated apparent competition: Immune 594 cells proliferate when their minimum energy-parasite requirement is met $(EP_1^*; gray dotted; EP_2^*; dashed)$. 595 The winner of apparent competition (higher I_i^*) supports more immune cells (here: less resistant P_i). Red 596 box: Conditions promoting coinfection.

597 Figure 5: Priority effects (founder control) in the 2PIEi model using assembly rules. (A-B) 598 Equilibrial densities along a gradient of nutrient supply (S) of (A) one parasite, P_1 (blue region, tiny dash, 599 left panel) or the other P2 (purple, large dash, right) or (B) leading to priority effects (grey) for a fixed 600 feeding rate of P_I ($f_{P_1} = 10.63$). (C-D) Comparison of energy (E) and immune (I) quantities with each 601 parasite singly provides insight into competitive outcomes. (C) Energy competition: Here, P_1 always remains the superior energy competitor with $(E_1^* < E_2^*)$ or without $(E_{I,I}^* < E_{2,I})$ immune cells. (D) Immune-602 603 mediated apparent competition: Immune cells invade when the minimum energy-parasite requirement of 604 the immune cells is met $(EP_1^*; gray dotted; EP_2^*; dashed)$. In the region of priority effects, more resistant 605 P_2 wins via apparent competition (since $I_2^* > I_1^*$). Grey box: Conditions promoting priority effects.

Figure 6: Interpretation of coinfection (coexistence) and priority effects (alternate stable states) in
 the 2PIEi model using nullclines. (A) Sensitivity to their niche: Densities of energy (E) and immune cells

608 (*I*) at which parasites 1 (P_1) and 2 (P_2) show zero growth ($dP_1/dt = 0$). Combinations below these nullclines 609 (higher energy, less immune attack) fall in the fundamental niche of each parasite. (B) *Impacts on their* 610 *niche:* densities of parasite 1 (P_1) and 2 (P_2) below the energy nullcline (dE/dt = 0, yellow) or above the 611 immune (dI/dt = 0, green) nullcline lead to increases in both. At a given *S*, the outcomes of within-host 612 competition can also be visualized at the intersection of (C,F) parasite or (E,H) energy-immune cell 613 nullclines involving the (D, G) equilibrium denoting coinfection (black, closed circle: C,D,E) or priority 614 effects (white, open circle: F,G,H). (See text and Appendix S1, Tables S1,2 for more details).

615 Figure 7: A mechanistic framework for within-host parasite competition, linking feedback, trait 616 ratios and key traits in the 2PIEi model. (A) Feedback: Coinfection v priority effects is governed by the 617 sum of level four feedback loops (F_4) ; from L-R, two destabilizing, positive (+), interspecific competition 618 loops and two stabilizing, negative (-), intraspecific competition ones. P_i benefits from an increase in its 619 density (positive feedback, interspecific) as it (i) directly "fuels" or (ii) indirectly "starves" immune cells 620 but is restrained (negative feedback, intraspecific) by (iii) and (iv) " P_i is attacked, P_i eats" loops (the product 621 of binary I-P_i and P_j -E consumer-resource-like loops). (B) Differences in ratios of key traits: With some 622 rearrangement, these feedback loops correspond to ratios of key quantities made of key traits. These ratios 623 encapsulate differences in how competing parasites are (B) sensitive to $(\alpha_1 - \alpha_2)$ and have (C) impacts on 624 $(\varepsilon_1 - \varepsilon_2)$ immune cells and energy. Those differences in ratios can be written proportional to and measured 625 either as a combination of (D) traits or as (E) traits and minimal requirements. For instance, (E) α_i is 626 proportional to the product of killing rate of immune cell on parasite $j(f_{IP_i})$ and the minimal energy 627 requirement of parasite $j(E_i^*)$, and ε_i is proportional to the per capita 'consumption' of energy by parasite j, $f_{IP_i}(I)$, times minimal energy and parasite needed for immune activation (EP_i^*). 628

Figure 8: Coinfected or not? Assuming a competition-resistance trade-off between competitors, (A) the 2D bifurcation plot of resource supply to host resources (S) and feeding rate of parasite P_1 on host energy (f_{P_1}) captures divergent infection outcomes in 2PIEi (as in Fig. 3): single infection either via priority effects (grey), or competitive exclusion (blue: P_1 wins; purple: P_2 wins), coinfection (orange), or no infection (yellow). (B) *Priority effects* in mice parasitized by strains of a bacterium (*B. burgdorferi*), where the first invader excludes the later. (C) *Competitive exclusion:* A malarial strain (P_2 ; purple) is excluded by one (P_3) or *coinfects* (coexists) with another strain (P_1 ; blue). (D-F) *Coinfection community structure:* Coinfecting parasites can shift relative and absolute abundance within hosts with increasing resource supply (S). (D) In 2PIEi, increasing nutrient supply favours more resistant P_2 over the superior P_1 . (E) Empirically, shifts in community structure arise in mice coinfected by two species of gastrointestinal worms (case I) and a mosquito larvae infected by a microsporidian (case II). (F) Re-examination of all outcomes of competition from Case II suggests that variation in infection outcomes could arise when individuals assimilate different amounts of food. Small deviations among individuals could drive large variation in infection outcomes.

Figure 9: *Coinfection burden:* The burden of parasites that infect a host can depend on baseline energy allocation to immunity (a_b). (A) vs. (B): Higher a_b squeezes parameter space for coinfection (2PIEi, induced *v* 2PIEc, constitutive immunity, respectively). (C-D) It also reduces density of competing parasites. Reduced burden allows the host to maintain slightly more energy for other metabolic work, potentially improving host health. (E) Increasing a_b favours the superior energy competitor (more resistant P_2) over the superior apparent competitor (P_1), eventually excluding it. By implication, differences in a_b can lead to varying (co)infection burden among hosts.

Table 1. *A guide to future coinfection experiments:* The outcomes of within-host competition can be tested by (A) measuring species' invasion growth rates via phenomenological models (e.g., Lotka-Volterra), or (B-D) via parameterizing a model mechanistic model like those outlined in Fig. 2 - 7. Questions' outline fundamental queries surrounding causes and consequences of divergent infection outcomes. 'Mechanisms' summarize the theory, and 'Relationships' shows equations or correlations that can connect theory to experiments. 'Empirical Tests' provides examples of how those relationships can be tested across a range of systems. Figure created using BioRender.com.

656 **REFERENCES**

657

- Amarasekare, P. (2002). Interference competition and species coexistence. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 269(1509), 2541-2550.
- 660
- Budischak, S. A., Sakamoto, K., Megow, L. C., Cummings, K. R., Urban Jr, J. F., & Ezenwa, V. O. (2015).

Resource limitation alters the consequences of co-infection for both hosts and parasites. *International Journal for Parasitology*, 45(7), 455-463.

664

Budischak, S. A., Wiria, A. E., Hamid, F., Wammes, L. J., Kaisar, M. M., van Lieshout, L., ... & Graham,

A. L. (2018). Competing for blood: the ecology of parasite resource competition in human malaria–
helminth co-infections. *Ecology letters*, 21(4), 536-545.

668

- Cassat, J. E., Moore, J. L., Wilson, K. J., Stark, Z., Prentice, B. M., Van de Plas, R., ... & Skaar, E. P.
 (2018). Integrated molecular imaging reveals tissue heterogeneity driving host-pathogen
 interactions. *Science translational medicine*, *10*(432), eaan6361.
- 672
- 673 Cervi, L., MacDonald, A. S., Kane, C., Dzierszinski, F., & Pearce, E. J. (2004). Cutting edge: dendritic
 674 cells copulsed with microbial and helminth antigens undergo modified maturation, segregate the antigens
 675 to distinct intracellular compartments, and concurrently induce microbe-specific Th1 and helminth-specific
 676 Th2 responses. *The Journal of Immunology*, *172*(4), 2016-2020.
- 677
- 678 Chen, C. C., Louie, S., McCormick, B., Walker, W. A., & Shi, H. N. (2005). Concurrent infection with an
 679 intestinal helminth parasite impairs host resistance to enteric Citrobacter rodentium and enhances
 680 Citrobacter-induced colitis in mice. *Infection and immunity*, 73(9), 5468-5481.
- 681

682 Clay, P. A., Cortez, M. H., Duffy, M. A., & Rudolf, V. H. (2019). Priority effects within coinfected hosts

683 can drive unexpected population-scale patterns of parasite prevalence. *Oikos*, *128*(4), 571-583.

684	
685	Cressler, C. E., Nelson, W. A., Day, T., & McCauley, E. (2014). Disentangling the interaction among host
686	resources, the immune system and pathogens. Ecology letters, 17(3), 284-293.
687	
688	De Roode, J. C., Helinski, M. E., Anwar, M. A., & Read, A. F. (2005). Dynamics of multiple infection and
689	within-host competition in genetically diverse malaria infections. The American Naturalist, 166(5), 531-
690	542.
691	
692	Devevey, G., Dang, T., Graves, C. J., Murray, S., & Brisson, D. (2015). First arrived takes all: inhibitory
693	priority effects dominate competition between co-infecting Borrelia burgdorferi strains. BMC
694	<i>microbiology</i> , <i>15</i> (1), 61.
695	
696	Ezeamama, A. E., Friedman, J. F., Olveda, R. M., Acosta, L. P., Kurtis, J. D., Mor, V., & McGarvey, S. T.
697	(2005). Functional significance of low-intensity polyparasite helminth infections in anemia. The Journal of
698	infectious diseases, 192(12), 2160-2170.
699	
700	Ezenwa, V. O. (2021). Co-infection and nutrition: integrating ecological and epidemiological perspectives.
701	In Nutrition and Infectious Diseases (pp. 411-428). Humana, Cham.
702	
703	Ezenwa, V. O., Etienne, R. S., Luikart, G., Beja-Pereira, A., & Jolles, A. E. (2010). Hidden consequences
704	of living in a wormy world: nematode-induced immune suppression facilitates tuberculosis invasion in
705	African buffalo. The American Naturalist, 176(5), 613-624.
706	
707	Ezenwa, V. O., & Jolles, A. E. (2011). From host immunity to pathogen invasion: the effects of helminth
708	coinfection on the dynamics of microparasites. Integrative and comparative biology, 51(4), 540-551.
709	
710	Fellous, S., & Koella, J. C. (2009). Infectious dose affects the outcome of the within-host competition
711	between parasites. The American Naturalist, 173(6), E177-E184.

7	1	2
1	I	L

712	
713	Fenton, A., & Perkins, S. E. (2010). Applying predator-prey theory to modelling immune-mediated, within-
714	host interspecific parasite interactions. Parasitology, 137(6), 1027-1038.
715	
716	Freckleton, R. P., & Watkinson, A. R. (2000). Designs for greenhouse studies of interactions between
717	plants: an analytical perspective. Journal of Ecology, 88(3), 386-391.
718	
719	Fuess, L. E., Weber, J. N., den Haan, S., Steinel, N. C., Shim, K. C., & Bolnick, D. I. (2021). Between-
720	population differences in constitutive and infection-induced gene expression in threespine
721	stickleback. Molecular Ecology, 30(24), 6791-6805.
722	
723	Galli, F., Varani, M., Lauri, C., Silveri, G. G., Onofrio, L., & Signore, A. (2021). Immune cell labelling and
724	tracking: implications for adoptive cell transfer therapies. EJNMMI radiopharmacy and chemistry, 6, 1-19.
725	
726	Graham, A. L. (2008). Ecological rules governing helminth-microparasite coinfection. Proceedings of the
727	National Academy of Sciences, 105(2), 566-570.
728	
729	Griffiths, E. C., Pedersen, A. B., Fenton, A., & Petchey, O. L. (2011). The nature and consequences of
730	coinfection in humans. Journal of Infection, 63(3), 200-206.
731	
732	Griffiths, E. C., Fairlie-Clarke, K., Allen, J. E., Metcalf, C. J. E., & Graham, A. L. (2015). Bottom-up
733	regulation of malaria population dynamics in mice co-infected with lung-migratory nematodes. Ecology
734	letters, 18(12), 1387-1396.
735	
736	Grainger, T. N., Levine, J. M., & Gilbert, B. (2019). The invasion criterion: a common currency for
737	ecological research. Trends in ecology & evolution, 34(10), 925-935.
738	

Gruner, D. S., Kolekar, A., McLaughlin, J. P., & Strong, D. R. (2009). Host resistance reverses the outcome 739

Guzmán-Franco, A. W., Clark, S. J., Alderson, P. G., & Pell, J. K. (2009). Competition and co-existence of

- 740 of competition between microparasites. *Ecology*, *90*(7), 1721-1728.
- 741

- Zoophthora radicans and Pandora blunckii, two co-occurring fungal pathogens of the diamondback moth,
 Plutella xylostella. *Mycological research*, *113*(11), 1312-1321.
 Halliday, F. W., Umbanhowar, J., & Mitchell, C. E. (2018). A host immune hormone modifies parasite
 species interactions and epidemics: insights from a field manipulation. *Proceedings of the Royal Society B*, *285*(1890), 20182075.
- 749
- Halliday, F. W., Penczykowski, R. M., Barrès, B., Eck, J. L., Numminen, E., & Laine, A. L. (2020).
- Facilitative priority effects drive parasite assembly under coinfection. *Nature Ecology & Evolution*, 4(11),
 1510-1521.
- 753
- Hart, S. P., Freckleton, R. P., & Levine, J. M. (2018). How to quantify competitive ability. *Journal of Ecology*, *106*(5), 1902-1909.
- 756
- Hite, J. L., & Cressler, C. E. (2018). Resource-driven changes to host population stability alter the evolution
 of virulence and transmission. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373(1745), 20170087.
- 760
- Holt, R. D., Grover, J., & Tilman, D. (1994). Simple rules for interspecific dominance in systems with
 exploitative and apparent competition. *The American Naturalist*, *144*(5), 741-771.
- 763
- Holt, R. D. (1977). Predation, apparent competition, and the structure of prey communities. *Theoretical population biology*, *12*(2), 197-229.
- 766
- 767 Karvonen, A., Jokela, J., & Laine, A. L. (2019). Importance of sequence and timing in parasite

768 coinfections. *Trends in parasitology*, *35*(2), 109-118.

- Karvonen, A., Savolainen, M., Seppälä, O., & Valtonen, E. T. (2006). Dynamics of Diplostomum
 spathaceum infection in snail hosts at a fish farm. *Parasitology Research*, *99*(4), 341-345.
- 772
- 773 Kendig, A. E., Borer, E. T., Boak, E. N., Picard, T. C., & Seabloom, E. W. (2020). Host nutrition mediates
- interactions between plant viruses, altering transmission and predicted disease spread. *Ecology*, *101*(11),
 e03155.
- 776
- 777 Lacroix, C., Seabloom, E. W., & Borer, E. T. (2014). Environmental nutrient supply alters prevalence and
- 778 weakens competitive interactions among coinfecting viruses. *New Phytologist*, 204(2), 424-433.
- 779
- Merrill, T. E. S., & Cáceres, C. E. (2018). Within-host complexity of a plankton-parasite
 interaction. *Ecology*, 99(12), 2864-2867.
- 782
- Leibold, M. A. (1996). A graphical model of keystone predators in food webs: trophic regulation of
 abundance, incidence, and diversity patterns in communities. *The American Naturalist*, 147(5), 784-812.
- 785
- Lively, C. M., de Roode, J. C., Duffy, M. A., Graham, A. L., & Koskella, B. (2014). Interesting open
 questions in disease ecology and evolution. *The American Naturalist*, *184*(S1), S1-S8.
- 788
- Lochmiller, R. L., & Deerenberg, C. (2000). Trade-offs in evolutionary immunology: just what is the cost
 of immunity?. *Oikos*, *88*(1), 87-98.
- 791
- 792 Lotka, A. J. (1925). *Elements of physical biology*. Williams & Wilkins.
- 793
- Mideo, N., Alizon, S., & Day, T. (2008). Linking within-and between-host dynamics in the evolutionary
- repidemiology of infectious diseases. *Trends in ecology & evolution*, 23(9), 511-517.

796	
797	Moret, Y., & Schmid-Hempel, P. (2000). Survival for immunity: the price of immune system activation for
798	bumblebee workers. Science, 290(5494), 1166-1168.
799	
800	Pásztor, L., Botta-Dukát, Z., Magyar, G., Czárán, T., & Meszéna, G. (2016). Theory-based ecology: a
801	Darwinian approach. Oxford University Press.
802	
803	Puccia, C. J., & Levins, R. (1991). Qualitative modeling in ecology: loop analysis, signed digraphs, and
804	time averaging. In Qualitative simulation modeling and analysis (pp. 119-143). New York, NY: Springer
805	New York.
806	
807	Råberg, L., De Roode, J. C., Bell, A. S., Stamou, P., Gray, D., & Read, A. F. (2006). The role of immune-
808	mediated apparent competition in genetically diverse malaria infections. The American Naturalist, 168(1),
809	41-53.
810	
811	Ramesh, A., & Hall, S. R. (2023). Niche theory for within-host parasite dynamics: Analogies to food web
812	modules via feedback loops. Ecology Letters, 26(3), 351-368.
813	
814	Restif, O., & Graham, A. L. (2015). Within-host dynamics of infection: from ecological insights to
815	evolutionary predictions. Philosophical Transactions of the Royal Society B: Biological
816	Sciences, 370(1675), 20140304.
817	
818	
819	Smith, V. H. (2014). Nutrient supply differentially alters the dynamics of co-infecting phytoviruses. New
820	Phytologist, 204(2), 265-267.
821	
822	Susi, H., Barres, B., Vale, P. F., & Laine, A. L. (2015). Co-infection alters population dynamics of
823	infectious disease. Nat Commun 6: 5975.

824

- Tilman, D. (1982). *Resource competition and community structure* (No. 17). Princeton university press.
 826
- 827 Tombak, K. J., Budischak, S. A., Hauck, S., Martinez, L. A., & Rubenstein, D. I. (2020). The non-invasive
- 828 measurement of faecal immunoglobulin in African equids. International Journal for Parasitology:
- 829 *Parasites and Wildlife*, *12*, 105-112.
- 830
- 831 Vogels, C. B., Rückert, C., Cavany, S. M., Perkins, T. A., Ebel, G. D., & Grubaugh, N. D. (2019). Arbovirus
- 832 coinfection and co-transmission: A neglected public health concern?. *PLoS biology*, *17*(1), e3000130.

833

834 Volterra, V. (1926). Fluctuations in the abundance of a species considered
835 mathematically. *Nature*, 118(2972), 558-560.

836

Zuk, M., & Stoehr, A. M. (2002). Immune defense and host life history. *The American Naturalist*, *160*(S4),
S9-S22.





Within-host models of parasite competition



2 Parasite – host immune cells - 2PI Apparent competition B. P.

Parasite competition along a single niche dimension, Je., (A) energy-mediated exploitative competition (2PE), or (B) immune-mediated apparent competition (2PI), ensures exclusion when parasites grow exponentially. Adding direct interference competition (grey box) enables coirrifection and priority effects.



(C) Exploitation of two energy resources (2P2E) in resource ratio competition, or (D) apparent competition for two types of immune cells, i.e., 'immune ratio' (2P2I), enables coninfection and priority effects. Production of immune cells can interfere with each other (grey box).

2 Parasite - immune cells - host energy



Exploitative and immunemediated apparent competition enables coinfection and priority effects (2PIE).

(E) Compared to hosts with solely induced immunity (2PIE), (F) constitutive immunity (via fixed energy allocation: 2PIE) reduces opportunities for coinfection and parasite burden (see text).









B. Impacts of parasites on host niche









Empirical pattern: (E) Nutrient supply (S) can shift 'community structure' of parasites within hosts

Consistent explanation: (D) 2PE predicts shifts in relative and abouted subndance lowershift more resistant pravatis (F2) with hirds 15 in successful confections. (F) Replayed C clase II for all holes exposed to both parasites, captures divergent outcomes: no infection or desance. 79, engine infection[®] Relay at low supply (F(S)), P₂, single infection[®] Relay at his parallel supply (S4), confection where parallel excellse of species are negatively correlated across nutriest supply as predicted by (D) model. (Reploiting of Case /: Budischak et al 2015; Case /: Relate & Koelas 2009)





Table 1

A. When

competing narasites

cointers?

can

Quantian Machaniem

Relationship

Empirical Tests

in general successful coinfection occurs when intra is creater than interspecific corrotition captured by simple phenomenological models like Lotka-Volterra competition

 $\frac{\partial}{\partial t} = p_1(1 - q_2 B - q_2 B)B$



A response surface design can evaluate growth rates estimated buistra, and inter-specific compation coefficients

1) Using a response a stare design later a host with an individual parasite grown it alone with a ranne of



increase (e.g. menonerosites that multiply within hosts) intramacife (a.) and intermedite (a.) coefficients 4) Quantify proveth rates and validate results with models fit to time series data within-hosts (e.g., LV, Eleverton-Holt modali: eae Gouper 2009. Grainpar 2019.

D Milest cointection?

. Priority offects

Correction exchasion

If the leas resistant competent for anarous (without or without immune cells) while the resistant parasite is best apparent competitor offects



When nemates cointest dose does not change final outcome. despite varying initial P, and constant initial P., With priority effects, sufficiently high initial density determines wirner.

L Testing priority effects: 1) Set up a dose-dependent intection assay where dose of one parasite is varied To Measure operative departure quer time and/or immune rails and rutritional status to delineate mechanism (s.o. superior resource v resistant competitor as in Fig. 4)

L Testing competitive exclusion: 1) The above exceriment can also yield exclusion. 2) Alternately, povern competitive ability, e.g., foraging rates of parasites. To Measure pairwise competition of two or more parasite strains with different feeding rates over time (Eig 7C)

C. How does nutrient enrichment mediate etmotion?

Nutrient supply to hosts mediates densities of issmunity, and hence. relative and absolute depaition of correction. Increasing nutrient supply favours the resistant but superior but supprise areasent competitor (P.)





parasites within boats from no intection -> P, wins -> coinfection -> P. wins

1) Raise hosts that are i) singly infected, and ii) coinfected along a nutrient supplement gradient (e.g., food envicted with nitrogen, iron, etc.).

2) Measure probability of infection outcomes (Fig. 8C), and/or paraste densities, immune cells and nutritional status over time (Budlechak 2015).



31 Wth single infection data measure minimum resource requirements of parasites and immune cells that establish the energy-immune trade-off (Fig. 4D-E) between competing

d) Alternately, field observations could correlate parasite densities to host resource or immune measurements to infer mechanism: see: Tombok 2021, Ezerwa 2010

D. How does cours libertion immunity hinder oninfaction?

Lesser baseline alcostion of anamy in immunity (a.) -> higher immune resconse. Higher immune nerasile bunten (P.'+ Ps' that allows the host to maintain alightly higher energy (F) for office matchells made



Constar immuna cell dansity (due to tell -> L conocturity for coinfection → 1 and excludes registant Ps freeing up energy for metabolic needs. Investment in immunity without compensatory intake leads to host death.

Effect of constitutive immunity on burden: Expose hosts with differences in constitutive immunity i.e. differences in immune resistance to multiple neresite strains. Houts should have a higher probability of coinfections y single infections with no / low constitutive immunity.

Cost of immune maintenance: Evonse hosts to inactivated parasites or stimulate immune system (e.g. LPS beads) under starvation. Lack of compensatory nutritional intake can affect host survival, conditio and reproduction. Differences between control maintenance, are: Moret & Schmid-Herroel 2000