

1 **Research Letter**

2 **Temporal changes in sleep parameters and body mass index after using the Pokémon**

3 **Sleep app**

4

5 **Masao Iwagami<sup>1,2+</sup>, Jaehoon Seol<sup>1,3,4+</sup>, Masashi Yanagisawa<sup>1,5</sup>**

6 <sup>1</sup>International Institute for Integrative Sleep Medicine (IIIS), University of Tsukuba, Tsukuba,

7 Ibaraki, Japan

8 <sup>2</sup>Institute of Medicine, University of Tsukuba, Tsukuba, Ibaraki, Japan

9 <sup>3</sup>Institute of Health and Sport Sciences, University of Tsukuba, Tsukuba, Ibaraki, Japan

10 <sup>4</sup>Department of Frailty Research, Center for Gerontology and Social Science, National Center

11 for Geriatrics and Gerontology, Obu, Aichi, Japan

12 <sup>5</sup>S'UIMIN Inc., Shibuya, Tokyo, Japan

13 <sup>+</sup>Co-first authors (contributed equally)

14

15 **Corresponding author:**

16 Masashi Yanagisawa

17 International Institute for Integrative Sleep Medicine (IIIS), University of Tsukuba, 1-1-1

18 Tennodai, Tsukuba, Ibaraki, 305-8575, Japan

19 Email: [yanagisawa.masa.fu@u.tsukuba.ac.jp](mailto:yanagisawa.masa.fu@u.tsukuba.ac.jp)

## 20    **Abstract**

21    Pokémon Sleep is a new sleep-tracking game app released in July 2023. We examined  
 22    whether sleep parameters would improve with the use of the Pokémon Sleep app, and  
 23    whether participants' improvement status was associated with changes in their body mass  
 24    index (BMI). We analyzed data from 2,063 volunteer Japanese participants (mean age 38.3 ±  
 25    10.7 years, female 82.1%) who used both the Pokémon Sleep and Asken (a commonly-used  
 26    app in Japan to record daily BMI) for ≥90 days. During the 90 days, improvement was seen  
 27    in total sleep time (TST) for 44.8% of participants, sleep latency for 18.1%, and percentage  
 28    of wakefulness after sleep onset for 24.4%. BMI tended to decrease faster in participants with  
 29    improved TST and sleep latency. In conclusion, the Pokémon Sleep app can improve some  
 30    sleep parameters, and by association, lead to improvement in BMI.

## 31 Introduction

32 In the era of digital health, a variety of mobile apps have been developed to record daily  
33 health-related parameters (e.g., step counts and screen time), with some including game  
34 elements to nudge users towards better health behaviors. Among these apps are a small  
35 number designed to track sleep behaviors and characteristics; however, their influence on  
36 users' behavioral and clinical outcomes remains unknown [1, 2].

37 Pokémon Sleep (The Pokémon Company, Tokyo, Japan) is a sleep-tracking app  
38 released in July 2023 in multiple countries, including Japan and the United States. Compared  
39 to previous sleep-tracking apps [1, 2], Pokémon Sleep has a stronger game element,  
40 rewarding users with points for longer sleep and the ability to raise Pokémon. Therefore, it  
41 can be hypothesized that key sleep parameters, such as total sleep time (TST), sleep latency  
42 (the duration from the start time in bed to sleep onset), and wakefulness after sleep onset  
43 (WASO), will improve after starting Pokémon Sleep. Further, we hypothesized that people  
44 whose sleep improved after starting Pokémon Sleep would also have improved health  
45 indicators, specifically, body mass index (BMI).

46 To examine these hypotheses, we conducted a retrospective observational study of  
47 data from users of both Pokémon Sleep and Asken (asken, Inc., Tokyo, Japan), an app  
48 commonly used in Japan to record daily diets and BMI [3].

49

## 50     **Methods**

51     Details of the Pokémon Sleep and Asken apps are described elsewhere [3, 4]. In brief,  
 52     Pokémon Sleep tracks and records daily sleep characteristics using the accelerometer  
 53     functions of mobile phones or smart devices named Pokémon GO Plus+. Users are instructed  
 54     to keep the app open and place the device by their pillow, or other location where it will  
 55     detect only the user's body movements. Using algorithms established by Cole and Kripke [5],  
 56     Pokémon Sleep distinguishes between periods of sleep and wakefulness. From these records,  
 57     we calculated daily TST, sleep latency, and the proportion of WASO (%WASO) during TST.  
 58     Asken is a calorie and nutrition tracker that also monitors daily BMI.

59             In-app messages were sent to all Asken users from January 19 through 31, 2024,  
 60     asking those who used both Pokémon Sleep and Asken for permission to use their stored data.  
 61     Respondents were asked to provide their Pokémon Sleep user IDs in their electronic informed  
 62     consent so that the Pokémon Sleep and Asken data were linkable. The study was approved by  
 63     the Institutional Review Board of Sapporo Yurinokai Hospital, Japan.

64             Using the retrieved data, we identified when each participant had started using  
 65     Pokémon Sleep ("day 0"), and included those continuing it until day 89, with or without gaps.  
 66     Persons were excluded if they had not used Asken during the 90 days of Pokémon Sleep  
 67     recordings, or did not have BMI recordings at days 0 and 89 (i.e., the first and last dates). For

gaps or days when the participants did not use Pokémon Sleep or Asken, missing data were imputed linearly from the records immediately before and after the day.

In statistical analysis, we summarized the basic characteristics of participants including age, sex, and baseline BMI. We also cross-sectionally estimated cubic spline curves for each sleep parameter (days 0–6 average) with BMI on day 0, based on a generalized additive model. Second, we plotted the average trend of each sleep parameter from days 0 to 89. For each sleep characteristic, we then classified participants into two groups, improved/not improved, based on the slope calculated for individuals during the 90 days. We regarded a slope  $>0$  as an improvement in TST, and a slope  $<0$  as an improvement in sleep latency and %WASO. Finally, for each sleep parameter, we plotted the average change in BMI from days 0 to 89 by group. Repeated-measures analysis of variance (ANOVA) at days 0, 29, 59, and 89 was used to test whether the temporal trend in BMI differed between the groups, adjusting for age and sex.

All analyses were performed using Python (version 3.12.2), which was built into Visual Studio Code (version 1.87.2). Statistical significance was set at  $P < 0.05$  (2-tailed).

## Results

Of 6,052 responses to our request, 3,934 participants used Pokémon Sleep for  $\geq 90$  days, and 2,063 participants were included for the analysis (**Supplementary Figure 1**). Of the 90 days,

the mean number of days spent on Pokémon Sleep and Asken was 64.1 (standard deviation [SD] 16.6 days) and 59.6 (SD 28.6 days), respectively. Participants' baseline characteristics are shown in **Supplementary Table 1**. In the cross-sectional analysis (**Supplementary Figure 2**), longer TST (up to around 8.5 hours) and shorter sleep latency were associated with lower BMI.

On average, the TST increased from days 0 to 89 by approximately 0.5 hours, whereas sleep latency and %WASO were nearly constant (**Supplementary Figure 3**). However, calculating the slope for individuals, 924 (44.8%), 374 (18.1%), and 504 (24.4 %) participants were classified as “improved” for TST, sleep latency, and %WASO, respectively. Participants' baseline characteristics by improvement status are shown in **Supplementary Table 2**.

**Figure 1** shows the temporal changes in BMI by improvement status groups. Regardless of the groups, participants' BMI tended to decrease during the 90 days. However, participants with improved TST (**panel A**) showed a steeper and more sustained decline in BMI than did the non-improved group, although the difference was not statistically significant in the repeated-measures ANOVA ( $P=0.107$ ). The mean change in BMI at day 89 was -0.38 and -0.19, respectively (unadjusted difference=-0.19, 95% confidence interval [CI]: -0.40–0.03; age-sex-adjusted difference=-0.25, 95% CI: -0.51–0.01). Participants with improved sleep latency (**panel B**) showed a significantly faster decline in BMI than the non-

improved group ( $P=0.030$ ). The mean change in BMI at day 89 was -0.51 and -0.23, respectively (unadjusted difference=-0.28, 95% CI: -0.56–0.00; age-sex-adjusted difference=-0.36, 95% CI: -0.69–0.02). There was no significant difference ( $P=0.851$ ) in the amount of BMI change by %WASO improvement status (**panel C**).

## Discussion

In this study of 2,063 volunteer participants using both Pokémon Sleep and Asken for  $\geq 90$  days, improvement was seen in TST for 44.8% of participants, sleep latency for 18.1%, and %WASO for 24.4%. Participants' BMI tended to decrease during the 90 days, most likely because Asken users are typically motivated to control their diets and decrease their body weight. However, BMI tended to decrease faster in participants with improved TST (without a statistical significance) and sleep latency (with a statistical significance). Potential mechanisms may include improvement of appetite-related hormones (e.g., ghrelin and leptin), or changes in energy metabolism and balance [6, 7].

This study had several limitations. The Pokémon Sleep app was the first to use the accelerometer functions of mobile phones to measure sleep on the bed. To our knowledge, its validity in distinguishing between sleep and wakefulness has not been rigorously tested in comparison to polysomnography or actigraphy. Second, those who used both Pokémon Sleep

and Asken for  $\geq 90$  days may be highly health-conscious and the generalizability of our findings to more general populations is thus unknown.

Despite these limitations, to our knowledge, this is the first evidence in the general population that a sleep-tracking app can improve some sleep parameters, and by association, lead to improvement in physical health parameters (i.e., BMI). Further studies are warranted to further examine whether the use of sleep-tracking apps can improve sleep and physical health parameters, and ultimately, their long-term health.

## Acknowledgements

We express our gratitude to the personnel of The Pokémon Company (Tokyo, Japan), asken Inc. (Tokyo, Japan), and S'UIMIN Inc. (Tokyo, Japan) for their contributions to data preparation. We also thank Editage ([www.editage.com](http://www.editage.com)) for the English language editing.

## Data Availability Statement

The data underlying this article cannot be shared publicly for the privacy of individuals that participated in the study. The data may be shared on reasonable request to the corresponding author.

## Disclosure Statement



143 M.Y. was paid by The Pokémon Company (Tokyo, Japan) for consultation in developing the  
144 Pokémon Sleep app. M.I. and J.S. declare no conflicts of interest.

145

## 146 **References**

147 1. Robbins R, Seixas A, Walton Masters L, et al. Sleep tracking: A systematic review of the  
148 research using commercially available technology. *Curr Sleep Med Rep*.

149 2019;5(3):156-163. doi: [10.1007/s40675-019-00150-1](https://doi.org/10.1007/s40675-019-00150-1).

150 2. Behar J, Roebuck A, Domingos JS, Geder E, Clifford GD. A review of current sleep

151 screening applications for smartphones. *Physiol Meas*. 2013;34(7):R29-R46. doi:

152 [10.1088/0967-3334/34/7/R29](https://doi.org/10.1088/0967-3334/34/7/R29).

153 3. Matsuzaki E, Michie M, Kawabata T. Validity of nutrient intakes derived from an Internet

154 Website dish-based dietary record for self-management of weight among Japanese

155 women. *Nutrients*. 2017;9(10). doi: [10.3390/nu9101058](https://doi.org/10.3390/nu9101058).

156 4. The Pokémon Company. Pokémon sleep. [https://support.pokemon.com/hc/en-](https://support.pokemon.com/hc/en-us/categories/360005098972-Pok%C3%A9mon-Sleep)

157 [us/categories/360005098972-Pok%C3%A9mon-Sleep](https://support.pokemon.com/hc/en-us/categories/360005098972-Pok%C3%A9mon-Sleep). Accessed May 29, 2024.

158 5. Cole RJ, Kripke DF, Gruen W, Mullaney DJ, Gillin JC. Automatic sleep/wake

159 identification from wrist activity. *Sleep*. 1992;15(5):461-469. doi:

160 [10.1093/sleep/15.5.461](https://doi.org/10.1093/sleep/15.5.461).

6. Stich FM, Huwiler S, D'Hulst G, Lustenberger C. The potential role of sleep in promoting a healthy body composition: Underlying mechanisms determining muscle, fat, and bone mass and their association with sleep. *Neuroendocrinology*. 2022;112(7):673-701. doi: [10.1159/000518691](https://doi.org/10.1159/000518691).
7. St-Onge MP, Cherta-Murillo A, Darimont C, Mantantzis K, Martin FP, Owen L. The interrelationship between sleep, diet, and glucose metabolism. *Sleep Med Rev*. 2023;69:101788. doi: [10.1016/j.smrv.2023.101788](https://doi.org/10.1016/j.smrv.2023.101788).

## Figure legends

**Figure 1. Temporal changes in body mass index by improvement status of total sleep time (panel A), sleep latency (panel B), and % wakefulness after sleep onset (panel C)**

BMI, body mass index; %WASO, percentage of wakefulness after sleep onset; SD, standard deviation.

## Supplemental files


**Supplementary Figure 1. Participant flow chart**

**Supplementary Table 1. Baseline characteristics of study participants**

**Supplementary Figure 2. Cross-sectional associations between sleep parameters and body mass index at baseline**


180     **Supplementary Figure 3. Temporal changes in sleep parameters, overall and by**  
181     **improvement status**  
182     **Supplementary Table 2. Baseline characteristics of study participants by improvement**  
183     **status for sleep parameters**

(A) Subgroup by improvement status of total sleep time




Mean change in BMI from day 0 (SD)	Day 29	Day 59	Day 89
Not improved (n=1,139)	-0.11 (0.60)	-0.19 (0.93)	-0.19 (2.43)
Improved (n=924)	-0.14 (0.53)	-0.25 (1.08)	-0.38 (2.55)

**(B) Subgroup by improvement status of sleep latency**



Mean change in BMI from day 0 (SD)	Day 29	Day 59	Day 89
Not improved (n=1,689)	-0.11 (0.58)	-0.20 (0.97)	-0.23 (2.48)
Improved (n=374)	-0.14 (0.53)	-0.29 (1.07)	-0.51 (2.51)

### (C) Subgroup by improvement status of %WASO



Mean change in BMI from day 0 (SD)	Day 29	Day 59	Day 89
Not improved (n=1,559)	-0.11 (0.60)	-0.21 (1.02)	-0.28 (2.49)
Improved (n=504)	-0.16 (0.49)	-0.24 (0.87)	-0.27 (2.50)