

Genomic insight into the *Comamonas kerstersii* isolated from diarrheal patients in Bangladesh

Noshin Ibnat Rib ^{1,2}, Fariza Shams ², Sakib Abrar Hossain ¹, Abdus Sadique ¹, Jahidul Alam ¹, Pronoy Debnath^{1,2}, Arman Hossain ^{1,2}, Aura Rahman ², Syeda Naushin Tabassum ^{1,2}, Tahrima Saiha Huq ², Maqsud Hossain ^{1,2*}

¹ NSU Genome Research Institute (NGRI), North South University

² Department of Biochemistry and Microbiology, North South University

***Corresponding author:**

Maqsud Hossain: muhammad.maqsud@northsouth.edu

Abstract

This study marks the first identification and genomic characterization of *Comamonas kerstersii* isolates from diarrheal patients in Bangladesh. *C. kerstersii* is a gram-negative opportunistic bacterium which has also been reported in several clinical samples, including urine, blood, stool, peritoneal liquid, sputum. We carried out the whole genome sequencing of three *C. kerstersii* isolates for the first time in Bangladesh with an aim to analyze genomic features in 3 different strains isolated from Bangladesh. Seven AMR genes, aph(6)-Id, aph(3'')-Ib, mph(E), mph(F), msr(E), sul2, and tet(A), have been observed in all the strains. We identified several virulence factors including copper homeostasis genes, type IV pili and type VI secretion system. phage regions and IncQ1 plasmid were also detected.

Keywords: *Comamonas kerstersii*, emerging bacteria, Bangladesh, diarrheal patient

Introduction

The increasing number of persistent infections caused by relatively unknown non-fermenting gram-negative rods is a crucial source of concern in healthcare environments (Ryan and Pembroke 2018). These emerging pathogens, which include species from the *Comamonas* genus, are opportunistic and pose severe risk to global health (Ryan et al. 2022). One of the little-known species in the *Comamonas* genus is *Comamonas kerstersii* (Rong et al. 2022). *C. kerstersii* was first recognized as a distinct species in 2003, alongside *Comamonas aquatica* and *Comamonas terrigena* (Wauters et al. 2003a; Opota et al. 2014a; Almuzara et al. 2018; Rong et al. 2022). This species is a part of the rRNA homology group III and defined by the presence of a unique 23S rRNA gene and a distinct 16S rRNA gene sequence (DE VOS et al. 1985; Wauters et al. 2003a; Willems and De Vos 2006). *C. kerstersii* was initially included in the *C. terrigena* complex (Almuzara et al. 2017). Willems, et al. (1991) studied the taxonomy of the genus *Comamonas* and suggested that *C. terrigena* contains 3 different DNA group. Based on this information, Wauters, et al. (2003) studied the species and suggested that it comprise three different species based on 16S rRNA sequencing and DNA-DNA hybridization. They renamed DNA group II of *C. terrigena* as *C. aquatica* and DNA group III as *C. kerstersii*.

C. kerstersii species are non-fermenting, gram-negative, rod shaped bacillus (Kumar et al.; Opota et al. 2014b; Voitenkova et al. 2018; Zhou et al. 2018; Almuzara et al. 2018; Wu et al. 2018; Liu et al. 2020b, a; Onya et al. 2020a, b; S et al. 2020; Farfán-Cano et al. 2021; Rong et al. 2022). 24 hours of incubation in ambient aerobic environment shows the growth of the species (Zhou et al. 2018). Better observation can be done if incubated for 48 hours and usually colonies have a diameter of 1-2 mm and seem translucent, smooth-surfaced, and nonpigmented (Zhou et al. 2018; Palacio et al. 2020). *C. kerstersii* has previously been considered nonpathogenic (Jiang et al., 2018; Liu et al., 2020b; Opota et al., 2014b; Rong et al., 2022; Voitenkova et al., 2018). It is extensively found in soil, plants, and water with the ability to withstand aquatic environment (Farfán-Cano et al. 2021; Mandal et al. 2022; Rong et al. 2022; Wang et al. 2024). But since 2013, the species have been found in several clinical samples like abdominal fluid, peritoneal fluid, blood, urine, stool and recently in sputum (Almuzara et al. 2013, 2017, 2018; Biswas et al. 2014; Opota et al. 2014b; Palacio et al. 2020; Liu et al. 2020a; Farfán-Cano et al. 2021; Mandal et al. 2022; Wang et al. 2024). Several cases of intra-abdominal including appendicitis, peritonitis, diverticulosis, and few extra abdominal cases including urinary tract infection, bacteremia and pneumonia have been reported in number of papers (Farfán-Cano et al. 2021; Ryan et al. 2022; Wang et al. 2024). In most cases the infections associated with *C. kerstersii* were polymicrobial (Farfán-Cano et al. 2021; Bennani et al. 2022; Ryan et al. 2022)

Jiang et al (2018) studied *C. kerstersii* 8943 and concluded in their study that the genome of *C. kerstersii* is circular with chromosome measuring 3,547,915 base pairs. The genome size is notably smallest among the *Comamonas* species studied aligning with previous findings. *C. kerstersii* 8943 has been identified to have numerous resistance genes, including tetA, strB, sul1, blaOXA-1, strA, sul2, catB3, and floR (Jiang et al. 2018). Another study shows that *C. kerstersii* is resistance to Ciprofloxacin (Almuzara et al. 2013). The strain was also resistant to penicillin and ampicillin, and it tested positive for the *strA* and *strB* genes, which confer streptomycin resistance (Jiang et al. 2018) while another study reported resistance to ampicillin (Pavone et al. 2021). Genetic analysis of the strain *C. kerstersii* 3132976 revealed three β -lactamase genes (CDSs H8N02_05890, H8N02_08740, and H8N02_17110) and among them H8N02_08740 and H8N02_17110 belong to class A and C β -lactamases, respectively (Moser et al. 2021). In a recent paper, it reported that the *C. kerstersii* strain 121606 was resistant to almost all of the β -lactam class of antibiotic indicating that it has a extensive drug resistance phenotype (Wang et al. 2024).

Species of the genus *Comamonas* are associated with certain human infection and has diversified virulence factors (Barbaro et al. 1987; Mandal et al. 2022). To date, Wu, et.al. (2018) one of the pioneers who studied virulence of *C. kerstersii* along with other species of the genus. The researchers discovered that *C. aquatica*, *C. terrigena*, and *C. kerstersii* constitute a unique cluster with common virulence features, such as genes for bacterial motility, adhesion, and chemotaxis signaling (Wu et al. 2018). They have adhesin biosynthesis genes like *IlpA* and *Hsp60*, as well as biofilm-forming components, hemolysin genes and share stress proteins like as *ClpEP* and *SodB* for cellular defense (Wu et al. 2018).

Given its widespread distribution and increasing detection in clinical samples, it is critical to investigate the genomic characteristics and pathogenic potential of *C. kerstersii*, especially in regions with limited data. This study aims to bridge that knowledge gap by presenting the first comprehensive genomic analysis of *C. kerstersii* isolates from diarrheal patients in Bangladesh. It also provides new insights into the antimicrobial resistance and potential virulence factors of these isolates offering new insights into their antimicrobial resistance and potential virulence factors from Bangladesh.

Material and Methods

Sample Collection and Isolation

From March to November 2020, 17 stool samples were collected from the Dhaka Central International Medical College & Hospital (DCIMCH) to isolate *Vibrio cholerae*. The samples were carried in 5ml Cary Blair transport medium at 4°C and then incubated at 37° in 3ml Alkaline peptone water (APW) for

enrichment. From the enriched culture, tellurite taurocholate gelatin agar (TTGA) and thiosulfate citrate bile-salts sucrose (TCBS) agar plates were inoculated and incubated overnight. Yellow colonies from TCBS and black-centred colonies from TTGA, presumptively identified by their morphology, were further streaked on gelatin agar (GA). Subsequent Gram staining and routine biochemical tests were performed for further characterization.

Genome Assembly and Annotation

Isolates were inoculated in LB broth and grown at 37°C overnight for DNA extraction. Genomic DNA was extracted using QIAamp® DNA Mini Kit (QIAGEN, Hilden, Germany) according to manufacturer's protocol. Library preparation and sequencing of 3 selected strains were carried out at Genome Research Institute of North South University (NGRI), Bangladesh. High molecular weight genomic DNA of the strains was used to prepare libraries using a Nextera DNA Library Prep Kit and employed for 250-bp paired-end whole genome sequencing with Illumina® MiSeq platform following the instruction provided by the manufacturer. High-quality paired-end Illumina sequencing data ($Q \geq 30$) were obtained for the bioinformatic analysis. Quality control was performed on the sequencing data to ensure the integrity and accuracy of the information. SPAdes genome assembly software (version 3.15.5) (Bankevich et al. 2012) was used for the *de novo* assembly of the genome with particular filters employed to lower the amount of mismatches and short indels, improving the quality of the generated contigs. To decipher the genetic content and functional elements encoded within the assembled contigs, the PROKKA pipeline (version 45) (Seemann 2014) was employed for annotation. Rapid Annotation using Subsystem Technology (RAST) (Aziz et al. 2008), an automated service for annotating genomes of bacteria and archaea, was also employed for annotation.

Genomic Feature determination

Nine genomes of *C. kerstersii* were obtained from the National Center for Biotechnology Information (NCBI). The obtained genomes were annotated using the Prokka pipeline (Seemann 2014). The core genome alignment was generated using roary (Page et al. 2015) and then used in FastTree (Price et al. 2009) to construct a midpoint rooting phylogenetic tree and visualized using FigTree. Assembled genomes were visualized using the BLAST ring image generator (BRIG) (Alikhan et al. 2011). Antimicrobial resistance genes were identified utilizing the Center for Genomic Epidemiology (CGE) based tool ResFinder (version 4.1) (Bortolaia et al. 2020). In this study, Virulence Factor Database (VFDB) (<http://www.mgc.ac.cn/VFs/>) has been used to identify and analyze virulence factors associated with the studied strains (Liu et al. 2019). Identification of plasmids were conducted employing PlasmidFinder (Carattoli et al. 2014) and phage regions using PHASTEST (Wishart et al. 2023) and Phigaro (Starikova et al. 2020) respectively.

Result

Bacterial Identification

Of the 17 presumptive isolates, 14 demonstrated gelatin hydrolysis when tested on gelatin agar (GA) plates. The remaining three isolates did not exhibit gelatin hydrolysis, suggesting a different enzymatic profile. These three strains (NG13, NG14, and NG17) also produced a biochemical profile consistent with *Comamonas* species rather than *Vibrio cholera* (Table 1).

Genomic Features

We conducted next-generation sequencing with the genomic DNA isolated from the three putative *Comamonas* strains NG13, NG14, and NG17. The draft genomes of strains NG13, NG14 and NG17 resulted in 50, 68, 81 contigs, respectively. The size of the assembled genomes of the 3 strains were 3.4 Mbp with approximately 59% GC content. NG13 and NG17 possess 94 tRNAs, NG14 possesses 97 tRNAs. All three strains contain 7 rRNAs and 1 tmRNA. The genomic features of the strains are summarized in Table 2.

Phylogeny Analysis

A core genome-based phylogenetic tree consisting of 12 *C. kerstersii* strains was constructed, which is shown in Figure 2. The tree shows two distinct clusters - Cluster A and Cluster B.

According to the phylogenetic tree, three *C. kerstersii* strains under study - NG13, NG14 and NG17, shared the same ancestor and belonged to Cluster A, with CP020121 being the closest relative. CP020121, also named as *C. kerstersii* 8943, was isolated from the effluent of peritoneal dialysis (Jiang et al. 2018), highlighting the close relationship among human-originated isolates. All these strains share features like ampicillin resistance, the presence of *sul2* and *tetA* genes conferring resistance to sulfonamide and tetracycline respectively. sulfonamides(Jiang et al. 2018).

The CP060413 strain, from the other cluster- Cluster B, was isolated from rectal swab (Moser et al. 2021) like the strains of the study. Even being from the same clinical source, The CP060413 branched distantly. This strain did not have any resistance genes other than 2 class A and 1 class C beta-lactamase genes.

The *C. kerstersii* strains used in the phylogenetic analysis were isolated either from human feces or as members of the gut microbiota. The presence of various resistance genes across different strains underscores the role of horizontal gene transfer in enabling adaptation to diverse environments.

Identification of antimicrobial resistance genes (ARG)

Whole genome sequencing revealed the presence of multiple antibiotic resistance genes in all three *C. kerstersii* strains using ResFinder (Table 3). The tet(A) gene, which potentially conferred resistance by exporting tetracycline out of the cell in exchange for protons, was identified (Nguyen et al. 2014). The strains possessed mph(E), mph(F), and msr (E) genes. The mph(E) and mph(F) encode macrolide phosphotransferase enzymes that likely inactivate macrolides through phosphorylation (Leclercq 2002). The msr(E) gene may confer resistance to macrolides, lincosamides, and streptogramin B (Kadlec et al. 2011). The presence of the aph(3'')-Ib and aph(6)-Id genes suggests resistance to streptomycin. These genes are often linked and encode enzymes that inactivate streptomycin (Navas et al. 2016). The presence of sul2 gene confer resistance to sulfonamides.

We found macA, macB and tolC encoding for macrolide export protein MacA, macrolide export ATP-binding/permease protein MacB and outer membrane protein TolC, respectively. This MacAB-TolC comprise tripartite efflux system which is described to contribute in resistance to macrolide class of antibiotics in *E. coli* and other Gram-negative pathogenic bacteria (Kobayashi et al. 2001). This complex is also responsible for transportation of, protoporphyrin (Turlin et al. 2014) and lipopolysaccharides (Lu and Zgurskaya 2013) and virulence factors i.e. enterotoxin II (Yamanaka et al. 2008). Besides these, a number of multidrug resistance protein encoding genes are also present in the genomes of *C. kerstersii* strains (Supplementary Table 1).

Adaptive Mechanism and Virulence

C. kerstersii can adapt to diverse environments, and annotating the genomes revealed some of the genes that enable them to withstand environmental stressors, including polyphosphate kinase (ppk). Polyphosphate (polyP) has previously been linked to osmotic and nutritional stress response in *E. coli* and bacterial growth, survival, and pathogenicity in other bacteria (Ault-Riché et al. 1998; Rao et al. 2009). The presence of this enzyme explains how *C. kerstersii* is equipped to withstand osmotic stresses in various aquatic environments like blood, urine, or peritoneal fluid. Polyphosphate reserves in *Vibrio cholerae* improve the bacterium's resistance to environmental stressors, especially when there is a shortage of phosphate (Jahid et al. 2006). The fact that *C. kerstersii* has polyphosphate kinase indicates that how it is adapted to survive in low-phosphate conditions, like those found in excrement in this study. PolyP has also been reported to be related with virulence including motility and biofilm formation in few studies (McMeechan et al. 2007; Candon et al. 2007; Lv et al. 2022).

The *C. kerstersii* strains of the study possess *osmY*, a periplasmic protein which is strongly induced by hyperosmotic stimuli. Like *E. coli*, *C. kerstersii* may withstand external osmotic pressure like in bladder or colon and Yim and Villarejo (1992) report that *E. coli* express *osmY* proteins in such conditions. The outer membrane protein A (*ompA*) precursor of *C. kerstersii* may enable it to withstand high osmolarity as the protein equips *E. coli* to survive in acidic environment, high osmolarity, and pooled human serum (Wang 2002). *OmpA* has also been associated with virulence factors like adhesion, invasion, intracellular survival, host defense evasion, and these activities are especially important in the urogenital, respiratory, or nervous system diseases (Confer and Ayalew 2013). The inner membrane protein required for strong colonization of the gut is encoded by the *cvpA* gene (Warr et al. 2020), and the presence of this gene in *C. kerstersii* strains of the study explains its adaptability in gut of the diarrheal patients. Additionally, a colicin V producing gene (*cvpA*), crucial for gut colonization, is present in *C. kerstersii* strains (Gilson et al. 1990; Warr et al. 2020) and the presence of this gene in *C. kerstersii* strains of the study explains its adaptability in gut of the diarrheal patients.

Analysis using the Virulence Factor Database (VFDB) identified key proteins potentially associated with pathogenicity in all three *C. kerstersii* strains (**Table 4**). Notable findings include proteins involved with type IV pili (T4P), type II secretion systems (T2SS), type VI secretion system (T6SS), general secretion pathways, and flagella. Notably, both NG13 and NG14 strains encoded proteins involved in T4P biogenesis, including PilB and PilF. Strain NG17 possessed a PilJ protein. T4P play a crucial role in pilus formation, adhesion, and bacterial motility, all of which contribute to pathogenesis. T4P play an important role in pilus formation, adhesion, and dynamics (Melville and Craig 2013).

The presence of genes encoding proteins involved in flagellar morphogenesis, assembly, and chemotaxis (Supplementary Table 1), emphasize the role of mobility in bacterial pathogenic strategies. Bacteria navigate their environments using a combination of motility and chemotaxis. Motility allows them to actively seek out nutrients, while chemotaxis acts like a "chemical GPS," guiding them toward favorable conditions and allows bacteria to rapidly adapt to changing environmental conditions (Schauer et al. 2018). In pathogenic bacteria, chemotaxis contributes to host invasion and infection by allowing the bacteria to sense and move towards host tissues and nutrients (Colin et al. 2021).

In general, there are noticeably few classical virulence determinants revealed by VFDB analysis. A thorough examination of the *C. kerstersii* genome, however, revealed a number of toxin coding sequences (CDS), including zona occludens toxin (*zot*) protein-encoding sequences that may play virulence-related activities. Another example is *apxIB*, which encodes Toxin RTX-I translocation ATP-binding protein, involved in the transport of the toxin RTX-I as well as that of RTX-II. RTX (Repeats-in-toxin) toxins are glycine- and aspartate-rich repeats containing exoproteins secreted by type I secretion system of Gram-

negative bacteria(Linhartova et al. 2015) . One classical RTX protein is hemolysin which is encoded by *tlyA* in the genomes of *C. kerstersii* strains. The members of the RTX toxin family are important virulence factors of many Gram-negative bacteria (Linhartova et al. 2015). Another gene is *ratA* which encodes Ribosome association toxin RatA. This protein is the toxic component of type II toxin-antitoxin (TA) system. In *E.coli* association of RatA protein with 50S subunit of ribosome prevents the binding of small subunit 30S, thereby preventing the formation of 70S ribosomes leading to decrease in protein translation (Zhang and Inouye 2011).

The *C. kerstersii* strains possess genes that encode proteins involved in alginate biosynthesis pathway, Phosphomannomutase/phosphoglucomutase, Poly(beta-D-mannuronate) C5 epimerase 1, Poly(beta-D-mannuronate) C5 epimerase 5 encoded by *algC*, *algE*, *algE1*, respectively. A negative regulator of alginate synthesis called alginate biosynthesis sensor protein KinB, encoding gene *kinB* is also present in the genome. *AlgC* protein plays a vital role in producing alginate, LPS and rhamnolipid. three important *P. aeruginosa* virulence-associated factors Alginate producing *P. aeruginosa* strains have a mucoid phenotype and are protected from antibiotics and other host defense components (Olvera et al. 1999).

Clp protease (caseinolytic protease; ClpP) is a highly conserved serine protease found in many bacteria. This ATP-dependent protease functions as a two-component complex consisting of a regulatory ATPase subunit and a proteolytic subunit. While the proteolytic subunit possesses some intrinsic catalytic activity, both components are essential for optimal enzymatic function in the presence of ATP. We identified genes encoding various components of the Clp protease complex, including *clpP* and *clpP1* (putative Clp proteolytic core), *clpA*, *clpC*, and *clpX* (Clp-ATPases), and *clpS* (adaptor protein). Clp proteases play crucial roles in various cellular processes, including the degradation of misfolded proteins, regulation of short-lived proteins, and housekeeping removal of dysfunctional proteins. They also control cell growth and target DNA-binding proteins from starved cells (Frees et al. 2007; Meyer and Baker 2011). ClpP has also been linked to the tight regulation of virulence genes in both Gram positive and Gram negative pathogens such as *Staphylococcus aureus*, *Listeria monocytogenes* and *Salmonella typhimurium*. Therefore, the presence of Clp protease complex encoding genes indicate the contribution of this system in the pathogenic lifestyle of the *C. kerstersii* strains (Frees et al. 2005, 2007).

Iron is often limited in host environments as part of the innate immune response, so bacteria need to actively scavenge iron from sources like hemoglobin. This iron homeostasis is closely linked to bacterial

virulence, stress resistance, and overall fitness. We examined the *C. kerstersii* genomes and found that a number of genes are present to encode enzymes and proteins involved in iron uptake mechanisms. Enterobactin is a siderophore that can extract iron from host iron-binding proteins like transferrin. Haemoglobin-haptoglobin binding proteins allow bacteria to retrieve heme iron from haemoglobin, a major iron source in the host (Chen et al. 2018). Iron acquisition regulators like Fur control the expression of iron uptake systems in response to iron availability, allowing bacteria to tightly regulate iron uptake to avoid toxicity while ensuring sufficient iron for growth (Chen et al. 2018) All these proteins are key adaptations that enable *C. kerstersii* to efficiently scavenge iron from the host, a critical nutrient for growth and virulence. In addition, several genes encode key enzymes in the heme biosynthesis pathway that is essential for bacterial growth, metabolism, and virulence. Targeting these enzymes could be a potential antimicrobial strategy.

A cluster of genes involved in copper homeostasis was found in the annotated genomes of *C. kerstersii* strains (Table 5). Li et al. mentioned that a crucial factor in the pathophysiology of bacteria is copper homeostasis, which involves the complex interaction of bacterial survival strategies and host defenses. To fight invasive pathogens, host macrophages deliberately create a copper overdose environment, exposing a host-driven defense mechanism (2019). Copper homeostasis-related genes, such as *copA* (efflux system), *copZ* (copper chaperone), *cueO* (copper oxidase), *copS* (sensor protein), and *copR* (transcriptional activator), were present in all three of the *C. kerstersii* strains included in this study. Notably, studies have shown that loss of the copper efflux pathway (CopA) in certain bacteria, such as *S. pneumoniae*, reduces virulence during infection (Begg 2019). This discovery implies that the pathogenic potential of *C. kerstersii* may depend on its capacity to maintain copper homeostasis.

Secretion systems

Protein export systems are responsible for regulating the secretion of different types of proteins across the cell envelope and thus has an important role in the colonization of bacteria living in various niches. *C. kerstersii* genomes were examined for the protein export and secretion system genes. In *Comamonas* species, the Sec (secretion) and Tat (twin-arginine translocation) pathways are essential for protein transport across the cytoplasmic membrane (Ma et al. 2009). We found *SecA*, *SecB*, *SecYEG* encoding genes, although these genes are not present in a cluster. In contrast, *tatABC* genes encoding the components of the Tat apparatus, are present in a cluster (Supplementary table 1). The Sec pathway typically handles the translocation of unfolded proteins, whereas the Tat pathway is specialized for transporting fully folded proteins. Most proteins transported by the Sec and Tat pathways remain within the

periplasm or inner membrane and may be transported in the extracellular environment through type II or type V secretion systems (Green and Meccas 2016)

The genes *epsE*, *gspF* and *outD* encode components of type II secretion system and genes to encode Chaperone-Usher pathway (supplementary table 1). The chaperone–usher (CU) pathway is the most widespread method for assembling adhesive surface pili on the surface of Gram-negative bacteria. For uropathogenic *Escherichia coli* and *Pseudomonas aeruginosa*, these adhesive pili mediated localization and host colonization and biofilm formation, are considered as important virulence factors (Vallet et al. 2001; Green and Meccas 2016).

Plasmids and Prophages

In each of the three strains used in this investigation, PlasmidFinder located a 157bp long fragment identical to plasmids of IncQ family. 100% identity was found in the region between 5741 and 6536 base pairs (bp) in the plasmid when it was compared using BLAST against the *E. coli* plasmid RSF1010 (Accession no. M28829). The distinct strand-displacement replication mechanism of the IncQ-family of plasmids allows them to work in a broad range of bacterial hosts and their high mobilization potential enables these plasmids to exhibit great promiscuity (Loftie-Eaton and Rawlings 2012). Several studies have mentioned plasmid IncQ to be multidrug resistant carrying *tetA*, Sul or carbapenem resistance gene *blaKPC-2* (Oliva et al. 2017; Fuga et al. 2021) although the IncQ1 plasmid of the studied strain do not contain any resistance genes.

PHASTEST and Phigaro analysis identified prophage regions in all three *C. kerstersii* strains (Figures 3 and 4). Each prophage region was approximately 63.6 kb in size with a GC content of ~58% and showed similarity to *Pseudomonas* phage F116 of the Podoviridae family. These regions encoded proteins involved in phage replication (integrase, repressor, terminases), regulation (regulatory protein), and lysis (lysis proteins, tail proteins, holin) and contained a significant number of hypothetical proteins.

Interestingly, Phigaro identified an additional prophage region belonging to the Siphoviridae family alongside the Podoviridae region in each strain (Figure 4a). This region encoded phage assembly proteins (tail, portal, coat proteins). Multiple prophage regions and a significant number of hypothetical proteins within them suggest potential roles in *C. kerstersii* fitness. Further investigation of these hypothetical proteins may provide valuable insights into their specific functions.

Discussion

C. kerstersii is becoming a prominent clinical and environmental pathogen due to its broad illness spectrum, ubiquity, adaptability in various environments and resistance capabilities. These characteristics, which indicate that *C. kerstersii* is highly diverse, call for a deeper comprehension of how the species has adapted to fill various niches. *C. kerstersii* came to light in Bangladesh for the first time; thus, genomic analysis and understanding the behaviour of this bacterium is crucial for mitigating potential public health risks associated with its infections.

Phylogenetic tree construction with the other available nine strains of *C. kerstersii* reveals its clonal nature, diversity and adaptability to different niches. Clustering of NG13, NG14, and NG17 together demonstrate their clonal nature, explaining the identical genotypical features.

The strains of the study have been isolated from clinical samples. However, they did not show significant virulence factors other than T4P and T6SS. It is plausible that these strains have acquired the virulent characteristics from the other species that may have coexisted in the infection milieu. The toxin protein zot is responsible for diarrhea or the holin protein present in the phage region can also induce the pathogenicity. The enhanced release of extracellular toxins in *E. coli*, including *SheA* and *StxI*, has been linked to phage-encoded holins (Vidor et al. 2022). According to a number of investigations, holins most likely also have a significant function in the production of biofilms (Saier and Reddy 2015). For instance, it is known that the *Staphylococcus aureus* *cid* and *lrg* operons regulate cell lysis and the release of genomic DNA, which eventually becomes a structural element of the biofilm matrix and are thus implicated in the production of biofilms (Saier and Reddy 2015). Alternatively, they may aid in the infection process through systems not generally linked to virulence but through resistance mechanisms like regulation of copper homeostasis.

As a relatively new opportunistic pathogen, *C. kerstersii* has received little attention in scientific research. Only two papers have explored and reported the genomic analysis of this species thus far suggesting that the genomic microbial databases might have incomplete information on *C. kerstersii*. This lack of information may have made it difficult to identify every gene connected to the organism. The presence of many hypothetical proteins has been discovered by the annotated data of the studied strains. These proteins are hypothetical gene products, the roles of which are neither entirely understood nor confirmed by experiments. Even though they are speculative, these potential proteins can provide important information about the genetic composition of *C. kerstersii* and could be useful research targets to clarify their functions in pathogenicity and antibiotic resistance.

Conflict of interest

All authors declare no conflict of interest

Ethical Approval

All isolates analyzed in this study were obtained for diagnostic purposes from a local tertiary hospital. All experiments and methodologies were conducted in compliance with relevant guidelines and regulations. The North South University (NSU) Institutional Review Board (IRB) / Ethical Review Committee (ERC) approved all experimental protocols, under protocol No. CTRG NSU-RP-20-043.

Data Availability

The 3 *Comamonas kerstersii* genome sequences and analysis from this study have been submitted to GenBank database, with accession numbers: JBEBZE000000000 (NG17), JBEBZF000000000 (NG13), and JBEBZG000000000(NG14).

References

- Alikhan NF, Petty NK, Ben Zakour NL, Beatson SA (2011) BLAST Ring Image Generator (BRIG): Simple prokaryote genome comparisons. *BMC Genomics* 12:1–10. <https://doi.org/10.1186/1471-2164-12-402/TABLES/3>
- Almuzara M, Barberis C, Veiga F, et al (2017) Unusual presentations of *Comamonas kerstersii* infection. *New Microbes New Infect* 19:91–95. <https://doi.org/10.1016/J.NMNI.2017.07.003>
- Almuzara M, Cittadini R, Estraviz ML, et al (2018) First report of *Comamonas kerstersii* causing urinary tract infection. *New Microbes New Infect* 24:4–7. <https://doi.org/10.1016/J.NMNI.2018.03.003>
- Almuzara MN, Cittadini R, Ocampo CV, et al (2013) Intra-Abdominal Infections Due to *Comamonas kerstersii*. *J Clin Microbiol* 51:1998. <https://doi.org/10.1128/JCM.00659-13>
- Ault-Riché D, Fraley CD, Tzeng CM, Kornberg A (1998) Novel Assay Reveals Multiple Pathways Regulating Stress-Induced Accumulations of Inorganic Polyphosphate in *Escherichia coli*. *J Bacteriol* 180:1841. <https://doi.org/10.1128/JB.180.7.1841-1847.1998>
- Aziz RK, Bartels D, Best A, et al (2008) The RAST Server: Rapid annotations using subsystems technology. *BMC Genomics* 9:. <https://doi.org/10.1186/1471-2164-9-75>
- Bankevich A, Nurk S, Antipov D, et al (2012) SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. *Journal of Computational Biology* 19:455. <https://doi.org/10.1089/CMB.2012.0021>
- Barbaro DJ, Mackowiak PA, Barth SS, Southern PM (1987) *Pseudomonas testosteroni* Infections: Eighteen Recent Cases and a Review of the Literature. *Rev Infect Dis* 9:124–129. <https://doi.org/10.1093/CLINIDS/9.1.124>
- Begg SL (2019) The role of metal ions in the virulence and viability of bacterial pathogens. *Biochem Soc Trans* 47:77–87
- Bennani H, El Ouarradi A, Hanchi AL, Soraa N (2022) A young child with acute perforated appendicitis due to *Comamonas kerstersii*: a rare case report. *Pan African Medical Journal* 41:. <https://doi.org/10.11604/pamj.2022.41.186.29615>
- Biswas JS, Fitchett J, O’Hara G (2014) *Comamonas kerstersii* and the Perforated Appendix. *J Clin Microbiol* 52:3134. <https://doi.org/10.1128/JCM.00909-14>
- Bortolaia V, Kaas RS, Ruppe E, et al (2020) ResFinder 4.0 for predictions of phenotypes from genotypes. *Journal of Antimicrobial Chemotherapy* 75:3491–3500. <https://doi.org/10.1093/JAC/DKAA345>

- Candon HL, Allan BJ, Fraley CD, Gaynor EC (2007) Polyphosphate Kinase 1 Is a Pathogenesis Determinant in *Campylobacter jejuni*. *J Bacteriol* 189:8099. <https://doi.org/10.1128/JB.01037-07>
- Carattoli A, Zankari E, Garcíá-Fernández A, et al (2014) In Silico detection and typing of plasmids using plasmidfinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother* 58:3895–3903. https://doi.org/10.1128/AAC.02412-14/SUPPL_FILE/ZAC007143013SO1.PDF
- Chen S, Yang D, Wen Y, et al (2018) Dysregulated hemolysin liberates bacterial outer membrane vesicles for cytosolic lipopolysaccharide sensing. *PLoS Pathog* 14:. <https://doi.org/10.1371/JOURNAL.PPAT.1007240>
- Colin R, Ni B, Laganenka L, Sourjik V (2021) Multiple functions of flagellar motility and chemotaxis in bacterial physiology. *FEMS Microbiol Rev* 45:1–19. <https://doi.org/10.1093/FEMSRE/FUAB038>
- Confer AW, Ayalew S (2013) The OmpA family of proteins: Roles in bacterial pathogenesis and immunity. *Vet Microbiol* 163:207–222
- DE VOS P, KERSTERS K, FALSEN E, et al (1985) *Comamonas* Davis and Park 1962 gen. nov., nom. rev. emend., and *Comamonas terrigena* Hugh 1962 sp. nov., nom. rev. *Int J Syst Bacteriol* 35:443–453. <https://doi.org/10.1099/00207713-35-4-443>
- Farfán-Cano GG, Sarmiento-Bobadilla JA, Jara León EA, et al (2021) *Comamonas kerstersii* strains on inpatients with acute appendicitis: review of literature and case report. *InterAmerican Journal of Medicine and Health* 4:. <https://doi.org/10.31005/iajmh.v4i.165>
- Frees D, Savijoki K, Varmanen P, Ingmer H (2007) Clp ATPases and ClpP proteolytic complexes regulate vital biological processes in low GC, Gram-positive bacteria. *Mol Microbiol* 63:1285–1295. <https://doi.org/10.1111/J.1365-2958.2007.05598.X>
- Frees D, Sørensen K, Ingmer H (2005) Global Virulence Regulation in *Staphylococcus aureus*: Pinpointing the Roles of ClpP and ClpX in the sar/agr Regulatory Network. *Infect Immun* 73:8100. <https://doi.org/10.1128/IAI.73.12.8100-8108.2005>
- Fuga B, Cerdeira L, Moura Q, et al (2021) Genomic data reveals the emergence of an IncQ1 small plasmid carrying blaKPC-2 in *Escherichia coli* of the pandemic sequence type 648. *J Glob Antimicrob Resist* 25:8. <https://doi.org/10.1016/J.JGAR.2021.02.014>
- Gilson L, Mahanty HK, Kolter R (1990) Genetic analysis of an MDR-like export system: the secretion of colicin V. *EMBO J* 9:3875–3884. <https://doi.org/10.1002/j.1460-2075.1990.tb07606.x>
- Green ER, Meccas J (2016) Bacterial Secretion Systems: An Overview. *Microbiol Spectr* 4:. <https://doi.org/10.1128/MICROBIOLSPEC.VMBF-0012-2015/ASSET/7FA9CACF-7AA0-4E78-8D18-243390487674/ASSETS/GRAPHIC/VMBF-0012-2015-FIG1B.GIF>
- Jiang X, Liu W, Zheng B (2018) Complete genome sequencing of *Comamonas kerstersii* 8943, a causative agent for peritonitis. *Sci Data* 5:180222. <https://doi.org/10.1038/sdata.2018.222>
- Kadlec K, Michael GB, Sweeney MT, et al (2011) Molecular Basis of Macrolide, Triamylide, and Lincosamide Resistance in *Pasteurella multocida* from Bovine Respiratory Disease. *Antimicrob Agents Chemother* 55:2475–2477. <https://doi.org/10.1128/AAC.00092-11>
- Kobayashi N, Nishino K, Yamaguchi A (2001) Novel Macrolide-Specific ABC-Type Efflux Transporter in *Escherichia coli*. *J Bacteriol* 183:5639. <https://doi.org/10.1128/JB.183.19.5639-5644.2001>
- Kumar S, Yadav M, Anwer R, Sehrawat N Isolation and Characterization of *E. coli* and *Comamonas kerstersii* from Chicken Litter Samples from North India

- Leclercq R (2002) Mechanisms of Resistance to Macrolides and Lincosamides: Nature of the Resistance Elements and Their Clinical Implications. *Clinical Infectious Diseases* 34:482–492. <https://doi.org/10.1086/324626>
- Li C, Li Y, Ding C Molecular Sciences The Role of Copper Homeostasis at the Host-Pathogen Axis: From Bacteria to Fungi. <https://doi.org/10.3390/ijms20010175>
- Linhartova I, Osicka R, Bumba L, et al (2015) RTX Toxins: A Review. In: *Microbial Toxins*. Springer Netherlands, pp 1–29
- Liu B, Zheng D, Jin Q, et al (2019) VFDB 2019: a comparative pathogenomic platform with an interactive web interface. *Nucleic Acids Res* 47:D687. <https://doi.org/10.1093/NAR/GKY1080>
- Liu X jiao, Qiao X wen, Huang T min, et al (2020a) *Comamonas kerstersii* bacteremia. *Med Mal Infect* 50:288–290. <https://doi.org/10.1016/J.MEDMAL.2019.12.005>
- Liu X, Qiao X, Huang T, et al (2020b) *Comamonas kerstersii* bacteremia. *Med Mal Infect* 50:288–290. <https://doi.org/https://doi.org/10.1016/j.medmal.2019.12.005>
- Loftie-Eaton W, Rawlings DE (2012) Diversity, biology and evolution of IncQ-family plasmids. *Plasmid* 67:15–34. <https://doi.org/10.1016/J.PLASMID.2011.10.001>
- Lu S, Zgurskaya HI (2013) MacA, a periplasmic membrane fusion protein of the macrolide transporter MacAB-TolC, binds lipopolysaccharide core specifically and with high affinity. *J Bacteriol* 195:4865–4872. <https://doi.org/10.1128/JB.00756-13/ASSET/AEE61305-EF1E-4D95-9C88-046E8AC2F3F0/ASSETS/GRAPHIC/ZJB9990928590003.JPEG>
- Lv H, Zhou Y, Liu B, et al (2022) Polyphosphate Kinase Is Required for the Processes of Virulence and Persistence in *Acinetobacter baumannii*. *Microbiol Spectr* 10:. <https://doi.org/10.1128/SPECTRUM.01230-22>
- Ma YF, Zhang Y, Zhang JY, et al (2009) The Complete Genome of *Comamonas testosteroni* Reveals Its Genetic Adaptations to Changing Environments. *Appl Environ Microbiol* 75:6812. <https://doi.org/10.1128/AEM.00933-09>
- Mandal J, Kannambath R, Mohan P, Stanley P (2022) Isolation of *Comamonas* Species from the Stool Samples of Patients with Underlying Gastrointestinal Pathology. *Journal of Gastrointestinal Infections* 11:47–50. <https://doi.org/10.5005/jp-journals-10068-3049>
- McMeechan A, Lovell MA, Cogan TA, et al (2007) Inactivation of *ppk* differentially affects virulence and disrupts ATP homeostasis in *Salmonella enterica* serovars Typhimurium and Gallinarum. *Res Microbiol* 158:79–85. <https://doi.org/10.1016/J.RESMIC.2006.10.008>
- Melville S, Craig L (2013) Type IV Pili in Gram-Positive Bacteria. *Microbiol Mol Biol Rev* 77:323. <https://doi.org/10.1128/MMBR.00063-12>
- Meyer AS, Baker TA (2011) Proteolysis in the *Escherichia coli* heat shock response: A player at many levels. *Curr Opin Microbiol* 14:194. <https://doi.org/10.1016/J.MIB.2011.02.001>
- Moser AI, Campos-Madueno EI, Keller PM, Endimiani A (2021) Complete Genome Sequence of a Third- and Fourth-Generation Cephalosporin-Resistant *Comamonas kerstersii* Isolate. *Microbiol Resour Announc* 10:. <https://doi.org/10.1128/MRA.00391-21>
- Navas J, Fernández-Martínez M, Salas C, et al (2016) Susceptibility to Aminoglycosides and Distribution of *aph* and *aac(3)-XI* Genes among *Corynebacterium striatum* Clinical Isolates. *PLoS One* 11:. <https://doi.org/10.1371/JOURNAL.PONE.0167856>
- Nguyen F, Starosta AL, Arenz S, et al (2014) Tetracycline antibiotics and resistance mechanisms. *Biol Chem* 395:559–575. https://doi.org/10.1515/HSZ-2013-0292/ASSET/GRAPHIC/HSZ-2013-0292_FIG7.JPG

- Oliva M, Monno R, D'Addabbo P, et al (2017) A novel group of IncQ1 plasmids conferring multidrug resistance. *Plasmid* 89:22–26. <https://doi.org/10.1016/J.PLASMID.2016.11.005>
- Olvera C, Goldberg JB, Snchez R, Sobern-Chvez G (1999) The *Pseudomonas aeruginosa* algC gene product participates in rhamnolipid biosynthesis. *FEMS Microbiol Lett* 179:85–90. <https://doi.org/10.1111/J.1574-6968.1999.TB08712.X>
- Onya O, Barbara N, Giorgio Z, et al (2020a) Bacteremia Caused by *Comamonas kerstersii* in a Patient with Diverticulosis. *J Clin Microbiol* 52:1009–1012. <https://doi.org/10.1128/jcm.02942-13>
- Onya O, Barbara N, Giorgio Z, et al (2020b) Bacteremia Caused by *Comamonas kerstersii* in a Patient with Diverticulosis. *J Clin Microbiol* 52:1009–1012. <https://doi.org/10.1128/jcm.02942-13>
- Opota O, Ney B, Zanetti G, et al (2014a) Bacteremia caused by *Comamonas kerstersii* in a patient with diverticulosis. *J Clin Microbiol* 52:1009–1012. <https://doi.org/10.1128/JCM.02942-13>
- Opota O, Ney B, Zanetti G, et al (2014b) Bacteremia caused by *Comamonas kerstersii* in a patient with diverticulosis. *J Clin Microbiol* 52:1009–1012. <https://doi.org/10.1128/JCM.02942-13/ASSET/5F1BCB07-4DDA-4365-B8E0-F477E0B0363C/ASSETS/GRAPHIC/ZJM9990931920001.JPEG>
- Page AJ, Cummins CA, Hunt M, et al (2015) Roary: rapid large-scale prokaryote pan genome analysis. *Bioinformatics* 31:3691–3693. <https://doi.org/10.1093/BIOINFORMATICS/BTV421>
- Palacio R, Cornejo C, Seija V (2020) *Comamonas kerstersii*. *Revista Chilena de Infectologia* 37:147–148. <https://doi.org/10.4067/S0716-10182020000200147>
- Pavone S, Rinoldo R, Albini E, et al (2021) First report of urinary tract infection caused by *Comamonas kerstersii* in a goat. *BMC Vet Res* 17:1–5. <https://doi.org/10.1186/S12917-021-02840-X/FIGURES/2>
- Price MN, Dehal PS, Arkin AP (2009) FastTree: Computing Large Minimum Evolution Trees with Profiles instead of a Distance Matrix. *Mol Biol Evol* 26:1641–1650. <https://doi.org/10.1093/MOLBEV/MSP077>
- Rao NN, Gomez-Garca MR, Kornberg A (2009) Inorganic polyphosphate: Essential for growth and survival. *Annu Rev Biochem* 78:605–647
- Rong K, Delport J, AlMutawa F (2022) *Comamonas kerstersii* Bacteremia of Unknown Origin. *Case Rep Infect Dis* 2022:1–3. <https://doi.org/10.1155/2022/1129832>
- Ryan MP, Pembroke JT (2018) *Brevundimonas* spp: Emerging global opportunistic pathogens. <https://doi.org/10.1080/21505594.2017.1419116>
- Ryan MP, Sevjahova L, Gorman R, White S (2022) The Emergence of the Genus *Comamonas* as Important Opportunistic Pathogens. *Pathogens* 11:.. <https://doi.org/10.3390/PATHOGENS11091032>
- S BJ, Joseph F, Geraldine O (2020) *Comamonas kerstersii* and the Perforated Appendix. *J Clin Microbiol* 52:3134. <https://doi.org/10.1128/jcm.00909-14>
- Saier MH, Reddy BL (2015) Holins in Bacteria, Eukaryotes, and Archaea: Multifunctional Xenologues with Potential Biotechnological and Biomedical Applications. *J Bacteriol* 197:7. <https://doi.org/10.1128/JB.02046-14>
- Schauer O, Mostaghaci B, Colin R, et al (2018) Motility and chemotaxis of bacteria-driven microswimmers fabricated using antigen 43-mediated biotin display. *Scientific Reports* 2018 8:1 8:1–11. <https://doi.org/10.1038/s41598-018-28102-9>

- Seemann T (2014) Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/BIOINFORMATICS/BTU153>
- Starikova E V., Tikhonova PO, Prianichnikov NA, et al (2020) Phigaro: High-throughput prophage sequence annotation. *Bioinformatics* 36:3882–3884. <https://doi.org/10.1093/bioinformatics/btaa250>
- Turlin E, Heuck G, Simões Brandão MI, et al (2014) Protoporphyrin (PPIX) efflux by the MacAB-TolC pump in *Escherichia coli*. *Microbiologyopen* 3:849–859. <https://doi.org/10.1002/MBO3.203>
- Vallet I, Olson JW, Lory S, et al (2001) The chaperone/usher pathways of *Pseudomonas aeruginosa*: Identification of fimbrial gene clusters (cup) and their involvement in biofilm formation. *Proc Natl Acad Sci U S A* 98:6911–6916. <https://doi.org/10.1073/PNAS.111551898/ASSET/9A1A9FE0-6816-46E3-BA77-7AC0A2E32776/ASSETS/GRAPHIC/PQ1115518005.JPEG>
- Vidor CJ, Hamiot A, Wisniewski J, et al (2022) A Highly Specific Holin-Mediated Mechanism Facilitates the Secretion of Lethal Toxin TcsL in *Paenibacillus sordellii*. *Toxins (Basel)* 14:. <https://doi.org/10.3390/TOXINS14020124/S1>
- Voitenkova E V., Matveeva ZN, Makarova MA, et al (2018) Difficulties in identification of *Comamonas kerstersii* strains isolated from intestinal microbiota of residents of republic of Guinea and Russian federation. *Russian Journal of Infection and Immunity* 8:163–168. <https://doi.org/10.15789/2220-7619-2018-2-163-168>
- Wang M, Wang H, Chen S, et al (2024) Clinical Case Report Medicine ® Extra-abdominal infections caused by *Comamonas kerstersii* Case report. <https://doi.org/10.1097/MD.00000000000037161>
- Wang Y (2002) The function of OmpA in *Escherichia coli*. *Biochem Biophys Res Commun* 292:396–401. <https://doi.org/10.1006/bbrc.2002.6657>
- Warr AR, Giorgio RT, Waldora MK (2020) Genetic analysis of the role of the conserved inner membrane protein CvpA in enterohemorrhagic *Escherichia coli* resistance to deoxycholate. *J Bacteriol* 203:661–681. <https://doi.org/10.1128/JB.00661-20>
- Wauters G, De Baere T, Willems A, et al (2003a) Description of *Comamonas aquatica* comb. nov. and *Comamonas kerstersii* sp. nov. for two subgroups of *Comamonas terrigena* and emended description of *Comamonas terrigena*. *Int J Syst Evol Microbiol* 53:859–862. <https://doi.org/10.1099/ijs.0.02450-0>
- Wauters G, De Baere T, Willems A, et al (2003b) Description of *Comamonas aquatica* comb. nov. and *Comamonas kerstersii* sp. nov. for two subgroups of *Comamonas terrigena* and emended description of *Comamonas terrigena*. *Int J Syst Evol Microbiol* 53:859–862. <https://doi.org/10.1099/ijs.0.02450-0>
- Willems A, De Vos P (2006) *Comamonas*. *The Prokaryotes* 723–736. https://doi.org/10.1007/0-387-30745-1_31
- Willems A, Pot B, Falsen E, et al (1991) Polyphasic taxonomic study of the emended genus *Comamonas*: Relationship to *Aquaspirillum aquaticum*, E. Falsen group 10, and other clinical isolates. *Int J Syst Bacteriol* 41:427–444. <https://doi.org/10.1099/00207713-41-3-427/CITE/REFWORKS>
- Wishart DS, Han S, Saha S, et al (2023) PHASTEST: faster than PHASTER, better than PHAST. *Nucleic Acids Res* 51:W443–W450. <https://doi.org/10.1093/NAR/GKAD382>

- Wu Y, Zaiden N, Cao B (2018) The core- and pan-genomic analyses of the genus *Comamonas*: From environmental adaptation to potential virulence. *Front Microbiol* 9:
<https://doi.org/10.3389/fmicb.2018.03096>
- Yamanaka H, Kobayashi H, Takahashi E, Okamoto K (2008) MacAB Is Involved in the Secretion of *Escherichia coli* Heat-Stable Enterotoxin II. *J Bacteriol* 190:7693.
<https://doi.org/10.1128/JB.00853-08>
- Yim HH, Villarejo M (1992) *osmY*, a New Hyperosmotically Inducible Gene, Encodes a Periplasmic Protein in *Escherichia coli*
- Zhang Y, Inouye M (2011) RatA (YfjG), An *Escherichia Coli* Toxin, Inhibits 70S Ribosome Association to Block Translation Initiation. *Mol Microbiol* 79:1418.
<https://doi.org/10.1111/J.1365-2958.2010.07506.X>
- Zhou YH, Ma HX, Dong ZY, Shen MH (2018) *Comamonas kerstersii* bacteremia in a patient with acute perforated appendicitis: A rare case report. *Medicine* 97:.
<https://doi.org/10.1097/MD.00000000000009296>

Tables

Table 1. Biochemical test results of the three *C. kerstersii* isolates

Test Name	NG13	NG14	NG17
Oxidase	+	+	+
Catalase	+	+	+
Motility	+	+	+
Gelatin Hydrolysis	-	-	-
Nitrate Reduction	+	+	+
Trypsin Hydrolysis	+	+	+
Acid from Glucose	-	-	-
Urease	-	-	-
H ₂ S Production	-	-	-
Indole	-	-	-

Table 2. Genome assembly and annotation statistics of the three *C. kerstersii* isolates

	NG13	NG14	NG17
Genome Size (appx)	34, 806, 663	34, 90, 070	34, 93, 044
No. of contigs	50	68	81
GC (%)	59.60	59.60	59.59
No. of CDS	3211	3215	3213
No. of tRNA	94	97	94
No. of rRNA	7	7	7
No. of tmRNA	1	1	1
N50 (bp)	181861	284992	221534

Table 3. Antimicrobial resistance genes associated with the three strains of *C. kerstersii*

Strain ID	Resistance Gene	Phenotype	Antibiotic	Coverage (%)	Identity (%)	Alignment Length/Gene Length
NG13, NG14, NG17	aph(6)-Id	Aminoglycoside resistance	streptomycin	100	100	837/837
	aph(3'')-Ib	Aminoglycoside resistance	streptomycin	100	100	804/804
	mph(E)	Macrolide resistance	erythromycin	100	100	885/885
	mph(F)	Macrolide resistance	erythromycin	100	100	900/900
	msr(E)	Macrolide, Lincosamide and Streptogramin B resistance	Erythromycin, azithromycin, Pristinamycin ia, virginiamycin s, quinupristin	100	100	1476/1476
	Sul2	Sulphonamide resistance	sulfamethoxazole	100	100	816/816
	tet(A)	Tetracycline resistance	Doxycycline, tetracycline	100	100	1275/1275

Table 4. Virulence profile of *C. kerstersii* strains

Strain ID	Virulence Factors	Length	E-value	Overall Identity
NG13	VFG000112 (pilB) type 4 fimbrial biogenesis protein PilB [Type IV pili] [<i>Pseudomonas aeruginosa</i> PAO1]	1701	2e-18	88.8% (166/187)
	VFG000233 (pilF) type IV pilus assembly ATPase protein [Type IV pili] [<i>Neisseria meningitidis</i> MC58]	1677	9e-17	80.9% (169/209)
	VFG013777 (pilB) type II secretion system protein E:General secretory system II, protein E, N-terminal [Type IV pili biosynthesis] [<i>Pseudomonas syringae</i> pv. <i>syringae</i> B728a]	1695	6e-18	85.4% (105/123)
	VFG014984 (algW) AlgW protein [Alginate regulation] [<i>Pseudomonas aeruginosa</i> PAO1]	1170	2e-18	83.6% (127/152)
	VFG025878 (tsr) methyl-accepting chemotaxis protein [Flagella] [<i>Burkholderia cenocepacia</i> J2315]	1959	9e-14	82.6% (119/144)
	VFG030697 (sugC) ABC transporter, ATP-binding protein SugC [Trehalose-recycling ABC transporter] [<i>Mycobacterium intracellulare</i> MOTT-02]	1179	1e-15	85.6% (95/111)
	VFG038316 (tapD/pilD) prepilin peptidase [exe] [<i>Aeromonas hydrophila</i> subsp. <i>hydrophila</i> ATCC 7966]	873	1e-15	87.4% (83/95)
	VFG043040 (tar/cheM) methyl accepting chemotaxis protein II [peritrichous flagella] [<i>Salmonella enterica</i> subsp. <i>enterica</i> serovar Typhimurium str. LT2]	1662	4e-19	85.6% (107/125)
	VFG000182 (xcpR) general secretion pathway protein E [xcp secretion system] [<i>Pseudomonas aeruginosa</i> PAO1]	1509	2e-09	85.9% (134/156)
	VFG002048 (gspE) general secretion pathway protein E [T2SS] [<i>Shigella dysenteriae</i> Sd197]	1494	6e-07	93.0% (40/43)
	VFG000114 (xcpA/pilD) type 4 prepilin peptidase PilD [Type IV pili] [<i>Pseudomonas aeruginosa</i> PAO1]	873	6e-07	90.2% (46/51)
	VFG043209 (cheD) methyl-accepting chemotaxis protein CheD [peritrichous flagella] [<i>Yersinia enterocolitica</i> subsp. <i>enterocolitica</i> 8081]	1674	9e-12	82.7% (191/231)
	NGCRVN 14	VFG036295 (pilF) type IV pilus assembly protein [Type IV pili] [<i>Neisseria lactamica</i> 020-06]	1677	2e-21
VFG019728 (pilB) type 4 fimbrial biogenesis protein PilB [Type IV pili biosynthesis] [<i>Pseudomonas aeruginosa</i> LESB58]		1701	9e-21	85.4% (229/268)
VFG026971 (sigA/rpoV) RNA polymerase sigma factor [Sigma A] [<i>Mycobacterium tuberculosis</i> CAS/NITR204]		1134	9e-24	84.2% (139/165)
VFG038432 (tapB) type IV-A pilus assembly ATPase PilB [Tap type IV pili] [<i>Aeromonas hydrophila</i> ML09-119]		1707	5e-19	88.2% (150/170)
VFG043040 (tar/cheM) methyl accepting chemotaxis protein II [peritrichous flagella] [<i>Salmonella enterica</i> subsp. <i>enterica</i> serovar Typhimurium str. LT2]		1662	5e-19	85.6% (107/125)

	VFG014984 (algW) AlgW protein [Alginate regulation] [Pseudomonas aeruginosa PAO1]	1170	2e-18	83.6% (127/152)
	VFG030705 (sugC) sugar-transport ATP-binding protein ABC transporter SugC [Trehalose-recycling ABC transporter] [Mycobacterium liflandii 128FXT]	1188	3e-17	86.4% (95/110)
	VFG043209 (cheD) methyl-accepting chemotaxis protein CheD [peritrichous flagella] [Yersinia enterocolitica subsp. enterocolitica 8081]	1674	9e-12	82.7% (191/231)
	VFG002529 (tsr) methyl-accepting chemotaxis protein I [Flagella] [Burkholderia pseudomallei K96243]	2022	1e-10	81.6% (115/141)
	VFG000182 (xcpR) general secretion pathway protein E [xcp secretion system] [Pseudomonas aeruginosa PAO1]	1509	2e-09	85.9% (134/156)
	VFG002048 (gspE) general secretion pathway protein E [T2SS] [Shigella dysenteriae Sd197]	1494	6e-07	93.0% (40/43)
	VFG000114 (xcpA/pilD) type 4 prepilin peptidase PilD [Type IV pili] [Pseudomonas aeruginosa PAO1]	873	6e-07	90.2% (46/51)
NGCRVN 17	VFG002076 (clpV1) type VI secretion system AAA+ family ATPase [HSI-I] [Pseudomonas aeruginosa PAO1]	2709	8e-08	86.8% (59/68)
	VFG043092 (tar/cheM) methyl-accepting chemotaxis protein II [peritrichous flagella] [Escherichia coli O157:H7 str. EDL933]	1662	3e-22	88.2% (97/110)
	VFG021055 (mgtB) Magnesium transport ATPase, P-type 2 [Mg ²⁺ transport] [Salmonella enterica subsp. enterica serovar Enteritidis str. P125109]	2727	5e-21	79.8% (233/292)
	VFG014045 (pilJ) type IV pilus biogenesis protein PilJ [Type IV pili twitching motility related proteins] [Pseudomonas syringae pv. tomato str. DC3000]	2043	2e-17	87.3% (89/102)
	VFG025879 (tsr) methyl-accepting chemotaxis sensory transducer [Flagella] [Burkholderia cenocepacia MC0-3]	1974	7e-14	82.6% (119/144)
	VFG046110 (ECVR50_2114) putative inner membrane ABC-transporter [Yersiniabactin siderophore] [Escherichia coli VR50]	1803	3e-13	84.0% (100/119)
	VFG034062 (ybtP) ABC transporter ATP-binding/permease rprotein [Yersiniabactin siderophore] [Escherichia coli O44:H18 042]	1803	3e-13	84.0% (100/119)
	VFG001231 (chpA) still frameshift probable component of chemotactic signal transduction system [Type IV pili] [Pseudomonas aeruginosa PAO1]	7419	8e-08	95.0% (38/40)

Table 5. Genes related to copper homeostasis in *C. kerstersii* strains

Strain ID	Group Alias all Copper related functions	Abbrev.	Functional Role
NG13, NG14, NG17	Copper related functions	CRTR	Cu(I)-responsive transcriptional regulator
	Copper related functions	copZ	Copper chaperone
	Copper related functions	clfA	Multidrug resistance transporter, Bcr/CflA family
	Copper related functions	MO	Multicopper oxidase
	Copper related functions	BCO	Blue copper oxidase CueO precursor
	Copper related functions	CTO	Copper tolerance protein
	Copper related functions	copG	CopG protein
	Copper related functions	CRB	Copper resistance protein B
	Copper related functions	TR	Negative transcriptional regulator-copper transport operon
	Copper related functions	copS	Sensor protein copS
	Copper related functions	copR	Transcriptional activator protein CopR
	Copper related functions	copCp	Copper resistance protein C precursor

Figures

Fig. 1 BRIG-obtained circular genomic maps of *C. kerstersii* strains. The homology rate is indicated by color saturation; blanks indicate no resemblance. The Blue circle represents the reference genome *C. kerstersii* 8943.

Fig. 2 Mid-point rooted phylogenetic tree of publicly available *C. kerstersii* strains and sequenced isolates from Bangladesh based on their core genome alignment. NG13, NG14 and NG17 belonging to CLADE A clustering together indicates their clonal character.

Fig. 3 An intact phage region resembling *Pseudomonas* phage F116, identified using PHASTEST.

Fig. 4 (a) A phage region belonging to Siphoviridae family and (b) Podoviridae family.