1	"It's his cheerfulness that gives me hope": A Qualitative Analysis of Access to Pediatric
2	Cancer Care in Northern Tanzania
3	Authors and Affiliations:
4	Madeline Metcalf ¹ , Happiness D. Kajoka ² , Esther Majaliwa ^{2,3} , Anna Tupetz ^{1,4} , Catherine A.
5	Staton ^{1,4} , João Ricardo Vissoci ^{1,4} , Pamela Espinoza ¹ , Cesia Cotache-Condor ^{1,5} , Henry E. Rice ^{1,6} ,
6	Blandina T. Mmbaga ^{2,7} , Emily R. Smith ^{1,4}
7	
8	¹ Duke Global Health Institute, Duke University, Durham, NC, USA; ² Kilimanjaro Christian
9	Medical University College, Moshi, Tanzania; ³ Pediatric Hematology and Oncology Services,
10	Kilimanjaro Christian Medical Centre, Moshi, Tanzania; ⁴ Department of Emergency Medicine,
11	Duke University Medical Center, Durham, North Carolina, USA; ⁵ Duke Center for Global
12	Surgery and Health Equity, Duke University, Durham, North Carolina, USA; 6Division of
13	Pediatric Surgery, Department of Surgery, Duke University Medical Center, Durham, North
14	Carolina, USA; ⁷ Kilimanjaro Clinical Research Institute, Kilimanjaro Christian Medical Centre,
15	Moshi, Tanzania
16	
17	Word count: 5310
18	Corresponding author:
19	Madeline A. Metcalf
20	Duke Global Health Institute, Duke University, 310 Trent Hall, Durham, NC 27705
21	madeline.metcalf@duke.edu +1 208 305 8501

22 ABSTRACT

23 <u>Background:</u>

Pediatric cancer is a significant and growing burden in low- and middle-income countries. The
objective of this project was to describe the factors influencing access to pediatric cancer care in
Northern Tanzania using the Three Delays Model.

27

28 <u>Methods:</u>

29 This was a cross-sectional qualitative study conducted between June and September 2023 at

30 Kilimanjaro Christian Medical Centre (KCMC). Parents and caregivers of children obtaining

31 pediatric cancer care at KCMC were approached for participation in in-depth interviews (IDIs)

32 and a demographic survey. All IDIs were facilitated in Swahili by a bilingual research

33 coordinator. Thirteen IDIs and surveys were completed during the study period. Analysis utilized

34 inductive and deductive coding approaches to identify dominant themes and sub-themes

35 impacting access to pediatric oncology care.

36

37 <u>Results:</u>

Participants reported significant financial barriers to accessing pediatric cancer care along the entire care continuum. Early delays were impacted by waiting for symptoms to resolve or worsen. The most substantial delays resulted from health infrastructure at mid-level health facilities, misdiagnoses, and delayed referral to KCMC for treatment. Participants did not describe delays after arrival to KCMC and rather offered perspective on their child's cancer diagnosis, their concerns while obtaining care, and their hopes for the future. Financial support provided by the Tanzanian government was the only facilitator noted by participants.

45

46 <u>Conclusions:</u>

- 47 We suggest targeted interventions including 1) empowerment of CHWs and local traditional
- 48 healers to advocate for earlier care seeking behavior, 2) implementation of clinical structures and
- 49 training at intermediary medical centers aimed at earlier referral to a treatment facility, 3)
- 50 incorporation of support and education initiatives for families of children with a cancer
- 51 diagnosis. Lastly, we suggest that national health plans include pediatric cancer care.

52 INTRODUCTION

53	The Lancet Oncology Commission on Sustainable Care for Children with Cancer estimates that
54	there will be 13.7 million newly diagnosed cases of pediatric cancer between 2020 and 2050,
55	nearly 90% of which will occur in low or lower-middle income countries (LMICs).(1) Further,
56	the Commission calculates that with the current global investment, diagnostic, and treatment
57	provision trends, 11.1 million children will experience a cancer-attributable death during this 30
58	year period, with 84% of these children residing in LMICs.(1) Considering the World Health
59	Organization (WHO) Global Initiative for Childhood Cancer's 2030 goal of 60% 5-year survival
60	for pediatric cancer patients worldwide, significant scale-up in the financing, accessibility, and
61	availability of quality pediatric oncology services is required.(1-3)
62	
63	In Tanzania, a LMIC in East Africa, several factors impact families' ability to obtain care for
64	their child diagnosed with cancer such as health education, transportation, household finances,
65	and traditional medicine.(4,5) Tanzania has three health centers with the capacity to provide
66	comprehensive cancer care to children, including diagnostic, laboratory, and treatment services.
67	Further, six licensed pediatric oncologists serve the entire country with a population of more than
68	33 million children.(6) Recent research conducted at Kilimanjaro Christian Medical Centre
69	(KCMC), one of the country's three tertiary level centers with cancer care capacity, reports that
70	over 75% of pediatric cancer patients are diagnosed with stage III or IV cancer, suggesting
71	notable delays in access to care.(7)
72	
73	Initially developed for examining maternal mortality in low-income settings, "Three Delays

74 Model" has been used to classify barriers and facilitators to maternal, emergency, and neonatal

75 care in Tanzania. (4,5,8,9) However, the model has yet to be qualitatively applied to pediatric 76 cancer care in the country. The Three Delays Model organizes barriers and facilitators to 77 healthcare service procurement into three groups including: 1) delays in seeking care, 2) delays 78 in reaching care, and 3) delays in receiving care.(9) The initial decision to seek care may be 79 impacted by factors such as a household's socioeconomic status, health education, cultural or 80 spiritual beliefs, and lack of geographic proximity or transportation to a health center.(4) For 81 pediatric cancer, the second delay is associated with higher rates of morbidity and mortality and 82 can be influenced by health service capacity and referral systems at smaller health centers.(2,10)83 This second delay can also be impacted by similar sociocultural factors described in the first 84 delay.(4,10) The final delay is concerned with delays experienced once reaching the definitive medical center where diagnosis and treatment services can be provided. While not always 85 86 contained within the framework, we have included financial influences within each of these delays to highlight the economic toll of pediatric cancer care. 87

88

Although there is a growing body of global epidemiological literature outlining the prevalence and incidence trends of pediatric cancer,(7,11,12) there remains a gap in qualitative research that illustrates the experience of families in obtaining oncology care. To understand the factors influencing increased mortality and limited access to pediatric cancer care, we gathered caregiver perspectives of the journey to care with the Three Delays Model, beginning from the identification of initial symptoms and extending beyond obtaining care to include post-treatment plans.

96

97 METHODS

98 Study Design

99 This cross-sectional qualitative study used data collected from in-depth interviews (IDIs) with
100 caregivers of children obtaining pediatric oncology care at KCMC between June and September
101 2023.
102

103 Research Team and Reflexivity

104 The research team involved in data collection and analysis included a research coordinator (HK),

105 KCMC's pediatric oncologist (EM), two graduate students (MM, PE), a postdoctoral fellow

106 (AT) and an epidemiologist and principal investigator of the study (ERS). All IDIs were

107 facilitated by the bilingual, female research coordinator (HK) based at KCMC and an employed

108 member of the research team. The research coordinator is professionally trained as a medical

109 doctor in Tanzania but was not involved in any patient care or known to the participants prior to

enrollment. HK was trained in qualitative methods by AT prior to the commencement of the

111 study.

112

113 Setting

114 Located within the Kilimanjaro region of Tanzania, KCMC is classified as a tertiary level

medical center, serving nearly 11 million patients each year.(13) The KCMC pediatric cancer

116 ward serves an estimated 100-150 patients a year and currently hosts one oncologist (EM) who is

responsible for the diagnosis and treatment of all pediatric cancer cases.(14)

118

119 Participant Screening and Enrollment

120 Using purposive sampling methods, potential participants were approached for study 121 participation upon presentation for care at the KCMC pediatric oncology ward between June and 122 September 2023. Screening and enrollment of study participants was led by the local research 123 coordinator (HK) and clinical staff (EM). Potential participants were approached based on the 124 following inclusion criteria: 1) patient was 14 years old or younger, 2) patient was actively 125 obtaining treatment for a diagnosed cancer condition, 3) caregiver was at least 18 years old, 4) 126 caregiver spoke Swahili or English (fluent languages of the research coordinator). Participants 127 were excluded from the study if they had cognitive problems that would inhibit the caregiver's 128 ability to participate or if the caregiver declined participation. Due to differences in treatment 129 provision, inclusion was limited to patients between 0-14 years of age to report solely on 130 children and early adolescents rather than older adolescents and young adults. During screening, 131 two potential interviewees declined participation.

132

133 Data Collection Procedures

134 All interviews took place in a quiet, private location in the KCMC pediatric oncology ward. The 135 interviewer's name and role within the study were disclosed to participants prior to the interview. 136 However, no additional information was shared regarding interviewer bias, assumptions, or 137 personal interests in the study. After completing the written informed consent procedures, the research coordinator completed a short paper survey with the participant to collect household 138 139 demographic data such as relationship to the child, child's age and sex, urban/rural home 140 community designation, ethnic and religious affiliation, education level, occupation, household 141 and medical expenses, household monthly income, and health insurance status.

142

143 Interviews were performed using a semi-structured interview guide containing open-ended 144 questions and probes based on the Three Delays Model. The interview guide was developed by 145 our team (CCC, MM) based on the conceptual framework and an existing interview guide 146 developed by the Harvard Medical School Program in Global Surgery and Social Change.(10,15) 147 We pilot tested the interview guide with two caregivers of children who were in remission from 148 their cancer diagnosis and obtaining follow up services at KCMC. Based on the review and 149 discussion of the pilot transcripts, final amendments were made to the guide. Mainly, questions 150 were rephrased to be linguistically appropriate for the study population. The interview guide was 151 developed in English and later translated to Swahili by the research coordinator (HK) and 152 validated by the bilingual pediatric oncologist and co-principal investigator at KCMC (EM). 153 154 In addition to interviewer field notes, all interviews were audio recorded for transcription and 155 translation. The research coordinator first transcribed interviews verbatim in Swahili and then 156 translated the interviews into English. As an additional quality check, a KCMC research assistant 157 not involved in the facilitation of the interviews compared the Swahili and English transcripts for

158 translation validity. No repeat interviews were conducted, and transcripts were not returned to 159 the participants for comments and/or corrections. Interviews lasted between 30-60 minutes and 160 participants received no compensation for their participation. Audio recordings, Swahili 161 interview transcripts, and final English transcript translations were uploaded to a password-162 protected and encrypted electronic data repository available to only the qualitative study team for 163 analysis. Paper surveys were stored in a locked file cabinet separate from all informed consent 164 and screening documents at KCMC until the demographic data had been de-identified and 165 transferred to Duke University via the electronic data repository. Caregiver IDIs were conducted

until the research team unanimously agreed that data saturation had been met, defined as theabsence of new codes after three consecutive interviews.

168 Analysis

169 After four interviews had been completed, the primary qualitative analyst (MM) coded the

170 interviews using NVivo 12 software (Lumivero, Denver, CO, USA).(16) Utilizing both inductive

and deductive coding approaches, prevalent themes were identified within the Three Delays

172 Model and a codebook was created to guide the analysis of the proceeding interviews. To

173 maintain analytical validity, the codebook was discussed weekly between the analyst (MM),

174 research coordinator (HK), and Duke University principal investigator (ERS). Changes were

175 made to the identified themes and sub-themes as appropriate. No revisions were made after the

176 10th interview and all interviews were recoded with the final version of the codebook. For

analytical rigor, a random sample of 6 interviews were coded by the KCMC research coordinator

178 (HK) and a graduate research assistant (PE). A coding comparison query resulted in >90% coder

agreeability between the three coders.

180

181 Ethical Considerations

Written informed consent was obtained, and participants received a hard copy of the consent document in their preferred language (Swahili or English) prior to the start of the interview. Data storage and study procedures were approved by the Duke University and KCMC institutional review boards (IRBs) and the Tanzania National Institute for Medical Research (NIMR) prior to the commencement of the study. We utilized the Consolidated Criteria for Reporting Qualitative Research (COREQ) guidelines which is included as supplemental information with this manuscript.(17)

189

190 **RESULTS**

191 Demographic Characteristics

- 192 Between June and September 2023, a total of 13 caregiver IDIs and complimentary demographic
- surveys were completed. Majority of the interviewed participants identified themselves as the
- 194 child's mother (61.5%) and more than 90% of interviewees designated their household's home
- 195 community as rural (Table 1). The most common household size including both adults and
- 196 children living in the home was between 2-6 individuals (53.8%), with the most common
- religion being Christianity (53.8%). Of the 13 pediatric patients, 7 were male (53.8%) and 6 were
- 198 female (46.2%). Three of these patients were between the ages of 0-2 years of age, four were
- between the ages of 3-6 years, two children were within 7-10 years, and four children were
- between the ages of 11-14 years. None of the interviewees reported having health insurance.

201

203 Table 1: Study Population Demographics

	Total
Characteristic	% (n)
	100.0 (13)
Interviewee's Relationship to Child	
Mother	61.5 (8)
Father	7.7 (1)
Sibling	15.4 (2)
Uncle	15.4 (2)
Child's Age (Years)	
0-2	23.1 (3)
3-6	30.8 (4)
7-10	15.4 (2)
11-14	30.8 (4)
Child's Sex	
Male	53.8 (7)
Female	46.2 (6)
Rural-Urban Designation	
Rural	92.3 (12)
Urban	7.7 (1)
Household Size	
2-6	53.8 (7)
7-11	38.5 (5)
12+	7.7 (1)
Caregiver Completed Education	
No Formal Education	7.7 (1)
Primary	84.6 (11)
Secondary	7.7 (1)
College	0.0 (0)
Religion	
Christian	53.8 (7)
Muslim	30.8 (4)
None	15.4 (2)
Occupation of Main Household Provider	
Skilled Employment	7.7 (1)
Farming and Livestock	69.2 (9)
Self Employed	23.1 (3)
Insurance Status	
Yes	0.0.(0)

205

206 Qualitative Results

207 Organized within the Three Delays Model, Figure 1 outlines the thematic codes included in this

- 208 manuscript. In addition to outlining the delays experienced in seeking, reaching, and receiving
- 209 care, participants provided contextual background regarding their journey to care at KCMC.
- 210

Three Delays Model				
Seeking Care ——	Reaching Care	→ Receiving Care		
<u>Delays:</u>	<u>Delays:</u>	<u>Delays:</u>		
Waiting for Worsening SymptomsFinances	 Traditional Medicine Hospital Infrastructure Referral System Travel Finances 	• Finances		
Contextual Factors:	Contextual Factors:	Contextual Factors:		
Initial Symptoms	• Care Received Prior to KCMC	 Care Received at KCMC Caregiver Concerns Treatment Success 		

211 *Figure 1: Thematic codes presented in this manuscript.*

212

Analysis suggested a duality between delays discussed by IDI participants and delays perceived by the research team. IDI responses suggested that families experienced delays stemming from traditional medicine, finances (across all three delays), waiting for worsening symptoms, and hospital infrastructure. The qualitative research team agreed with all participant delays noted above, however, suggested the inclusion of referral systems and travel as additional barriers.

219 Delay 1 - Seeking Care

220 Initial symptoms

221 Caregivers outlined the symptoms that motivated care seeking behavior. One participant detailed

some of the symptoms noticed within their child before seeking care, "At first, it looked like he

had some sort of squinting or eye misalignment, but it wasn't squinting. It was actually his eye

224 protruding because it was inside, and it started coming out." (IDI Participant 6) Many

225 participants suggested the initial local diagnosis "child of the eye."

226

227 Another respondent discussed the changes over time in the formation of their child's abdomen

stating, "I began to notice small lump appearing in the abdomen. As time went on, the lump grew

229 *larger, and the whole abdomen was distended.* " (IDI Participant 8)

230

231 Waiting for worsening symptoms

232 Delays in initiation of care seeking behavior were often a result of waiting for the child's

233 condition to worsen or resolve over time. One mother discussed how they delayed care seeking

behavior for over six months with hope that the condition would heal. The mother stated, "I

thought it would go away on its own, that's why I waited because I had never seen anything like

it before. So, I thought it would resolve itself, and I waited but rather it grew." (IDI Participant

237 6)

238

Another participant explained how their child's condition recurred despite medicine provided at
home, "*I would give him medicine, and he would seem better, so he would go back to school. It*went on like that, back and forth. He would get better, then worse. But then, he got really sick,
all of his nostrils were like rotten. The sores were like rotting wounds." (IDI Participant 12)

243

244 Finances

245	More than half of participants discussed how household finances resulted in an early delay to
246	seeking care. Caregivers explained how they had to sell land or possessions to initiate care
247	seeking. A sibling describes this within their household, "The trees they sold were trees they had
248	planted on their property, and when they faced financial difficulties, they decided to sell them to
249	get money for treatment." (IDI Participant 10) Another participant described how they had to
250	"throw" or sell possessions to collect enough money for transportation to the first medical
251	center. "I had to carry a bag of rice and throw it so that we could get transportation to the
252	healthcare centerI had some roosters, and I threw them, and that's what helped me to travel to
253	Ifakara." (IDI Participant 2)
254	
255	A participant noted how government support has served as a financial facilitator in seeking care
256	for their child. "We were financially constrained, and when the child fell ill, we didn't have the
257	means to seek immediate medical attention. Fortunately, the government stepped in and brought
258	me hereIf it weren't for the government's support, I don't know what would have happened."
259	(IDI Participant 1) No additional examples of government or social support schemes for
260	financing medical expenses were provided.
261	
262	Delay 2 - Reaching Care
263	Traditional medicine
264	Once the care seeking process began, cultural and spiritual beliefs suggesting traditional

265 medicine techniques influenced when caregivers sought care from health centers. Many

266 participants noted how traditional medicine was used as first aid before opting to seek additional

267	medical care. One participant explained how traditional remedies have proven effective within
268	their community, "Yes, there are some people who have used them and found them helpful But
269	for me, they didn't help." (IDI Participant 1)
270	
271	Participants also discussed how economic factors impacted their decision to use traditional
272	medicines, often noting the "fear" associated with the costs of medical treatment, particularly in
273	rural areas (IDI Participant 2). "The other factor that has contributed is finances because you
274	find the child falls suddenly ill and you don't have anything in your pocket, you say, let's try
275	traditional medicine since even other children in the village have used it and recovered." (IDI
276	Participant 3)
277	
278	Care received prior to KCMC
279	Many participants explained how their child's condition had been misinterpreted at intermediary
280	health facilities and been prescribed ineffective medications prior to KCMC. One mother
281	explained how their child had repeatedly been treated for malaria before obtaining their cancer
282	diagnosis, "So, whenever he had a fever at night, we would rush him to the hospital. They would
283	say it's just a fever or sometimes say he has severe malaria." (IDI Participant 1) Most often,
284	children were prescribed pain medications which failed to provide long-term relief or treatment
285	for the child's condition. "It was about two months that he was using those medications. Then I
286	went to Ifakara, and they also gave me some medications just for soothing." (IDI Participant 2)
287	
288	A minority of respondents discussed how initial testing was done at smaller health facilities
289	which led to their final arrival at KCMC, but diagnostic services were never mentioned at these

290 medical centers. One participant explained this experience, "When we reached the regional

- 291 *hospital, they tested him for liver disease, took an X-ray, did an ultrasound, but the medication*
- *didn't help either. That's when they told us to come here (KCMC).* " (IDI Participant 3)
- 293

294 Hospital infrastructure

295 More than half of respondents suggested that workforce, laboratory, diagnostic, and treatment

296 capacity for pediatric patients at smaller health centers contributed to delays in arrival to KCMC.

297 One mother explained the experience of not being able to obtain testing services for their child at

298 multiple health facilities, "When you go to a hospital, they just touch the child briefly and say

they can't help, or they just look at him and say they can't do anything, go somewhere else. You

300 move to another hospital, and they say the same thing. Meanwhile, the child's condition is

301 *troubling, and he's crying. As a mother, you feel bad, and sometimes you even start crying out of*

302 pity for your child... They just look at the child without conducting proper tests." (IDI

303 Participant 1)

304

Another respondent explained their perspective regarding the capacity to provide quality and sufficient care at some medical centers and how this impacts the child's health status, "*Well, you know, when I started at those smaller hospitals, what I realized is that you could have a patient that is okay and later on their disease worsens just because they lack that specific expertise or equipment.*" (IDI Participant 11)

³¹¹ Referral system

312 Respondents described how they were referred to multiple mid-level facilities before KCMC, 313 outlining how the stepwise referral system within the country contributed to delays. While these 314 referrals may have been linked to other delays in diagnoses and infrastructure outlined above, the 315 system itself presented challenges in obtaining timely care. One participant explained the 316 challenges related to delayed referrals, "Perhaps one challenge is that when you arrive at a 317 hospital, they should give you a referral early because sometimes you arrive, and they perform 318 tests and tests. You find you have spent like 400,000 shillings and the child has not even 319 improved. Later on, when you're already broke that's when they give you a referral." (IDI 320 Participant 3) 321

Despite participants commonly noting multiple week- or month-long delays in being referred to
KCMC, caregivers expressed overall satisfaction with the referral system, focusing on how the
system had helped their family. *"I haven't seen any issue because this is where the treatment is. Even if I had continued to want to remain there and there was no treatment so when they brought me here, it wasn't a problem for me because I saw and believed that my child would get better here.* " (IDI Participant 8)

328

329 Travel

330 Caregivers and children endured long journeys to receive care at KCMC. Travel time varied

between participants, ranging from three hours to two days. One participant explained the

emotional and physical challenges of traveling with a sick child, *"It was difficult for me to travel"*

333 with the child in public transport the wound was still fresh, and I was carrying the child in my

334 *arms.* " (IDI Participant 4) Caregivers also expressed variability in the travel experience

depending on the type of transportation and familiarity with the area, "I traveled alone. I didn't 335 336 know Moshi and I didn't know anyone here...I arrived in Moshi at 2 o'clock at night! When I got 337 here at 2am I didn't have any familiarity, I didn't know anyone and there was nobody there. They 338 dropped me off along the way. I sat with the child on the roadside." (IDI Participant 5) 339 Finances 340 341 For the second delay, finances played a key role in the provision of care prior to reaching 342 KCMC. Households were generally required to pay out-of-pocket (OOP) for all medical 343 expenses related to their child's illness at intermediary hospitals before obtaining a KCMC 344 referral. A participant explained how financial concerns of paying for medical care prohibited 345 them from attending the clinic stating, "There was a month when I couldn't go. They told me I 346 needed to bring 240,000 shillings for the child to undergo a test. The date arrived, but I hadn't 347 received the money.." (IDI Participant 8) 348 349 Similar to the first delay, more than half of participants described how they had to sell 350 possessions or borrow money to continue seeking care. An interviewee explained how they both 351 borrowed money and sold livestock to pay for medical expenses, "I borrowed money from 352 someone, and when I returned home, I sold some items at home to pay them back. I also sold 353 some of my cows to continue with my child's treatment." (IDI Participant 4) Another participant 354 suggested how households can end up with "nothing" after selling these possessions, "If you 355 have cows, you sell them. If you have something, you sell it. It all ends, and you find that nothing 356 *is left.*" (IDI Participant 12)

357

358 Delay 3 - Receiving Care

359 Care received at KCMC

360 Participants discussed the care they had received at KCMC, often describing how the child was

- 361 diagnosed and their perspective of what treatment was being provided. One participant outlined
- the initial care provided by KCMC clinical staff, *"they started running tests on his head, and the"*
- 363 *next day, they told me that he needed surgery to prevent it from spreading to the other eye. He*
- had the surgery, and I stayed there for three days before coming here to the cancer ward." (IDI
- 365 Participant 6)

366

367 Although most caregivers opted to begin treatment immediately for their child at KCMC,

368 potential delays presented for families who returned home before initiating treatment. One

369 mother describes her experience after obtaining her child's cancer diagnosis, *"When we came*

- 370 *here to KCMC in July, the eye was not that swollen, and the eye had not come out and the*
- 371 examination revealed the cancer... After they gave me the results, I went back home...I tried

372 covering the eye that had no disease, and I gave the child something to look at, but I realized he

373 *couldn't see it nor anything else. That's when I made up my mind to come back."* (IDI Participant

374 5)

375

376 Caregiver concerns

Participants described some of their worries about being away from home such as familial
relationships, household economic implications of obtaining care, and the health trajectory of
their child. A participant describes their worries about the additional children at home and how
being away from home may be impacting the children's school attendance, *"Yes, worrying is a*

381	must because we left our young children at home. They go to school on their own, and if I'm not
382	there, they might only go for two or three days a week, or sometimes they don't go at all." (IDI
383	Participant 3) Another respondent expressed the sense of hopelessness that can accompany a
384	cancer diagnosis. "There is no worse illness than this. You need great support. I see people here
385	laughing and appearing well-fed, and I feel like something is wrong in their brains this is not a
386	place to laugh; it's not a safe place. I'd prefer being in jail than here because in jail, you'll
387	eventually get out alive, but here, it's by grace." (IDI Participant 12)
388	
389	Nearly half of participants described how they drew upon their religious or spiritual beliefs to
390	manage their worries and find a sense of comfort and "grace" (IDI Participant 11). One
391	participant stated, "I just pray to God, rely on him, and ask God for strength. When I pray and
392	lean on God, the fear diminishes." (IDI Participant 4) Further, a child's uncle suggests how the
393	pride they feel in reaching care has lessened their concerns, "I don't have any worries because I
394	have saved a life. I only think about the child I have saved." (IDI Participant 2)
395	
396	Treatment success
397	Despite concerns, participants expressed a sense of hope in their perceived effectiveness of the
398	treatment provided at KCMC. Nearly all participants felt like their child would complete the
399	treatment plan, rarely noting additional delays that may prevent them continuing to receive
400	services. Caregivers felt reassured by the progress in their child's condition, which motivated
401	them to continue with the treatment plan. One respondent said, "just seeing the child playing
402	gives me some peace of mind." (IDI Participant 5) Another participant expressed their
403	satisfaction with KCMC as a medical facility compared to another hospital, "KCMC is the

404 biggest and the best treatment here... So, if anyone has a problem, I will tell them to directly
405 come to KCMC for salvation." (IDI Participant 2)

406 Finances

407 Finally, participants described how finances impacted their ability to continue receiving

408 oncology care and avoid treatment abandonment. When asked if they experienced any stress

409 regarding paying for oncology care, one mother noted how the philanthropic financing available

410 at KCMC allows her to not have financial worries stating, "Not at all, because they told me that

411 *the child's treatment costs here at the hospital are covered. They told me there are no expenses,*

412 *my cost is my time.*" (IDI Participant 7) Overall, participants expressed a sense of relief in

413 knowing they would be able to continue treatment for their child without incurring catastrophic

414 medical expenses. However, before arriving to KCMC, respondents were unaware of this

415 financing option and often noted how the perceived price of treatment in Moshi was

416 *"incomprehensible"* (IDI Participant 2). One participant said, *"I knew that it might involve a lot*

417 of money because we are used to hearing that when you hear about KCMC, you know you need

418 *to be well-prepared. They say the tests themselves are expensive, and just entering the hospital*

419 *itself is costly, unlike other hospitals.*" (IDI Participant 3)

420

While this did not necessarily cause a delay in the continuation of care, participants discussed
how they had to forego their normal employment or livelihood activities regardless of the free
treatment provided. One mother explained how the lost income will impact her family, *"It affects us significantly because my job was a great source of income... It's the biggest challenge that I see affecting our lives."* (IDI Participant 4)

426

427 DISCUSSION

428 To the best of our knowledge, this is the first paper to qualitatively apply the Three Delays 429 Model to pediatric cancer care in Northern Tanzania. Most notably, we report the most 430 substantial delays occur once families have sought care and experience barriers related to the 431 Tanzanian referral system and health infrastructure capacity. While global health research has 432 pushed for decentralized healthcare expansion in LMICs, health system capacity must 433 concurrently advance to support the appropriate delivery of pediatric cancer care. Overall, an 434 increase in community and clinical resources aimed at reducing financial constraints and 435 ensuring quality, timely access to pediatric oncology care is required in Tanzania. Further, we 436 recommend that future interventions extend beyond basic access to care to include holistic 437 strategies that aid families in completing full treatment regimes.

438

439 Respondents described initial delays due to waiting for the child's condition to resolve or 440 deteriorate before seeking medical care. Although caregivers expressed how they identified 441 symptoms and changes in the child's behavior, unlike injurious or emergency conditions, early 442 signs of pediatric cancer can be difficult to discern. Due to the complexities of recognizing signs 443 of pediatric cancer, we recommend that community health workers (CHWs) and traditional 444 healers be trained to identify and suggest when further medical treatment should be sought. For 445 example, a study conducted in Cameroon developed an early recognition and referral training for 446 primary healthcare workers, including traditional healers, and found that healthcare workers 447 knowledge of early childhood cancer symptoms increased by 52% after completing the training 448 program.(18) The WHO Global Strategy on Human Resources for Health further suggests the 449 value of CHWs, and urges countries to embrace CHWs as diverse and sustainable primary

healthcare professionals.(19,20) Building on the current literature and global recommendations,
future interventions should engage CHWs and traditional healers in cost-effective and culturally
reflective pediatric cancer training and outreach, thus limiting delays at the household level.

454 Out of all the barriers mentioned, health system barriers were the most often discussed among 455 respondents. While the stepwise referral system in Tanzania has benefits for preventing the 456 treatment of mild conditions at high-level facilities, our study suggests that the referral system 457 hindered patients from obtaining care in a timely manner. Keating et al. report that referral 458 delays for severe pediatric illnesses put children at greater risk for mortality in Tanzania, which 459 we believe to be applicable to pediatric cancer.(4) Additionally, the costs associated with 460 misdiagnosis and ineffective treatments were considerable and prohibited some participants from 461 accessing referrals promptly. While caregiver's did not perceive this referral system to have 462 hindered the overall treatment or prognosis of their child, the global pediatric oncology literature 463 strongly supports the necessity of prompt diagnosis for treatment effectiveness, minimizing out-464 of-pocket (OOP) costs, and reducing the suffering accompanying a cancer diagnosis.(21) 465 Households in LMICs, such as Tanzania, are more likely to finance medical expenses by selling 466 assets, using personal savings, and borrowing money, impacting the long-term socioeconomic 467 status of the household.(22) These financial strategies are supported by the data we present and 468 are exacerbated by the lack of health insurance in our study sample. Smith et al. has highlighted 469 the cost-effectiveness, economic growth, and equity building that investment in global pediatric 470 surgical care produces, principles relevant to global pediatric cancer care. (23) Therefore, 471 progressive financial protection schemes such as individual-level support for financing indirect 472 medical expenses such as lodging, food, and transportation and macro-level interventions such as

473 the inclusion of pediatric oncology diagnosis and treatment in universal health coverage (UHC) 474 plans must be considered. In addition, it will be critical that the capacity to recognize children 475 with potential cancer conditions improves at smaller health facilities. To minimize disease 476 progression and household OOPs costs, workers within these facilities should be trained to 477 identify concerning symptoms of pediatric cancer and refer these patients to a definitive 478 treatment facility. Efforts addressing these concerns have proven successful in Tanzania with 479 Muhimbili National Hospital reporting an 8% treatment abandonment rate for acute leukemia 480 after accommodation was provided for families of children obtaining cancer care in addition to 481 infrastructure advancements.(24,25) 482 483 While the severity of cancer was well known, respondents expressed varying thoughts 484 surrounding the causes of cancer, how cancer may reoccur, treatment strategies, and follow up 485 care. Participants also suggested the importance of spirituality and support networks for coping with their child's diagnosis. The feelings described by participants represent the necessity of 486 487 clear health education and tangible strategies for managing the emotions that accompany a 488 cancer diagnosis. The National Cancer Institute (NIC) has developed six elements of patient-489 centered communication in cancer care including 1) fostering healing relationships, 2) 490 responding to emotions, 3) exchanging information, 4) making decisions, 5) managing 491 uncertainty, and 6) enabling self-management. (26) While these elements have mostly been 492 applied to HICs, a review has recommended the NCI guidelines as a baseline for further research and intervention in LMICs.(27) For example, cancer support groups have been implemented in 493 494 numerous settings to support families in HICs. However, similar programs have yet to be widely 495 incorporated in LMICs. Participant responses detailing worries about being away from home and

the confusion that accompanied their child's cancer diagnosis suggests that future childhood
cancer interventions could focus on delivering comprehensible health education and providing
supportive spaces for managing fears and uncertainty. Due to treatment abandonment concerns,
this extension of health education is especially valuable in pediatric oncology settings where
routine follow-up care is necessary.

501

502 While these results will inform future interventions aimed at increasing the quality and 503 accessibility of pediatric cancer care in Tanzania, our study contains important limitations. Like 504 any qualitative research study, these findings are not intended to be representative of the entire 505 pediatric oncology patient population. Rather, this study intends to detail the journey to pediatric 506 cancer care and identify potential opportunities for intervention that support timely and quality 507 service provision. Second, while the thematic codebook underwent routine discussion and 508 adaptation based on the perspectives of research team members based at KCMC, not all 509 interviews were doubled coded, and agreeability was determined by a random sample of 6 510 interviews. While this is a notable limitation, no major discrepancies in coding outcomes were 511 noted. between Lastly, we report perspectives of only households who received care at KCMC, 512 excluding the perspectives of families who were unsuccessful in reaching a final destination for 513 oncology treatment and therefore may have experienced more significant barriers to accessing 514 care.

515

516 **CONCLUSION**

517 In this qualitative study, we report on the journey to pediatric cancer care beginning at the518 household level and extending beyond care received at KCMC to describe stressors and plans for

- the future. Our recommendations include 1) empowerment of CHWs and local traditional healers
- 520 to advocate for earlier care seeking behavior at the community level, 2) implementation of
- 521 financial protection, clinical structures, and training at intermediary medical centers aimed at
- 522 earlier referral to a definitive treatment facility, 3) incorporation of support and health education
- 523 initiatives for families of children with a cancer diagnosis. Most importantly, in accordance with
- the United Nation's target of achieving universal health coverage (UHC) worldwide by
- 525 2030,(28) future national health plans should include pediatric cancer care.

- 526 *Funding:* This research was supported by the Duke Global Health Institute and the National
- 527 Institutes of Health K01
- 528 *Competing Interests:* The authors declare no competing interests
- 529

530 Author Contributions

- 531 Conceptualization: MM, ERS, HER, PE, CC-C
- 532 Methodology: MM, ERS, AT, PE
- 533 Formal Analysis and Investigation: MM, ERS, HK, AT, PE
- 534 Data Collection: MM, HK, EM
- 535 Writing Original Draft Preparation: MM, ERS
- 536 Writing Review and Editing: MM, AT, JRV, CAS, ERS, HER, CC-C
- 537 Funding Acquisition: ERS, CAS, EM, BTM
- 538 Supervision: ERS, EM, BTM, CAS

539 REFERENCES

- Atun R, Bhakta N, Denburg A, Frazier AL, Friedrich P, Gupta S, et al. Sustainable care for children with cancer: a Lancet Oncology Commission. Lancet Oncol. 2020 Apr 1;21(4):e185–224.
- The Global Initiative for Childhood Cancer [Internet]. [cited 2023 Feb 16]. Available from: https://www.who.int/initiatives/the-global-initiative-for-childhood-cancer
- Farmer P, Frenk J, Knaul FM, Shulman LN, Alleyne G, Armstrong L, et al. Expansion of
 cancer care and control in countries of low and middle income: a call to action. The Lancet.
 2010 Oct 2;376(9747):1186–93.
- 548 4. Keating EM, Sakita F, Mmbaga BT, Amiri I, Nkini G, Rent S, et al. Three delays model
 549 applied to pediatric injury care seeking in Northern Tanzania: A mixed methods study. PLOS
 550 Glob Public Health. 2022 Aug 25;2(8):e0000657.
- 5. Hosaka L, Tupetz A, Sakita FM, Shayo F, Staton C, Mmbaga BT, et al. A qualitative
 assessment of stakeholder perspectives on barriers and facilitators to emergency care delays
 in Northern Tanzania through the Three Delays. Afr J Emerg Med. 2023 Sep;13(3):191–8.
- 554 6. UNICEF DATA [Internet]. [cited 2023 Aug 31]. United Republic of Tanzania (TZA) 555 Demographics, Health & Infant Mortality. Available from:
 556 https://data.unicef.org/country/tza/
- 557 7. Majaliwa E, Smith ER, Cotache-Condor C, Rice H, Gwanika Y, Canick J, et al. Childhood
 558 and Adolescent Cancer Care at a Tertiary Hospital in Northern Tanzania: A Retrospective
 559 Study. JCO Glob Oncol. 2023 Sep;(9):e2200263.
- Greenspan JA, Chebet JJ, Mpembeni R, Mosha I, Mpunga M, Winch PJ, et al. Men's roles in care seeking for maternal and newborn health: a qualitative study applying the three delays model to male involvement in Morogoro Region, Tanzania. BMC Pregnancy Childbirth.
 2019 Aug 13;19(1):293.
- 564 9. Thaddeus S, Maine D. Too far to walk: maternal mortality in context. Soc Sci Med 1982.
 565 1994 Apr;38(8):1091–110.
- 10. Cotache-Condor C, Kantety V, Grimm A, Williamson J, Landrum KR, Schroeder K, et al.
 Determinants of delayed childhood cancer care in low- and middle-income countries: A
 systematic review. Pediatr Blood Cancer. 2023 Mar;70(3):e30175.
- 569 11. Steliarova-Foucher E, Colombet M, Ries LAG, Moreno F, Dolya A, Bray F, et al.
 570 International incidence of childhood cancer, 2001–10: a population-based registry study.
 571 Lancet Oncol. 2017 Jun 1;18(6):719–31.
- Johnston WT, Erdmann F, Newton R, Steliarova-Foucher E, Schüz J, Roman E. Childhood
 cancer: Estimating regional and global incidence. Cancer Epidemiol. 2021 Apr 1;71:101662.
- 13. About Us [Internet]. [cited 2023 Jan 23]. Available from: https://www.kcmc.ac.tz/about

- 575 14. Cancer Care [Internet]. [cited 2023 Sep 9]. Available from: https://www.kcmc.ac.tz/cancer 576 care
- 577 15. pgssc [Internet]. [cited 2024 Apr 9]. Old_National Surgical Planning. Available from: 578 https://www.pgssc.org/national-surgical-planning
- 579 16. Lumivero [Internet]. [cited 2023 Oct 26]. NVivo. Available from:
 580 https://lumivero.com/products/nvivo/
- 581 17. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research
 582 (COREQ): a 32-item checklist for interviews and focus groups. Int J Qual Health Care J Int
 583 Soc Qual Health Care. 2007 Dec;19(6):349–57.
- 18. Afungchwi GM, Hesseling PB, Kouya F, Enow SA, Kruger M. The outcome and cost of a capacity-building training programme on the early recognition and referral of childhood cancer for healthcare workers in North-West Cameroon. Nurs Open. 2020;7(6):2029–38.
- 587 19. Global strategy on human resources for health: Workforce 2030 [Internet]. [cited 2024 Jun 26]. Available from: https://www.who.int/publications/i/item/9789241511131
- 589 20. Community-based health workers (CHWs) [Internet]. [cited 2024 Jun 26]. Available from:
 590 https://www.who.int/teams/health-workforce/community
- 591 21. Childhood cancer [Internet]. [cited 2023 Sep 19]. Available from:
 592 https://www.who.int/news-room/fact-sheets/detail/cancer-in-children
- 593 22. Donkor A, Atuwo-Ampoh VD, Yakanu F, Torgbenu E, Ameyaw EK, Kitson-Mills D, et al.
 594 Financial toxicity of cancer care in low- and middle-income countries: a systematic review
 595 and meta-analysis. Support Care Cancer. 2022 Sep 1;30(9):7159–90.
- 596 23. Smith ER, Concepcion TL, Niemeier KJ, Ademuyiwa AO. Is Global Pediatric Surgery a
 597 Good Investment? World J Surg. 2019 Jun 1;43(6):1450–5.
- 598 24. Schroeder K, Saxton A, McDade J, Chao C, Masalu N, Chao C, et al. Pediatric Cancer in
 599 Northern Tanzania: Evaluation of Diagnosis, Treatment, and Outcomes. J Glob Oncol. 2017
 600 Jun 9;4:JGO.2016.009027.
- Kersten E, Scanlan P, DuBois SG, Matthay KK. Current treatment and outcome for
 childhood acute leukemia in Tanzania. Pediatr Blood Cancer. 2013;60(12):2047–53.
- Epstein RM, Street Jr. RL. Patient-centered communication in cancer care: Promoting
 healing and reducing suffering. 2007;
- Graetz DE, Garza M, Rodriguez-Galindo C, Mack JW. Pediatric cancer communication in
 low- and middle-income countries: A scoping review. Cancer. 2020;126(23):5030–9.
- 607 28. Universal health coverage (UHC) [Internet]. [cited 2023 Feb 5]. Available from:
 608 https://www.who.int/news-room/fact-sheets/detail/universal-health-coverage-(uhc)