1	
2	
3	
4	Relationship between midwifery density and midwifery regulatory environments
5	in International Confederation of Midwives member countries
6	
7	
8	Emma Virginia Clark ^{1*} , Marianna LaNoue ¹ , Kate Clouse ¹ , Alexandra Zuber ² , Jeremy Neal ¹
9	
10	
11	¹ School of Nursing, Vanderbilt University, Nashville, Tennessee, United States of America
12	² Ata Health Strategies LLC, Washington, District of Columbia, United States of America
13	
14	
15	*Corresponding Author:
16	E-mail: <u>emma.v.clark@vanderbilt.edu</u> (EVC)
17	

18

19 Abstract

20 Background

- 21 Midwives are a cost-effective solution for reducing preventable maternal death, but barriers to
- 22 their recruitment, training, deployment, and retention exist. Improved midwifery regulation is
- proposed to address these challenges through activities related to education and qualification,
- 24 licensure, registration/re-licensure, and scope and conduct of practice. However, no
- 25 quantitative analysis of the associations between midwifery regulation and workforce density in
- low- and middle-income countries has been conducted, limiting actionable insights for
- 27 policymakers. The objective of this study was to evaluate the relationship between midwifery
- regulatory environments and midwifery workforce densities (midwives per 10,000 population)
- 29 in International Confederation of Midwives member countries.
- 30

31 Methods

- 32 The study used secondary data from the 2019 Global Midwives' Associations Map Survey to
- perform a cross-sectional analysis of 103 International Confederation of Midwives member
- 34 countries. To operationalize the strength of a country's midwifery regulatory environment, we
- 35 used a previously developed Midwifery Regulatory Environment Index. We conducted
- 36 multivariable linear regression analysis to evaluate the relationship between this index and
- 37 country-level midwifery density, available from the National Health Workforce Accounts.

38 Results

- 39 Midwifery regulatory environments are significantly associated with midwifery workforce
- 40 densities across countries. For every 1-point increase in the Midwifery Regulatory Environment
- 41 Index (indicating activities in at least one additional regulatory domain), midwifery density
- 42 increased by 0.66 midwives per 10,000 population (p = 0.011). This effect decreased to 0.41
- 43 midwives per 10,000 population yet remained significant even after controlling for country level
- 44 income, per capita health spending, income inequality, and human development index (p =
- 45 0.04).

46 Conclusion

- 47 Stronger regulatory environments are associated with higher densities of midwives, even when
- 48 controlling for potential confounding factors such as country-level income and health spending.
- 49 This relationship suggests that policymakers should invest in midwifery regulation to improve
- 50 maternal health outcomes.

51 Introduction

52

53	The current global sexual, reproductive, maternal, newborn, and adolescent health
54	(SRMNAH) workforce is insufficient, meeting only 75% of the global need for essential services
55	[1]. This workforce provides care across the reproductive lifespan, including preconception
56	(e.g., offering modern contraceptive methods), early pregnancy (e.g., post-abortion care,
57	ectopic pregnancy management), antenatal (e.g., screening and managing hypertensive
58	disorders, diabetes, and malaria), intrapartum (e.g., parenteral administration of
59	anticonvulsants, uterotonics, and antibiotics, manual removal of placenta), and postpartum
60	(e.g., promoting breastfeeding) periods [2]. In low-income countries, only up to 41% of the
61	need for these essential services is met [1].
62	Midwives are increasingly recognized as a cost-effective means for delivering SRMNAH
63	services that improve outcomes and reduce preventable maternal and newborn mortality [3-5],
64	yet an estimated 900,000 more midwives are needed to meet global service delivery needs [1].
65	The availability of a country's health workforce to provide essential SRMNAH services in any
66	given geographical area can be estimated using national workforce density reports, with low-
67	and middle-income countries (LMICs) having the greatest workforce shortfall [6]. Numerous
68	regulatory barriers at the country level can contribute to this shortfall, hindering the
69	recruitment, training, deployment, and retention of midwives. These health professional
70	regulation barriers include unclear and non-standardized role definitions, limited scopes of
71	practice misaligned with international standards, varying educational and clinical standards for
72	entry to practice, and weak licensure and registration authorities and systems [7,8].

73	Health professional regulation establishes mechanisms and standards that recognize
74	and delineate the roles of health professionals, providing flexibility to adapt to complex and
75	evolving health system contexts [9]. According to the World Health Organization (WHO),
76	regulation is a priority component of advancing the health workforce, and policymakers should
77	consider it a strategic tool for addressing workforce imbalances and other challenges to ensure
78	adequate training and availability of health professionals [10]. Specifically, midwifery regulation
79	defines criteria and processes outlined in legislation to identify what a midwife is and to
80	describe their scope of practice [11]. The aim of regulation is to promote regulatory
81	mechanisms that protect the public (women and families) by ensuring that safe and competent
82	midwives provide high standards of midwifery care to every woman and baby through such
83	actions as producing autonomous, high-quality practice [11].
84	The presence of distinct regulatory structures and processes for midwifery is crucial to
85	enable autonomous midwifery practice and maintain midwifery standards, both of which are
86	essential for providing high-quality care to mothers and infants [11]. Midwifery regulation can
87	be categorized into the following regulatory domains, which are derived from the International
88	Confederation of Midwives (ICM) Global Standards for Midwives [11] and adapted through
89	additional research [12-14].
90	• Overarching Regulatory Policy and Legislation: Enacting legislation that recognizes
91	midwifery as a profession distinct from nursing and establishing a dedicated regulatory
92	body with specifically for midwives or with separate and distinct policies and processes

93 for midwives.

94	• Education and Qualification: Establishing standards for midwifery education, curricula,
95	and, occasionally, accreditation of educational institutions to develop necessary
96	competence for practice.
97	• Licensure: Developing and enforcing criteria, standards, and processes for initial entry
98	into practice.
99	Registration/Re-Licensure: Establishing continuing medical education or other
100	requirements for demonstrating continued competence and maintaining a current
101	database of midwives, including their registration/licensure status, to support workforce
102	planning, and consumer transparency.
103	• Scope and Conduct of Practice: Clearly defining the scope of practice of a midwife
104	consistent with ICM's definition of a midwife and establishing mechanisms for handling
105	complaints and disciplinary action.
106	A strong midwifery regulatory environment is one in which midwifery is recognized as a distinct
107	profession, and a specific regulatory body, led by midwives, exists. Alternatively, a distinct track
108	for midwifery regulation within a nursing board or council may suffice. This empowered body,
109	established through regulatory policies, has statutory authority and effectively implements and
110	oversees activities across the domains noted above. Conversely, a weak regulatory
111	environment is characterized by the absence, inadequacy, or non-functionality of regulatory
112	activities and mechanisms.
113	Midwifery density is the number of midwives per 10,000 population in a specified
114	geographic area. Density data are critical for assessing workforce availability, recruitment,
115	deployment, and retention. The WHO and global professional associations have emphasized

the significance of regulation in advancing the midwifery workforce, resulting in a growing body
of research aiming to formally establish linkages between regulation and midwifery workforce
densities. Predominantly conducted in high-income countries, these studies have explored how
regulations constraining midwifery practice can lead to a reduction in the midwifery workforce
[15-17].

Research in the United States has underscored that strong regulatory environments are 121 122 related to higher midwifery workforce densities [18,19]. However, limited evidence exists on 123 this relationship in LMICs, where research mostly focuses on documenting midwifery workforce 124 challenges and suggesting that improved regulation might alleviate these challenges, without exploration of the underlying mechanisms. For example, a scoping review in Kenya [20] and a 125 126 descriptive review in Nigeria [21] effectively highlight how poor job satisfaction and low 127 retention rates contribute to midwifery shortages, with the suggestion that weak regulatory 128 environments may be a driving factor. However, similar to other LMIC-focused research, these studies lacked a quantitative analysis of associations between regulation and workforce, 129 130 limiting actionable insights for policymakers. This study aims to evaluate the relationship between midwifery regulatory 131 environments and midwifery workforce densities in ICM member countries, addressing a 132 133 critical gap in the literature. We hypothesize that countries with stronger midwifery regulatory 134 environments will have higher densities of midwives.

135 Methods

This study was conducted using a cross-sectional design and secondary data analysis of multiple publicly available databases. Approval for this study was obtained from the Vanderbilt University Medical Center Institutional Review Board.

139 Data and Variables

140 The independent variable for this study was the Midwifery Regulatory Environment (MRE) Index, a composite score our team developed to operationalize the strength of a 141 country's midwifery regulatory environment. Data used in developing this index were drawn 142 143 from the ICM Global Midwives' Association Map Survey. For this survey, ICM collected data 144 from country midwifery associations with support from the United Nations Population Fund (UNFPA) Country Office and relevant government organizations between October and 145 146 December 2019 [22]. The survey included four subsections on midwifery: association, education, leadership, and regulation. To ensure data quality, ICM used a multi-step process 147 148 including an initial review with country follow-up for clarifications and completeness, internal logic, and validity; review of the revised and validated data by country midwifery associations; 149 and a letter of validation from an official for each survey section or confirmation that data were 150 publicly available from reputable sources [22]. 151

To develop the MRE Index, we began by identifying relevant data from the Map Survey for inclusion in the composite score. The regulatory sub-section contained 38 questions assessing a country's regulatory environment [23]. We eliminated sixteen questions for freetext answers and/or not directly relating to at least one regulatory domain. The remaining 22 questions were consolidated and organized into 12 items, each associating with one of the five essential domains noted above [overarching regulatory policy and legislation (4 items);

education and qualification (0 items); licensure (4 items); registration/re-licensure (2 items); and scope and conduct of practice (2 items)]. Because the education and qualification domain was not reflected in the regulatory sub-section questions, one question from the education subsection of the Map Survey about the national curriculum for midwifery education was included as a final item for MRE Index development purposes. This resulted in a total of 13 items for inclusion in calculation of the final composite MRE Index.

Five different potential scoring methodologies for aggregating the 13 items into a single 164 165 MRE Index were developed, and the characterization that best fit the data and had the highest 166 predictive ability for each of three outcomes (low birthweight prevalence, stillbirth rate, MMR) was selected. While all characterizations performed well, the "Aggregated Domain Scoring" 167 characterization proved to be the best. In this approach, the 13 items are grouped into five 168 169 categories, each representing a regulatory domain: overarching regulatory policy and 170 legislation, education and qualification, licensure, registration/re-licensure, or scope and conduct of practice. One point is assigned to each domain with at least one "yes" response to 171 the item(s) within that domain, allowing for a total score between zero (no regulatory activities 172 in any domain) and five (regulatory activities in all five domains). Detailed methodology 173 regarding development of the MRE Index is published in a forthcoming manuscript. 174 175 The dependent variable for this study was the density of midwives at the country level, 176 defined as the number of midwives per 10,000 population. These data were obtained from WHO's National Health Workforce Accounts (NHWA), which facilitates data reporting on key 177 workforce indicators to improve the availability, guality, and use of health workforce data with 178 179 the goal of achieving universal health coverage [24]. WHO's 194 member states submit health

180	workforce data annually to the NHWA. These data sources include health workforce registries
181	or databases, aggregated data from health facilities such as routine administrative records and
182	District Health Information System census or surveys, professional council, chamber, and/or
183	association registers, and labor force surveys [25]. The NHWA uses population estimates
184	provided by the United Nations (UN) Statistics Division, drawn from civil registration and vital
185	statistics systems, population censuses or registers, population registers, and household
186	surveys. If other methodologies are used, WHO recalculates densities using the UN Statistics
187	Division population estimates described above to harmonize densities [25].
188	Covariate data were obtained from publicly available databases including the United
189	Nations Development Program's (UNDP) Human Development Index (HDI) [26] and WHO's
190	Global Health Expenditure database [27], and the World Bank's Gini index [28] and income
191	group databases [29]. Four control variables were selected to reduce potential bias related to
192	overall allocation and distribution of resources within countries. These included two continuous
193	variables: per capita health spending in United States dollars and the Gini index, which
194	measures the extent to which the distribution of income or consumption among individuals or
195	households within an economy deviates from a perfectly equal distribution [28]. We also used
196	two categorical variables. A country's income group is assigned annually by the World Bank
197	based on the gross national income (GNI) per capita of the previous year, expressed in United
198	States dollars. In 2022, countries were classified as low income (GNI per capita less than
199	\$1,085), lower-middle income (GNI per capita \$1,086-4,255), upper-middle income (GNI per
200	capita \$4,256-13,205), and high income (GNI per capita greater than \$13,205) [29]. The HDI, the
201	final control variable, is a composite measure of average achievement in three key dimensions

of human life: a long and healthy life measured by life expectancy at birth; education, measured
by mean years of schooling and expected years of schooling; and a decent standard of living,
measured by gross national income per capita in Purchasing Power Parity (PPP) international
dollars. It is reported on a scale from 0 to 1.00, with ≥ 0.800 classified as "very high" human
development, 0.700-0.799 as high, 0.550-0.699 as medium, and < 0.550 as low [26].

207 Data Screening and Analysis

One hundred fifteen countries submitted complete responses to the regulatory 208 209 subsection of the ICM Global Midwives' Association Map Survey, which was used for calculating 210 the MRE Index. Three countries were missing density information, and nine others were missing data for one or more covariates, leaving a total of 103 countries for analyses. We transformed 211 212 two highly-skewed variables, midwifery workforce density and per capita health spending, by 213 selecting the positive square root transformation which had the best effect on skewness and 214 kurtosis. We used multivariable linear regression to test for associations between the MRE 215 Index and midwifery density, adjusting for the covariates. Statistical significance was set a priori 216 at 0.05. SPSS Statistics 27 was used to conduct the analysis.

217 **Results**

MRE Indices across countries ranged from 2 to 5, with an average score of 4.36 (SD = 0.88). Midwifery workforce density ranged from 0.11 midwives per 10,000 population in Burundi to 20.46 midwives per 10,000 population in Ireland. The median density was 2.41 midwives per 10,000 population (interquartile range [IQR] 1.43, 5.10). Table 1 presents descriptive summaries of MRE Index, midwifery workforce density, and control variables used in the regression analysis (i.e., per capita health spending, Gini index, income group, and HDI).

224	Table 1. Descriptive statistical summaries	s of country characteristics (N = 103)
-----	--	--

	Mean	SD
MRE Index	4.36	0.88
Gini Index (1-100)	37.39	7.48
	Median	IQR
Health Spending (USD)	201.06	58.5, 1278.18
Density of Midwives (per 10,000 population)	2.41	1.43, 5.10
	%	
Human Development Index		
Low	22.0	
Medium	28.0	
High	17.0	
Very High	33.0	
Income Level		
Low	25.0	
Low-Middle	29.0	
Upper-Middle	17.0	
High	29.0	

225

226 Correlations among the explanatory variables indicate that all control variables were 227 significantly associated with each other, with health spending, Gini index, and income group 228 demonstrating very strong associations (Table 2). However, MRE Index was not significantly 229 associated with any of the covariates.

	MRE Index	Health Spending ^a	Income Group	Huma Developi Index
Health Spending ^a	-0.097	-		
Income Group	1.19	155.037***	-	
Human Development Index	1.42	101.55***	181.055***	-
Gini Index	0.003	0.304*	9.35***	8.10**

230 Table 2: Correlations among MRE Index and control variables (N= 103)

Chi square for categorical variables, Person r for continuous variables, one-way ANOVA for
 mixed categorical (>2 categories) & continuous variables.

249 * = p < 0.05, ** = p < 0.01, *** p < 0.001.

250 ^a transformed variable

251 252

Results of univariate and multivariate regression analyses are shown in Table 3. The

253 multiple regression of all explanatory variables with midwifery workforce density was

significant (Multiple *R* = 0.405, p = 0.010), explaining approximately 12% of the variance. While

255 all explanatory variables showed significant univariate correlations with midwifery workforce

density, only health spending and the MRE Index remained significant after adjusting for health

257 spending, HDI, Gini index, and income level.

258 Since transformed variables were used for midwifery workforce density and per capita

- 259 health spending, beta values were calculated from non-transformed variables to aid
- 260 interpretation. These values demonstrate that for every 1-point increase in the MRE Index
- 261 (indicating activities in at least one additional regulatory domain), midwifery density increased
- by 0.66 midwives per 10,000 population (p = 0.011). This effect decreased to 0.41 midwives per

- 263 10,000 population when accounting for health spending, income group, inequality, and HDI, yet
- the MRE Index remained significantly associated with midwifery workforce density (p = 0.04).

265	Table 3: Summary of univariate and multivariate associations with midwifery density (N =
266	103)

Characteristic	r	<i>p-</i> value	beta ^c	SE	95% Wald Confidence Interval		<i>p-</i> value
					Lower	Upper	
MRE Index	-0.24	0.011	0.41	.013	-0.37	1.19	0.04
Health Spending (USD)	0.31	<0.001	0.00	0.00	0.00	0.00	0.03
Income Group ^a	5.00	0.003					
2-1			-1.51	1.39	-4.24	1.23	0.28
3-1			-0.93	1.89	-4.63	2.77	0.28
4-1			0.39	2.15	-3.82	4.61	0.39
Human Development Index ^b	6.24	<0.001					
2-1			3.89	1.43	1.09	6.68	0.05
3-1			2.16	1.92	-1.61	5.92	0.59
4-1			1.46	1.97	-2.39	5.31	0.97
Gini Index	0.29	0.002	-0.02	0.05	-0.12	0.09	0.12

267 Multiple R = 0.405, p = 0.010; $R^2 = 0.164$ (Adjusted $R^2 = 0.120$)

^aIncome Group: 1 = Low; 2 = Lower Middle; 3 = Upper Middle; 4 = High; all income group values
 are relative to low-income group.

^bHuman Development Group: 1 = Low; 2 = Middle; 3 = High; 4 = Very High.

²⁷¹ ^cBeta obtained from model with untransformed variables for interpretation purposes.

272

273 **Discussion**

This study aimed to determine if there is a relationship between midwifery regulatory 274 275 environments and density of midwives in ICM member countries. The findings support our hypothesis that there is a positive association between the midwifery regulatory environments 276 277 and midwife density; stronger regulatory environments are associated with higher densities. 278 This relationship remains significant even when controlling for potential confounding health 279 system and country factors. The sample included a relatively balanced representation of low-, 280 lower-middle, upper-middle, and high-income countries, as well as countries with varying levels 281 of human development, although there were slightly more high-income and high human 282 development countries. Research such as the WHO's "Midwives' Voices, Midwives' Realities" consultation 283 highlights numerous issues that affect density. These issues include inadequate education, 284 uncertainty about midwifery's role in the health system, restrictions on practice that are 285 286 inconsistent with the legal scope of practice, and overwork leading to burnout and attrition [30]. Each regulatory domain -- overarching regulatory policy and legislation, education and 287 288 qualification, licensing, registration/re-licensure, and scope and conduct of practice – contains aspects that can influence midwifery density, with significant cross-over and reinforcement 289 290 between domains. For example, in the overarching regulatory policy and legislation domain, 291 Mattison et al. found when health system decision-makers did not recognize midwifery as an 292 autonomous profession, such as through distinct regulatory bodies, midwifery was excluded from maternal health-related political agendas [31]. This omission led to downstream impacts 293 such as lack of clarity about midwives' specific roles in the health system and inappropriate 294

deployment and remuneration. These factors significantly affect the desirability of midwifery as 295 296 a profession and influence how many people choose to enter and remain in the profession. Regulation in the education and gualification domain helps standardize education 297 298 requirements for midwives, ensuring they meet at least minimum education standards and are 299 prepared for their defined scope of practice. Aligning these standards with international norms further strengthens credibility, making the profession more attractive and drawing in more and 300 301 stronger candidates. Administrative aspects of regulation, such as licensure and registration, 302 can improve density by providing data on the number of midwives working in the country, 303 distinguishing between active, inactive, and retired midwives, and identifying their locations. This information is crucial for forecasting supply and analyzing it alongside demand data to 304 305 understand labor market dynamics and gaps. Such insights support appropriate investments in 306 midwifery training institutions and efforts to expand midwifery scope of practice, ensuring an 307 adequate number of midwives are available to provide needed services. Lastly, activities within the scope and conduct of practice domain are fundamental aspects of regulatory environments 308 309 [12] and major drivers of midwifery workforce density. Well-defined scopes of practice that promote strong interprofessional collaboration have been shown to attract and retain 310 311 healthcare workers by clarifying task division, enhancing support, and managing overall 312 workload [32,33]. A qualitative systematic review including studies from 23 LMICs found that 313 workplace relationships between healthcare professionals that are built on trust contribute to social interactions and cooperation among healthcare workers, impacting their intrinsic 314 motivation, retention, performance, and quality of care [34]. There is also evidence suggesting 315 316 that midwives who practice autonomously – defined as the authority to make decisions and act

freely based on their knowledge base- experience increased occupational satisfaction, 317 318 improved professional identity, and reduced burnout [4,35,36]. Qualitative research on policymakers' and regulators' perspectives from both strong and weak regulatory environments 319 320 could identify additional domains or domain activities not considered in this research. Research 321 tracking health systems that are in the process of enhancing regulatory mechanisms for midwives, such as Lao People's Democratic Republic [37], could also be evaluated using 322 approaches that account for the complexity of health systems, such as difference-in-323 324 differences. It is also critical to ascertain if the relationship between midwifery workforce 325 density and regulation holds at a sub-national level, as remote and rural areas often have much lower densities than urban areas and national averages, leading to poorer maternal, newborn, 326 and child health outcomes [38]. While the NHWA has the capacity to report sub-national 327 328 densities, only two countries in the study sample reported this information. As sub-national 329 density data becomes more available, or in countries where it is currently available, this analysis could influence overall and specific domain investment in midwifery regulation. 330

331 Limitations

This study had several limitations. As a secondary data analysis, it is limited by the quality of the datasets used. The data quality and validation processes for the main data sources are discussed, with quality variation being a generally accepted limitation of secondary data sources. Our regression analyses only included countries that are ICM members and participated in the ICM mapping survey, potentially introducing bias toward countries with more midwives and more sophisticated regulatory systems. However, many countries which are not ICM members do not have a midwifery cadre, mitigating some of this bias. Additional research is also needed to demonstrate a causal relationship between regulatory environments
and midwifery workforce density to maximize policymaker interest in regulatory investments.
Our findings only state an association, not causation, and these associations may be influenced
by other factors in the health system or country governance, such as other investments in
maternal and newborn health (e.g., abolition of user fees for MNH services).

344 **Conclusion**

345 Stronger midwifery regulatory environments, characterized by regulatory policies and legislation, education and qualification standards, licensure, registration/re-licensure processes, 346 and well-defined scopes and conduct of practice, are associated with higher densities of 347 midwives globally. This likely results from the positive impact of regulation on the profession's 348 349 ability to attract, develop, and retain midwives due to enhanced professional identity, 350 autonomy, and credibility. These findings underscore the importance of investing in midwifery regulation as a critical component for ensuring sufficient midwifery workforce and, 351 consequently, improved maternal, newborn, and child health outcomes. However, additional 352 research is needed to better understand how specific regulatory factors influence the pipeline 353 of midwifery students, as well as the recruitment and retention of midwives, to establish 354 355 causality and identify the role individual domains play in driving density.

356 **References:**

357 358 359 360	1.	United Nations Population Fund, World Health Organization, International Confederation of Midwives. The state of the world's midwifery 2021. 2021 [cited 6 March 24]. Available from: <u>https://www.unfpa.org/sowmy-webappendices</u> .
361 362	2.	World Health Organization. The global strategy for women's, children's, and adolescents' health (2016-2030). 2018.
363		https://iris.who.int/bitstream/handle/10665/276423/A71 19Rev1-en.pdf?sequence=1
364		
365	3.	Nove A, Friberg IK, de Bernis L, McConville, F, Moran, A Najjemba M, et al. Potential
366		impact of midwives in preventing and reducing maternal and neonatal mortality and
367		stillbirths: a Lives Saved Tool modeling study. Lancet Glob Health. 2021;9(1):e24-e32.
368	_	
369	4.	Homer, CS. Models of maternity care: evidence for midwifery continuity of care. Med J
370		Aust. 2016;205(8):370-374.
371	_	
372	5.	Sandall J, Soltani H, Gates S, Shennan A, Devane D. Midwife-led continuity models
3/3		versus other models of care for childbearing women. Cochrane Database Syst Rev.
374		2016(4): CD004667.
3/5	c	World Health Organization Clobal Health Observatory, 142020 SP 1.2, density of
370	0.	neuronal neuronal and midwives per 10,000 penulation, 2024 [sited 20 March 24]
3// 270		Available from: https://www.wbo.int/data/gbo/indicator.motadata.rogistry/imr
270		dotails/7766
380		
380	7	Renfrew MI McEadden & Bastos MH Campbell I Channon AA Cheung NE et al
382	7.	Midwifery and quality care: findings from a new evidence-informed framework for
383		maternal and newborn care Lancet 2014: 384(9948) 1129–1145
384		
385	8.	Filby A. McConville F. Portela A. What prevents quality midwifery care? A systematic
386		mapping of barriers in low and middle income countries from the provider
387		perspective. <i>PloS one</i> , 2016; <i>11</i> (5), e0153391.
388		
389	9.	Nkowane AM, Wheeler E. Partnerships and collaboration: A means to effective
390		regulation and practice for nursing and midwifery professions in the African region.
391		Africa J Midwifery. 2014;8(2):3-3.
392		
393	10.	WHO. Transforming and scaling up health professionals' education and training: World
394		Health Organization guidelines 2013. 2013 [cited 31 May 24]. Available from:
395		https://www.who.int/publications/i/item/transforming-and-scaling-up-health-
396		professionals%E2%80%99-education-and-training.
397		

398 399 400	 International Confederation of Midwives. Global standards for midwifery regulation. 2011 [cited 31 May 24]. Available from: https://internationalmidwives.org/resources/global-standards-for-midwifery-
400	regulation/.
402 403	12. Kennedy HP, Myers-Ciecko JA, Carr KC, Breedlove G, Bailey T, Farrell MV, et al. United
404 405	States model midwifery legislation and regulation: development of a consensus document. J Midwifery Womens Health. 2018;63(6):652-659.
406 407 408	13. Jefferson K, Bouchard ME, Summers L. The regulation of professional midwifery in the United States. J Nurs Regul. 2021;11(4):26-38.
409	
410 411 412 413 414 415 416	14. Kreutzberg A, Reichebner, C, Maier, CB, Destrebecq, F & Panteli, D. Regulating the input: health professions. In: Busse R, Klazinga N, Panteli D, et al., editors. Improving healthcare quality in Europe: characteristics, effectiveness and implementation of different strategies (Health Policy Series, No. 53.). Copenhagen (Denmark): European Observatory on Health Systems and Policies; 2019 [cited 31 May 24]. <u>Available from: https://www.ncbi.nlm.nih.gov/books/NBK549267/</u>
410 417 418 419 420	 Ranchoff BL, Declercq ER. The scope of midwifery practice regulations and the availability of the certified nurse-midwifery and certified midwifery workforce, 2012- 2016.J Midwifery Womens Health.2020;65(1):119-130.
421 422 423 424	 Hoehn-Velasco L, Jolles D, Silverio-Murillo A, Plemmons A. Health outcomes and provider choice under independent practice for certified nurse-midwives. SSRN Electronic Journal. 2022:3878127.
425 426 427	17. Markowitz S, Adams EK, Lewitt MJ, Dunlop AL. Competitive effects of scope of practice restrictions: public health or public harm? J Health Econ. 2017;55:201-218.
428 429 430 431	 Vanderlaan J. Midwifery workforce density moderates the association between independent practice and pregnancy outcomes. J Midwifery Womens Health. 2023;68(5):588-595.
432 433 434 435	 Vedam S, Stoll K, MacDorman M, Declercq E, Cramer R, Cheyney M, Fisher T, Butt E, Yang YT, & Powell Kennedy H (2018). Mapping integration of midwives across the United States: Impact on access, equity, and outcomes. PloS one. 2018;13(2), e0192523–e0192523.
436 437 438 439	20. Kimani RW, Gatimu SM. Nursing and midwifery education, regulation and workforce in Kenya: A scoping review. Int Nurs Rev. 2023;70(3):444-455. doi:10.1111/inr.12840
440 441	21. Adegoke AA, Atiyaye FB, Abubakar AS, Auta A & Aboda A. Job satisfaction and retention of midwives in rural Nigeria. Midwifery. 2015; <i>31</i> (10), 946–956.

442	
443	22. United Nations Population Fund, World Health Organization, International
444	Confederation of Midwives. The state of the world's midwifery 2021: webappendices.
445	2021 [Cited 6 March 24]. Available from: https://www.unfpa.org/sowmy-
446	webappendices.
447	
448	23. International Confederation of Midwives, Global midwives' associations mapping
449	survey, 2019 [cited 5 March 24]. Available from:
450	https://experience.arcgis.com/experience/c60a4fch935a47fabcac76cdf025583h
451	
451 //52	24 World Health Organization, National Health Workforce Account Data Platform: 2024
452	[cited 5 June 2024] Available from: https://apps.who.int/phwaportal/
455	
434 AEE	25 World Health Organization, National health workforce accounts: a handbook, 2017
455	23. World Health Organization. National field workforce accounts. a fiandbook. 2017.
450	Geneva. Wond Health Organization.
457	26 United Nations Development Dreament Lunger Development Index 2022 [sited 9 March
458	26. United Nations Development Program. Human Development index. <u>2022 [cited 8 March</u>]
459	24]. Available from: https://hdr.undp.org/data-center/human-development-
460	<u>index#/indicies/HDI</u> .
461	
462	27. World Health Organization. Global health expenditure database. 2024 [cited 7 March
463	24]. Available from: <u>https://apps.who.int/nha/database</u> .
464	
465	28. The World Bank. Gini Index Data. 2024 [cited 8 March 24]. <u>Available from:</u>
466	https://data.worldbank.org/indicator/SI.POV.GINI.
467	
468	29. The World Bank. Country Income Group. 2024 [cited 8 March 24]. Available from:
469	https://datahelpdesk.worldbank.org/knowledgebase/topics/19280-country-
470	<u>classification</u> .
471	
472	30. World Health Organization. Midwives' voices, midwives' realities: find-ings from a global
473	consultation on providing quality midwifery care. October 2016 [cited 20 March 24].
474	http://www.who.int/maternal_child_adolescent/documents/midwives-voices-
475	<u>realities/en/</u> .
476	
477	31. Mattison C, Bourret K, Hebert E, Leshabari S, Kabeya A, Achiga P, Robinson J, Darling E.
478	Health systems factors impacting the integration of midwifery: an evidence-informed
479	framework on strengthening midwifery associations. BMJ Glob Health. 2021: 6(6).
480	e004850.
481	
482	32. Soewondo P. Puiisubekti R. Prastvani AW. Rahmavanti NM. Interprofessional
483	collaborative practice and health workers retention at remote primary health care: case
484	study from nusantara sebat team-based program. Kesmas (Denok), 2022-17(2)
485	
.00	

486 487	33. Belaid L, Dagenais C, Moha M, Ridde V. Understanding the factors affecting the attraction and retention of health professionals in rural and remote areas: a mixed-
188	method study in Niger, Hum Resour Health, 2017;15(1), 60–60
489	
490	34. Okello DRO. Gilson L. Exploring the influence of trust relationships on motivation in the
491	health sector: a systematic review. Hum Resour Health, 2015;13(1), 16–16.
492	
493	35. West M, Bailey S, Williams E. The courage of compassion: supporting nurses and
494	midwives to deliver high quality care. The King's Fund. Sep. 2020 [cited 8 Mar 24].
495	Available at: https://www.kingsfund.org.uk/publications/courage-compassion-
496	supporting-nursesmidwives
497	
498	36. Fenwick J, Sidebotham M, Gamble J, Creedy DK. The emotional and professional
499	wellbeing of Australian midwives: A comparison between those providing continuity of
500	midwifery care and those not providing continuity. Women Birth. 2018;31(1):38-43.
501	
502	37. Kubota S, Ando M, Murray J, et al. A regulatory gap analysis of midwifery to deliver
503	essential reproductive, maternal, newborn, child and adolescent health services in Lao
504	People's Democratic Republic. Lancet Reg Health West Pac. 2023;43:100960. Published
505	2023 Dec 5.
506	
507	38. Saralegui-Gainza A, Soto-Ruiz N, Escalada-Hernández P, Arregui-Azagra A, García-Vivar C,
508	San Martín-Rodríguez L. Density of nurses and midwives in sub-Saharan Africa: Trends
509	analysis over the period 2004–2016. J Nurs Manag. 2022;30(8), 3922–3932.
510	