

Prevalence and factors associated with incompleteness of measles and rubella vaccine routine immunization among children aged 18-59 months in Kilimanjaro region, Tanzania

David M. Gichogo¹, Abdul A. Mbezi¹, Sasi K. Sasi¹, Vailet Magogo¹, Edwin J. Shewiyo^{2,3}
William Nkenguye^{2,3}, Florian Tinuga⁴, Sia E. Msuya^{1,3,5}

Affiliations

¹ Institute of Public Health, Department of Community Health, Kilimanjaro Christian Medical University College (KCMUCo), P.O. Box 2240 Moshi, Tanzania.

² Kilimanjaro Clinical Research Institute (KCRI), Box 2236, Moshi, Tanzania

³ Institute of Public Health, Department of Epidemiology and Biostatistics, Kilimanjaro Christian Medical University College, P.O. Box 2240 Moshi, Tanzania.

⁴ Immunization and Vaccine Development Program, Ministry of Health, Dodoma, Tanzania.

⁵ Department of Community Health, Kilimanjaro Christian Medical Centre (KCMC), P.O. Box 3010, Moshi, Tanzania.

ABSTRACT

Background: In Tanzania Measles and Rubella (MR) vaccination coverage of 90% in 90% of the regions was achieved in 2018. However, there are councils lagging in MR1 coverage, 38 (19%) councils out of 195 had <90% coverage by the end of 2019 while MR2 coverage is struggling with 99 (51%) of councils having a coverage of <90% at the end of 2019. Kilimanjaro region is among the regions with some councils having MR coverage of <80%. There is a need for information on why a significant proportion of eligible children are not receiving the second dose of MR vaccine.

Objective: To determine prevalence and factors associated with incompleteness of MR vaccination among children aged 18-59 months in two districts of Kilimanjaro region.

Methodology: This study was a cross-sectional study conducted in Moshi urban and rural councils, two out of the seven councils in Kilimanjaro region. The study population was children aged 18-59 months whose parents/ caregivers have been the residents of the respective districts for at least 3 past years. Interviews using questionnaires were used for data collection. Data was entered and analyzed using SPSS version 20.

Results: A total of 415 children aged 18-59 months were enrolled. The proportion of children with incomplete MR2 vaccination was 14.2%. Only 33% of the 415 caregivers knew MR vaccine is given at 9 and 18 months and 24% reported unavailability of vaccine at their scheduled visit. Children from Moshi rural council (AOR=2.53, 95% CI =1.36-4.73) had higher odds of MR incompleteness. Lower odds of MR vaccine incompleteness were among caregivers who were aware on the time for MR vaccination; 9 and 18 months (AOR=0.27, 95% CI=0.10-0.73), caregivers who had knowledge that MR vaccine protects measles, and those who had health promotion talk on the vaccination day.

Conclusion: Nearly one in ten (14.2%) children do not complete the recommended two doses of MR vaccines. Strategies to improve awareness and knowledge on timing, frequency and advantage of MR and other vaccines is needed among parents/ caregivers in this setting. Further, improvement in ordering and supply of vaccines at health facilities needs to be improved.

Keywords: Measles, Rubella, MR vaccine, Vaccination, Children, Prevalence, Tanzania

BACKGROUND

Measles and rubella, both viral infections, are of significant public health concern due to their potential for widespread transmission and severe health consequences, particularly among vulnerable populations such as young children and pregnant women. Globally, measles remains a leading cause of vaccine-preventable deaths among children, with over 140,000 deaths reported in 2018 alone(1). Sub-Saharan Africa, in particular, bears a disproportionate share of the measles burden, with an estimated 155,000 measles-related deaths occurring in the region in 2018 alone(1).

The introduction of measles-rubella (MR) vaccines has played a crucial role in reducing the burden of these diseases. Through widespread vaccination efforts, substantial declines in measles-related deaths have been observed globally. For instance, between 2000 and 2018, measles vaccination prevented an estimated 23.2 million deaths worldwide, with most of the deaths averted in the African region(2). However, despite these successes, outbreaks of measles persist, particularly in regions with inadequate vaccination coverage.

Tanzania, like many other countries in sub-Saharan Africa, has reported measles outbreaks, despite efforts to control the disease through vaccination. In Tanzania, MR vaccination is administered to children under five years in two doses, with the first dose given at 9 months and the second dose at 18 months, according to the Tanzanian routine vaccination schedule(3). Tanzania has observed an increase in number of measles cases from 2019, with about 12,253 cases reported between 2019 – 2022(4). Children aged 1-4 years had the highest proportion of laboratory confirmed cases compared to other groups. Michael et al (2024) also showed that, MR unvaccinated children had 2.6 higher odds of getting measles outbreak compared to fully vaccinated children. Currently Tanzania is number 9 leading country with measles outbreak(5).

Suboptimal measles vaccination coverage is the main cause of the reported outbreaks by increasing the pool of susceptible children over time. The 2019 data from the Integrated Vaccination Data (IVD) program have shown that despite the availability of the vaccine, there is an incomplete uptake of MR vaccination, with the second dose facing particular challenges. Tanzania MR coverage of 90% in 90% of the regions was achieved in 2018 according to DHIS-2 data. By the

end of 2019, coverage of MR1 was high (> 95%), but 19% (38 out of 195) of councils had coverage of < 90% (6). There are many councils lagging when it comes to the second dose of MR (MR2), with 51% (99 out of 195) of councils having a coverage of <90% at the end of 2019 representing a significant gap in vaccination coverage(4). Studies from Dar es Salaam and Mtwara have shown 17% and 55% of children under five had not received the second dose of MR vaccine(7,8).

In Asia studies have shown vaccine availability, negative perception on vaccines and poor knowledge on vaccination schedule to be factors associated with incompleteness of MR vaccine(9,10). Literature from East Africa reported both health system and caregivers' characters to influence completion of MR vaccination. Awareness of MR vaccine among caregivers, vaccine hesitancy driven by misinformation or cultural beliefs, myths, negative attitudes, and poverty contribute to the incompleteness of the MR vaccination schedule(11). Logistical barriers, such as limited access to vaccination services in remote or rural areas, and distance to facilities also influenced completion of MR vaccine(4,12).

Against this backdrop, this study aims to investigate the prevalence and factors associated with the incompleteness of MR vaccine routine immunization among children aged 18-59 months in Kilimanjaro region, Tanzania. The region is among those with MR2 vaccine coverage of < 90% in several councils. By identifying barriers to vaccination uptake and elucidating the context-specific challenges faced by communities, this research seeks to inform evidence-based strategies for enhancing vaccination coverage and advancing progress towards Sustainable Development Goal 3.8, which focuses on achieving universal health coverage, including access to essential vaccines, for all individuals by 2030.

Specific Objectives:

1. To determine the prevalence of MR2 vaccine incompleteness among children aged 18-59 months.
2. To identify the factors associated with the incompleteness of MR2 vaccination.
3. To assess caregivers' awareness and knowledge regarding MR vaccination.

METHODS

Study Design

This was a community based cross-sectional study conducted from June to August 2022.

Setting

The study was conducted in two out of seven councils of Kilimanjaro region, namely Moshi municipal council and Moshi district council. The region is situated in northern Tanzania with approximate population of 1,861,934 and 224,780 children U5. Moshi municipal council has a total population of 221,733 people and 25,280 children U5, while Moshi DC has a population of 535,803 people and 61,762 children aged U5 years (Census, 2022). The two districts were selected because they had low coverage where MR2 immunization coverage of <80% according to the Immunization Coverage update for performance of routine immunization of January to August 2021 in Tanzania. Main economic activities done in the areas are tourism, agriculture and manufacturing activities.

Study Population

The study population was children aged 18-59 months. Parents especially mothers and/ or caretaker of the child-aged 18-59 months provided the information. The study included children whose parents or caretakers were the residents of the two selected councils. Residency in this study was defined as having an address in the selected council and having lived in the area for the past three years prior to the study.

Sample size and sampling

Multi-staged sampling technique was used to access the study participants. From the list of 53 wards, 21 wards in Moshi municipal and 32 wards from Moshi DC respectively, a sample of 6 wards 3 wards from each district with their respective population of children under 5 years were randomly selected. Random sampling was employed for each selected ward. The sample size for each selected ward was obtained.

From each selected ward, all streets within the ward were listed and 7 streets were drawn randomly from Moshi municipal and 2 villages from Moshi DC from the selected wards. The households were randomly selected and from the households, a caregiver or parent was interviewed. In situations where there was more than one child with 18-59 months of age

all were interviewed.

Data Collection

Face-to-face interviews with a questionnaire was used at household level to collect information from parents and caretakers of the children aged 18-59 months. The questionnaire was uploaded into Kobo toolbox, an online data kit. Prior to the interview, informed consent was sought from the study participants then the interview proceeded among the consented caregivers. Interviews lasted between 15 and 20 minutes depending on the respondents' ability to respond to posed questions.

Study variables

Dependent variable

Dependent variable of this study was incompleteness of MR2 vaccination according to the immunization card and by report speech.

Independent variables

The independent variables of this study included; caretakers' individual factors (such as fear of adverse events following vaccination, awareness on the need for MR second dose, attending antenatal and postnatal care), health system factors (such as absenteeism of health worker at the health facility, vaccine availability, existence of outreach services, opening days, time of vaccination session, long waiting time), caretaker' socio-demographic factors (such as occupational status, education level, economic status, marital status and family size), child factors (such as age, sex, child birth rank, orphan hood) and community factors (such as travelling distance to health facility, means of transport and geographical accessibility to health facility).

Data Analysis

The collected questionnaires from Kobo toolbox were checked manually for its completeness, and after reviewing them the data was transferred and analyzed using SPSS version 20. Data was cleaned by running the frequency of each variable. Categorical variables were summarized into frequency and percent tables. Numeric variables were summarized using mean/ median and their respective measures of dispersion. Bivariate and multivariable analyses were conducted to identify independent factors associated with the incompleteness of MR2 vaccine. Crude Odds Ratios (COR) and Adjusted Odds Ratios (AOR) and 95% Confidence Intervals (CI) were calculated. Variables with p-value of < 10% in bivariate analysis were entered into multivariable logistic regression models.

RESULTS

A total of 425 children aged 18-59 months were approached. A total of 415 (97.6%) children were enrolled, 10 children were not included because the caregivers were not around at the time of interview.

Background characteristics of the participants

The median age of the 415 children was 35 months, (IQR 24-46) and 52% were males. Most of the informants were mothers (86.0%) with a median age of 28 years (IQR 25-34). About half (50.4%) of the informants had secondary education or higher and most of them were married (65.5%). The common occupational activity of informants was informal employment (47.7%) where the majority earned less than 50,000 Tanzanian shillings per month (56.6%), (**Table 1**).

Table 1: Social demographic characteristics of children aged 18-59 months and caregivers in Moshi urban and rural districts, Kilimanjaro region (N=415)

Variable	Frequency	Percentage
District of enrollment		
Moshi urban	226	54.5
Moshi rural	189	45.5
Age of the child(months)		
Median (IQR)	35(24-46)	
18-24	107	25.8
25-36	128	30.8
37-48	104	25.1
49-59	76	18.3
Sex of the child		
Male	214	51.6
Female	201	48.4
Informant		

Mother	357	86.0
Father	11	2.7
Other relatives	47	11.3
Informant's age		
Median (IQR)	28(25-34)	
18-24	86	20.7
25-34	232	55.9
35-44	50	12.1
45+	47	11.3
Marital status of the informant		
Not married	143	34.5
Married	272	65.5
Educational level of the informant		
Primary level and below	206	49.6
Secondary level and above	209	50.4
Occupation of the informant		
Non employed	143	34.5
Informally employed	198	47.7
Formally employed	43	10.4
Others	31	7.1
Average income of the informant (TZS)		
≤50000	235	56.6
>50000	180	43.4
Number of people in the household total		
≤5	330	79.5
>5	85	20.5
Number of children in the household		
≤4	401	96.6
>4	14	3.4
Distance to health facility		
<2km	340	81.9
>2km	75	18.1
Obstacles to reach the health facility		
Yes	38	9.2
No	377	90.8

Health facility characteristics and Caregivers' awareness and knowledge on MR vaccination

Majority (82%) of the 415 parents/ caregivers reported to live less than 2 kilometers to a health facility that provides vaccination services. About 1.9% of 415 parents/caregivers of the enrolled children reported that they did not find the health worker at the vaccination post on the scheduled date of vaccination and about quarter (24.1%) reported they attended for vaccination, but the vaccines were out of stock, (**Table 2**).

Nearly one in ten (11.6%) of the 415 parents/ caregivers had no awareness of the MR vaccine. Of 367 who were aware, many of the caregivers (58.9%) did not know the time of MR

vaccination, and only 33% knew MR vaccine is given twice, at 9 and 18 months as per vaccination schedule in Tanzania (Table 2). Few (37.9%) were aware of the importance of MR2 vaccine, (Table 2).

Prevalence of MR vaccine incompleteness

A total of 59 children out of 415 (14.2%) had not completed MR2 at the time of the study. Out of 415 children enrolled in the study, caregivers of 216 children had vaccination cards, 21 children (9.7%) were not vaccinated with MR2. (Table 2).

Table 2: Background characteristics of health facilities and awareness, knowledge of caregivers on MR vaccination in Moshi urban and rural districts, Kilimanjaro region (N=415)

Variable	Frequency	Percentage
Awareness of MR vaccine		
Yes	367	88.4
No	48	11.6
Knowledge on time of MR vaccination (n=367)		
9 months	31	8.4
9 and 18 months	120	32.7
I don't know	216	58.9
Awareness of importance of MR2 vaccine (n=367)		
Yes	139	37.9
No	228	62.1
Measles as a vaccine protected disease		
Yes	366	88.2
No	49	11.8
Vaccination status by card (n=216)		
MR2		
Yes	195	90.3
No	21	9.7
Vaccination status by recall (did not have card, n=199)		
MR2		
Yes	161	80.9
No	38	19.1
MR2 status by card and recall (N=415)		
Yes	356	85.8
No	59	14.2
Absenteeism of health worker on vaccination day		
No	407	98.1
Yes	8	1.9
Unavailability of the vaccine on vaccination day		
Yes	100	24.1
No	315	75.9
Promotion talk on vaccines when attending for PNC		
Yes	168	40.5
No	247	59.5

PNC: Postnatal care

Factors associated with MR2 incompleteness

Children enrolled in Moshi district council were three times more likely to be unvaccinated for MR2 vaccine compared to children enrolled in Moshi urban (COR=3.20, 95% CI=1.77-5.79). Likewise male children were more likely to be unvaccinated compared to female children (COR=2.02 95% CI=1.13-3.59), (Table 3).

Table 3: Bivariate analysis of socio-demographic characteristics associated with incompleteness of MR2 vaccine in Kilimanjaro region

Variable	N	Unvaccinated with MR2, n (%)	COR (95% CI)	P-Value
District of enrollment				
Moshi urban	226	18 (7.9)	1	
Moshi rural	189	41 (21.7)	3.20 (1.77-5.79)	<0.001
Education level of the caregiver				
None/Primary level	206	29 (14.1)	1	
Secondary level and above	209	30 (14.3)	1.02 (0.59-1.77)	0.936
Occupation level of the caregiver				
Non employed	143	22 (22.4)	1	
Employed	270	37 (13.7)	0.86 (0.49-1.53)	0.620
Economic status of the caregiver*				
≤50000	229	34 (14.8)	1	
>50000	180	25 (13.9)	1.92 (0.53-1.62)	0.784
Marital status of the caregiver				
Not married	143	25 (17.9)	1	
Married	272	34 (12.5)	0.67 (0.38-1.18)	0.169
Total number of people in the household				
≤5	330	42 (12.7)	1	
>5	85	17 (20.0)	1.71 (0.92-3.19)	0.090
Number of children in the household				
≤4	401	56 (14.0)	1	
>4	14	3 (21.4)	1.68 (0.45-6.21)	0.437
Age of the child (months)				
18-24	107	17 (15.9)	1.12 (0.49-2.54)	0.793
25-36	128	13 (10.2)	0.67 (0.28-1.58)	0.357
37-48	104	18 (17.3)	1.24 (0.55-2.80)	0.610
49-59	76	11 (14.5)	1	
Sex of the child				
Male	214	39 (18.2)	2.02 (1.13-3.59)	0.017
Female	201	20 (10.0)	1	
Birth rank of the child				
1-3	390	56 (14.4)	1.23 (0.35-4.24)	0.744
4-6	25	3 (12.0)	1	
Caregiver's relationship with the child				
Mother	357	47 (13.2)	1	
Father	11	2 (18.2)	1.47 (0.31-6.99)	0.632
Other relatives	47	10 (21.3)	1.78 (0.83-3.82)	0.138

*Amount in Tanzanian Shilling

Caregivers who did not receive counseling or health promotion talks postnatal care had less odds of completing MR2 vaccination than others. Caregivers who were not aware on importance and schedule of MR vaccination had 4 times higher odds of having a child with incomplete MR vaccination than others (COR=4.38, 95% CI=1.91-10.05), (Table 4).

Table 4: Bivariate analysis of the health facility, caregivers' and community factors associated with the incompleteness of MR2 vaccine

Variable	N	Unvaccinated with MR2, n (%)	COR (95% CI)	P-Value
Awareness of MR vaccine				
Yes	367	50 (13.6)	0.68 (0.31-1.50)	0.341
No	48	9 (18.8)	1	
Knowledge on time of MR vaccination (n=367)				
9 months	31	13 (41.9)	1	
9 and 18 months	120	13 (10.8)	0.17 (0.07-0.42)	<0.001
I don't know	216	24 (11.1)	0.17 (0.08-0.40)	<0.001
Awareness of importance of MR2 vaccine (n=367)				
Yes	139	7 (5.0)	1	
No	228	43 (18.8)	4.38 (1.91-10.05)	<0.001
Measles as a vaccine protected disease				
Yes	366	47 (12.8)	1	
No	49	12 (24.5)	2.20 (1.07-4.52)	0.032
Distance to nearest health facility				
<2km	340	43 (12.6)	1	
>2km	75	16 (21.3)	1.87 (0.99-3.55)	0.054
Obstacles to reach the nearest health facility				
Yes	38	7 (18.4)	1	
No	377	52 (13.8)	0.71 (0.30-1.69)	0.438
Absenteeism of health worker on vaccination day				
No	407	57 (14.0)	1	
Yes	8	2 (25.0)	2.05 (0.40-10.39)	0.388
Unavailability of the vaccine on vaccination day				
Yes	100	20 (20.0)	1.77 (0.98-3.20)	0.060
No	315	39 (12.4)	1	
Promotion talk on vaccine when attending for PNC				
Yes	168	15 (8.9)	1	
No	247	44 (17.8)	2.21 (1.18-4.41)	0.012

Multivariable analysis of the factors associated with the incompleteness of MR2 vaccine.

In multivariable analysis, children enrolled in Moshi rural were two times more likely to be unvaccinated with MR2 vaccine compared to those enrolled in Moshi urban (AOR=2.53, 95% CI =1.36-4.73). Likewise, awareness on the time for MR vaccination, children with caregivers who were aware that ages for MR vaccination are 9 and 18 months and those who were not aware were less likely to be unvaccinated with MR2 vaccine compared to those who were aware of only 9 months as the age for MR vaccination (AOR=0.27, 95% CI=0.10-0.73), (AOR=0.17, 95% CI=0.07-0.44) respectively (Table 5).

Table 5: Multivariable logistic regression on factors associated with the incompleteness of MR2 vaccine

Variable	Adjusted OR (95% CI)	P-Value
District of enrollment		
Moshi municipal	1	
Moshi DC	2.53 (1.36- 4.73)	0.004
Sex of the child		
Male	1	
Female	0.51 (0.27-0.97)	0.040
Measles as a vaccine protected disease		
Yes	1	
No	1.56 (0.61-3.96)	0.350
Knowledge on time of MR vaccination (n=415)		
9 months	1	
9 and 18 months	0.27 (0.10-0.73)	0.010
I don't know	0.17(0.07-0.44)	0.001
Others	0.25(0.09-0.73)	0.010
Awareness of the importance of MR2 vaccine (n=367)		
Yes	1	
No	2.89 (1.17-7.16)	0.022
Promotion talk on vaccine when attending PNC		
Yes		
No	1.24 (0.61-2.51)	0.541

Discussion

Our study found that 14.2% of children had not completed the MR2 vaccination, with an overall vaccination coverage of 85.8%. This rate is higher than the national average of 74% and exceeds the coverage reported in community studies conducted in southern rural Tanzania (44.2%) and urban Tanzania (83%)(8,13). The higher MR2 coverage observed in our study can be attributed to multiple immunization campaigns conducted across various districts in the Kilimanjaro region, which have enhanced access to vaccination services for children in the community.

However, significant disparities in vaccination coverage exist between rural and urban areas. Children in rural areas were found to have twice the odds of missing MR2 vaccination compared to their urban counterparts. This disparity is critical, considering that the majority of the children in the Kilimanjaro region reside in rural areas, thus highlighting significant gaps in immunization coverage. This finding is consistent with studies from other low-resource settings, where geographic and socio-economic barriers often impede access to vaccination services(14,15).

Our study also identified caregiver awareness and knowledge as significant factors influencing MR2 vaccination coverage. Caregivers who were aware of the importance of MR2 vaccination and knew the correct vaccination schedule were significantly less likely to have children who missed the MR2 dose. This aligns with previous research in Tanzania and other settings, which underscores the importance of parental education and awareness in improving vaccination uptake(16,17).The high level of unawareness observed in our study, with 62.1% of caregivers unaware of the importance of the MR2 vaccine and 58.9% unaware of the vaccination schedule, underscores the need for enhanced health education initiatives.

The role of postnatal care (PNC) visits in improving vaccination coverage is evident in our findings. However, about 10% of caregivers reported not receiving counseling or health promotion on vaccination during their PNC visits. This represents a missed opportunity to educate caregivers on the importance of complete immunization. Studies have shown that integrating immunization education into routine PNC services can significantly improve vaccine uptake(18,19).

In terms of health system factors, the availability of vaccines and healthcare worker presence at vaccination posts were crucial. Approximately 24.1% of caregivers reported vaccine stockouts during scheduled visits, which is a significant barrier to vaccination completion. This issue is echoed in other studies from low- and middle-income countries (LMICs), where logistical challenges in vaccine supply chains hinder immunization efforts(20,21).

Furthermore, our study's findings suggest that community-level interventions and improved healthcare infrastructure are necessary to address these gaps. Ensuring consistent vaccine availability and enhancing health worker training and motivation are essential steps toward achieving higher vaccination coverage. Additionally, targeted outreach programs in rural areas can help bridge the coverage gap between urban and rural populations.

The findings from our study align with the broader literature on vaccine coverage and highlight the multifaceted nature of immunization challenges. Studies have shown that comprehensive approaches, combining health education, improved healthcare delivery, and robust supply chain management, are essential for enhancing vaccination rates(22,23).

For example, a study conducted in Nigeria found that improved training for health workers and enhanced community outreach programs significantly increased vaccination rates in rural areas(14). Similarly, research in Ethiopia indicated that integrating health education programs into routine health services improved caregiver knowledge and vaccine uptake(24). In India, addressing logistical barriers by ensuring consistent vaccine supply and reducing travel distances to health facilities was shown to improve vaccination coverage(25).

In summary, our study highlights the significant gaps in MR2 vaccination coverage in the Kilimanjaro region, with a particular focus on rural areas. Enhancing caregiver awareness through health education, ensuring vaccine availability, and addressing logistical barriers are crucial for improving MR2 vaccination rates. These findings underscore the need for multifaceted interventions to achieve higher vaccination coverage and better health outcomes for children in Tanzania.

CONCLUSION

The study aimed at assessing the prevalence and factors associated with incompleteness of MR2 vaccine among children aged 18-59 months whereas a proportion of (14.2%) of children had not received MR2 dose. Knowledge on MR vaccine, awareness on the importance of MR2 dose, awareness of measles as a disease protected by vaccination and health promotion talks significantly contributed to the incompleteness of MR2 dose. Intervention measures thus should aim at improving community education at all levels to reduce the burden of unvaccinated children.

RECOMMENDATION

With regard to improving MR2 coverage in Kilimanjaro region, the following recommendations should be considered so as to attain the recommended coverage of >95% by 2025.

- More health promotion talks should be offered regarding postnatal vaccinations whereas caregivers should be educated on the importance of vaccines and the proper schedule for administration of all vaccines for their children. Also, these talks should be extended to the community level so as to reach out even those caregivers who find no importance of vaccinations.
- Health care workers should consider screening vaccination status of children who visit health facilities for other purposes out of vaccination and consider vaccinating those who have missed as per Tanzania immunization guideline.
- The government together with the Ministry of Health could also make efforts to make vaccination knowledge more available in as many different platforms as possible like mass media, social media, religious centers and many more so as to reach a lot of people with this knowledge.

Ethics statement: The ethical clearance was obtained from the Ethical Committee of the Kilimanjaro Christian Medical University College Research Ethics and Review Committee (CRERC) in Moshi (UG 53/2022). Permission was obtained from the President's Office,

Regional Administration, and Local Government (POLARG), from the regional administrative secretary and regional medical officer office. Purpose of the study, benefit, right to refuse participation or leaving the study was explained to each participant before the interview. The study participants were informed that the data will be kept confidential by using codes and not personal identifiers such as names and is meant only for the purpose of study. Verbal consent was regarded as sufficient to be used in the study.

Patient Consent for Publication: Not Applicable

Acknowledgement: We are grateful to all participants who participated in one way of this work and the data collection team

REFERENCES

1. CDCGlobal. Centers for Disease Control and Prevention. 2023 [cited 2023 Apr 12]. Fast Facts on Global Measles, Rubella, and Congenital Rubella Syndrome. Available from: <https://www.cdc.gov/globalhealth/measles/data/fast-facts-global-measles-rubella.html>
2. Patel MK, Dumolard L, Nedelec Y, Sodha SV, Steulet C, Gacic-Dobo M, et al. Progress Toward Regional Measles Elimination — Worldwide, 2000–2018. *Morb Mortal Wkly Rep* [Internet]. 2019 Dec 6 [cited 2024 Jun 13];68(48):1105–11. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6897527/>
3. Rabaan AA, Mutair AA, Alhumaid S, Garout M, Alsubki RA, Alshahrani FS, et al. Updates on Measles Incidence and Eradication: Emphasis on the Immunological Aspects of Measles Infection. *Medicina (Mex)* [Internet]. 2022 May 20 [cited 2024 Jun 13];58(5):680. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9147347/>
4. Michael F, Mirambo MM, Misinzo G, Minzi O, Beyanga M, Mujuni D, et al. Trends of measles in Tanzania: A 5-year review of case-based surveillance data, 2018-2022. *Int J Infect Dis* [Internet]. 2024 Feb [cited 2024 Jun 13];139:176–82. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10784152/>
5. Thomas JA, Paul IK, Jubilate A, Kavishe Z Joseph, Kazinza M, Abdi SSH, et al. Interventions and recommendations to mitigate measles outbreak in Tanzania. *Ann Med Surg* [Internet]. 2022 Nov 18 [cited 2024 Jun 13];84:104964. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9682328/>

6. Hayman DTS. Measles vaccination in an increasingly immunized and developed world. *Hum Vaccines Immunother* [Internet]. 2018 Sep 19 [cited 2024 Jun 13];15(1):28–33. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6363159/>
7. Sangeda RZ, James D, Mariki H, Mbwambo ME, Mwenesi ME, Nyaki H, et al. Childhood vaccination trends during 2019 to 2022 in Tanzania and the impact of the COVID-19 pandemic. *Hum Vaccines Immunother* [Internet]. [cited 2024 Jun 13];20(1):2356342. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC11123454/>
8. Vasudevan L, Baumgartner JN, Moses S, Ngadaya E, Mfinanga SG, Ostermann J. Parental concerns and uptake of childhood vaccines in rural Tanzania – a mixed methods study. *BMC Public Health* [Internet]. 2020 Oct 20 [cited 2024 Jun 13];20(1):1573. Available from: <https://doi.org/10.1186/s12889-020-09598-1>
9. Bari MdS, Hossain MdJ, Ahmmed F, Sarker MdMR, Khandokar L, Chaitly AP, et al. Knowledge, Perception, and Willingness towards Immunization among Bangladeshi Population during COVID-19 Vaccine Rolling Period. *Vaccines* [Internet]. 2021 Dec 7 [cited 2024 Jun 13];9(12):1449. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8706921/>
10. Adisu MA, Bogale WA, Alemu TG. Second dose of measles-containing vaccine coverage and associated factors among children aged 24–36 months in Gondar city, Central Gondar, Northwest Ethiopia, 2023. *Front Public Health* [Internet]. 2024 May 2 [cited 2024 Jun 13];12:1364865. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC11097900/>
11. Danso SE, Frimpong A, Seneadza NAH, Ofori MF. Knowledge, attitudes, and practices of caregivers on childhood immunization in Okaikoi sub-metro of Accra, Ghana. *Front Public Health* [Internet]. 2023 Sep 15 [cited 2024 Jun 13];11:1230492. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10540614/>
12. Addis M, Mekonnen W, Estifanos AS. Health system barriers to the first dose of measles immunization in Ethiopia: a qualitative study. *BMC Public Health* [Internet]. 2024 Mar 1 [cited 2024 Jun 13];24:665. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10908078/>
13. Magodi R, Mmbaga EJ, Massaga J, Lyimo D, Mphuru A, Abade A. Factors associated with non-uptake of measles-rubella vaccine second dose among children under five years in Mtwara district council, Tanzania, 2017. *Pan Afr Med J* [Internet]. 2019 May 29 [cited 2023 Apr 12];33:67. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6689852/>
14. Adedokun ST, Uthman OA, Adekanmbi VT, Wiysonge CS. Incomplete childhood immunization in Nigeria: a multilevel analysis of individual and contextual factors. *BMC Public Health* [Internet]. 2017 Mar 8 [cited 2024 Jun 13];17:236. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5341359/>
15. Newton P, Smith DM. Factors influencing uptake of measles, mumps and rubella (MMR) immunization in site-dwelling Gypsy, Roma and Traveller (G&T) communities: a qualitative

- study of G&T parents' beliefs and experiences. *Child Care Health Dev.* 2017 Jul;43(4):504–10.
16. Perceptions and experiences of labour companionship: a qualitative evidence synthesis - PMC [Internet]. [cited 2024 Jun 13]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6422112/>
 17. Oku A, Oyo-Ita A, Glenton C, Fretheim A, Ames H, Muloliwa A, et al. Perceptions and experiences of childhood vaccination communication strategies among caregivers and health workers in Nigeria: A qualitative study. *PLoS ONE* [Internet]. 2017 [cited 2024 Jun 13];12(11). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5678719/>
 18. Ibnouf A, Van Den Borne H, Maarse J. Factors influencing immunisation coverage among children under five years of age in Khartoum State, Sudan. *South Afr Fam Pract* [Internet]. 2007 Sep [cited 2024 Jun 13];49(8):14–14f. Available from: <https://www.tandfonline.com/doi/full/10.1080/20786204.2007.10873611>
 19. Brown KF, Kroll JS, Hudson MJ, Ramsay M, Green J, Long SJ, et al. Factors underlying parental decisions about combination childhood vaccinations including MMR: A systematic review. *Vaccine* [Internet]. 2010 Jun 11 [cited 2024 Jun 13];28(26):4235–48. Available from: <https://www.sciencedirect.com/science/article/pii/S0264410X10005761>
 20. Cataldi JR, Kerns ME, O'Leary ST. Evidence-based strategies to increase vaccination uptake: a review. *Curr Opin Pediatr* [Internet]. 2020 Feb [cited 2024 Jun 13];32(1):151. Available from: https://journals.lww.com/co-pediatrics/abstract/2020/02000/evidence_based_strategies_to_increase_vaccination.21.aspx
 21. Dolan SB, Burstein R, Shearer JC, Bulula N, Lyons H, Carnahan E, et al. Changes in on-time vaccination following the introduction of an electronic immunization registry, Tanzania 2016-2018: interrupted time-series analysis. *BMC Health Serv Res* [Internet]. 2022 Sep 20 [cited 2024 Jun 13];22:1175. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9485799/>
 22. Why children are not vaccinated: a review of the grey literature | International Health | Oxford Academic [Internet]. [cited 2024 Jun 13]. Available from: <https://academic.oup.com/inthealth/article/4/4/229/672421?login=false>
 23. Arsenault C, Johri M, Nandi A, Mendoza Rodríguez JM, Hansen PM, Harper S. Country-level predictors of vaccination coverage and inequalities in Gavi-supported countries. *Vaccine* [Internet]. 2017 Apr 25 [cited 2024 Jun 13];35(18):2479–88. Available from: <https://www.sciencedirect.com/science/article/pii/S0264410X17303353>
 24. Factors associated with incomplete childhood immunization in Arbegona district, southern Ethiopia: a case – control study | BMC Public Health | Full Text [Internet]. [cited 2024 Jun 13]. Available from: <https://bmcpublichealth.biomedcentral.com/articles/10.1186/s12889-015-2678-1>

25. Mathew JL. Inequity in childhood immunization in India: a systematic review. *Indian Pediatr.* 2012 Mar;49(3):203–23.