

1 Sexual and Reproductive Health Needs of Women with Severe
2 Mental Illness in Low- and Middle-Income Countries: A Scoping
3 Review

4 Short Title: Sexual Health Needs of Women with Severe Mental Illness in Low- and
5 Middle-Income Countries

6

7 Shilpa Sisodia^{1¶}, Zara Hammond^{2¶}, Jo Leonardi-Bee^{1,3}, Charlotte Hanlon^{1,5}, Laura Asher^{1,6}

8

9 ¹ Nottingham Centre for Public Health and Epidemiology, School of Medicine, University of
10 Nottingham, Nottingham, United Kingdom

11 ² Placed with NHS England Midlands Regional Public Health Directorate, Nottingham, United
12 Kingdom

13 ³ Nottingham Centre for Evidence Based Healthcare, School of Medicine, University of Nottingham,
14 Nottingham, United Kingdom

15 ⁴ College of Health Sciences, School of Medicine, Department of Psychiatry, Addis Ababa University,
16 Addis Ababa, Ethiopia

17 ⁵ Centre for Global Mental Health, Health Service and Population Research Department, King's
18 College London, Institute of Psychiatry, Psychology and Neuroscience, London, UK

19 ⁶ Institute of Mental Health, University of Nottingham

20

21 ¶ These authors contributed equally to this work.

22

23 *Corresponding author:

24 Laura.Asher@nottingham.ac.uk (L.Asher)

25 Abstract

26 Background

27 This scoping review aimed to understand the extent and type of evidence in relation to sexual and
28 reproductive health needs of women with severe mental illness (SMI) in low- and middle-income
29 countries (LMIC) and to summarise those needs.

30 Methods

31 Inclusion criteria were 1) focus on sexual and reproductive health needs 2) women or girls with SMI,
32 professionals, caregivers of women with SMI and community members 3) study set in a LMIC 4) peer
33 reviewed literature (no restriction on study date or design). Studies were identified from
34 comprehensive searches of Medline, EMBASE, CINAHL and PsycINFO (to July 2023).

35 Results

36 The review included 100 papers. Most studies were cross-sectional and set in hospital outpatient
37 departments. We identified studies in only 19 of 140 LMIC countries and only 15 studies were set-in
38 low-income countries. Included studies often had multiple focus areas which were grouped into
39 themes of HIV (prevalence, risk behaviour and knowledge), other sexually transmitted infections
40 (STIs), sexual function, contraception use and family planning, sexual violence, fertility, pregnancy
41 and postpartum.

42 Included studies indicated women with SMI have worse outcomes and worse sexual and
43 reproductive health compared to both women without SMI and men with SMI. Women with SMI
44 were shown to have higher rates of HIV and low levels of contraception knowledge and use, with
45 little advice offered by professionals.

46 Conclusions

47 This review highlights the need for a greater diversity of study methodology, robustness of ethical
48 and consensual reporting when researching vulnerable populations and for further research on

49 interventions and models of care aimed at addressing stigma, discrimination and improving the
50 sexual and reproductive health of women with SMI. Future research should better represent the
51 breadth of LMIC, investigate cultural adaptability of interventions and consider sexual health needs
52 across the life course.

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67 Introduction

68 Good sexual and reproductive health is fundamental to a person's 'physical, mental, emotional and
69 social wellbeing' [1][2][3][4][5][84]. The United Nations Convention on the Rights of Persons with
70 Disabilities (CRPD) sets out that access to high quality, non-discriminatory preventative and
71 treatment services and the rights of individuals to make their own choices about their reproductive
72 and sexual health are human rights, which should not be compromised by disability due to mental,
73 physical, intellectual or sensory impairment [6].

74 Good sexual and reproductive health describes individuals achieving their full potential for
75 pleasurable and safe sexual experiences across the life course through being empowered to
76 maintain their own sexual and reproductive health [1][7]. People experiencing poor sexual and
77 reproductive health will have a greater need for health care to support them in reaching their
78 optimal sexual and reproductive health [8]. Need for sexual and reproductive health care relates to
79 the provision of safe, effective, accessible services, encompassing a broad range of activities
80 including but not limited to detection, treatment and management of sexual and reproductive
81 health-related conditions, comprehensive information and education on not only health risks but
82 also sexual function and satisfaction, support for fertility choices including contraceptive, antenatal,
83 childbirth, postnatal and abortion services, as well as sexual health across the life course including
84 during and after menopause [8][7][9]. All aspects of sexual health needs should be experienced free
85 from coercion, discrimination and violence [1].

86 High levels of unintended pregnancies, use of unsafe abortion methods, untreated sexually
87 transmitted infections (STIs) and poor access to sexual health services are just a handful of examples
88 illustrating that the sexual and reproductive health and healthcare needs of women living in low-
89 and middle-income countries are not well met [8][9][10][11]. The reasons behind this are complex
90 [12][10]. Beliefs around sexuality, reproductive health and gender are often closely underpinned by
91 social, cultural and political views [11][13][14]. For women with severe mental illness (SMI), the

92 compounding effect of stigma and exclusion, economic deprivation associated with mental health,
93 gender inequality, paucity of comprehensive mental and sexual health services in LMIC and
94 psychosocial disability provides a multiplicity of conditions which make them acutely vulnerable to
95 poor sexual health [14][11]. Many women with SMI are formally or informally subjected to the
96 assumption that they lack capacity to make decisions about their own sexual and reproductive
97 health [11][15]. This renders women with SMI vulnerable to human right abuses such as the
98 experience of forced or coercive reproductive interventions and being deprived of or unable to
99 access information and services related to sexuality and reproductive health, putting them at
100 significantly greater risk of sexual violence with often little opportunity for justice [15] [11].

101 This scoping review aimed to understand and quantify the extent and type of evidence available in
102 relation to the sexual and reproductive health needs of women with SMI in LMIC. Secondary to the
103 this, the scoping review also aimed to summarise the sexual and reproductive health needs of
104 women with SMI in LMIC.

105 Materials and Methods

106 This scoping review was conducted using the Joanna Briggs Institute (JBI) guidance for scoping
107 reviews [16] and reported according to the PRISMA extension for scoping reviews [17]. A pilot
108 literature search was conducted, showing a range of literature available on the research topic and no
109 similar review published or underway. A scoping review protocol was developed and published on
110 Open Science Framework on 2 June 2023, DOI: 10.17605/OSF.IO/P5YA8.

111 Inclusion criteria:

- 112 - Concept: Studies focusing on sexual and reproductive health (including but not limited to
113 family planning, contraception, pregnancy and the postpartum period, sexual violence, sex
114 work, STI, fertility, female orgasms, denial of sexual life), and;

115 Described need (the concept of “need” referred to any aspects of need including needs that
116 are met, unmet, quantification of need, nature of need, determinants of need and needs in
117 relation to services). Studies focusing solely on child sexual exploitation, parenting,
118 marriageability and marital relationships and gynaecological topics were excluded.

119 - Population: Studies involving women or girls (including transgender women) and non-binary
120 people assigned female at birth with SMI (including schizophrenia spectrum disorders,
121 bipolar disorder, psychosis and depression with psychotic features). In studies where
122 aggregated data were presented, studies were included if over 50% of participants had SMI
123 and if over 80% of participants were women. Studies involving families, community
124 members and professionals that interact with the above population were also included.

125 - Context: Studies conducted in a LMIC as defined by the World Bank. In studies where
126 aggregated data were presented, studies were included if over 50% of participants were
127 from LMIC (this was a post-hoc criterion applied to one study).

128 - Study Design: Peer reviewed qualitative and quantitative literature were included with no
129 restriction on study type.

130

131 Search terms were based around three elements: terms related to sexual and reproductive health,
132 terms related to SMI and terms related to LMIC. There was no specific search term related to
133 women to avoid excluding studies that recruited both men and women. A search for peer reviewed
134 studies was conducted on 23rd December 2022 (re-run 7th July 2023) on EMBASE using the EBSCO
135 platform and Medline CINAHL and PsycINFO using the Ovid platform. Due to the large amount of
136 peer reviewed literature identified, grey literature was not searched. There was no restriction on
137 study date or language.

138 Screening of Studies

139 Studies identified from the searches were collated in Mendeley, de-duplicated and transferred into
140 Rayyan for screening. One reviewer screened all articles with a second reviewer screening 20% of
141 articles. Full texts of the potentially eligible studies were screened by two independent reviewers.

142 Data Extraction

143 A data extraction form was developed in Microsoft Excel based on the standard JBI template. Two
144 reviewers each extracted data from 50% of the included studies. A third reviewer independently
145 extracted data from 10% of studies for quality assurance. Data were extracted on study publication
146 date, country, setting, study type, aim/purpose, total sample size, sample size of participants
147 relevant to the inclusion criteria, participant characteristics, methodology, focus of study, outcome
148 assessed and findings. Quantitative data were summarised and study findings were grouped
149 thematically.

150 Results


151 The search identified 3054 articles, which included 332 duplicates, thus resulting in 2722 articles for
152 title and abstract screening. Full text screening was carried out for 279 articles. An article published
153 after re-run of the database was identified and added, thus yielding 280 articles for full text
154 screening, of which 187 studies were excluded, primarily due to not meeting the eligibility criteria
155 (concept/described need/context/population) for inclusion or full text articles not being available.

156 A total of 88 studies were included from the database searches and a further 12 papers were
157 identified from citation searching. Therefore, the scoping review included 100 studies (Figure 1).
158 Included studies are summarised in Table 1.

159

160

161 Figure 1: PRISMA flow diagram



162

163

Author	Year	Country	Setting	Study type and methodology	Total sample size	Relevant sample size	Outcome measures
Abayomi[18]	2013	Nigeria	Psychiatric Hospital – inpatient and outpatient	Cross-sectional	102	28	Risky sexual behaviour (bespoke questions)
Abekah-Carter[19]	2022	Ghana	In the community	Qualitative	20	20	Sexual violence focus
Afe[20]	2016	Nigeria	Outpatient – Psychiatric Hospital	Cross-sectional	77	77	Sexual assault (bespoke questions)
Aboobaker [21]	2022	South Africa	Outpatient – Psychiatric Hospital	Cross-sectional	368	181	HIV blood test
Amr[22]	2012	Egypt	Outpatient – Psychiatric Hospital	Cross-sectional	98	37	Trauma Assessment for Adults – Brief Revised Version

							(TAA)
Aneja[23]	2020	India	Outpatient – Psychiatric Hospital	Case Study	1	1	n/a
Bagadia[24]	2020	India	Outpatient – Psychiatric Hospital	Qualitative (grounded theory)	42	42	Pregnancy focus
Ben[25]	2013	Tunisia	Outpatient – Psychiatric Hospital	Cross-sectional	61	61	Sexual Behaviour Questionnaire
Bhatia[26]	2004	India	Psychiatric hospital – inpatient and outpatient	Cross-sectional	368	108	Fertility (bespoke questions)
Bram[27]	2014	Tunisia	Inpatient – Psychiatric Hospital	Cross-sectional	38	16	Changes in Sexual Functioning Questionnaire
Bursalioglu [28]	2013	Turkey	Inpatient – Psychiatric Hospital	Cross-sectional	196	96	Personal Information Questionnaire (contraception use)
Caqueo-Urizar[29]	2018	Bolivia Peru Chile	Outpatient – Psychiatric Hospital	Cross-sectional	247	83	Schizophrenia Quality of Life Questionnaire (SqoL18)
Carey[30]	2007	India	Inpatient – Psychiatric Hospital	Cross-sectional	948	375	Blood test; HIV-Risk Screening Instrument (HRSI)
Carmo[31]	2013	Brazil	Psychiatric hospital – inpatient and outpatient	Cross-sectional	2087	1089	HCV seropositivity
Carmo[32]	2014	Brazil	Psychiatric hospital – inpatient and outpatient	Cross-sectional	2206	1147	HbsAg seropositivity
Ceylan[33]	2019	Turkey	Outpatient – Psychiatric Hospital Community	Cross-sectional	186	186	Attitudes, awareness and practices regarding reproductive health needs of people with schizophrenia (bespoke questions)
Chandra (a)[34]	2003	India	Inpatient – Psychiatric Hospital	Cross-sectional	146	146	Sexual Experiences Survey;

							HIV-Risk Screening Instrument
Chandra (b) [35]	2003	India	Inpatient – Psychiatric Hospital	Cross-Sectional	50	50	Sexual Experiences Survey;
Chandra (c) [36]	2003	India	Inpatient – Psychiatric Hospital	Cross-sectional	618	429	HIV-Risk Screening Instrument (HIS)
Chandra[37]	2006	India	Inpatient – Psychiatric Hospital	Cohort	39	19	HIV knowledge (bespoke questions)
Chopra[3]	1998	India	Inpatient – Psychiatric Hospital	Cross-sectional	59	30	HIV risk (bespoke questions); Blood test
Collins[38]	2006	South Africa	Inpatient – Psychiatric Hospital	Qualitative	46	46	Sexual violence focus
Collins[39]	2009	South Africa	Inpatient – Psychiatric Hospital	Cross-sectional	151	75	Blood test
Collins[40]	2001	South Africa	Mixed – inpatient and outpatient facilities (community and hospital)	Ethnography	56	56	Sexual violence, sexual function, contraception use focus
Correa[41]	2020	Colombia	Outpatient – Psychiatric Hospital	Cross-sectional	160	79	Contraception awareness (bespoke questions)
De Oliveira [42]	2012	Brazil	Mixed – inpatient and outpatient facilities (community and hospital)	Cross-sectional	2475	1277	Sexual violence (bespoke questions)
Desai[43]	2009	India	Outpatient – psychiatric hospital	Case Study	N/a	N/a	Ethical issues in pregnant women with SMI
Dogu[44]	2012	Turkey	Outpatient – Psychiatric Hospital	Cross-sectional	120	63	Sexual function (bespoke questions)
Dutra[45]	2014	Brazil	Psychiatric Hospital – inpatient and outpatient	Cross-sectional	2145	1129	Self-reported STI

Eroglu[46]	2020	Turkey	Outpatient – psychiatric hospital	Cross-sectional	117	58	Contraception use, gravidity and parity (bespoke questions)
Esan[47]	2018	Nigeria	Outpatient – Psychiatric Hospital	Cross-sectional	90	45	Arizona Sexual Experiences Questionnaire; New Sexual Satisfaction Scale
Fanta[48]	2018	Ethiopia	Outpatient – Psychiatric Hospital	Cross-sectional	422	132	Change in Sexual Functioning Questionnaire
Gebeyehu[49]	2021	Ethiopia	Outpatient – Psychiatric Hospital	Cross-sectional	223	113	Risky sexual behaviour (bespoke questions adapted from behavioural surveillance survey)
Ghebrehwet [50]	2020	Ethiopia	In the community	Qualitative (grounded theory)	39	39	Sexual violence focus)
Grover[51]	2019	India	Outpatient – Psychiatric Hospital	Cross-sectional	219	65	Contraception use and gravidity (bespoke questions)
Guimarães [52]	2010	Brazil	Psychiatric hospital – inpatient and outpatient	Cross-sectional	2475	1277	HIV risk behaviour (bespoke questions)
Guimarães[53]	2014	Brazil	Psychiatric hospital – inpatient and outpatient	Cross-sectional	2237	1161	HIV seroprevalence
Hall[54]	2019	Timor-Leste	In the community	Qualitative	85	85	Sexual violence focus
Halouani[55]	2018	Tunisia	Outpatient – Psychiatric Hospital	Case Control	32	32	Female Sexual Functioning Index (FSFI)
Hariri[56]	2009	Turkey	Outpatient – Psychiatric Hospital	Cross-sectional	360	174	Golombok Rust Inventory of Sexual Satisfaction
Henning[57]	2012	South Africa	Inpatient – Psychiatric Hospital	Cross-sectional	195	95	HIV rapid test; HIV ELISA test
Hocaoglu [58]	2014	Turkey	Outpatient – Psychiatric Hospital	Cross-sectional	190	38	Arizona Sexual Experiences Questionnaire

Hou[59]	2016	China	Primary Care	Cross-sectional	607	279	Arizona Sexual Experiences Questionnaire
Huang[60]	2019	China	Primary Care	Cross-sectional	720	272	Arizona Sexual Experiences Questionnaire
Incedere[61]	2017	Turkey	Outpatient – Psychiatric Hospital	Cross-sectional	200	116	Sexual violence (bespoke questions); Arizona Sexual Experiences Questionnaire
Joska[62]	2014	South Africa	Outpatient – Psychiatric Hospital	Cross-sectional	100	85	HIV appointment attendance (retrospective clinical record review)
Kazour[63]	2020	Lebanon	Inpatient – Psychiatric Hospital	Cross-sectional	60	30	Sexual Behaviour Questionnaire
Kesebir[64]	2014	Turkey	Outpatient – Psychiatric Hospital	Cross-sectional	57	28	Arizona Sexual Experiences Questionnaire; Golombok Rust Inventory of Sexual Satisfaction (GRIS)
Kumar[65]	2021	India	Outpatient – Psychiatric Hospital	Cross-sectional	57	57	Changes in Sexual Functioning Questionnaire
Loganathan [66]	2022	India	Outpatient – Psychiatric Hospital	Qualitative	200	82	Pregnancy focus
Lundberg [67]	2012	Uganda	Psychiatric hospital – inpatient and outpatient	Qualitative	20	13	Sexual function and sexual violence focus
Lundberg [68]	2015	Uganda	Inpatient – Psychiatric Hospital	Cross-sectional	602	343	World Health Organization Violence Against Women Instrument; HIV blood test
Lundberg [69]	2013	Uganda	Inpatient – Psychiatric Hospital	Cross-sectional	602	343	Blood test
Madziro – Ruwizhu[70]	2019	Zimbabwe	Outpatient – Psychiatric Hospital	Cross-sectional	270	142	Blood test
Magalhães [71]	2009	Brazil	Outpatient – Psychiatric	Cross-sectional	136	136	Contraception use (bespoke

			Hospital				questions)
Maling[72]	2011	Uganda	Inpatient – Psychiatric Hospital	Cross-sectional	272	116	Blood test
Mamabolo [73]	2012	South Africa	Inpatient – Psychiatric Hospital	Cross-sectional	113	36	Sexual violence (bespoke questions)
Marengo (a) [74]	2015	Argentina	Outpatient – Psychiatric Hospital	Cross-sectional	126	63	Contraception use (bespoke questions); London Measure of Unplanned Pregnancy (LMUP)
Marengo (b) [75]	2015	Argentina	Outpatient – Psychiatric Hospital	Cross-sectional	126	63	Contraception use (bespoke questions); HIV-risk Time Line Follow Back interview (TLFB)
Mashaphu [76]	2007	South Africa	Inpatient – Psychiatric Hospital	Cross-sectional	63	13	HIV Blood test
Matshoba[4]	2021	South Africa	Outpatient – Psychiatric Hospital	Cross-sectional	214	76	HIV knowledge and attitudes (bespoke questions)
Melo[5]	2010	Brazil	Psychiatric hospital – inpatient and outpatient	Cross-sectional	2475	1277	HIV knowledge (bespoke questions)
Mere[77]	2018	South Africa	Inpatient – Psychiatric Hospital	Cross-sectional	201	121	HIV status (retrospective clinical record review))
Mirsepassi [78]	2022	Iran	Outpatient – psychiatric hospital	Qualitative	21	4	Sexual function focus
Mpango[79]	2022	Uganda	Outpatient – Psychiatric Hospital	Cross-sectional	1201	654	Blood Tests (HIV and Syphilis)
Mwelase [80]	2023	South Africa	Inpatient – psychiatric hospital	Cross-sectional	294	92	HIV status (retrospective clinical record review)
Nakhli[81]	2014	Tunisia	Outpatient – Psychiatric Hospital	Cross-sectional	100	30	Arizona Sexual Experiences Scale
Nakigudde [82]	2013	Uganda	Outpatient – Psychiatric Hospital Community	Qualitative	23	8	Family planning and postpartum focus
Negash[83]	2019	Uganda	Outpatient –	Cross-	442	115	Risky sexual


			Psychiatric Hospital	sectional			behaviour (bespoke questions)
Obo[84]	2019	Ethiopia	Outpatient – Psychiatric Hospital	Cross-sectional	424	201	Risky sexual behaviour (bespoke questions)
Olisah[85]	2016	Nigeria	Outpatient – Psychiatric Hospital	Cross-sectional	255	133	International Index of Erectile Function (IIEF) Questionnaire: Female Sexual Function Index (FSFI)
Opondo[86]	2018	Botswana	Inpatient – Psychiatric Hospital	Cross-sectional	1482	482	HIV status (retrospective clinical record review)
Ozcan[87]	2014	Turkey	Inpatient – Psychiatric Hospital	Cross-sectional	292	292	Contraception use and knowledge, sexual violence (bespoke questions)
Ozcan[88]	2018	Turkey	Inpatient – Psychiatric Hospital Outpatient – Psychiatric Hospital Community	Cross-sectional and comparative descriptive	349	149	Prenatal Attachment Inventory; Maternal Attachment Scale
Pehlivanoglu[89]	2007	Turkey	Outpatient – psychiatric hospital	Cross-sectional	200	100	Contraception use and knowledge (bespoke questions)
Peixoto[2]	2014	Brazil	Psychiatric Hospital – Inpatient and outpatient	Cross-sectional	1475	791	Contraception use, sexual violence, HIV knowledge (bespoke questions, clinical data)
Pinto[90]	2007	Brazil	Inpatient – Psychiatric Hospital	Qualitative – Ethnography	88	30	HIV risk behaviour focus
Poreddi[91]	2021	India	Inpatient – Psychiatric Hospital	Qualitative	20	20	Sexual violence focus
Rezaie[92]	2020	Iran	Inpatient – Psychiatric Hospital	Qualitative	42	42	Pregnancy and sexual function focus
Sethuraman [93]	2019	India	Outpatient – psychiatric hospital	Cross-sectional	96	96	Modified National Family Health Survey-3
Shahvari [94]	2020	Iran	Inpatient –	Qualitative	21	21	HIV risk

			Psychiatric Hospital				behaviour and contraception use focus
Sibanyoni [95]	2022	South Africa	Psychiatric hospital – inpatient and outpatient	Qualitative	23	23	Pregnancy and family planning focus
Simiyon [96]	2016	India	Outpatient – Psychiatric Hospital	Cross-sectional	63	63	Marital Quality Scale Female Sexual Function Index (FSFI).
Singh[97]	2009	South Africa	Inpatient – Psychiatric Hospital	Cross-sectional	206	88	HIV Blood tests – ELISA
Souaiby [98]	2020	Lebanon	Outpatient – Psychiatric Hospital	Cross-sectional	95	13	Psychotropic-Related Sexual Dysfunction Questionnaire (PRSexDQ)
Souto[99]	2011	Brazil	Inpatient – Psychiatric hospital Outpatient – Psychiatric Hospital	Cross-sectional	2380	1232	Self-reported HIV testing
Terzian [100]	2006	Brazil	Outpatient – Psychiatric Hospital	Cross-sectional	167	65	Fertility rate
Tharoor [101]	2015	India	Outpatient – Psychiatric Hospital	Cross-sectional	136	67	Psychotropic Related Sexual Dysfunction Questionnaire (PRSexDQ-Salsex)
Tomruk [102]	2006	Turkey	Outpatient – Psychiatric Hospital	Cross-sectional	200	100	HIV risk behaviour (bespoke questions)
Tumwakire[103]	2022	Uganda	Inpatient – Psychiatric Hospital	Qualitative	14	14	Sexual function, contraception use and sexual violence focus
Tunde [104]	2013	Nigeria	Outpatient – Psychiatric Hospital	Cross-sectional	100	100	Family planning experience/knowledge (bespoke questions)
Wainberg [105]	2007	Brazil	Psychiatric hospital – inpatient and outpatient	Qualitative	88	30	Sexual function and contraception use focus
Wainberg [106]	2008	Brazil	Outpatient – Psychiatric	Cross-sectional	98	50	Sexual Risk Behaviour

			Hospital				Assessment Schedule (SERBAS)
Yang[107]	2023	China	Inpatient – Psychiatric Hospital Outpatient – Psychiatric Hospital	Qualitative – phenomenological	20	10	Sexual function focus
Yu[108]	2022	China	Psychiatric Hospital – inpatient and outpatient	Qualitative – phenomenological	15	15	Fertility, pregnancy, post-partum focus
Zerihun [109]	2020	Ethiopia	Outpatient – General Hospital (psych dept)	Cross-sectional	422	422	Gravidity, contraception use and knowledge (bespoke questions)
Zerihun(a) [110]	2021	Ethiopia	Outpatient - Psychiatric Hospital	Nested qualitative	16	16	Pregnancy, family planning, sexual violence focus
Zerihun(b) [111]	2021	Ethiopia	Outpatient - General Hospital (psych dept)	Cross-sectional	422	422	Adapted WHO Intimate Partner Violence tool
Zhang (a) [112]	2018	China	Inpatient - Psychiatric Hospital	Cross-sectional	87	33	Syphilis seroprevalence
Zhang (b) [113]	2018	China	Inpatient - Psychiatric Hospital	Cross-sectional	118	63	Arizona Sexual Experience Scale

164 Year of study

165 The oldest included study was from 1998 [3], whilst the most recently published were from 2023
 166 [80][107]. Over the 25 years that relevant studies were published, 62 were published within the last
 167 10 years. The highest number of studies were published in 2014 (n=10). Figure 2 displays the
 168 number of included studies by publication year.



169

170 Context

171 Of the 140 LMIC globally, only 20 countries (14%) were represented in the included studies. Figure 3
172 illustrates the breadth of countries in the review in proportion to the number of included studies
173 from each of those countries. All studies focused on a single country, except for one study which
174 included participants drawn from three South American countries. The most common continent for
175 included studies was Africa (n=36). Asia was the second most common (n=24), followed by the
176 Middle East (n=22) and South America (n=18).

177

178


179

180

181

182

183 **Figure 3: Countries displayed as a proportion of included studies.**



184
185

186

187 India was the country setting for the largest number of studies (n=17). Other countries with >10
188 included studies were Brazil (n=14), South Africa (n=13) and Turkey (n=12). This meant that 4 of the
189 140 (3%) LMIC globally accounted for 56% of all included studies.

190 The topic of studies carried out in India were heterogenous. Eight of the 14 studies carried out in
191 Brazil and 11 of the 13 studies carried out in South Africa focused on HIV. In contrast, most studies
192 set in Turkey were focused on sexuality and sexual function (n=6) and family planning (n=4).

193 The majority (n=7) of studies in Uganda were based around HIV and HIV risk behaviours. Four of the
194 six studies set in China were based around sexual function, three of which utilised the Arizona Sexual
195 Experiences (ASEX) Questionnaire [59][60][113]. All Tunisian studies (n=4) were focused on sexuality
196 and all Iranian studies (n=3) were qualitative.

197 Psychiatric hospital outpatient facilities were the most frequent setting for studies (n=46), followed
198 by psychiatric hospital inpatient settings (n=29). In addition, a proportion of studies utilised mixed
199 inpatient and outpatient psychiatric settings (n=20), five of which also incorporated participants

200 from the community. Two studies were conducted in primary care, both of which were based in
201 China. A small number of studies were based solely in the community and not in health facilities
202 (n=3).

203 Participants

204 A total of 38,510 participants were included in the studies, however some studies used the same
205 participants so the total likely overestimates the number included. Of the total number of
206 participants, 19,987 (52%) were relevant to the study population of this scoping review. Small
207 numbers of men and women without SMI may be included in this total in studies that were not
208 disaggregated by diagnosis or gender.

209 The most represented SMI diagnosis was schizophrenia. Twenty-eight studies focused solely on
210 schizophrenia and people with schizophrenia were included in a further 54 studies. People with a
211 diagnosis of bipolar disorder were included in 55 studies with an additional 8 studies exclusively
212 focused on people with bipolar disorder.

213 Generally, inclusion criteria of the studies restricted participation to those 18 years of age or older.
214 Only four studies [28][26][102][89] included women aged 15-years or above and one study included
215 those aged 16 or older [18]. The oldest recorded participant included in a study was 83 years.
216 Reporting of age ranges within studies was variable and some studies reported mean age of
217 participants only. Where age range was reported, it was noted that only a handful of studies
218 included women of menopausal age and older.

219 Participants without lived experience of SMI were included in 11 studies
220 [19][105][103][94][82][78][54][50][40][38][33]. These included psychiatric nurses, midwives,
221 administrators of mental health care facilities, caregivers or family members of people with SMI,
222 government decision makers, mental healthcare providers, and people living in communities that
223 include people with SMI.

224 Ethics

225 Included studies were searched for evidence of ethics approval, informed consent and assessment of
226 capacity to consent. Nineteen of the 100 included studies did not report whether ethical approval
227 had been obtained [23][25][36][35][34][37][43][44][51][55][65][66][71][81][102][90] [104][19].
228 Seventy-six studies explicitly stated that informed consent had been sought from participants but of
229 those, only 16 studies [31][32][37][36][42][52][57][60][67][70][5][2][99][110][111][113] included a
230 statement that capacity to consent had been assessed prior to participants being included in studies.
231 This means that almost a quarter of studies did not provide information in their published papers as
232 to whether consent had been obtained and most studies in the review (n=79) did not include
233 information as to whether capacity to consent had been assessed. A small number of papers (n=5)
234 did not require assessment of capacity to consent due to the nature of the study or the focus of the
235 study being participants without lived experience of SMI [19][112][40][38][33].

236 Of the studies that included participants aged 15 or 16 years and older (n=5), none explicitly stated
237 that capacity to consent had been assessed [18][28][89][97][102] and one study did not document
238 that consent had been obtained. All five studies did provide information indicating that ethics
239 approval had been obtained.

240 Methodology

241 Most studies (n=76) implemented a cross-sectional design. Other quantitative study designs were
242 one case control study [55] and one cohort study [37]. Questionnaires were the main data collection
243 tool used in these studies. A sizeable number of studies also used blood samples as a data source.

244 There were 22 qualitative studies included in the review, two of which used ethnographic
245 methodologies [40][90]. There were two case studies in the review [43][23]. Most of the qualitative
246 studies used semi-structured interviews (n=18). The range of sample sizes for the qualitative studies
247 was 14-200 participants.

248 Study Focus

249 Included studies frequently had multiple areas of focus. Focus areas were grouped into 6
250 overarching themes. The largest domains were HIV prevalence, risk behaviour and knowledge
251 (n=32), sexual function (n=30), contraception use and family planning (n=25), sexual violence (n=21),
252 fertility, pregnancy and the post-partum period (n=19 studies) and non-HIV STIs (n=11). The most
253 relevant results from each study have been reported in Appendix 2.

254 HIV (prevalence, risk-taking behaviour and knowledge)

255 All of the 14 studies that ascertained the prevalence of HIV reported it to be higher in women with
256 SMI than in men with SMI [21][30][39][80][53][57][69][70][72][76][86][97][77]; however, not all of
257 these studies were probabilistically sampled. Sixteen studies researched HIV risk behaviour
258 [18][34][36][3][40][49][53][68][75][83][2][90][94][102][106]. Results were varied, although the
259 majority reported HIV risk behaviour was generally higher among women with SMI than men with
260 SMI. Key findings from three qualitative studies [40][90][94] found mental healthcare providers 1.)
261 were more concerned about the avoidance of pregnancy than the avoidance of HIV, 2.) perceived
262 patients to be at high risk of HIV, and 3.) noted that patients may not be able to 'resist' high risk
263 sexual behaviour and may also lack the agency to refuse unprotected sexual intercourse.

264 Six studies assessed the knowledge of people with SMI about HIV/AIDS [37][3][4][5][102][2] with
265 inconsistent findings between the studies. Three of the studies reported that HIV knowledge in
266 women with SMI was lower than a comparator group of men with SMI [37][3] and a comparator
267 group of women without SMI [102]. Women with SMI had a higher knowledge score in one study
268 [5]. Two studies reported no significant difference in HIV knowledge score between men and women
269 with SMI [2][4]. Limited evidence from individual studies reported that a higher proportion of
270 women with SMI attended HIV appointments compared to men with SMI [62]. A study looking at
271 whether men and women with SMI had ever had testing for HIV reported no significant difference
272 [99].

273 STIs (not including HIV)

274 Most studies focused on non-HIV STIs used men with SMI as a comparator, with only one study
275 comparing to women without SMI [75]. The prevalence of a range of STIs were reported. One study
276 showed a similar prevalence of chlamydia between men and women[30]. In the three studies that
277 reported the prevalence of syphilis, one showed no difference [79] whilst the other two showed
278 prevalence to be higher in men [112][30]. A qualitative study noted the perception of mental
279 healthcare providers that rates of syphilis were higher in their hospital population compared to the
280 general population [40]. Two studies reported hepatitis B prevalence, both showed a higher
281 prevalence of hepatitis B in men with SMI than women with SMI. Similarly, a study of hepatitis C
282 prevalence showed a higher prevalence in men [30][32]. Prevalence of history of STI was reported by
283 five studies [18][42][45][75][32]. Generally, studies showed a similar prevalence for history of STI
284 between women and men but a study that compared women with SMI to women without SMI
285 showed a higher prevalence of STI in women with SMI.

286 The two qualitative studies included utilised healthcare provider staff participants only or in
287 combination with people with SMI [40][90]. One qualitative study suggested that mental healthcare
288 providers felt that SMI patients were at a higher risk of STI than the general population [90].

289

290 Sexuality and Sexual Function

291 A total of 30 studies focused on sexuality and sexual function. The most commonly used measure of
292 sexual function was the Arizona Sexual Experiences Scale (ASEX) which was used in eight of the
293 included studies [81][58][61][64][59][60][47][113]. Other measures of sexual function included The
294 Changes in Sexual Functioning Questionnaire (CSFQ) [48][27][65] and Female Sexual Functioning
295 Index (FSFI) [55][96][85], the Sexual Behaviour Questionnaire (SBQ)[25], the Golombok Rust
296 Inventory of Sexual Satisfaction (GRISS) [56][64], the Psychotropic-Related Sexual Dysfunction

297 Questionnaire (PRSexDQ)[101][98], the Sexual Risk Behaviour Assessment Schedule (SERBAS)[106],
298 the Schizophrenia Quality of Life Questionnaire (SQoL18) questionnaire, specifically the “sentimental
299 life score” component (measuring satisfaction with love life)[29]. One study used a bespoke scale
300 and found more women with SMI than men with SMI reported a problem in their sexual life [44].

301 Of the 18 studies that used standardised measures of sexual dysfunction and compared women with
302 SMI to men with SMI, ten studies reported higher levels of sexual dysfunction in women with SMI
303 than men with SMI [85][81][60][59][106][47][107][61][98][25]. Five studies reported no significant
304 difference between women with SMI and men with SMI [58][27][48][56][63]. Only one study
305 reported higher levels of sexual dysfunction in men with SMI [101] and another reported lower
306 levels of sexual satisfaction in men with SMI [29]. Limited evidence from one study indicated that
307 self-reported levels of sexual dysfunction varied by gender when type of medication was taken into
308 consideration [64].

309 Of those studies that did not use a male comparator, one study comparing groups of women with
310 SMI taking risperidone or olanzapine medication for SMI found both groups to have high levels of
311 sexual dysfunction [65], another study reported women with SMI to have higher levels of sexual
312 dysfunction compared to women without SMI [55]. A third study reported 70% of women with SMI
313 had high levels sexual dysfunction [96] although no comparator was used.

314 Eight qualitative studies relating to sexuality were identified [40][67][78][90][92][103][105][107].
315 Stigmatisation was consistently reported by women with SMI and that this was at times perpetuated
316 by healthcare providers. Examples given were perceptions that women with SMI lacked sexual needs
317 and the promotion of lifelong abstinence. Studies including healthcare providers reported that staff
318 caring for women with SMI agreed they have the same sexual desires as women without SMI. Both
319 healthcare providers and people with SMI felt that sexuality could be affected by mental illness, with
320 both increases and decreases in sexual activity described. Women with SMI described positive
321 experiences of sexuality to be beneficial to their mental health but also negative sexual experiences

322 to be a result of their illness. There was limited research indicateing women with SMI were keen to
323 learn more about the interaction between SMI and their sexual fulfilment. No studies reported
324 sexuality or sexual function needs and experiences of lesbian, gay, bisexual, transgender, intersex,
325 queer, asexual or other sexually or gender diverse LGBTQIA+ women with SMI.

326 Contraception and Family Planning

327 Twenty-five studies were based around contraception and family planning. The majority of these
328 (n=13) focused on contraception use [28][40][43][51][53][71][74][75][2][90][94][105][106]. Four
329 studies, which did not include a comparison, reported large proportions of women with SMI had
330 never used contraception, had not used contraception during their most recent sexual experiences
331 and had no intention of using contraception during sexual intercourse [87][109][71][93].

332 Of those studies that included a comparator, most studies used a comparator group of women with
333 SMI rather than men with SMI. Contraception use compared to women without SMI was mixed with
334 two studies stating no difference in contraceptive use [74][75], two studies stating women with SMI
335 were more likely to use contraception [46][28] (in particular ‘traditional’ methods such as
336 withdrawal and calendar methods) and one stating that women with SMI were less likely to use
337 contraception [89]. There were similar results for studies comparing women with SMI to men with
338 SMI; two studies found contraception use to be lower in women [51][2] whilst one study found it to
339 be lower in men [41].

340 Four qualitative studies were also identified on the topic of contraception [94][103][105][106].
341 Healthcare professionals perceived that women with SMI were commonly engaged in unprotected
342 sex and therefore prescribed reversible contraceptive methods for them. Women with SMI felt they
343 may be perceived negatively by partners for using condoms. When asked about knowledge of
344 different types of contraception, coitus interruptus, condoms and intrauterine devices (IUDs) were
345 generally the most cited.

346 A small number of studies looked at informed choices and knowledge when making contraception
347 decisions. One study assessed whether there were differences in participants being informed about
348 contraception based on gender and diagnosis of either schizophrenia or bipolar disorder [41]. A
349 further study looked at the proportion of women with SMI referred to a perinatal psychiatric clinic
350 and reported that health care providers had discussed contraception issues with them [43]. A
351 qualitative study in Uganda looked at the autonomy of women with SMI in making contraception
352 choices and noted that this was often absent, with consent taken from family members to force a
353 woman to use contraception if she refused [103].

354 Seven studies focused on family planning advice from professionals [41][33][87][93][82][110][104].
355 The focus of included studies was varied. Most studies looked at the quantity of provision or receipt
356 of family planning advice from the perspective of either healthcare professionals or patients. None
357 of the included studies assessed the quality of family planning advice provided. One study included a
358 comparison and found more women with SMI received family planning advice compared to men
359 with SMI [41]. In one study, 23.5% of healthcare professionals stated they gave family planning
360 advice to patients [33], in another 21.9% of people with SMI stated they received family planning
361 advice from professionals [87]. Two further studies concluded there was a lack of family planning
362 advice being provided to patients from healthcare professionals [93][104]. In a qualitative study,
363 caregivers and postpartum women with SMI felt that family planning education should be added to
364 an existing psychoeducation programme [82]. Two studies reported that women with SMI felt that
365 they would prefer to receive information about family planning from mental healthcare providers
366 [110][104] as they are knowledgeable about the impact of their health conditions on family planning
367 decision making.

368 Two studies (from India and Turkey) provided evidence of healthcare professionals making family
369 planning decisions for patients [24][33], with forced abortion being considered necessary by over
370 39% of health professionals working with people who have schizophrenia in one study [33]. Use of

371 sterilisation and abortion for failed contraception was reported in a qualitative study carried out in
372 South Africa, which looked at the perceptions and experiences of mental health care staff [40].
373 Another study found that family planning considerations affected desire to take medication and
374 were influenced by medication, with some women with SMI not wanting to take sodium valproate
375 due to teratogenicity risk [95].

376 Sexual Violence

377 Six studies [22][42][61][68][73][2] compared experiences of sexual violence in women with SMI to
378 men with SMI or women without SMI. Most studies reported lifetime experience of sexual violence.
379 Two studies reported experience of sexual violence from a partner or non-partner[68][87]. All but
380 one study found women with SMI to be more likely to have experienced sexual violence. An Egyptian
381 study reported men to be more likely to report sexual violence [22]. Six studies reported the
382 prevalence of sexual violence in populations of women with SMI, and this ranged from 10%-100%
383 [20][34][35][102][87][111].

384 There were nine qualitative studies [110][103][91][67][54][50][40][38][19] focusing on the theme of
385 sexual violence. Two thirds of included studies utilised non-patient participants either wholly or in
386 combination with people who have lived experience. Only three of the nine qualitative studies
387 included exclusively people with lived experience of SMI [110][67][91].

388 All nine studies reported the vulnerability of women with SMI to sexual violence. This included
389 women being raped by male patients, whilst inpatients in psychiatric facilities. In addition to reports
390 of sexual violence, participants reported difficulties accessing medical care after rape for homeless
391 women; rape resulting in unintended pregnancies and delayed discharge from medical care facilities
392 in women who experienced sexual violence at home.

393 Fertility, Pregnancy and Postpartum

394 Five studies were based around fertility [26][51][74][100][108]. These were varied in terms of the
395 outcome being studied. Three studies looked at conception and childlessness and found similar rates
396 between women with SMI and men with SMI or women without SMI [26][51][74]. A study carried
397 out in Brazil found lower fertility rates and fecundity in women with SMI compared to women in the
398 general population [100]. A qualitative study described difficulties in the fertility process for women
399 with SMI, alongside parenting being described as a rewarding process that aids recovery [108].

400 Fourteen studies focused on pregnancy [23][24][43][46][66][87][88][91][41][92][95][108] [109][110].
401 Notably, the only two case studies included in this review were based around decision-making
402 during pregnancy in the context of medication and abortion [23][43]. Seven studies mentioned the
403 impact of medication on pregnancy and/or the effect of withholding medication during pregnancy.
404 One qualitative study mentioned a higher risk perception related to impacts of medication on a
405 fetus compared to risk of unmedicated mental illness during pregnancy [110]. One Turkish study
406 included a comparator and found that women with SMI were less likely to have experienced
407 antenatal care during pregnancy, had a greater frequency of Caesarean birth and were subject to
408 more birth related trauma [88]. Two studies reported that high numbers of women with SMI
409 experience unplanned pregnancy (51.4% of women in one study [87] and 87.8% of women who had
410 experienced pregnancy in another [109]. A study that compared women with schizophrenia to
411 bipolar disorder [41] found those with bipolar disorder to have a lower age of first pregnancy (22
412 years for bipolar and 32 years for schizophrenia). Two studies reported pregnancy being discouraged
413 for women with SMI, 69% of Ethiopian community members felt that women with SMI should not
414 give birth in one study [110], whilst in India, there were reports of women being forced to abort or
415 coerced not to have children [66]. A comment in another Indian study reported instances of
416 pregnancy loss due to domestic violence [91]. One study found no significant difference in the
417 number of pregnancies or number of lost or interrupted pregnancies between for women with

418 bipolar disorder compared to controls but that a higher proportion of women with bipolar disorder
419 had a higher proportion of unplanned pregnancies [46].

420 Three studies were based in the postpartum period [88][82][108]. One study compared women with
421 SMI to women without SMI and found that women with SMI were less likely to breastfeed, more
422 likely to express concerns about infant care, experience childbirth related trauma and rely on others
423 to care for their babies [88]. Two studies mentioned breastfeeding: a Ugandan study discussed
424 instances of women with SMI being told not to breastfeed as mental illness could be passed onto an
425 infant through breastmilk [82] and some women with schizophrenia in a Chinese study stated that
426 breastfeeding could lead to a loss of routine for the women with SMI that could lead to an
427 exacerbation of illness [108].

428 Discussion

429 This review identified a substantial body of evidence relating to the sexual and reproductive health
430 needs of women with SMI in LMIC. The most studied domains of sexual and reproductive health
431 were HIV and sexual function, followed by contraception and family planning, sexual violence,
432 fertility, pregnancy and post-partum and finally non-HIV STIs. Age ranges were not reported for all
433 studies but where age range was known, the literature indicated a focus on adult women of
434 reproductive age. There were no studies specifically looking at sexual and reproductive health needs
435 of women with SMI during menopause. This could however be influenced by some of the studies
436 being conducted in countries where average life expectancy for women is close to expected age for
437 menopause [114] and that menopause was not an included term in the search strategy. Sexual and
438 reproductive health in girls and LGBTQIA+ women with SMI were also areas with no studies
439 identified by this review.

440 The review frequently found themes of women with SMI in LMIC having worse sexual and
441 reproductive health compared to both women without SMI and men with SMI. The increasing

442 number of studies over time may indicate this topic is receiving relatively more attention. However,
443 only a small number of LMICs (14%) were represented in this review, signifying that the sexual and
444 reproductive health needs of women with SMI remain underexplored in the majority of LMIC. Study
445 focus varied by world region in this review with Middle Eastern studies mainly focused on sexuality
446 and almost all South African studies focused on HIV. Schizophrenia was the most commonly
447 represented SMI and the most common study type was cross-sectional (n = 76), followed by
448 qualitative interview studies (n = 18).

449 A range of ethical concerns related to the sexual and reproductive health of women with SMI in
450 LMIC were identified. In studies with mental healthcare providers, there were reports of women
451 being forced to take medication and to undergo abortion and sterilisation. A 2019 Turkish study [33]
452 reported that 79% of healthcare providers felt that forced abortion and sterilisation was necessary.
453 A lack of agency was a theme throughout studies, with women reporting being pressured to not
454 have children or breastfeed, for fear their illness would pass into their offspring. Women were also
455 pressured by partners to avoid contraception. In addition, ethical approval, informed consent and
456 capacity assessment procedures were not documented for a substantial minority of studies. These
457 findings show that some women with SMI may not have their autonomy upheld and may be
458 vulnerable to harm from family members, partners, healthcare professionals and even researchers.

459 Availability of family planning support and contraception seemed to be mixed. Some women with
460 SMI may lack capacity to make decisions, such as those related to contraception. Contraception was
461 often given without their consent in these instances. However, one study mentioned that planning
462 for management of relapse of mental illness when a women was well would allow them to make
463 their own decisions about family planning, reflecting the principles of the CRPD in terms of making
464 decisions according to the best knowledge of an individuals' preferences [92].

465 Comparison to Previous Literature

466 Hughes et al. [115] completed a systematic review and meta-analysis of HIV, hepatitis B and
467 hepatitis C prevalence in people with SMI globally in 2016 (70). They found a higher prevalence of
468 these viral infections in people with SMI in countries with a low prevalence in the general population
469 and similar prevalence in countries with a high prevalence in the general population. This systematic
470 review included many studies that were also included in the current review. The authors identified
471 high levels of bias present in studies focusing on HIV prevalence in LMIC. They also commented on a
472 general theme of higher HIV prevalence in women compared to men in both high and low
473 prevalence countries, which is in keeping with findings from this scoping review. A 2016 systematic
474 review [117] found that 10% of women with SMI experienced sexual violence, and that this was
475 significantly higher than men with SMI and the general population. That review mainly included
476 studies from HIC but had similar findings to this review.

477 What This Review Adds

478 Throughout the results, the burden of sexual and reproductive health needs for women with SMI in
479 LMIC is clearly shown. Rates of STIs, HIV and risky sexual behaviour were high. A small number of
480 studies on educational programmes for HIV showed positive outcomes. The need for more
481 education on STIs and unwanted pregnancy prevention was identified. Studies based around
482 pregnancy were often focused on the use of medication in pregnancy, but a small amount of
483 research on pregnancy did state advantages (such as providing a purpose in life) and disadvantages
484 (exacerbation of mental illness) of pregnancy for women with SMI.

485 The link between sexual dysfunction and SMI was found to be complex with multiple factors such as
486 psychotropic medication, self-esteem and positive and negative symptoms of SMI to be involved. All
487 studies about sexual violence found that women with SMI were at risk of sexual violence, including
488 whilst admitted onto mixed inpatient wards, in the home environment or when homeless.

489 Limitations

490 Limitations were present in this scoping review. Many LMIC were not covered in this review, and
491 with the wide variation in contexts, cultures and geography, it is difficult to generalise these findings
492 to all LMIC. Most studies were of a cross-sectional design, therefore causality cannot be implied for a
493 majority of papers. As this is a scoping review, bias was not assessed. There may be high levels of
494 bias in some studies which could skew the conclusions of this review. Many studies had small sample
495 sizes and are considered alongside larger studies, although an effort was made to note studies that
496 were particularly small. Grey literature was excluded in this review due to the large amount of peer
497 reviewed literature available. Capturing unpublished literature may ensure inclusion of more LMICs.

498 Strengths

499 The strengths of this scoping review include a rigorous methodology with an iterative approach that
500 resulted in early decisions to refine the inclusion criteria. Initially, only studies that provided
501 disaggregated data by gender and diagnosis were included. It was felt that some useful data was
502 being excluded based on these criteria and therefore, the inclusion criteria was refined to include
503 aggregated data where at least 50% of participants had SMI or 80% were women. A broad definition
504 of sexual and reproductive health needs ensured that multiple areas of the review topic were
505 included and ensured a broad overview of this multifaced area. Double coding was utilised
506 throughout the review process to increase robustness of findings.

507 Implications

508 Implications for a subsequent review

509 This scoping review systematically identified literature on the sexual and reproductive health needs
510 of women in LMIC in relation to study type, focus of study, population, setting and outcome.
511 Subsequent synthesis of evidence from one or more of the focus areas from this review is planned.

512 This would enable the literature identified to be utilised to inform decision-making in policy and
513 practice.

514 There are challenges highlighted by this scoping review that will need to be addressed to progress to
515 full evidence synthesis. Study type will need careful consideration as many of the studies included in
516 this review were observational and may provide rich insights but be challenging to draw conclusions
517 from. This review highlighted that only a small proportion of LMIC countries are represented and a
518 future review will likely need to consider how applicable findings are for the culture and diversity of
519 other LMIC countries and consider the use of grey literature.

520 Implications for research and practice

521 There is a need for future research to better represent the breadth of LMIC countries and for high
522 quality research to be carried out. Other key gaps in the literature were identified by this review.
523 This included a need for greater focus on inclusion of people with lived experience as part of studies
524 looking at sexual violence, as well as ensuring that the ethical and consensual aspects are fully
525 considered and reported by researchers. These are both areas underscored by the recent work of
526 ‘The Lancet Commission on ending Stigma and Discrimination in Mental Health’, which outlined the
527 increased effectiveness of interventions developed in co-production with people who have lived
528 experience [118]. The Lancet Psychiatry has also recently specified that research submitted for
529 publication will be required to provide information on how people with lived experience have
530 shaped research priorities, the design of the study and interpretation and write up of findings [119]
531 with authors expected to acknowledge this as a limitation of their work where it has not been
532 carried out [119].

533 Marginalisation and social exclusion not only exacerbate access to mental and physical health care
534 but are often the catalyst that leads to basic human rights of people with mental health conditions
535 being contravened [118]. Future research would benefit from focussing on the effectiveness of
536 interventions that integrate physical and mental health, most notably those aimed at reducing

537 stigma and discrimination around mental health and sexual health needs, as well as culturally
538 appropriate adaptations [118].

539 Financial Disclosure Statement

540

541 CH and LA receive support from the National Institute for Health and Care Research (NIHR) through
542 the NIHR Global Health Research Group on Homelessness and Mental Health in Africa (NIHR134325)
543 and CH also receives support from the SPARK project (NIHR200842) using UK aid from the UK
544 Government. The views expressed in this publication are those of the authors and not necessarily
545 those of the NIHR or the Department of Health and Social Care. CH also receives support from WT
546 grants 222154/Z20/Z and 223615/Z/21/Z.

547

548 For the purposes of open access, the authors have applied a Creative Commons Attribution (CC BY)
549 licence to any Accepted Author Manuscript version arising from this submission.

550

552 References

- 553 [1] United Nations Population Fund. Sexual & reproductive health 2023.
554 <https://www.unfpa.org/sexual-reproductive-health> (accessed January 4, 2024).
- 555 [2] Peixoto ER de M, Barros FCR de, Guimarães MDC. Fatores associados com a prática do sexo
556 desprotegido entre homens e mulheres com transtornos mentais no Brasil. *Cad Saude Publica*
557 2014;30:1475–86. <https://doi.org/10.1590/0102-311X00029313>.
- 558 [3] Chopra MP, Eranti SSV, Prabha SC. HIV-Related Risk Behaviors Among Psychiatric Inpatients in
559 India. *Psychiatric Services* 1998;49.
- 560 [4] Matshoba T, Mashaphu S, Tomita A, Paruk S. HIV knowledge, attitudes and practices amongst
561 patients with severe mental illnesses and chronic medical illnesses in Durban, South Africa. *S*
562 *Afr J Psychiatr* 2021;27:1586.
563 <https://doi.org/https://dx.doi.org/10.4102/sajpspsychiatry.v27i0.1586>.
- 564 [5] Melo APS, Cesar CC, Acurcio F de A, Campos LN, Ceccato M das GB, Wainberg ML, et al.
565 Individual and treatment setting predictors of HIV/AIDS knowledge among psychiatric
566 patients and their implications in a national multisite study in Brazil. *Community Ment Health*
567 *J* 2010;46:505–16. <https://doi.org/https://dx.doi.org/10.1007/s10597-010-9303-7>.
- 568 [6] United Nations. Convention on the Rights of Persons with Disabilities (CRPD) | Division for
569 Inclusive Social Development (DISD) 2006.
570 [https://social.desa.un.org/issues/disability/crpd/convention-on-the-rights-of-persons-with-](https://social.desa.un.org/issues/disability/crpd/convention-on-the-rights-of-persons-with-disabilities-crpd)
571 [disabilities-crpd](https://social.desa.un.org/issues/disability/crpd/convention-on-the-rights-of-persons-with-disabilities-crpd) (accessed January 4, 2024).
- 572 [7] WHO. Sexual and Reproductive Health and Research (SRH) 2023.
573 [https://www.who.int/teams/sexual-and-reproductive-health-and-research-\(srh\)/areas-of-](https://www.who.int/teams/sexual-and-reproductive-health-and-research-(srh)/areas-of-work/sexual-health)
574 [work/sexual-health](https://www.who.int/teams/sexual-and-reproductive-health-and-research-(srh)/areas-of-work/sexual-health) (accessed January 4, 2024).
- 575 [8] Starrs AM, Ezeh AC, Barker G, Basu A, Bertrand JT, Blum R, et al. Accelerate progress—sexual
576 and reproductive health and rights for all: report of the Guttmacher– Lancet Commission. *The*
577 *Lancet* 2018;391:2642–92. [https://doi.org/10.1016/S0140-6736\(18\)30293-9](https://doi.org/10.1016/S0140-6736(18)30293-9).
- 578 [9] Guttmacher Institute. Investing in Sexual and Reproductive Health in Low- and Middle-
579 Income Countries | Guttmacher Institute 2020. [https://www.guttmacher.org/fact-](https://www.guttmacher.org/fact-sheet/investing-sexual-and-reproductive-health-low-and-middle-income-countries)
580 [sheet/investing-sexual-and-reproductive-health-low-and-middle-income-countries](https://www.guttmacher.org/fact-sheet/investing-sexual-and-reproductive-health-low-and-middle-income-countries) (accessed
581 January 4, 2024).
- 582 [10] Sedgh G, Ashford LS, Hussain R. Unmet Need for Contraception in Developing Countries:
583 Examining Women’s Reasons for Not Using a Method 2016.
- 584 [11] Women Enabled International. Women Enabled International Submission to OHCHR Report
585 on Mental Health and Human Rights 2016.
- 586 [12] Mitchell KR, Lewis R, O’Sullivan LF, Fortenberry JD. What is sexual wellbeing and why does it
587 matter for public health? *Lancet Public Health* 2021;6:e608–13.
588 [https://doi.org/10.1016/S2468-2667\(21\)00099-2](https://doi.org/10.1016/S2468-2667(21)00099-2).
- 589 [13] WHO. Transforming mental health for all 2022.

- 590 [14] Rathod S, Pinninti N, Irfan M, Gorczynski P, Rathod P, Gega L, et al. Mental Health Service
591 Provision in Low- and Middle-Income Countries. *Health Serv Insights*
592 2017;10:117863291769435. <https://doi.org/10.1177/1178632917694350>.
- 593 [15] WHO. Sexual health, human rights and the law 2015.
- 594 [16] Peters MDJ, Godfrey CM, Khalil H, McInerney P, Parker D, Soares CB. Guidance for conducting
595 systematic scoping reviews. *Int J Evid Based Healthc* 2015;13:141–6.
596 <https://doi.org/10.1097/XEB.000000000000050>.
- 597 [17] Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for
598 Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med* 2018;169:467–73.
599 <https://doi.org/10.7326/M18-0850>.
- 600 [18] Abayomi O, Adelufosi A, Adebayo P, Ighoroje M, Ajogbon D, Ogunwale A. HIV risk behavior in
601 persons with severe mental disorders in a psychiatric hospital in Ogun, Nigeria. *Ann Med*
602 *Health Sci Res* 2013;3:380. <https://doi.org/10.4103/2141-9248.117960>.
- 603 [19] Abekah-Carter K, Oti GO. Perspectives of community members on homeless people with
604 mental illness in Nsawam, Ghana. *International Journal of Social Psychiatry* 2022;68:196–202.
605 <https://doi.org/10.1177/0020764020984195>.
- 606 [20] Afe TO, Emedoh TC, Ogunsemi O, Adegbohun AA. Intimate partner violence, psychopathology
607 and the women with schizophrenia in an outpatient clinic South-South, Nigeria. *BMC*
608 *Psychiatry* 2016;16. <https://doi.org/10.1186/s12888-016-0898-2>.
- 609 [21] Aboobaker A, Zingela Z. Correlates and cascade of HIV care in patients with psychiatric
610 disorders in the Eastern Cape province, South Africa. *South African Journal of Psychiatry*
611 2022;28:a1753. <https://doi.org/https://dx.doi.org/10.4102/sajpsychiatry.v28i0.1753>.
- 612 [22] Amr M, El-Wasify M, Amin T, Roy A. Prevalence and correlates of physical and sexual assault
613 history in patients with schizophrenia. *Arab Journal of Psychiatry* 2012;23:22–9.
- 614 [23] Aneja J, Arora S. Pregnancy and severe mental illness: Confounding ethical doctrines. *Indian J*
615 *Med Ethics* 2020:133–9. <https://doi.org/10.20529/IJME.2020.037>.
- 616 [24] Bagadia A, Nanjundaswamy MH, Ganjekar S, Thippeswamy H, Desai G, Chandra PS. Factors
617 influencing decision-making around pregnancy among women with severe mental illness
618 (SMI): A qualitative study. *Int J Soc Psychiatry* 2020;66:792–8.
619 <https://doi.org/https://dx.doi.org/10.1177/0020764020925104>.
- 620 [25] Ben Mahmoud S, Zouari L, Dammak M, Ben Thabet J, Zouari N, Maalej M. Evaluation of
621 sexuality in 61 subjects suffering from chronic psychosis. *Sexologies: European Journal of*
622 *Sexology and Sexual Health / Revue Europeenne de Sexologie et de Sante Sexuelle*
623 2013;22:e59–63. <https://doi.org/https://dx.doi.org/10.1016/j.sexol.2012.08.002>.
- 624 [26] Bhatia T, Franzos MA, Wood JA, Nimgaonkar VL, Deshpande SN. Gender and procreation
625 among patients with schizophrenia. *Schizophr Res* 2004;68:387–94.
626 <https://doi.org/https://dx.doi.org/10.1016/j.schres.2003.08.009>.
- 627 [27] Bram N, Rafrafi R, Abdelghaffar W, Lakhali MH, Ouanes S, El Hechmi Z. Sexual dysfunctions in
628 Tunisian patients with schizophrenia. *Sexologies: European Journal of Sexology and Sexual*
629 *Health / Revue Europeenne de Sexologie et de Sante Sexuelle* 2014;23:e65–70.
630 <https://doi.org/https://dx.doi.org/10.1016/j.sexol.2014.05.004>.

- 631 [28] Bursalioglu FS, Aydin B, Yazici E. The correlation between psychiatric disorders and women's
632 lives. *Journal of Clinical and Diagnostic Research* 2013;7:695–9.
633 <https://doi.org/https://dx.doi.org/10.7860/JCDR/2013/5635.2884>.
- 634 [29] Caqueo-Urizar A, Fond G, Urzua A, Boyer Guillaume; ORCID: [https://orcid.org/0000-0003-](https://orcid.org/0000-0003-3249-2030)
635 3249-2030 AI - Urzua, Alfonso; ORCID: <https://orcid.org/0000-0002-0882-2194> LAI-F. Gender
636 differences in schizophrenia: A multicentric study from three Latin-America countries.
637 *Psychiatry Res* 2018;266:65–71.
638 <https://doi.org/https://dx.doi.org/10.1016/j.psychres.2018.05.032>.
- 639 [30] Carey MP, Ravi V, Chandra PS, Desai A, Neal DJ. Prevalence of HIV, hepatitis B, syphilis, and
640 chlamydia among adults seeking treatment for a mental disorder in southern India. *AIDS*
641 *Behav* 2007;11:289–97. <https://doi.org/10.1007/s10461-006-9134-2>.
- 642 [31] Carmo RA, Campos LN, Melo APS, Guimaraes MDC. Hepatitis C among patients with mental
643 illness in Brazil: an analysis of associated factors. *Gen Hosp Psychiatry* 2013;35:129–33.
644 <https://doi.org/https://dx.doi.org/10.1016/j.genhosppsy.2012.11.005>.
- 645 [32] Carmo RA, Melo APS, Dezanet LNC, De Oliveira HN, Cournos F, Guimarães MDC. Correlates of
646 hepatitis B among patients with mental illness in Brazil. *Gen Hosp Psychiatry* 2014;36:398–
647 405. <https://doi.org/10.1016/j.genhosppsy.2014.03.001>.
- 648 [33] Ceylan B, Kocoglu-Tanyer D. An Examination of Turkish Nurses' Attitudes, Awareness and
649 Practices Regarding Reproductive Health Needs of Individuals With Schizophrenia. *Issues*
650 *Ment Health Nurs* 2019;40:413–20. <https://doi.org/10.1080/01612840.2018.1509404>.
- 651 [34] Chandra PS, Carey MP, Carey KB, Shalinianant MP, Thomas T. Sexual Coercion and Abuse
652 Among Women with a Severe Mental Illness in India: An Exploratory Investigation. vol. 44.
653 2003.
- 654 [35] Chandra PS, Deepthivarma S, Carey MP, Carey KB, Shalinianant MP. A Cry from the Darkness:
655 Women with Severe Mental Illness in India Reveal Their Experiences with Sexual Coercion.
656 *Compr Psychiatry* 2003.
- 657 [36] Chandra PS, Carey MP, Carey KB, Rao P, Mbbs J, Thomas T. HIV risk behaviour among
658 psychiatric inpatients: results from a hospital-wide screening study in southern India. *Compr*
659 *Psychiatry* 2003.
- 660 [37] Chandra PS, Krishna VAS, Carey MP. Improving Knowledge about HIV and AIDS Among
661 Persons with a Severe Mental Illness in India. *Indian Journal of Soc Psychiatry* 2006;22:104–9.
- 662 [38] Collins PY, Holman AR, Freeman MC, Patel V. What is the relevance of mental health to
663 HIV/AIDS care and treatment programs in developing countries? A systematic review. *AIDS*
664 2006;20:1571–82.
- 665 [39] Collins PY, Berkman A, Mestry K, Pillai A. HIV prevalence among men and women admitted to
666 a South African public psychiatric hospital. *AIDS Care* 2009;21:863–7.
667 <https://doi.org/https://dx.doi.org/10.1080/09540120802626188>.
- 668 [40] Collins PY. Dual Taboos: Sexuality and Women with Severe Mental Illness in South Africa.
669 *Perceptions of Mental Health Care Providers*. vol. 5. 2001.
- 670 [41] Posada Correa AM, Andrade Carrillo RA, Suarez Vega DC, Gómez Cano S, Agudelo Arango LG,
671 Tabares Builes LF, et al. Sexual and Reproductive Health in Patients with Schizophrenia and

- 672 Bipolar Disorder. *Rev Colomb Psiquiatr* 2020;49:15–22.
673 <https://doi.org/10.1016/j.rcp.2018.04.007>.
- 674 [42] de Oliveira HN, Machado CJ, Guimaraes MDC. Factors associated with self-report of sexual
675 violence against men and women with mental disorders in Brazil. *Soc Psychiatry Psychiatr
676 Epidemiol* 2012;47:1567–79. <https://doi.org/https://dx.doi.org/10.1007/s00127-011-0463-8>.
- 677 [43] Desai G, Chandra P. Ethical issues in treating pregnant women with severe mental illness.
678 *Indian J Med Ethics* 2019;2.
- 679 [44] Dođu B, Güler J, Čýtak S, Altunkaynak Y, Alpay N, Cankatan Týp Merkezi Ö, et al. Bizofreni
680 Hastalarýnýn Cinsel Yařama Ýliřkin Öznel Dűbűncelerinin Cinsiyetler Arasý Karbýlařtırmasý
681 Gender Differences in Subjective Thoughts of Schizophrenia Patients About Their Sexuality.
682 *Klinik Psikiyatri* 2012;15:238–74.
- 683 [45] Dutra MRT, Campos LN, Guimaraes MDC. Sexually transmitted diseases among psychiatric
684 patients in Brazil. *Braz J Infect Dis* 2014;18:13–20.
685 <https://doi.org/https://dx.doi.org/10.1016/j.bjid.2013.04.004>.
- 686 [46] Eroglu MZ, Lus MG. Impulsivity, unplanned pregnancies, and contraception among women
687 with bipolar disorder. *Neuropsychiatr Dis Treat* 2020;16:407–14.
688 <https://doi.org/10.2147/NDT.S238887>.
- 689 [47] Esan O, Esan A. Sexual Dysfunction Among Patients With Schizophrenia in Southwest Nigeria.
690 *J Sex Marital Ther* 2018;44:657–66.
691 <https://doi.org/https://dx.doi.org/10.1080/0092623X.2018.1447055>.
- 692 [48] Fanta T, Haile K, Abebaw D, Assefa D, Hibdye G. Assessment of sexual dysfunction and
693 associated factors among patients with schizophrenia in Ethiopia, 2017. *BMC Psychiatry*
694 2018;18. <https://doi.org/10.1186/s12888-018-1738-3>.
- 695 [49] Gebeyehu DA, Mulatie M. Risky sexual behavior and its associated factors among patients
696 with severe mental disorder in University of Gondar Comprehensive Specialized Hospital,
697 2018. *BMC Psychiatry* 2021;21. <https://doi.org/10.1186/s12888-021-03054-z>.
- 698 [50] Ghebrehiwet S, Baul T, Restivo JL, Kelkile TS, Stevenson A, Gelaye B, et al. Gender-specific
699 experiences of serious mental illness in rural Ethiopia: A qualitative study. *Global Public
700 Health: An International Journal for Research, Policy and Practice* 2020;15:185–99.
701 <https://doi.org/https://dx.doi.org/10.1080/17441692.2019.1680723>.
- 702 [51] Grover S, Sahoo S, Chaudhary S, Chakrabarti S, Nehra R, Avasthi A. Gender differences, family
703 size and fertility rate among patients with bipolar disorder: A study from India. *Psychiatry Res*
704 2019;272:562–8. <https://doi.org/10.1016/j.psychres.2018.12.156>.
- 705 [52] Guimaraes MDC, McKinnon K, Campos LN, Melo APS, Wainberg M. HIV risk behavior of
706 psychiatric patients with mental illness: a sample of Brazilian patients. *Braz J Psychiatry*
707 2010;32:351–60.
- 708 [53] Guimaraes MDC, McKinnon K, Cournos F, Machado CJ, Melo APS, Campos LN, et al.
709 Correlates of HIV infection among patients with mental illness in Brazil. *AIDS Care*
710 2014;26:505–13. <https://doi.org/https://dx.doi.org/10.1080/09540121.2013.832722>.
- 711 [54] Hall T, Kakuma R, Palmer L, Minas H, Martins J, Kermode M. Social inclusion and exclusion of
712 people with mental illness in Timor-Leste: a qualitative investigation with multiple

- 713 stakeholders. BMC Public Health 2019;19:702.
714 <https://doi.org/https://dx.doi.org/10.1186/s12889-019-7042-4>.
- 715 [55] Halouani N, Ellouze S, Aloulou J, Charfeddine F, Aribi L, Amami O. Sexuality of schizophrenic
716 women in Tunisia: Case-control study. *Sexualite de la femme schizophrene en Tunisie*: Etude
717 cas-temoins 2018;27:211–6. <https://doi.org/https://dx.doi.org/10.1016/j.sexol.2017.12.003>.
- 718 [56] Hariri AG, Karadag F, Gurol DT, Aksoy UM, Tezcan AE. Sexual problems in a sample of the
719 Turkish psychiatric population. *Compr Psychiatry* 2009;50:353–60.
720 <https://doi.org/10.1016/j.comppsy.2008.09.009>.
- 721 [57] Henning MP, Krüger C, Fletcher L. HIV sero-positivity in recently admitted and long-term
722 psychiatric in-patients: Prevalence and diagnostic profile. *African Journal of Psychiatry (South
723 Africa)* 2012;15:47–53. <https://doi.org/10.4314/ajpsy.v15i1.7>.
- 724 [58] Hocaoglu C, Celik FH, Kandemir G, Guveli H, Bahceci B. Sexual dysfunction in outpatients with
725 schizophrenia in Turkey: A cross-sectional study. *Shanghai Arch Psychiatry* 2014;26:347–56.
- 726 [59] Hou C-L, Zang Y, Rosen RC, Cai M-Y, Li Y, Jia F-J, et al. Sexual dysfunction and its impact on
727 quality of life in Chinese patients with schizophrenia treated in primary care. *Compr
728 Psychiatry* 2016;65:116–21.
729 <https://doi.org/https://dx.doi.org/10.1016/j.comppsy.2015.11.002>.
- 730 [60] Huang Y-H, Hou C-L, Ng CH, Chen X, Wang Q-W, Huang Z-H, et al. Sexual dysfunction in
731 Chinese rural patients with schizophrenia. *BMC Psychiatry* 2019;19:218.
732 <https://doi.org/https://dx.doi.org/10.1186/s12888-019-2205-5>.
- 733 [61] Incedere A. Sexual Life and Associated Factors in Psychiatric Patients. *Sex Disabil* 2017;35:89–
734 106. <https://doi.org/https://dx.doi.org/10.1007/s11195-017-9475-y>.
- 735 [62] Joska JA, Obayemi AJ, Cararra H, Sorsdahl K. Severe mental illness and retention in anti-
736 retroviral care: a retrospective study. *AIDS Behav* 2014;18:1492–500.
737 <https://doi.org/https://dx.doi.org/10.1007/s10461-014-0709-z>.
- 738 [63] Kazour F, Obeid S, Hallit Francois; ORCID: <https://orcid.org/0000-0001-9337-4797> AI - Obeid,
739 Sahar; ORCID: <https://orcid.org/0000-0002-0735-4704> AI - Hallit, Souheil; ORCID:
740 <https://orcid.org/0000-0001-6918-5689> SAI-K. Sexual desire and emotional reactivity in
741 chronically hospitalized Lebanese patients with schizophrenia. *Perspect Psychiatr Care*
742 2020;56:502–7. <https://doi.org/https://dx.doi.org/10.1111/ppc.12455>.
- 743 [64] Kesebir S, Toprak B, Baykaran B, Hariri A, Bilici M. Comparison of sexual function and
744 hormonal parameters between mood stabilizer treatment modalities in bipolar disorder. *Iki
745 uclu bozuklukta duygudurum duzenleyici tedavi modaliteleri arasinda cinsel islev ve hormonal
746 degiskenlerin karsilastirilmesi* 2014;51:242–7.
747 <https://doi.org/https://dx.doi.org/10.4274/npa.y6838>.
- 748 [65] Kumar NS, Radhika MK, Suresh R, Uvais NA. Comparative Study of Sexual Side Effects in
749 Female Patients With Schizophrenia Receiving Risperidone or Olanzapine. *Primary Care
750 Companion for CNS Disorders* 2021;23. <https://doi.org/10.4088/PCC.20m02835>.
- 751 [66] Loganathan S, Murthy RS. Living with schizophrenia in India: Gender perspectives. *Transcult
752 Psychiatry* 2011;48:569–84. <https://doi.org/10.1177/1363461511418872>.

- 753 [67] Lundberg P, Johansson E, Okello E, Allebeck P, Thorson A. Sexual risk behaviours and sexual
754 abuse in persons with severe mental illness in Uganda: A qualitative study. *PLoS One* 2012;7.
755 <https://doi.org/10.1371/journal.pone.0029748>.
- 756 [68] Lundberg P, Nakasujja N, Musisi S, Thorson AE, Cantor-Graae E, Allebeck P. Sexual Risk
757 Behavior, Sexual Violence, and HIV in Persons With Severe Mental Illness in Uganda: Hospital-
758 Based Cross-Sectional Study and National Comparison Data. *Am J Public Health*
759 2015;105:1142–8. <https://doi.org/https://dx.doi.org/10.2105/AJPH.2014.302479>.
- 760 [69] Lundberg P, Nakasujja N, Musisi S, Thorson AE, Cantor-Graae E, Allebeck P. HIV prevalence in
761 persons with severe mental illness in Uganda: a cross-sectional hospital-based study. *Int J*
762 *Ment Health Syst* 2013;7:20. <https://doi.org/https://dx.doi.org/10.1186/1752-4458-7-20>.
- 763 [70] Madziro-Ruwizhu TB, Nyagura T, Takarinda K, Jack HE, Kajawu L, Mangezi W. HIV Infection in
764 Attendees of Psychiatric Clinics in Harare, Zimbabwe; Prevalence, Associated Factors and HIV
765 Care Uptake. *AIDS Behav* 2019;23:3471–81.
766 <https://doi.org/https://dx.doi.org/10.1007/s10461-019-02633-2>.
- 767 [71] Magalhaes PV da S, Kapczynski F, Kauer-Sant’Anna M. Use of contraceptive methods among
768 women treated for bipolar disorder. *Arch Womens Ment Health* 2009;12:183–5.
769 <https://doi.org/https://dx.doi.org/10.1007/s00737-009-0060-y>.
- 770 [72] Maling S, Todd J, Van Paal L, Grosskurth H, Kinyanda E. HIV-1 seroprevalence and risk factors
771 for HIV infection among first-time psychiatric admissions in Uganda. *AIDS Care - Psychological*
772 *and Socio-Medical Aspects of AIDS/HIV* 2011;23:171–8.
773 <https://doi.org/10.1080/09540121.2010.498939>.
- 774 [73] Mamabolo MM, Magagula TG, Kruger C, Fletcher L. A survey of risk behaviour for contracting
775 HIV among adult psychiatric patients. A South African study - Part 1. *Afr J Psychiatry*
776 (Johannesbg) 2012;15:329–34. <https://doi.org/https://dx.doi.org/10.4314/ajpsy.v15i5.40>.
- 777 [74] Marengo E, Martino DJ, Igoa A, Scápola M, Fassi G, Baamonde MU, et al. Unplanned
778 pregnancies and reproductive health among women with bipolar disorder. *J Affect Disord*
779 2015;178:201–5. <https://doi.org/10.1016/j.jad.2015.02.033>.
- 780 [75] Marengo E, Martino DJ, Igoa A, Fassi G, Scápola M, Urtueta Baamonde M, et al. Sexual risk
781 behaviors among women with bipolar disorder. *Psychiatry Res* 2015;230:835–8.
782 <https://doi.org/10.1016/j.psychres.2015.10.021>.
- 783 [76] Mashapu S, Mkize DL. HIV seropositivity in patients with first-episode psychosis. *The South*
784 *African Journal of Psychology* 2007;13:90–4.
- 785 [77] Mere SM, Paruk S. A chart review of human immunodeficiency virus status in patients
786 admitted with psychosis in Durban, South Africa. *South African Journal of Psychiatry* 2018;24.
787 <https://doi.org/10.4102/sajpsy.2018.24i0.1129>.
- 788 [78] Mirsepassi Z, Raisi F, Shahvari Z, FirooziKhojastefar R, Yahyavi ST. Evaluating family
789 knowledge about sexual health in patients with severe mental illness: a qualitative study in
790 Iran. *BMC Psychiatry* 2022;22:174. <https://doi.org/https://dx.doi.org/10.1186/s12888-022-03788-4>.
- 792 [79] Mpango RS, Ssembajjwe W, Rukundo GZ, Birungi C, Kalungi A, Gadown KD, et al. Physical and
793 psychiatric comorbidities among patients with severe mental illness as seen in Uganda. *Eur*

- 794 Arch Psychiatry Clin Neurosci 2022. [https://doi.org/https://dx.doi.org/10.1007/s00406-022-](https://doi.org/https://dx.doi.org/10.1007/s00406-022-01478-6)
795 01478-6.
- 796 [80] Mwelase MP, Ntlantsana V, Tomita A, Chiliza B, Paruk S. HIV prevalence and access to HIV
797 testing and care in patients with psychosis in South Africa. South African Journal of Psychiatry
798 2023;29. <https://doi.org/10.4102/sajpsychiatry.v29i0.1918>.
- 799 [81] Nakhli J, El Kissi Y, Bouhlel S, Amamou B, Nabli TA, Nasr S Ben, et al. Reliability and validity of
800 the Arizona Sexual Experiences Scale-Arabic version in Tunisian patients with schizophrenia.
801 Compr Psychiatry 2014;55:1473–7.
802 <https://doi.org/https://dx.doi.org/10.1016/j.comppsy.2014.04.006>.
- 803 [82] Nakigudde J, Ehnvall A, Mirembe F, Musisi S, Airaksinen E. An exploratory study on the
804 feasibility and appropriateness of family psychoeducation for postpartum women with
805 psychosis in Uganda. BMC Psychiatry 2013;13:131.
806 <https://doi.org/https://dx.doi.org/10.1186/1471-244X-13-131>.
- 807 [83] Negash B, Asmamewu B, Alemu WG. Risky sexual behaviors of schizophrenic patients: A
808 single center study in Ethiopia, 2018. BMC Res Notes 2019;12.
809 <https://doi.org/10.1186/s13104-019-4673-6>.
- 810 [84] Obo CS, Sori LM, Abegaz TM, Molla BT. Risky sexual behavior and associated factors among
811 patients with bipolar disorders in Ethiopia. BMC Psychiatry 2019;19:313.
812 <https://doi.org/https://dx.doi.org/10.1186/s12888-019-2313-2>.
- 813 [85] Olisah VO, Sheikh TL, Abah ER, Mahmud-Ajeigbe AF. Sociodemographic and clinical correlates
814 of sexual dysfunction among psychiatric outpatients receiving common psychotropic
815 medications in a Neuropsychiatric Hospital in Northern Nigeria. Niger J Clin Pract
816 2016;19:799–806. <https://doi.org/https://dx.doi.org/10.4103/1119-3077.180063>.
- 817 [86] Opondo PR, Ho-Foster AR, Ayugi J, Hatitchki B, Pumar M, Bilker WB, et al. HIV Prevalence
818 Among Hospitalized Patients at the Main Psychiatric Referral Hospital in Botswana. AIDS
819 Behav 2018;22:1503–16. <https://doi.org/https://dx.doi.org/10.1007/s10461-017-1878-3>.
- 820 [87] Özcan NK, Boyacıoğlu NE, Enginkaya S, Dinç H, Bilgin H. Reproductive health in women with
821 serious mental illnesses. J Clin Nurs 2014;23:1283–91. <https://doi.org/10.1111/jocn.12332>.
- 822 [88] Ozcan NK, Boyacioglu NE, Dikeç G, Dinc H, Enginkaya S, Tomruk N. Prenatal and Postnatal
823 Attachment Among Turkish Mothers Diagnosed with a Mental Health Disorder. Issues Ment
824 Health Nurs 2018;39:795–801.
825 <https://doi.org/https://dx.doi.org/10.1080/01612840.2018.1455773>.
- 826 [89] Pehlivanoglu K, Tanriover O, Tomruk N, Karamustafalioglu N, Oztekin E, Alpay N. Family
827 Planning Needs and Contraceptive Use in Female Psychiatric Outpatients. TURKISH JOURNAL
828 OF FAMILY MEDICINE AND PRIMARY CARE (TJFMPC) 2007;1.
- 829 [90] Pinto D de S, Mann CG, Wainberg M, Mattos P, Broxado de Oliveira S. Sexuality, vulnerability
830 to HIV, and mental health: an ethnographic study of psychiatric institutions. Cad Saúde
831 Pública 2007;23:2224–33. <https://doi.org/10.1590/s0102-311x2007000900030>.
- 832 [91] Poreddi V, Sai Nikhil Reddy S, Gandhi S, Marimuthu P, BadaMath S. “Unheard voices”:
833 Perceptions of women with mental illness on nurses screening routinely for domestic

834 violence: A qualitative analysis. *Invest Educ Enferm* 2021;39.
835 <https://doi.org/10.17533/udea.iee.v39n3e03>.

836 [92] Rezaie L, Phillips D. Post-discharge needs of Iranian women diagnosed with severe mental
837 illness: A qualitative study. *J Psychiatr Ment Health Nurs* 2020;27:752–62.
838 <https://doi.org/10.1111/jpm.12634>.

839 [93] Sethuraman B. Factors associated with contraceptive use in women with schizophrenia. *Arch
840 Womens Ment Health* 2019;22:705–6. [https://doi.org/https://dx.doi.org/10.1007/s00737-
841 019-00996-y](https://doi.org/https://dx.doi.org/10.1007/s00737-019-00996-y).

842 [94] Shahvari Z, Raisi F, Firoozi Khojastefar R, Yahyavi ST, Mirsepassi Z. Memos on Sexuality
843 Education Effectiveness; A Qualitative Study in Severe Mental Illness in Iran. *International
844 Journal of Sexual Health* 2020;32:254–66. <https://doi.org/10.1080/19317611.2020.1793047>.

845 [95] Sibanyoni AU, Joubert M. Are female bipolar patients of reproductive age aware of the
846 teratogenic risk of sodium valproate? A qualitative study. *South African Journal of Psychiatry*
847 2022;28:a1719. <https://doi.org/https://dx.doi.org/10.4102/sajpsychiatry.v28i0.1719>.

848 [96] Simiyon M, Chandra PS, Desai G. Sexual dysfunction among women with Schizophrenia-A
849 cross sectional study from India. *Asian J Psychiatr* 2016;24:93–8.
850 <https://doi.org/https://dx.doi.org/10.1016/j.ajp.2016.08.022>.

851 [97] Singh D, ChB M, Berkman A, Bresnahan M. Seroprevalence and HIV-associated factors among
852 adults with severe mental illness-a vulnerable population. *South African Medical Journal*
853 2009.

854 [98] Souaiby L, Kazour F, Zoghbi M, Bou Khalil R, Richa S. Sexual dysfunction in patients with
855 schizophrenia and schizoaffective disorder and its association with adherence to
856 antipsychotic medication. *Journal of Mental Health* 2020;29:623–30.
857 <https://doi.org/https://dx.doi.org/10.1080/09638237.2019.1581333>.

858 [99] Souto Melo AP, Machado CJ, Crosland Guimaraes MD. HIV testing in psychiatric patients in
859 Brazil. *J Acquir Immune Defic Syndr* 2011;57 Suppl 3:S157-63.
860 <https://doi.org/https://dx.doi.org/10.1097/QAI.0b013e31821e9be1>.

861 [100] Terzian ACC, Andreoli SB, Razzouk D, Chaves AC. Fertility and fecundity of an outpatient
862 sample with schizophrenia. *Revista Brasileira de Psiquiatria* 2006;28:305–7.
863 <https://doi.org/http://dx.doi.org/10.1590/S1516-44462006005000018>.

864 [101] Tharoor H, Kaliappan A, Gopal S. Sexual dysfunctions in schizophrenia: Professionals and
865 patients perspectives. *Indian J Psychiatry* 2015;57:85–7. [https://doi.org/10.4103/0019-
866 5545.148532](https://doi.org/10.4103/0019-5545.148532).

867 [102] Tomruk N, Tanriover O. HIGH RISK BEHAVIOURS FOR SEXUALLY TRANSMITTED DISEASES IN
868 FEMALE PSYCHIATRIC OUTPATIENTS IN TURKEY review View project Yaşlılıkta Beslenme ve
869 Diyet View project. 2006.

870 [103] Tumwakire E, Arnd H, Gavamukulya Y. A qualitative exploration of Ugandan mental health
871 care workers' perspectives and experiences on sexual and reproductive health of people
872 living with mental illness in Uganda. *BMC Public Health* 2022;22:1722.
873 <https://doi.org/https://dx.doi.org/10.1186/s12889-022-14128-2>.

- 874 [104] Tunde-Ayinmode MF. Current knowledge and pattern of use of family planning methods
875 among a severely ill female Nigerian psychiatric outpatients: Implication for existing service.
876 *Ann Afr Med* 2013.
- 877 [105] Wainberg ML, Alfredo Gonzalez M, McKinnon K, Elkington KS, Pinto D, Gruber Mann C.
878 Targeted ethnography as a critical step to inform cultural adaptations of HIV prevention
879 interventions for adults with severe mental illness. *Soc Sci Med* 2007;65:296–308.
880 <https://doi.org/https://dx.doi.org/10.1016/j.socscimed.2007.03.020>.
- 881 [106] Wainberg ML, McKinnon K, Elkington KS, Mattos PE, Mann CG, Pinto D de S, et al. HIV risk
882 behaviors among outpatients with severe mental illness in Rio de Janeiro, Brazil. *World*
883 *Psychiatry* 2008;7:166–72. [https://doi.org/http://dx.doi.org/10.1002/j.2051-](https://doi.org/http://dx.doi.org/10.1002/j.2051-5545.2008.tb00190.x)
884 [5545.2008.tb00190.x](https://doi.org/http://dx.doi.org/10.1002/j.2051-5545.2008.tb00190.x).
- 885 [107] Yang J wei, Yu K, Wang X qing, Wang Y, Zhang CC, Ma R, et al. Sexual needs of people with
886 schizophrenia: a descriptive phenomenological study. *BMC Psychiatry* 2023;23.
887 <https://doi.org/10.1186/s12888-023-04640-z>.
- 888 [108] Yu K, Wang Y, Wang X, Ma R, Li Y, Zhou Y. Experience of reproductive concerns in women
889 with schizophrenia: A descriptive phenomenological study. *Int J Nurs Stud* 2022;135:1–7.
890 <https://doi.org/https://dx.doi.org/10.1016/j.ijnurstu.2022.104343>.
- 891 [109] Zerihun T, Bekele D, Birhanu E, Worku Y, Deyesa N, Tesfaye M. Family planning awareness,
892 utilization and associated factors among women of reproductive age attending psychiatric
893 outpatient care, a cross- sectional study, Addis Ababa, Ethiopia. *PLoS One* 2020;15:e0238766.
894 <https://doi.org/https://dx.doi.org/10.1371/journal.pone.0238766>.
- 895 [110] Zerihun T, Sorsdahl K, Hanlon C. Family planning for women with severe mental illness in
896 rural Ethiopia: a qualitative study. *Reprod Health* 2021;18. [https://doi.org/10.1186/s12978-](https://doi.org/10.1186/s12978-021-01245-1)
897 [021-01245-1](https://doi.org/10.1186/s12978-021-01245-1).
- 898 [111] Zerihun T, Tesfaye M, Deyessa N, Bekele D. Intimate partner violence among reproductive-
899 age women with chronic mental illness attending a psychiatry outpatient department: Cross-
900 sectional facility-based study, Addis Ababa, Ethiopia. *BMJ Open* 2021;11.
901 <https://doi.org/10.1136/bmjopen-2020-045251>.
- 902 [112] Zhang Q, Xie JJ. Association between schizophrenia and syphilis: A retrospective study in
903 Xiamen, China. *BMC Psychiatry* 2018;18. <https://doi.org/10.1186/s12888-018-1869-6>.
- 904 [113] Zhang Y, Tang Z, Ruan Y, Huang C, Wu J, Lu Z, et al. Prolactin and Thyroid Stimulating
905 Hormone (TSH) Levels and Sexual Dysfunction in Patients with Schizophrenia Treated with
906 Conventional Antipsychotic Medication: A Cross-Sectional Study. *Med Sci Monit*
907 2018;24:9136–43. <https://doi.org/https://dx.doi.org/10.12659/MSM.913759>.
- 908 [114] The World Bank. Life Expectancy at Birth (Total) - Low and Middle Income Countries .
909 [https://DataWorldbankOrg/Indicator/SpDynLe00.in?End=2021&locations=xo&start=1960&vi](https://DataWorldbankOrg/Indicator/SpDynLe00.in?End=2021&locations=xo&start=1960&view=chart)
910 [ew=chart](https://DataWorldbankOrg/Indicator/SpDynLe00.in?End=2021&locations=xo&start=1960&view=chart) 2022.
- 911 [115] Hughes E, Bassi S, Gilbody S, Bland M, Martin F. Prevalence of HIV, hepatitis B, and hepatitis C
912 in people with severe mental illness: a systematic review and meta-analysis. *Lancet*
913 *Psychiatry* 2016;3:40–8. [https://doi.org/10.1016/S2215-0366\(15\)00357-0](https://doi.org/10.1016/S2215-0366(15)00357-0).

- 914 [116] Hughes E, Bassi S, Gilbody S, Bland M, Martin F. Prevalence of HIV, hepatitis B, and hepatitis C
915 in people with severe mental illness: a systematic review and meta-analysis. *Lancet*
916 *Psychiatry* 2016;3:40–8. [https://doi.org/10.1016/S2215-0366\(15\)00357-0](https://doi.org/10.1016/S2215-0366(15)00357-0).
- 917 [117] Khalifeh H, Oram S, Osborn D, Howard LM, Johnson S. Recent physical and sexual violence
918 against adults with severe mental illness: a systematic review and meta-analysis.
919 *International Review of Psychiatry* 2016;28:433–51.
920 <https://doi.org/10.1080/09540261.2016.1223608>.
- 921 [118] Thornicroft G, Sunkel C, Alikhon Aliev A, Baker S, Brohan E, el Chammay R, et al. The Lancet
922 Commission on ending stigma and discrimination in mental health. *The Lancet*
923 2022;400:1438–80. [https://doi.org/10.1016/S0140-6736\(22\)01470-](https://doi.org/10.1016/S0140-6736(22)01470-2/ATTACHMENT/89C56927-C832-4D5C-ACFB-15D3517E058C/MMC3.PDF)
924 [2/ATTACHMENT/89C56927-C832-4D5C-ACFB-15D3517E058C/MMC3.PDF](https://doi.org/10.1016/S0140-6736(22)01470-2/ATTACHMENT/89C56927-C832-4D5C-ACFB-15D3517E058C/MMC3.PDF).
- 925 [119] Davis S, Pinfold V, Catchpole J, Lovelock C, Senthil B, Kenny A. Reporting lived experience
926 work. *Lancet Psychiatry* 2024;11:8–9. [https://doi.org/10.1016/S2215-0366\(23\)00402-9](https://doi.org/10.1016/S2215-0366(23)00402-9).
- 927