

## **Research article: Reevaluation of the Bicycle Ergometry in the Diagnosis of Ischemic Heart Disease**

**Authors' list: Basheer Abdullah Marzoog<sup>1,\*</sup>, Peter Chomakhidze<sup>1</sup>, Philipp Kopylov<sup>1</sup>**

<sup>1</sup> World-Class Research Center «Digital Biodesign and Personalized Healthcare», I.M. Sechenov First Moscow State Medical University (Sechenov University), 119991 Moscow, Russia; postal address: Russia, Moscow, 8-2 Trubetskaya street, 119991.

\*Corresponding author: Basheer Abdullah Marzoog, MD, Ph.D. cardiology student at Sechenov First Moscow State Medical University at the World-Class Research Center "Digital biodesign and personalized healthcare". (marzug@mail.ru, +79969602820). Address: Russia, Moscow, 8-2 Trubetskaya street, 119991. Postal address: Russia, Moscow, 8-2 Trubetskaya street, 119991. ORCID: 0000-0001-5507-2413, Scopus ID: 57486338800

Competing interests: No competing interests regarding the publication.

## **Abstract**

**Background:** Ischemic heart disease (IHD) and related complications and outcomes remain the most frequently reported cardiovascular events. However, the diagnosis and prevention of ischemic heart disease primary depends on the clinical manifestations and electrocardiography changes.

**Objectives:** Patients with IHD are victims of the poor diagnostic methods and the prevention strategies.

**Aims:** To assess the diagnostic accuracy of the used bicycle ergometry in the diagnosis of ischemic heart disease based on the results of the stress computed myocardial perfusion (CTP) imaging with vasodilator test (adenosine triphosphate).

**Materials and methods:** A single center prospective, and non-randomized study included participations aged  $\geq 40$ . The study included 38 participants with vs without IHD confirmed by CTP. The participants done bicycle ergometry following Bruce protocol then performed CTP. For statistical analysis used the descriptive statistics, Pearson Correlation test, Student test. The Statistica 12 program used for the statistical test.

**Results:** The study included 38 participants included 19 (50 %) patients with positive stress induced myocardial perfusion defect on the CTP and 19 (50 %) with negative results. The mean age of the participants 58,77 years (std. div.  $\pm 9,664$ ). The bicycle ergometry specificity and sensitivity 61.11%, 57.89%, respectively. The diagnostic accuracy of the bicycle ergometry 59.46%.

**Conclusions:** The bicycle ergometry test characterized by severe limitations in terms of the IHD diagnosis.

**Others:** Recommended to combine additional diagnostic methods with the bicycle ergometry to improve it is diagnostic accuracy, such as single channel electrocardiography and exhaled breath analysis.

**Keywords:** Ischemic heart disease, Bicycle Ergometry, Stress Computed Tomography Myocardial Perfusion Imaging, Diagnosis

## **Introduction**

Ischemic heart disease (IHD) is one of the most common pathologies among the cardiovascular disease in terms of the mortality and morbidity <sup>[1]</sup>. This high prevalence of IHD in the community returns to the poor early detection and diagnostic methods of IHD. One of the most frequently used diagnostic method for individuals with clinical signs and symptoms of ischemic heart disease is the physical stress test, including bicycle ergometry <sup>[2]</sup>. The debate about the efficacy of this method is under evaluation since it is first use in clinical practice <sup>[3, 4]</sup>. However, this method is easy to use and safe as well as cost-effective. Therefore, this method is widespread as an elementary diagnostic method for IHD in compare to other expansive and invasive methods such as coronary angiography with Coronary Flow Reserve <sup>[5]</sup>.

The current paper revises the specificity and sensitivity of the bicycle ergometry for usage as an elementary tool for IHD diagnosis through performing research on a participant with specific inclusion and exclusion criteria (published on the clinical trial (NCT06181799)).

## **Materials and methods**

A non-randomized, single center, mini-invasive, parallel, cross-sectional, diagnostic, observational, case-control prospective cohort study performed at the University Clinical Hospital-1. The data of this study extracted from another clinical trial performed by Marzoog, the full design of the study published on [clinicaltrials.gov](https://clinicaltrials.gov) (NCT06181799) <sup>[6]</sup>. The study approved by the Sechenov University, Russia, from “Ethics Committee Requirement № 19-23 from 26.10.2023”.

The study included both males and females, and the age of the participants  $\geq 40$  years. The study participants passed exercise bicycle ergometry (on SCHILLER device; Bruce protocol) test to evaluate the response to physical activity. According to the results metabolic equivalent; Mets-BT (BT), the angina functional class (FC) in participants with positive physical stress test results determined, Where  $BT/Mets < 50/ < 4$  FC-III,  $BT/Mets 50-100/4-7$  FC-II,  $BT/Mets > 100/7$  FC-I. During the bicycle ergometry, the participants monitored with 12-lead ECG and manual blood pressure measurement, 1 measured at the end of each 2 minute.

The ergometry procedure discontinued if an increase in blood pressure  $\geq 220$  mmHg or horizontal or downsloping ST segment on the ECG  $\geq 1$  mm. Moreover, stop the procedure if the target heart rate (86% of the 220-age) is reached.

To perform the stressed myocardial perfusion computer tomography imaging, all the participants present results of the venous creatinine level, eGFR (estimated glomerular filtration rate) according to the 2021 CKD-EPI Creatinine > 30 ml/min/1,73 m<sup>2</sup>, according to the recommendation for using this formula by the National kidney foundation and the American Society of Nephrology [7-10].

The participants got catheterization in the basilar vein or the radial vein for injection of contrast and Natrii adenosine triphosphate (10 mg/1ml) to induce pharmacological stress test to the heart by increasing heart rate.

To prepare the Natrii adenosine triphosphate, an Adenosine triphosphate 3 ml dilute in 17 ml of Isotonic Sodium Chloride Solution 0.9%, the injected volume of the diluted drug in milliliters is calculated by body weight. For 1 dose, take 3 ml of adenosine triphosphate (3 ampoules of each 1 ml (10mg)) + 17 ml of isotonic solution of sodium chloride 0.9% in one syringe, 20 ml. For one patient, manually inject intravenously (IV) through the already inserted catheter at a rate of 300 µg/kg/2 minutes, depending on weight: 60 kg = 12 ml, 70 kg = 14 ml. 80 kg = 16 ml, 100 kg = 20 ml of the full dose.

Computer tomography (Canon with 640 slice, 0,5mm thickness) with contrast (Omnipaque, 350-100ml), inject the contrast, then make rest image for myocardial perfusion, then wait 20 minutes and then inject the Natrii adenosine triphosphate intravenously (10 mg/1ml) according to the body weight to cause pharmacological stress test to the heart during two minutes in to the catheter. Then make the second image of the myocardial perfusion after stress test immediately.

## Results

The descriptive statistic of the continuous variables demonstrated in table 1.

Table 1: Descriptive statistics of the presented sample.

Variable	Descriptive Statistics (All in primary data.stw)				
	Valid N	% Valid obs.	Mean	Minimum	Maximum
Age (years)	38	100.0000	58.77	42.05	77.94
Pulse rate (beat per minute)	38	100.0000	72.21	53.00	92.00
SBP (mmHg)	38	100.0000	122.47	100.00	155.00
DBP (mmHg)	38	100.0000	77.84	60.00	100.00
Body weight (Kg)	38	100.0000	75.92	52.50	118.30
Height (cm)	38	100.0000	171.29	148.00	190.00
BMI ((kg/m <sup>2</sup> )	38	100.0000	25.84	18.49	35.00
Goal heart rate (220-age)	38	100.0000	161.23	142.06	177.95
Max HR (beat per minute)	38	100.0000	148.08	108.00	199.00
Reached % on bicycle ergometry	38	100.0000	92.18	64.89	135.25
WT	38	100.0000	119.74	75.00	250.00
METs	38	100.0000	6.59	3.40	11.90
Creatinine (µmol/L)	38	100.0000	85.14	57.00	138.00
eGFR (2021 CKD-EPI Creatinine)	38	100.0000	82.04	45.40	107.00

The number of patients with normal body weight 17 (44.73684 %). overweight 15 (39.47368 %), and obesity first degree 6 (15.78947 %).

The categorical variables are presented as a relative and absolute values in table 2 (A-W)

Table 2: The table represents the categorical variables of the study in absolute and relative values.

2-A

Category	Frequency table: Smoking (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
No	33	33	86.84211	86.8421
Yes	5	38	13.15789	100.0000
Missing	0	38	0.00000	100.0000

2-B

Category	Frequency table: Concomitant disease (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
No	22	22	57.89474	57.8947
Yes	13	35	34.21053	92.1053
Missing	3	38	7.89474	100.0000

2-C

Category	Frequency table: Atherosclerosis of the coronary artery (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
No	19	19	50.00000	50.0000
Yes	19	38	50.00000	100.0000
Missing	0	38	0.00000	100.0000

2-D

Category	Frequency table: Hemodynamically significant (>60%) (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
No	33	33	86.84211	86.8421
Yes	5	38	13.15789	100.0000
Missing	0	38	0.00000	100.0000

2-E

Category	Frequency table: Myocardial perfusion defect after stress ATP (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
No	19	19	50.00000	50.0000

Category	Frequency table: Myocardial perfusion defect after stress ATP (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
Yes	19	38	50.00000	100.0000
Missing	0	38	0.00000	100.0000

2-G

Category	Frequency table: Myocardial perfusion defect before Stress ATP (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
Yes	13	13	34.21053	34.2105
No	25	38	65.78947	100.0000
Missing	0	38	0.00000	100.0000

2-H

Category	Frequency table: Atherosclerosis in other arteries (Yes/No) (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
No	18	18	47.36842	47.3684
Yes	15	33	39.47368	86.8421
Missing	5	38	13.15789	100.0000

2-I

Category	Frequency table: Atherosclerotic vascular (Namely) (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
Carotid, Brachiocephalic bifurcation	12	12	31.57895	31.5789
Carotid, Brachiocephalic bifurcation	2	14	5.26316	36.8421
Carotid	1	15	2.63158	39.4737
Missing	23	38	60.52632	100.0000

2-J

Category	Frequency table: Carotid (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
No	20	20	52.63158	52.6316
Yes	14	34	36.84211	89.4737
Missing	4	38	10.52632	100.0000

2-K

Category	Frequency table: Brachiocephalic (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
No	22	22	57.89474	57.8947

Category	Frequency table: Brachiocephalic (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
Yes	12	34	31.57895	89.4737
Missing	4	38	10.52632	100.0000

2-L

Category	Frequency table: Arterial Hypertension (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
Yes	21	21	55.26316	55.2632
No	17	38	44.73684	100.0000
Missing	0	38	0.00000	100.0000

2-M

Category	Frequency table: Stage AH (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
II	14	14	36.84211	36.8421
I	2	16	5.26316	42.1053
III	5	21	13.15789	55.2632
I	1	22	2.63158	57.8947
Missing	16	38	42.10526	100.0000

2-N

Category	Frequency table: Degree AH (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
III	5	5	13.15789	13.1579
II	5	10	13.15789	26.3158
I	11	21	28.94737	55.2632
II	1	22	2.63158	57.8947
Missing	16	38	42.10526	100.0000

2-O

Category	Frequency table: Risk of CVD (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
Moderate	14	14	36.84211	36.8421
High	10	24	26.31579	63.1579
Low	11	35	28.94737	92.1053
Very high	3	38	7.89474	100.0000
Missing	0	38	0.00000	100.0000

2-P

Category	Frequency table: SCAD (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent

Category	Frequency table: SCAD (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
No	18	18	47.36842	47.3684
Yes	3	21	7.89474	55.2632
Missing	17	38	44.73684	100.0000

## 2-Q

Category	Frequency table: Blood pressure reaction type (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
Normotonic	28	28	73.68421	73.6842
Mild hypertonic	2	30	5.26316	78.9474
Hypertonic	2	32	5.26316	84.2105
Asthenic	3	35	7.89474	92.1053
Hypotonic	3	38	7.89474	100.0000
Missing	0	38	0.00000	100.0000

## 2-R

Category	Frequency table: FC by WT (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
FC-II	7	7	18.42105	18.4211
FC-I	5	12	13.15789	31.5789
Missing	26	38	68.42105	100.0000

## 2-S

Category	Frequency table: FC by METs (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
FC-II	7	7	18.42105	18.4211
FC-I	4	11	10.52632	28.9474
FC-III	1	12	2.63158	31.5789
Missing	26	38	68.42105	100.0000

## 2-T

Category	Frequency table: Reaction type ( positive/negative) (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
Positive	12	12	31.57895	31.5789
Negative	17	29	44.73684	76.3158
Suspected	9	38	23.68421	100.0000
Missing	0	38	0.00000	100.0000

## 2-U

Category	Frequency table: Reason of discontinuation (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
Horizontal ST depression >1mm	6	6	15.78947	15.7895
Reach goal HR	32	38	84.21053	100.0000
Missing	0	38	0.00000	100.0000

2-V

Category	Frequency table: Tolerance to exertion (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
Close to high	4	4	10.52632	10.5263
Moderate	23	27	60.52632	71.0526
Very high	6	33	15.78947	86.8421
High	4	37	10.52632	97.3684
Low	1	38	2.63158	100.0000
Missing	0	38	0.00000	100.0000

2-W

Category	Frequency table: CKD stage (All in primary data.stw)			
	Count	Cumulative Count	Percent	Cumulative Percent
I	16	16	42.10526	42.1053
II	18	34	47.36842	89.4737
IIIa	4	38	10.52632	100.0000
Missing	0	38	0.00000	100.0000

The results of the student test of the sample represented in table 3 (A-M).

Table 3: The results of the student test of the sample represented. Only statistically significant difference represented in the table.

3-A

Variable	T-tests; Grouping: Gender (All in primary data.stw)					
	Group 1: F		Group 2: M		p	Valid N F
	Mean F	Mean M	t-value	df		
SBP rest	117.41	126.57	-2.40957	36	0.021210	17
Body weight	67.85	82.44	-3.58890	36	0.000981	17
Height	164.62	176.69	-4.83464	36	0.000025	17
WT	88.24	145.24	-4.82673	36	0.000025	17
METs	5.81	7.22	-2.24874	36	0.030743	17
Creatinine (µmol/L)	76.05	92.50	-3.65241	36	0.000820	17

Variable	T-tests; Grouping: Gender (All in primary data.stw)					
	Group 1: F		Group 2: M			

	Valid N M	Std.Dev. F	Std.Dev. M	F-ratio Variances	p Variances
SBP rest	21	12.654	10.782	1.377398	0.492763
Body weight	21	10.219	13.997	1.876169	0.205644
Height	21	8.795	6.600	1.775870	0.223898
WT	21	17.936	45.839	6.531701	0.000392
METs	21	1.206	2.358	3.822979	0.008835
Creatinine (µmol/L)	21	11.782	15.230	1.670764	0.301144

### 3-B

Variable	T-tests; Grouping: Concomitant disease (All in primary data.stw)					
	Group 1: No Group 2: Yes					
	Mean No	Mean Yes	t-value	df	p	Valid N No
Height	173.27	165.62	2.52705	33	0.016476	22
Reached %	88.58	97.49	-2.30415	33	0.027648	22
WT	130.68	98.08	2.13328	33	0.040428	22

Variable	T-tests; Grouping: Concomitant disease (All in primary data.stw)				
	Group 1: No Group 2: Yes				
	Valid N Yes	Std.Dev. No	Std.Dev. Yes	F-ratio Variances	p Variances
Height	13	8.062	9.622	1.424600	0.461098
Reached %	13	8.428	14.566	2.987445	0.027287
WT	13	52.288	21.558	5.882893	0.002912

### 3-C

Variable	T-tests; Grouping: Atherosclerosis of the coronary artery (All in primary data.stw)					
	Group 1: No Group 2: Yes					
	Mean No	Mean Yes	t-value	df	p	Valid N No
Age	53.13	64.41	-4.39659	36	0.000093	19
Goal heart rate	166.87	155.59	4.39659	36	0.000093	19
Creatinine (µmol/L)	79.40	90.89	-2.35327	36	0.024191	19

Variable	T-tests; Grouping: Atherosclerosis of the coronary artery (All in primary data.stw)				
	Group 1: No Group 2: Yes				
	Valid N Yes	Std.Dev. No	Std.Dev. Yes	F-ratio Variances	p Variances
Age	19	6.825	8.850	1.68168	0.279471
Goal heart rate	19	6.825	8.850	1.68168	0.279471
Creatinine (µmol/L)	19	12.642	17.114	1.83250	0.208473

### 3-C

Variable	T-tests; Grouping: Hemodynamically significant (>60%) (All in primary data.stw) Group 1: No Group 2: Yes					
	Mean No	Mean Yes	t-value	df	p	Valid N No
SBP rest	120.36	136.40	-2.96948	36	0.005282	33
Max HR	150.79	130.20	2.97336	36	0.005229	33

Variable	T-tests; Grouping: Hemodynamically significant (>60%) (All in primary data.stw) Group 1: No Group 2: Yes				
	Valid N Yes	Std.Dev. No	Std.Dev. Yes	F-ratio Variances	p Variances
SBP rest	5	10.911	13.686	1.573274	0.410528
Max HR	5	13.713	19.215	1.963335	0.247939

### 3-E

Variable	T-tests; Grouping: Myocardial perfusion defect before Stress ATP (All in primary data.stw) Group 1: Yes Group 2: No					
	Mean Yes	Mean No	t-value	df	p	Valid N Yes
Max HR	139.15	152.72	-2.70213	36	0.010442	13
Reached %	86.04	95.37	-2.46004	36	0.018825	13

Variable	T-tests; Grouping: Myocardial perfusion defect before Stress ATP (All in primary data.stw) Group 1: Yes Group 2: No				
	Valid N No	Std.Dev. Yes	Std.Dev. No	F-ratio Variances	p Variances
Max HR	25	14.899	14.573	1.045192	0.886020
Reached %	25	9.231	11.919	1.667134	0.357847

### 3-F

Variable	T-tests; Grouping: Atherosclerosis in other arteries (Yes/No) (All in primary data.stw) Group 1: No Group 2: Yes					
	Mean No	Mean Yes	t-value	df	p	Valid N No
Age	52.16	65.61	-5.29700	31	0.000009	18
Goal heart rate	167.84	154.39	5.29700	31	0.000009	18
METs	7.61	5.63	3.03459	31	0.004844	18

Variable	T-tests; Grouping: Atherosclerosis in other arteries (Yes/No) (All in primary data.stw) Group 1: No Group 2: Yes				
	Valid N Yes	Std.Dev. No	Std.Dev. Yes	F-ratio Variances	p Variances
Age	15	6.804	7.776	1.30590	0.594317
Goal heart rate	15	6.804	7.776	1.30590	0.594317

Variable	T-tests; Grouping: Atherosclerosis in other arteries (Yes/No) (All in primary data.stw) Group 1: No Group 2: Yes				
	Valid N Yes	Std.Dev. No	Std.Dev. Yes	F-ratio Variances	p Variances
METs	15	2.452	0.635	14.89954	0.000007

### 3-G

Variable	T-tests; Grouping: Atherosclerotic vascular (Namely) (All in primary data.stw) Group 1: Carotid, Brachiocephalic bifurcation Group 2: Carotid, Brachiocephalic bifurcation	
	Mean Carotid, Brachiocephalic bifurcation	Mean Carotid, Brachiocephalic bifurcation
Max HR	139.33	184.00
Reached %	90.37	124.61

Variable	T-tests; Grouping: Atherosclerotic vascular (Namely) (All in primary data.stw) Group 1: Carotid, Brachiocephalic bifurcation Group 2: Carotid, Brachiocephalic bifurcation			
	t-value	df	p	Valid N Carotid, Brachiocephalic bifurcation
Max HR	-5.07001	12	0.000275	12
Reached %	-5.00824	12	0.000305	12

### 3-H

Variable	T-tests; Grouping: Atherosclerotic vascular (Namely) (All in primary data.stw) Group 1: Carotid, Brachiocephalic bifurcation Group 2: Carotid, Brachiocephalic bifurcation	
	Valid N Carotid, Brachiocephalic bifurcation	Std.Dev. Carotid, Brachiocephalic bifurcation
Max HR	2	10.210
Reached %	2	8.173

Variable	T-tests; Grouping: Atherosclerotic vascular (Namely) (All in primary data.stw) Group 1: Carotid, Brachiocephalic bifurcation Group 2: Carotid, Brachiocephalic bifurcation		
	Std.Dev. Carotid, Brachiocephalic bifurcation	F-ratio Variances	p Variances
Max HR	21.2132	4.317	0.123886
Reached %	15.0526	3.392	0.185215

### 3-I

Variable	T-tests; Grouping: Carotid (All in primary data.stw) Group 1: No Group 2: Yes					
	Mean No	Mean Yes	t-value	df	p	Valid N No
Age	53.33	66.08	-4.77434	32	0.000038	20

Variable	T-tests; Grouping: Carotid (All in primary data.stw)					
	Group 1: No Group 2: Yes					
	Mean No	Mean Yes	t-value	df	p	Valid N No
Goal heart rate	166.67	153.92	4.77434	32	0.000038	20
METs	7.35	5.56	2.60444	32	0.013842	20

Variable	T-tests; Grouping: Carotid (All in primary data.stw)				
	Group 1: No Group 2: Yes				
	Valid N Yes	Std.Dev. No	Std.Dev. Yes	F-ratio Variances	p Variances
Age	14	7.532	7.845	1.08495	0.849210
Goal heart rate	14	7.532	7.845	1.08495	0.849210
METs	14	2.508	0.597	17.65041	0.000005

### 3-J

Variable	T-tests; Grouping: Brachiocephalic (All in primary data.stw)					
	Group 1: No Group 2: Yes					
	Mean No	Mean Yes	t-value	df	p	Valid N No
Age	54.18	66.65	-4.38832	32	0.000116	22
Goal heart rate	165.82	153.35	4.38832	32	0.000116	22
METs	7.18	5.56	2.24411	32	0.031872	22

Variable	T-tests; Grouping: Brachiocephalic (All in primary data.stw)				
	Group 1: No Group 2: Yes				
	Valid N Yes	Std.Dev. No	Std.Dev. Yes	F-ratio Variances	p Variances
Age	12	8.165	7.432	1.20683	0.769942
Goal heart rate	12	8.165	7.432	1.20683	0.769942
METs	12	2.452	0.587	17.45301	0.000023

### 3-K

Variable	T-tests; Grouping: Arterial Hypertension (All in primary data.stw)					
	Group 1: Yes Group 2: No					
	Mean Yes	Mean No	t-value	df	p	Valid N Yes
Age	62.13	54.62	2.55288	36	0.015067	21
SBP rest	126.00	118.12	2.03149	36	0.049637	21
Goal heart rate	157.87	165.38	-2.55288	36	0.015067	21
METs	5.90	7.44	-2.45597	36	0.019008	21
eGFR (2021 CKD-EPI Creatinine)	77.32	87.88	-2.16746	36	0.036892	21

Variable	T-tests; Grouping: Arterial Hypertension (All in primary data.stw)				
	Group 1: Yes Group 2: No				
	Valid N No	Std.Dev. Yes	Std.Dev. No	F-ratio Variances	p Variances
Age	17	8.924	9.128	1.04632	0.910887
SBP rest	17	12.235	11.450	1.14179	0.797035
Goal heart rate	17	8.924	9.128	1.04632	0.910887
METs	17	1.407	2.395	2.89929	0.026188

Variable	T-tests; Grouping: Arterial Hypertension (All in primary data.stw) Group 1: Yes Group 2: No				
	Valid N No	Std.Dev. Yes	Std.Dev. No	F-ratio Variances	p Variances
eGFR (2021 CKD-EPI Creatinine)	17	15.615	14.046	1.23599	0.674001

3-L

Variable	T-tests; Grouping: Reaction type ( positive/negative) (All in primary data.stw) Group 1: Positive Group 2: Negative					
	Mean Positive	Mean Negative	t-value	df	p	Valid N Positive
WT	106.25	144.12	-2.18459	27	0.037776	12

Variable	T-tests; Grouping: Reaction type ( positive/negative) (All in primary data.stw) Group 1: Positive Group 2: Negative				
	Valid N Negative	Std.Dev. Positive	Std.Dev. Negative	F-ratio Variances	p Variances
WT	17	37.119	51.181	1.901152	0.283098

3-M

Variable	T-tests; Grouping: Reason of discontinuation (All in primary data.stw) Group 1: Horizontal ST depression >1mm Group 2: Reach goal HR			
	Mean Horizontal ST depression >1mm	Mean Reach goal HR	t-value	df
Creatinine (µmol/L)	98.62	82.62	2.39629	36

Variable	T-tests; Grouping: Reason of discontinuation (All in primary data.stw) Group 1: Horizontal ST depression >1mm Group 2: Reach goal HR		
	p	Valid N Horizontal ST depression >1mm	Valid N Reach goal HR
Creatinine (µmol/L)	0.021881	6	32

Variable	T-tests; Grouping: Reason of discontinuation (All in primary data.stw) Group 1: Horizontal ST depression >1mm Group 2: Reach goal HR		
	Std.Dev. Horizontal ST depression >1mm	Std.Dev. Reach goal HR	F-ratio Variances
Creatinine (µmol/L)	23.934	13.007	3.385970

Variable	T-tests; Grouping: Reason of discontinuation (All in primary data.stw) Group 1: Horizontal ST depression >1mm Group 2: Reach goal HR

	p Variances
Creatinine (μmol/L)	0.029651

Interestingly, the sensitivity and specificity of the bicycle ergometry showed a very low level and not sufficient for confirming the diagnosis of ischemic heart disease. (Table 4)

Table 4: The specificity and sensitivity of the bicycle ergometry. (\*) These values are dependent on disease prevalence.

Statistic	Value	95% CI
Sensitivity	57.89%	33.50% to 79.75%
Specificity	61.11%	35.75% to 82.70%
Positive Likelihood Ratio	1.49	0.74 to 2.98
Negative Likelihood Ratio	0.69	0.36 to 1.31
Disease prevalence (*)	51.35%	34.40% to 68.08%
Positive Predictive Value (*)	61.11%	43.96% to 75.89%
Negative Predictive Value (*)	57.89%	41.95% to 72.35%
Accuracy (*)	59.46%	42.10% to 75.25%

All the correlation results between the continuous variables presented in supplementary material. (Supplementary file 1)

### Discussion

Interestingly, on the CTP test, 19 (50 %) participants with coronary artery atherosclerosis, 5 (26 %) of them with significant atherosclerosis (stenosis >60 %). Furthermore, out of the 19 patients with significant atherosclerosis of the coronary artery, 11 (57 %) had myocardial perfusion defect after stress test with ATP.

Out of the 11 patients with positive myocardial perfusion defect after the stress test, 4 (36 %) participants with positive bicycle ergometry test.

More interesting findings, in the whole sample there is 19 (50 %) participants with positive myocardial perfusion defect, and only 5 (26 %) of them with significant coronary artery stenosis. Out of the 19 positive myocardial perfusion defect, 11 (57 %) with positive/suspected results on the bicycle ergometry.

2 (5.26 %) participants out of the 38 with positive results on the bicycle ergometry with heart rate < 85 % of the required goal heart rate (220- age). In other participants with positive (11)/suspected (8) results on the bicycle ergometry with heart rate > 85 % of the required heart rate goal.

In the whole sample there is 12 (31.58 %) positive, 9 (23.68 %) suspected, and 17 (44.47 %) negative on the bicycle ergometry. However, on the CTP, out of the 12 positive bicycle ergometry, only 5 (41.67 %) with positive myocardial perfusion defect after stress test with

ATP. Moreover, out of the 9 suspected participants, 6 (66.67 %) had positive myocardial perfusion defect after stress test with ATP.

The presented results showed that bicycle ergometry suffers from severe limitations in terms of specificity and sensitivity in the diagnosis of IHD. Additional methods are required to improve the diagnostic accuracy of the bicycle ergometry such as the exhaled breath analysis and single channel electrocardiography<sup>[11]</sup>.

### **Conclusions**

On the light of the presented results, the diagnosis of IHD using bicycle ergometry independently is ultra extremely not sufficient and further investigations required to improve the diagnostic accuracy of the bicycle ergometry. As well as to innovate novel methods for early detection of IHD through the application of the current advances in technologies such as the use of artificial intelligence (machine learning models) in the interpretation of the electrocardiography monitoring for patient at home bases using their watches or phones<sup>[6, 12]</sup>. Additionally, supplementary methods can be combined with the bicycle ergometry to enhance its diagnostic accuracy.

### **List of abbreviations**

IHD; Ischemic heart disease, CTP; stress computed tomography myocardial perfusion imaging

### **Decelerations**

1. Ethics approval and consent to participate: The study approved by the Sechenov University, Russia, from “Ethics Committee Requirement № 19-23 from 26.10.2023”.
2. The study is under consideration by the [clinicaltrials.gov](https://clinicaltrials.gov) for registration.
3. Consent for publication: Written informed consent was obtained from the participants for publication of study results and any accompanying images.
4. Availability of data and materials: applicable on reasonable request.
5. Competing interests: The authors declare that they have no competing interests regarding publication.
6. Funding’s: The work of Philipp Kopylov and Peter Chomakhidze was financed by the government assignment 1023022600020-6 «Application of mass spectrometry and exhaled air emission spectrometry for cardiovascular risk stratification».
7. Authors' contributions: **Basheer Abdullah Marzoog**, is the writer, researcher, collected and analyzed data, and revised the manuscript, **Peter Chomakhidze and Philipp Kopylov** revised the manuscript. All authors have read and approved the manuscript.
8. Acknowledgments: not applicable

9. Authors' information: **Basheer Abdullah Marzoog**, World-Class Research Center «Digital Biodesign and Personalized Healthcare», I.M. Sechenov First Moscow State Medical University (Sechenov University), 119991 Moscow, Russia; postal address: Russia, Moscow, 8-2 Trubetskaya street, 119991. (marzug@mail.ru, +79969602820). ORCID: 0000-0001-5507-2413. Scopus ID: 57486338800. **Peter Chomakhidze**, World-Class Research Center «Digital Biodesign and Personalized Healthcare», I.M. Sechenov First Moscow State Medical University (Sechenov University), 119991 Moscow, Russia; postal address: Russia, Moscow, 8-2 Trubetskaya street, 119991. ORCID: 0000-0003-1485-6072. Email: m.ba.m@bk.ru. **Philipp Kopylov**, World-Class Research Center «Digital Biodesign and Personalized Healthcare», I.M. Sechenov First Moscow State Medical University (Sechenov University), 119991 Moscow, Russia; postal address: Russia, Moscow, 8-2 Trubetskaya street, 119991. ORCID: 0000-0002-4535-8685. Email: kopylovf\_yu@staff.sechenov.ru.
10. The paper has not been submitted elsewhere

## References

- [1] Tsao, C. W.; Aday, A. W.; Almarzooq, Z. I.; Anderson, C. A. M.; Arora, P.; Avery, C. L.; Baker-Smith, C. M.; Beaton, A. Z.; Boehme, A. K.; Buxton, A. E.; et al. Heart Disease and Stroke Statistics - 2023 Update: A Report from the American Heart Association. *Circulation*, **2023**, *147* (8), E93–E621. <https://doi.org/10.1161/CIR.0000000000001123>.
- [2] Yeates, K.; Lohfeld, L.; Sleeth, J.; Morales, F.; Rajkotia, Y.; Ogedegbe, O. A Global Perspective on Cardiovascular Disease in Vulnerable Populations. *Can. J. Cardiol.*, **2015**, *31* (9), 1081–1093. <https://doi.org/10.1016/j.cjca.2015.06.035>.
- [3] Ofori, E. K.; Frimpong, E.; Ademiluyi, A.; Olawale, O. A. Ergometer Cycling Improves the Ambulatory Function and Cardiovascular Fitness of Stroke Patients—a Randomized Controlled Trial. *J. Phys. Ther. Sci.*, **2019**, *31* (3), 211–216. <https://doi.org/10.1589/jpts.28.211>.
- [4] Myers, J.; Arena, R.; Franklin, B.; Pina, I.; Kraus, W. E.; McInnis, K.; Balady, G. J. Recommendations for Clinical Exercise Laboratories. *Circulation*, **2009**, *119* (24), 3144–3161. <https://doi.org/10.1161/CIRCULATIONAHA.109.192520>.
- [5] van der Wall, E. E. Alternative Stress Methods in the Evaluation of Coronary Artery Disease. In *Nuclear Cardiology and Cardiac Magnetic Resonance*; Springer Netherlands: Dordrecht, 1992; pp 199–214. <https://doi.org/10.1007/978-94-011-2826->

1\_11.

- [6] Marzoog, B. A.; Chomakhidze, P.; Kopylov, P. The Biomarkers of Exhaled Breath & Single Channel Electrocardiography in the Diagnosis of Ischemic Heart Disease. *New Emirates Med. J.*, **2024**.
- [7] Cockcroft, D. W.; Gault, M. H. Prediction of Creatinine Clearance from Serum Creatinine. *Nephron*, **1976**, *16* (1), 31–41. <https://doi.org/10.1159/000180580>.
- [8] Winter, M. A.; Guhr, K. N.; Berg, G. M. Impact of Various Body Weights and Serum Creatinine Concentrations on the Bias and Accuracy of the Cockcroft-Gault Equation. *Pharmacotherapy*, **2012**, *32* (7), 604–612. <https://doi.org/10.1002/j.1875-9114.2012.01098.x>.
- [9] Brown, D. L.; Masselink, A. J.; Lalla, C. D. Functional Range of Creatinine Clearance for Renal Drug Dosing: A Practical Solution to the Controversy of Which Weight to Use in the Cockcroft-Gault Equation. *Ann. Pharmacother.*, **2013**, *47* (7–8), 1039–1044. <https://doi.org/10.1345/aph.1S176>.
- [10] Delgado, C.; Baweja, M.; Crews, D. C.; Eneanya, N. D.; Gadegbeku, C. A.; Inker, L. A.; Mendu, M. L.; Miller, W. G.; Moxey-Mims, M. M.; Roberts, G. V.; et al. A Unifying Approach for GFR Estimation: Recommendations of the NKF-ASN Task Force on Reassessing the Inclusion of Race in Diagnosing Kidney Disease. *Am. J. Kidney Dis.*, **2022**, *79* (2), 268-288.e1. <https://doi.org/10.1053/j.ajkd.2021.08.003>.
- [11] Marzoog, B. A. Breathomics Detect the Cardiovascular Disease: Delusion or Dilution of the Metabolomic Signature. *Curr. Cardiol. Rev.*, **2024**, *20*. <https://doi.org/10.2174/011573403X283768240124065853>.
- [12] Marzoog, B. A.; Averina, D. Nicotinamide Mononucleotide in the Context of Myocardocyte Longevity. *Curr. Aging Sci.*, **2023**, *17*. <https://doi.org/10.2174/0118746098266041231212105020>.