More

1 National Implementation Trial of BeUpstandingTM: An Online Initiative for Workers to Sit Less and Move

2

3

4 Authors

Name	Institutional Address(es) and Affiliations
Genevieve N. Healy	The University of Queensland, School of Human Movement and Nutrition
	Sciences, Brisbane, Queensland, Australia
Ana D. Goode	The University of Queensland, School of Human Movement and Nutrition
	Sciences, Brisbane, Queensland, Australia
Lisa Ulyate	The University of Queensland, School of Human Movement and Nutrition
	Sciences, Brisbane, Queensland, Australia
Alison Abbott	Work Health Design (retired), Workplace Health and Safety Qld, Office of
	Industrial Relations
David Dunstan	1. Physical Activity Laboratory, Baker Heart and Diabetes Institute. 75
	Commercial Road, Melbourne, VIC Australia
	2. Institute for Physical Activity and Nutrition, Deakin University,
	Melbourne, VIC Australia
Elizabeth G. Eakin	The University of Queensland, School of Public Health, Brisbane,
	Queensland, Australia
Nicholas D. Gilson	The University of Queensland, School of Human Movement and Nutrition
	Science, Brisbane, Queensland, Australia
Lynn Gunning	Comcare, GPO Box 9905, Canberra ACT 2601
Jodie Jetann	The University of Queensland, School of Human Movement and Nutrition
	Sciences, Brisbane, Queensland, Australia
Anthony D.	Deakin University, Institute for Health Transformation, Determinants of
LaMontegness preprint reports	new cashtacil insite a Stock being certified by Abby tealing and should not be used to guide clinical practice.

Marj Moodie	Deakin University, Institute for Health Transformation, Deakin Health		
	Economics, Melbourne, VIC, Australia		
Samantha Mulcahy	The University of Queensland, Health and Wellbeing Centre for Research		
	Innovation, School of Human Movement and Nutrition Science, Brisbane,		
	Queensland, Australia		
Neville Owen	1. Centre for Urban Transitions, Swinburne University of Technology,		
	Hawthorn, Victoria, Australia		
	2. Physical Activity Laboratory, Baker Heart & Diabetes Institute, 75		
	Commercial Road, Melbourne 3004, Victoria, Australia		
Trevor Shilton	1. Curtin University, School of Public Health, Kent Street, Bentley, Western		
	Australia		
	2. World Heart Federation		
Leanne Sweeny	Work Health Design, Workplace Health and Safety Qld, Office of Industrial		
	Relations		
Leon Straker	School of Allied Health, Curtin University, Perth, Australia		
Elisabeth A.H. Winkler	The University of Queensland, School of Human Movement and Nutrition		
	Sciences, Brisbane, Queensland, Australia		

5

6 Corresponding author

- 7 Genevieve N. Healy
- 8 The University of Queensland, School of Human Movement and Nutrition Sciences
- 9 St Lucia, Brisbane, Queensland, 4072
- 10 Email: <u>g.healy@uq.edu.au</u>
- 11

12 Abstract

Background: The online BeUpstandingTM program is an eight-week workplace-delivered intervention for
desk-based workers to raise awareness of the benefits of sitting less and moving more and build a
supportive culture for change. A workplace representative (the "champion") delivers the program, which
includes a workshop where teams collectively choose their sit less/move more strategies. A toolkit
provides the champion with a step-by-step guide and associated resources to support program uptake,
delivery, and evaluation. Here we report on the main findings from the Australian national implementation
trial of BeUpstanding.

20 Methods: Recruitment (12/06/2019 to 30/09/2021) was supported by five policy and practice partners,

21 with desk-based work teams from across Australia targeted. Effectiveness was measured via a single arm,

22 repeated-measures trial. Data were collected via online surveys, toolkit analytics, and telephone calls with

champions. The RE-AIM framework guided evaluation, with adoption/reach (number and characteristics);

24 effectiveness (primary: self-reported workplace sitting time); implementation (completion of core

components; costs); and, maintenance intentions reported here. Linear mixed models, correcting for

26 cluster, were used for effectiveness, with reach, adoption, implementation, and maintenance outcomes

27 described.

28 **Results:** Of the 1640 website users who signed-up to BeUpstanding during the recruitment period, 233 were eligible, 198 (85%) provided preliminary consent, and 118 (50.6%) champions consented and started 29 30 the trial, with 94% (n=111 champions) completing. Trial participation was from across Australia and 31 across industries, and reached 2,761 staff, with 2,248 participating in the staff survey(s): 65% female, 64% university educated; 16.9% from non-English speaking background. The program effectively changed 32 33 workplace sitting (-38.5 [95%CI -46.0 to -28.7] minutes/8-hour workday) and all outcomes targeted by BeUpstanding (behaviours and culture), with small-to-moderate statistically-significant effects observed. 34 All participating teams (n=94) completed at least 5/7 core steps; 72.4% completed all seven. Most 35 36 champions spent (72%) or > (10%) per team member; most (67/70.96%) intended to continue or repeat the program. 37

- 38 **Conclusions:** BeUpstanding can be adopted and successfully implemented by a range of workplaces, reach
- 39 a diversity of staff, and be effective at creating a supportive culture for teams of desk-based workers to sit
- 40 less and move more. Learnings will inform optimisation of the program for longer-term sustainability.
- 41
- 42 Trial registration: ACTRN12617000682347.
- 43 The trial was prospectively registered on the 12th May, 2017 (ACTRN12617000682347), prior to the soft
- 44 launch of the program, with the last update on the 11th June, 2019 prior to the commencement of
- 45 recruitment to the trial on the 12th June, 2019.
- 46

47 Keywords

48 Implementation trial; workplace; sitting; health promotion; activity; health and safety; public health;

- 49 occupational; web-based; RE-AIM
- 50
- 51

52 Introduction

53	High levels of sedentary time can be associated with poor health, wellbeing and productivity outcomes and
54	indicators (1) with detrimental impacts particularly pronounced in those who are also physically
55	inactive.(2, 3) The desk-based workplace has been identified as a key target setting for intervening on
56	prolonged sedentary time,(4) with systematic review evidence concluding that interventions addressing
57	prolonged workplace sedentary time can be both effective and acceptable.(5-9) Recent economic analyses
58	have also indicated that reductions in sedentary time, to the levels achieved through workplace
59	interventions, could result in Australian healthcare cost savings of \$39 million per year.(10) Given this
60	evidence, and the associated identification of workplace sedentary time as an emergent workplace health
61	and safety issue,(11) there is demand from occupational policy and practice partners for an evidence-based,
62	low-cost/no-cost, scalable solution to support workers and organisations to reduce prolonged sedentary
63	time.(12)
64	The online BeUpstanding TM program (13) is based on a strong foundation of research evidence, including

from randomised controlled trials (14) and best-practice approaches to workplace health promotion.(15)

66 The 8-week program, targeted at desk-based work teams and using a champion-led approach, has a 67 theoretical basis in social cognitive theory (16) and a social-ecologic model.(17, 18) The program is 68 designed to raise awareness of the benefits of sitting less and moving more and create a culture of 69 sustainable change through teams collectively identifying and promoting strategies to support these

70 behaviours that are suitable for their work team and context.(19)

BeUpstanding has been iteratively developed and tested across multiple phases in collaboration with end
users and key policy and practice stakeholders, (12, 20-23) and guided by the RE-AIM (reach,

ramework.(24) Use of this framework is relevant

74 for informing broader dissemination as it guides design and evaluation of programs in applied

contexts.(25) This study reports on the adoption, reach, effectiveness, selected implementation, and

76 maintenance intent outcomes from the Australian national implementation trial (Phase 4) of BeUpstanding.

77 Cost-benefit analyses, and detailed implementation, maintenance, engagement and process analyses, will

- be reported separately. Importantly, the implementation trial, which started in mid-June 2019, occurred in
 the context of major events in Australia and globally, namely the Australian Black Summer bushfires (July
 2019 to March 2020) and the COVID-19 pandemic (March 2020--), which caused major disruptions to
 ways of working.(26, 27) Findings from this evaluation are intended to inform further optimisation of the
- 82 BeUpstanding program prior to Phase 5 (dissemination).
- 83
- 84 Methods
- 85 Study design

Detailed methods for this implementation trial have been previously reported.(19) In brief, a single-arm 86 87 design was used for evaluation of effectiveness, with repeated cross-sectional evaluations at pre-program (0 weeks), end-of-program (\approx 8 weeks; primary endpoint), and at 9 months post-program (\approx 12 months post-88 sign-up). Only the pre-program and end-of-program data are reported herein, with data reporting according 89 to the TIDieR (28) and TREND (29) checklists (Additional Files 1 and 2). The implementation trial was 90 91 funded by a National Health and Medical Research Council (NHMRC) of Australia Partnership Project Grant (#1149936), which included cash and/or in-kind support from the five partners (Safe Work Australia. 92 Comcare, Queensland Office of Industrial Relations, The Victorian Health Promotion Foundation 93 94 (VicHealth), and Healthier Workplace Western Australia). These organisations are responsible for 95 developing, implementing and/or promoting workplace health, safety and wellbeing in Australian workplaces. Ethical approval was from The University of Queensland Human Research Ethics Committee 96 (approval #2016001743). Recruitment into the trial ceased on the 30th September, 2021, noting the online 97 98 program remained available for anyone to access and use. Only data from users who signed-up during the recruitment period are reported here. 99

100

101 **Participants and recruitment**

Promotional efforts, co-ordinated by a detailed marketing and promotional plan developed by the research
team and external marketing and communication experts, were made by the five partners to direct potential

users to the BeUpstanding website (13) and the implementation trial. Multiple promotional channels were
used, including social media, web links, email listservers, newsletters, workplace health promotion and
occupational health networks, conferences, and workshops. Recruitment efforts were aimed at champions
and decision makers of desk-based employees from a wide cross-section of industries, especially including
the five sectors identified as priorities by the partners (regional, call centre, small business, blue-collar, and
government).

110

Recruitment into the trial was a multi-step process. The BeUpstanding online program was freely available 111 online for the entire trial period and beyond, without any requirement to be eligible or participate in the 112 trial. A user could register (by completing preliminary details of themselves and their organisation) and 113 fully sign-up to access the toolkit and associated resources by completing a champion profile survey and 114 providing informed consent. Preliminary eligibility for the trial was ascertained from these two data 115 sources (initial registration; champion profile survey), with eligibility being: not having previously run the 116 BeUpstanding program; team size of at least five staff; and, most of the team do predominantly desk-based 117 work. Those potentially eligible were contacted by telephone (call 1) by the research team to confirm 118 criteria and ascertain the final eligibility requirements. Namely, that a workplace representative/employee 119 signing up to the toolkit could perform the duties of a workplace champion and planned to run the program 120 within the recruitment window. Potential champions were made aware that the main feature of taking part 121 122 in the trial (as distinct from using the program independently) was they were committing to the evaluation steps in the BeUpstanding program and would receive expert health coaching support to deliver the 123 program via email and phone calls. 124

125

All confirmed eligible champion participants were invited to participate in the trial, except where recruitment quota had been reached for a particular priority sector (see sample size) or a limit was reached for coaching availability to be allocated to a single organisation. Consent for the trial was in addition to that provided as part of the website sign-up process. The program was intended to be delivered to all staff

7

130 in participating teams, with no exclusion criteria at the staff level. Staff provided informed consent as a

131 step for gaining entry to the staff surveys.

132

BeUpstanding program – national implementation trial version

A detailed description of the BeUpstanding program is reported elsewhere.(19) In brief, the program was 134 designed to be run by the workplace champion across three phases (plan, do, review), each with associated 135 tasks for the champion to complete. The accompanying toolkit provided information and training for the 136 champion on the purpose of each phase and associated resources to support implementation, with the seven 137 core components highlighted (starred) to indicate their importance. The most critical was the staff 138 information and consultation workshop (or equivalent), where staff were provided information about the 139 benefits of sitting less/moving more and collectively chose three (or more) team strategies they would use 140 to sit less/move more as a team. Champions were encouraged to promote these strategies over eight weeks 141 via the posters and emails provided in the toolkit. This participative approach meant each intervention 142 program was unique for each team. In line with public, occupational, and clinical guidelines,(30-32) 143 behavioural targets were for workers to achieve a 50:50 split between sitting and upright activities during 144 work hours and to alternate sitting/upright posture at least every 30 minutes. Increased incidental 145 movement throughout the day was also encouraged through the move more messaging. No incentives were 146 provided as part of the trial. The main distinguishing feature of the implementation trial (as compared with 147 148 the program in general) is that the research team both supported (and collected further data on) program implementation, via email support and telephone calls at five key timepoints: 1) recruitment; 2) obtaining 149 150 consent; 3) at program initiation, following the staff consultation; 4) at end of program; and, 5) after approximately 12 months (maintenance). Project staff providing this support all had a minimal Masters 151 level qualification and were experienced in motivational interviewing for behaviour change. 152

153

Due to the major shift to hybrid work resulting from COVID-19,(33) the toolkit and resources were audited and modified, and new resources were developed (e.g., see Additional File 3) during the trial to ensure the program was suitable for delivery for all desk-workers, no matter where they were working. These new

- resources went online in July 2020. Additional measures to capture work-from-home arrangements (staff
 completed) and the impact of COVID-19 on the workplace (champion completed) were also added (May
 2020).
- 160

161 **Data collection**

Detailed description of the data collection process is provided in the protocol paper.(19) Data were 162 primarily collected via the dedicated, stand-alone BeUpstanding website, using surveys and toolkit 163 analytics, with the project management team collecting and confirming implementation data via the 164 telephone check-ins. Each toolkit user was required to complete the registration survey and champion 165 profile survey as part of the sign-up process. Each champion was requested to complete a workplace audit 166 before delivering the program and then a program completion survey after program delivery. Champions 167 were also responsible for sending their bespoke links for the staff surveys (pre-program, post-program) to 168 all staff in their team. For each survey, an anonymous staff identifier was constructed based on responses 169 to three questions (month of birth; first initial of mothers first name; and, last three digits of mobile phone). 170 This identifier, coupled with the cluster identification number, was used to ensure, per combined work 171 team (cluster), each staff member responded only once and to match pre- and post-program responses. 172 There was no blinding. 173

174

175 Outcomes and Measures

176 The outcomes according to the RE-AIM framework are outlined below.

177

178 Adoption

Adoption was chiefly described in terms of uptake of the toolkit (n and % unlocking the toolkit), trial participation (n and %), the organisational, workplace, team and champion characteristics of those who participated in the trial, and how they had heard of BeUpstanding (in order to adopt it). Organisational characteristics were reported by champions, reconciled across champions from the same organisation, and checked against public records. Workplace characteristics were permitted to diverge between different

184	champions from the same workplace. Champions were classified by project staff according to their role
185	delivering the program to a team (overseer and primary champion / primary champion / co-champion) or
186	supporting other aspects of BeUpstanding (overseer only / decision maker / other role). Further detailed
187	information relevant to understanding adoption were the predictors of non-adoption (comparison of
188	participants with non-participants), reasons given for not participating or withdrawing, and reasons for
189	adopting the program, focusing specifically on what the champions said that they aimed to achieve by
190	running BeUpstanding (open ended).

191

192 *Reach*

193 Reach outcomes included: the number of staff in participating teams (as estimated by the champion, and as

verified through staff surveys), and the characteristics of staff who took part in the evaluation (by

responding to one or both staff surveys).

196

194

197 *Implementation*

Implementation outcomes reported are completion rates (yes/no) of the seven core program steps, and champion-reported costs they incurred (reported in the program completion survey) or expected to incur (reported in the planning stage implementation check, shortly after selecting team strategies) through running BeUpstanding.

202

203 *Effectiveness*

Program effectiveness was assessed by the difference between pre-and post-program staff survey responses regarding 13 indicators of the program's impact on workplace behaviour and workplace culture targeted directly by the program and 13 measures of productivity and health and wellbeing that were expected might improve as a result of the program. The primary outcome for effectiveness was workplace sitting, measured by the Occupational Sitting and Physical Activity Questionnaire (OSPAQ).(34) The post-staff survey also captured staff perceptions of the program and its impact, and adverse events. When described,

adverse events were classified by two researchers as to whether they constituted adverse events and

- 211 whether they could reasonably have resulted from the program. Program satisfaction was collected from
- staff in post-program surveys and from team champions in program completion surveys.
- 213

214 Maintenance intentions

- 215 The maintenance intentions of the champions of completing teams were ascertained by coding and
- reviewing the responses to an open text question of "What are your plans now in relation to
- 217 BeUpstanding?" asked during program completion calls. Their detailed actual maintenance of the program
- and longer-term outcomes will be addressed in detail in subsequent papers.
- 219

220 Sample size (primary effectiveness outcome)

As described elsewhere,(19) using assumptions informed by earlier iterations of the program (SD 90,

r=0.5, ICC=0.1, post-attrition n/workplace = 5), it was determined that 47 to 62 workplaces (235 to 310)

staff) were required to provide 80% to 90% power to detect a change in workplace sitting of at least 20

224 minutes per 8 hours of work (i.e., 4% of the workday) with 5% two-tailed significance. Consequently, the

target sample size was at least 50 workplaces per priority sector to permit subgroup analysis.

226

227 Statistical Analysis

Most of the reach, adoption, implementation and maintenance outcomes are described using descriptive 228 statistics. Additionally, logistic regression was used to compare the organisational, workplace, team, and 229 230 personal characteristics of those who included in the trial (trial adopters) versus non-participants (nonadopters), as well as those who completed the post-program evaluation of their team versus those who did 231 not. Effectiveness outcomes (continuous) were compared between pre- and post-program staff surveys 232 using linear mixed models, correcting for cluster (workplace) as a random intercept, or as random slopes, 233 where likelihood ratio tests indicated the random slopes model was superior. Unadjusted models were 234 reported in addition to the main findings, which adjusted for potential confounding due to different staff 235 and teams providing data at each survey. Specifically, these were characteristics with a p < 0.2 association 236

237	either with teams being part of the post-program evaluation or not (tested by logistic regression models), or
238	with staff members providing outcome data at only the pre-program survey (n=1481) versus only the post-
239	program survey (n=397) (tested by mixed logistic regression models, correcting for workplace cluster as a
240	random intercept). To test the sensitivity of conclusions to missing data handling, confounder-adjusted
241	models were also reported with missing data imputed, using multiple imputation by chained equations. In
242	addition to the analytic variables, imputation models contained auxiliary variables (additional variables
243	predicting outcomes at either pre- or post- program surveys at p<0.2 in backwards eliminations) to help
244	improve prediction of missing outcomes. A further sensitivity analysis excluded staff survey responses that
245	had been collected after the program had already started. Analyses were performed in STATA version 18,
246	with the final date of data extraction 15 May 2024. Significance was set at p<0.05 (two-tailed).
247	
248	Results
249	For ease of reporting, RE-AIM outcomes are presented these in the order of Adoption, Reach,
250	Implementation, Effectiveness, and Maintenance intentions.
250 251	Implementation, Effectiveness, and Maintenance intentions.
250 251 252	Implementation, Effectiveness, and Maintenance intentions. INSERT FIGURE 1 ABOUT HERE
250 251 252 253	Implementation, Effectiveness, and Maintenance intentions. INSERT FIGURE 1 ABOUT HERE
250 251 252 253 254	Implementation, Effectiveness, and Maintenance intentions. INSERT FIGURE 1 ABOUT HERE Adoption
250 251 252 253 254 255	Implementation, Effectiveness, and Maintenance intentions. INSERT FIGURE 1 ABOUT HERE Adoption The participant flow diagram (Figure 1) shows that out of the 1640 website users who signed up during the
250 251 252 253 254 255 256	Implementation, Effectiveness, and Maintenance intentions. INSERT FIGURE 1 ABOUT HERE Adoption The participant flow diagram (Figure 1) shows that out of the 1640 website users who signed up during the trial recruitment period, 847 fully registered and unlocked the toolkit, and 233 were determined to be
250 251 252 253 254 255 256 257	Implementation, Effectiveness, and Maintenance intentions. INSERT FIGURE 1 ABOUT HERE Adoption The participant flow diagram (Figure 1) shows that out of the 1640 website users who signed up during the trial recruitment period, 847 fully registered and unlocked the toolkit, and 233 were determined to be eligible for the trial, with ineligibility mostly due to not being contactable or not planning to run the
250 251 252 253 254 255 256 257 258	Implementation, Effectiveness, and Maintenance intentions. INSERT FIGURE 1 ABOUT HERE Adoption The participant flow diagram (Figure 1) shows that out of the 1640 website users who signed up during the trial recruitment period, 847 fully registered and unlocked the toolkit, and 233 were determined to be eligible for the trial, with ineligibility mostly due to not being contactable or not planning to run the program either at all or in the requisite timeframe. In total, 82 potentially would have been eligible if the
250 251 252 253 254 255 256 257 258 259	Implementation, Effectiveness, and Maintenance intentions. INSERT FIGURE 1 ABOUT HERE Adoption The participant flow diagram (Figure 1) shows that out of the 1640 website users who signed up during the trial recruitment period, 847 fully registered and unlocked the toolkit, and 233 were determined to be eligible for the trial, with ineligibility mostly due to not being contactable or not planning to run the program either at all or in the requisite timeframe. In total, 82 potentially would have been eligible if the recruitment quota for their sector(s) was considered not already met. Out of the 233 eligible users, the
250 251 252 253 254 255 256 257 258 259 260	Implementation, Effectiveness, and Maintenance intentions. INSERT FIGURE 1 ABOUT HERE Adoption The participant flow diagram (Figure 1) shows that out of the 1640 website users who signed up during the trial recruitment period, 847 fully registered and unlocked the toolkit, and 233 were determined to be eligible for the trial, with ineligibility mostly due to not being contactable or not planning to run the program either at all or in the requisite timeframe. In total, 82 potentially would have been eligible if the recruitment quota for their sector(s) was considered not already met. Out of the 233 eligible users, the preliminary consent rate was 85.0% (n=198), which dropped to 50.6% (n=118) after excluding those who
250 251 252 253 254 255 256 257 258 259 260 261	Implementation, Effectiveness, and Maintenance intentions. INSERT FIGURE 1 ABOUT HERE Adoption The participant flow diagram (Figure 1) shows that out of the 1640 website users who signed up during the trial recruitment period, 847 fully registered and unlocked the toolkit, and 233 were determined to be eligible for the trial, with ineligibility mostly due to not being contactable or not planning to run the program either at all or in the requisite timeframe. In total, 82 potentially would have been eligible if the recruitment quota for their sector(s) was considered not already met. Out of the 233 eligible users, the preliminary consent rate was 85.0% (n=198), which dropped to 50.6% (n=118) after excluding those who preliminarily seemed eligible and consented but withdrew before starting the program. Completion rates

most common reasons for non-participation prior to preliminary consent were reasons related to not being
able to run the program, or lack of interest in the trial. Reasons for not participating after preliminarily
consenting were primarily due to changing circumstances at either the organisation or champion level.
Withdrawing after having started the program was rare and most often due to it being a difficult time in the

267 organisation (n=5/7).

268

Almost all trial respondents (n=116) indicated what they aimed to achieve by adopting the program, with 269 270 results categorised under 22 themes (Additional File 4). The most common aim was to improve either physical health and/or wellbeing (50.9%), with changing sitting / physical activity levels (37.9%) also 271 often mentioned. There was some evidence that aims were associated with trial participation (Additional 272 File 5). Reporting an aim to improve culture (OR=1.95, 95%CI: 1.09, 3.48), connection (OR=1.85, 273 95%CI: 1.02, 3.35), and/or productivity (OR=2.78, 95%CI: 1.47, 5.24) were associated with significantly 274 higher odds of participation. Reporting a general aim to improve health was associated with a significantly 275 lower odds of participation (OR=0.30, 95%CI: 0.09, 0.98). 276

277

Table 1 shows the characteristics of teams participating in the trial and their champions. The sample size 278 target was met in terms of total numbers but not in any of the subgroups, due in part to the COVID-19 279 pandemic. Teams were recruited from every priority sector: 48 teams (51.1%) from the public sector; 24 280 teams (25.5%) from small-medium enterprises (11 (11.75%) small business); 33 teams (35.1%) from blue-281 collar workplaces; 32 teams (34.0%) with regional staff; and, six teams with call-centre staff (6.4%). Team 282 size was highly varied and averaged 29.4 (SD=32.1) members. Overall, 13 of the 19 standard industry 283 classifications in Australia (35) had some representation in the trial. The most common industry of the 284 organisations of participating teams was Health Care and Social Assistance (n=16, 17.0%) and 285 Professional / Technical and Scientific Services (n=16, 17.0%). Teams were recruited from every state and 286 territory of Australia and were predominantly located in Queensland (n=43, 45.7%). Champions' work 287 locations varied widely in socioeconomic status, with a mean (SD) state percentile of 67.6 (26.7) in terms 288 of socioeconomic index of advantage and disadvantage. Champions had a mean (SD) age of 42.9 (11.0) 289

- years. Many were female (75.9%), in middle (38.8%) or senior (15.5%) management, had a health and
 safety role (65.5%), and workplace health promotion training (56.9%). Less than half (36.2%) had prior
- 292 experience delivering workplace health promotion programs.

(1.52, 95% CI: 1.09, 2.12 per unit change efficacy).

- 293
- 294

INSERT TABLE 1 ABOUT HERE

- 295
- Additional File 5 shows the odds of participation (adopting the program) by these characteristics.

Significant predictors included location (p<0.001, with highest participation in Oueensland, and lowest 297 participation in WA/NT/TAS), sector (p=0.001, with highest participation by public followed by non-profit 298 then private sector), and COVID-19 impact level (p<0.001, with highest participation in high followed by 299 general followed by no impact). Predictors significantly associated with lower odds of participation were: 300 having an occupational health and safety role (0.60, 95% CI: 0.39, 0.91); inclusion of call centre staff 301 (0.38, 95%CI: 0.16, 0.89); small-medium enterprise (0.50, 95%CI: 0.32, 0.79); and, hearing of the program 302 303 through an article/publication/newsletter (0.56. 95% CI: 0.32, 0.98). Predictors significantly associated with higher odds of participation were: hearing of the program through a seminar/presentation/conference 304 (1.97, 95%CI: 1.07, 3.64) or a colleague (2.29, 95%CI: 1.48, 3.52); and, higher organisational readiness 305

307

306

308 Reach outcomes

Based on estimates from participating champions, the program reached 2761 staff in the trial, with 2248 309 able to be verified via responding to at least the identifier questions in one or both staff surveys. Staff were 310 311 on average (mean (SD)) aged 42.9 (11.4) years, worked 38.2 (9.0) hours per week, and exercised for 3.1 (2.2) days per week prior to BeUpstanding, with 27.3% of staff meeting the recommended five or more 312 days of exercise per week (Table 2). More than half of staff were female (65.0%), in full-time employment 313 314 (82.1%), university educated (64.4%), non-managerial employees (68.5%), and in the highest occupational skill category (50.3%). A very small minority were shift-workers (2.1%), and 16.9% of staff were from a 315 non-English speaking background. 316

317

- 318

INSERT TABLE 2 ABOUT HERE

319

320 Implementation outcomes

The champions had completed all core steps for most of the participating teams (72.4%) and most of the 321 322 completing teams (76.4%). The steps with the lowest completion rates were the workplace audit, strategy selection, and the program completion survey (Table 3). During their planning stage implementation 323 check-in calls, shortly after selecting their team strategies, 80 champions reported on any costs they had 324 incurred or expected to incur in order to support or enact their team strategies. In total, 35 cost items were 325 reported, the most common of which was sit-stand desks or similar (n=14). Other costs included: 326 accessories to assist with standing at the desk (Bluetooth headsets (n=3); anti-fatigue mats (n=1)), aids for 327 prompting (reminder/lockout software (n=4); timer (n=1)), enhancements to communal areas (standing 328 benches/tables (n=3); communal bins (n=1)), equipment (n=2) and subscriptions (n=1) for physical activity 329 330 at work; and, miscellaneous other (pedometers (n=1); undescribed equipment (n=2); sporting goods gift cards (n=1)). In the program completion survey, 72 champions reported what was spent in total to run the 331 program for their team, with the majority reporting spending 0 (n=52, 72.2%, Table 4). Sit-stand desks 332 333 had been purchased for every team where the amount spent was >\$1000.

- 334
- 335

INSERT TABLES 3 AND 4 ABOUT HERE

336

337 Effectiveness outcomes

338 Outcomes targeted by the intervention

Baseline levels of the staff-reported program outcomes and their changes over the program are shown in
Table 5. The main models adjusted for characteristics that might differ between those providing data at
post- and pre-program surveys, based on participation in the post-program evaluation by the team
(Additional File 6) or staff member (Additional File 7). These models (Figure 2) indicated significant

343	improvements in all outcomes directly targeted by the program: the primary outcome (self-reported work
344	sitting); work standing and moving; the extent to which sitting was accumulated in prolonged bouts; the
345	alignment between preferred and actual sitting, standing and moving at work; and, all measures of
346	workplace culture. The magnitude of these effects on average were mostly small-to-moderate, with the
347	smallest effects seen for moving, moving alignment, and perceived control over sitting. Workplaces varied
348	significantly in their changes in activity at work and sitting in prolonged bouts, with tests for random
349	slopes reaching statistical significance (Additional File 8).

350

351

352

INSERT FIGURE 2 AND TABLE 5 ABOUT HERE

Conclusions regarding workplace behaviours from the main adjusted models were mostly robust to modelling choices, except that moving results strengthened slightly and became significant in adjusted models and then further strengthened with imputation. Results concerning culture were unaffected by statistical adjustment but were sensitive to missing data handling, whereas results from imputation models were often weaker and sometimes no longer statistically significant. Excluding the pre-program surveys that had been collected after the program commenced had no meaningful impact on findings (Additional File 9).

360

361 Other outcomes

Of the 13 outcomes concerning work productivity and health, significant improvements were seen in self-362 rated job performance, job satisfaction, energy, self-rated physical health, and musculoskeletal discomfort 363 364 in the upper body, lower back, and lower body, with improvements ranging in magnitude from very small to small (Figure 2; Table 5). Other changes were consistently all very small, and not statistically 365 significant. Random slopes tests were not statistically significant for any of these outcomes (Additional 366 File 8). Findings were similar to those from unadjusted models, except that adjustment brought results for 367 engagement even closer to the null and the previously significant change in self-rated psychological health 368 became borderline significant. There were no meaningful changes in outcomes excluding late pre-program 369

370	surveys (Additional File 9). Many findings, especially for work-related and productivity outcomes, were
371	highly sensitive to missing data, where multiple imputation results only indicated significant changes (all
372	improvements) in energy, self-rated physical health, and musculoskeletal discomfort in the upper body and
373	lower back, all of which were somewhat smaller than they had been in the evaluable case analyses. In
374	imputed models, effects for lower body discomfort were somewhat attenuated in magnitude and no longer
375	statistically significant, while effects for estimates of changes in self-rated job performance and job
376	satisfaction were completely attenuated. Effect sizes became further away from the null for sick days,
377	stress, and creativity, but remained very small and did not reach statistical significance.

378

379 Priority sectors

The changes in the primary outcome (work sitting) overall amounted on average to -38.5 (-46.0, -31.0) 380 min/8 h for the trial in general. Changes were mostly also present within each of the priority sectors: -28.2 381 (-39.7 to -16.6) min/8-h in small-medium enterprises (based on 24 workplaces and 496 staff); -35.8 (-49.4 382 to -22.1) min/8-h in blue-collar workplaces (based on 26 workplaces and 772 staff), -41.7 (-52.5 to -31.0) 383 min/8-h in public-sector workplaces (based on 40 workplaces and 954 staff); and, -45.2 (-60.0 to -30.3) 384 385 min/8-h in teams that included regional staff (based on 26 workplaces and 626 staff). Changes in the teams that included call-centre staff were -46.3 (-126.4 to 33.9) min/8-h and did not reach statistical significance 386 with the small sample of 3 workplaces and 97 staff. 387

388

389 Adverse events and staff perceptions of the program

Most (n=604, 96.0%) of the 629 staff responding to the staff follow-up surveys did not experience any adverse events. Only 7 staff (1.1%) reported an adverse event that was confirmed by research staff, with six of these (1.0%) potentially being attributable to the study, and a further 18 staff (2.9%) reported that they experienced some adverse event that could not be verified as they did not describe it. The six events that could have arisen from program participation were: new or exacerbation of musculoskeletal pain or injury (n=4), specifically to the hip from standing propped on a stool, to the back from "overdoing things",

```
397 guilty about leaving the desk (n=1) and feeling followed by their supervisor (n=1).
```

398

399	The majority of staff reported positive impacts on outcomes targeted directly by BeUpstanding (culture,
400	knowledge, attitudes, awareness), with few reporting negative impacts (Figure 3). Staff provided more
401	mixed responses in relation to the impact of the program on activity outside work, with very few staff
402	(n=11, 1.7%) perceiving a negative impact and the rest of the staff being almost equally divided between
403	perceiving a negligible impact (51.1%) or a positive impact (47.2%).

- 404
- 405

INSERT FIGURE 3 ABOUT HERE

406

407 **Program satisfaction**

Staff (n=654) reported on their satisfaction with the program. Excluding those who indicated they were not aware of BeUpstanding or responded not applicable (n=35), most staff (n=477, 77.1%) indicated they enjoyed BeUpstanding, some were uncertain (n=115, 18.6%), and a minority of staff said they did not enjoy participating (n=27, 4.4%). Champions also reported on their satisfaction with the program in postcompletion surveys (n=72). When asked whether the program met expectations, most said yes (n=57, 79.2%) or somewhat (n=13, 18.1%) and only two (2.8%) said no. Similarly, most said they would (n=63, 87.5%) or maybe would (n=8, 11.1%) recommend the program and only one said they would not (1.4%).

416 Maintenance intentions

The 8-week BeUpstanding program was completed by 89 of the 94 teams that were part of the trial. Upon completion, project staff were able to contact 70 of their team champions and ask about their planned next steps concerning BeUpstanding. Most responses to the open-ended question indicated the champions intended to continue the program (n=59, 84.3%) or repeat it (n=8, 11.4%), while very few indicated that

- 421 the champion intended to stop or pause the program (n=1, 1.4%), including to restart it with different staff
- 422 (n=1, 1.4%), and one champion intended to check with management (n=1, 1.4%).

plantar fasciitis, and lower leg and lower back pain; and, adverse work impacts (n=2), specifically feeling

423

424 **Discussion**

This study reports the main outcomes, covering the five dimensions of the RE-AIM framework, of the first national implementation trial of a workplace health promotion initiative specifically supporting teams of desk-based workers to sit less and move more (BeUpstanding). This trial was the fourth phase in the research-to-practice translation process from an intervention that had demonstrated effectiveness in the context of a cluster randomised controlled trial.(14)

430

431 Adoption and Reach

While there was interest in the program more broadly, with over 600 organisations unlocking the toolkit 432 during the trial period, adoption of the program as part of the research trial was more modest, with 433 approximately half of eligible workplaces running the program. This needs to be understood in the context 434 of the BeUpstanding program being available for free online without participation in the trial, and two 435 significant emergencies in Australia that occurred during the trial's recruitment period, specifically the 436 COVID-19 pandemic and the 2019-20 bushfires. Reported reasons for not adopting the program were 437 mostly regarding it being a difficult time in the organisation (e.g., availability of personnel) or lack of 438 interest in the research trial. Only rarely did withdrawal occur after initial consent due to an issue with the 439 program. As might be expected, a higher level of organisational readiness, specifically change efficacy, 440 441 was significantly associated with higher odds of adoption into the trial while conversely having an occupational health and safety role was associated with lower odds of adoption. One interpretation (among 442 many) is that champions who do not feel their workplace is well placed to run the program may require 443 additional supports (beyond what was provided in the implementation trial) to adopt the program, while 444 conversely those in occupational health and safety roles may have felt confident to run the program 445 446 independently outside of the trial context without support. Other reasons could include that occupational health and safety staff had competing priorities especially in the COVID-19 pandemic, and unready 447 workplaces may have wanted to avoid the oversight of a trial. 448

449

19

Workplaces and staff took part from every partner-identified priority sector, from every Australian state 450 and territory (including rural/regional locations), from 13 of 19 ANZIC industries,(35) and with diverse 451 workplace and worker characteristics. Some groups were easier to reach than others, with limited 452 participation from shift workers, from those not in fulltime employment, and with lower-skilled 453 occupations. While this may align with our specific focus on desk-based workers, it is important in future 454 research to understand how this program and others like it may work in these groups, as well as other 455 underrepresented or untested key populations, such as people with disabilities. Adoption was significantly 456 lower for workplaces outside the public sector, of small-medium size, and especially in teams that included 457 call-centre staff (despite deliberate recruitment efforts). Staff in call centres or contact centres generally 458 have higher rates of sedentary time than general office workers.(36) but typically also have reduced 459 autonomy to break up their sedentary time due to their job tasks and job demands.(37) In line with previous 460 recommendations,(38) it will be important to work with relevant stakeholders to co-design appropriate 461 approaches for this particular setting. 462

463

464 *Implementation*

Champions were provided automated feedback via the toolkit that was engaging and meaningful and had a 465 streamlined online user experience that highlighted the core components. As part of the trial, they also 466 received support from the research team at key stages: all factors that had been added and optimised based 467 468 on learnings from the version used by the early adopters.(21) In the previous phase of this research-topractice translation (Phase 3), the seven core steps were completed for only 5% of teams, due in part to 469 470 retention and in part to progressing through without completing core steps.(20) These modifications appear to have resulted in a much more complete implementation of the program, with most teams (>72%) 471 completing all seven core steps, and none completing fewer than five. Most workplaces implemented the 472 program in a no or low-cost manner, and only a minority spent significant sums of money, usually on sit-473 474 stand workstations.

475

476 Effectiveness

The program was significantly effective in relation to the primary outcome (workplace sitting), behaviours 477 478 and culture (which were the main targets of the program), and also for some indicators of productivity, health and wellbeing. The average decrease in self-reported work sitting of -38.5 (95%CI -46.0 to -31.0) 479 minutes per 8-hour workday appeared to have co-occurred mainly with increased standing, and also a 480 smaller amount of increased moving. Notably the extent of change strongly resembled the average effect 481 on sedentary behaviour of 38 minutes reduction per workday (95% CI -47.3 to -28.7) achieved by 482 sedentary behaviour interventions in office workers (across all intervention types combined) as reported by 483 484 a recent systematic review and meta-analysis.(39) Given the timing of the trial, and the substantial shift to hybrid work resulting from the COVID-19 pandemic, (33) these findings also provide some evidence that 485 BeUpstanding is suitable for work from home and hybrid work contexts. A pre-post design was required 486 for feasibility reasons, leaving open the possibility of other explanations for changes over time, especially 487 in light of the pandemic. Notably however, the controlled studies that informed the BeUpstanding program 488 development had not observed any meaningful degree of changes in control groups.(14, 40, 41) It is not 489 490 likely that the pandemic produced the improvements, considering COVID-19 saw an increase in sedentary time and a reduction in physical activity in desk workers(42), and the average change was -43 minutes per 491 8=hour workday (-9.0%, 95% CI: -12.0% to -5.9%) in the pre-pandemic evaluation of the program's 492 early adopters.(20) 493

494

495 Findings concerning workplace sitting, other workplace behaviours and workplace culture were robust to 496 issues of potentially selective evaluation, being largely unaltered in sensitivity analyses. Where selective 497 participation may have been more of an issue was in gauging the impact of the program on productivity 498 and health, where changes were typically smaller after accounting for missing data, but notably were still present for energy, self-rated physical health, and upper body and lower back musculoskeletal discomfort. 499 The benefits seen to at least some of the productivity and health indicators adds to the growing evidence 500 501 from field-based studies that interventions aimed at reducing workplace sedentary time, particularly those where sedentary time is predominantly replaced with standing, are not detrimental to indicators of work 502 performance.(43) The musculoskeletal findings in particular add to the currently limited evidence base in 503

this area.(17) Although these reductions were small on average, they are potentially important given that work-related musculoskeletal disorders are the leading work health and safety issue in Australia in terms of both frequency and costs,(44) and were not necessarily small for all participants. Future detailed evaluation will explore whether changes are possibly enhanced based on factors such as pre-existing discomfort, and nuanced aspects of the behaviour changes, such as the reallocation of sitting to standing versus to moving, the prolonged or interrupted nature of sedentary accumulation, and the specific behavioural strategies that were adopted.

511

A key goal of the implementation trial was to evaluate effectiveness in not just overall but also within key 512 sectors identified *a priori*.(19) Here, significant improvements in workplace sitting were seen within 513 almost all sectoral subgroups. The sole exception was that changes in teams with call centre staff were 514 substantial -46.3 minutes per 8-hour workday (95%CI: -126.4 to 33.9) but inconclusive with wide 515 confidence intervals as a consequence of only three call centres participating in the evaluation. This finding 516 adds to a known paucity of evidence, as indicated by a recent scoping review(45) that called for further 517 evidence on the effectiveness, acceptability and feasibility of health-promoting interventions (including for 518 addressing sedentary behaviour) conducted in the call-centre setting. Minimal adverse events were 519 recorded and were of mostly a similar nature (musculoskeletal) as have been reported in similar 520 programs.(46, 47) 521

522

523 Satisfaction and Maintenance intentions

Satisfaction was generally high for both staff and champions and most indicated they would participate again in BeUpstanding. Although further evidence is needed,(48) shifting culture may be the key for sustainable change in workplace sedentary behaviour. The BeUpstanding program heavily targeted culture, and significantly improved many cultural aspects. Following BeUpstanding, most champions intended to continue on with the program, often also intending adaptation -- an important feature of sustainable delivery.(49) Many particularly mentioned intentions to expand to either more staff and/or embedding the program into standard practice, which is consistent with the intent of BeUpstanding. Findings from the 12-

month follow-up champions (to be reported elsewhere), will provide further insights into workplacechanges following BeUpstanding.

533

534 Strengths and limitations

Strengths of the study include: the wide-ranging participation from workplaces across Australia and from a 535 wide variety of industries and all priority sectors identified by the policy and practice partners; the data 536 collection across the RE-AIM framework, enabling understanding of how the program is working in 537 practice; and, the data collection on a range of measures, including those relevant to organisations (e.g., job 538 satisfaction). Limitations include the single-group evaluation design, which was necessary for feasibility 539 reasons, and the large workforce turnover, reflective of the broader job mobility occurring during this 540 period.(33) To some extent this was mitigated by combining longitudinal and repeated cross-sectional 541 542 designs, and using statistics that account for repeated observations but do not require them. Staff could be included in the evaluation whether they were part of the team and participating in the evaluation in either 543 one or both evaluations. Another limitation was the measures used are self-report, or proxy-reported by 544 champions, and both responses concerning the organisation, the program, and the sending out of staff 545 surveys all relied on the workplace champions. While necessary for the scalable online program, this does 546 547 have significant potential for measurement error and bias. Importantly, staff perspectives were collected as well as the champion perspectives, and many staff findings were robust to missing data handling. Single-548 549 item or minimal item measures were chosen where possible in order to minimise participant burden in line with our preliminary work informing the trial. (21) One possibility for better gauging the impact of this 550 program and others like it on productivity and health is to continue the broad, brief evaluation model but 551 add optional evaluation tools with more detailed assessments. These may have measurement properties 552 more suitable to detecting changes. 553

554

555 Conclusion

556	Findings from this Australian national implementation trial found that the workplace champion-delivered
557	BeUpstanding program was effective in supporting teams of desk-based workers to sit less and move more
558	and in creating a supportive culture for this change. It also demonstrated that the program can be taken up
559	and implemented by a wide range of users from across Australia and across industries. Learnings from the
560	trial are being used to inform the next iteration of the program (BeUpstanding 2.0), which in turn will
561	inform Phase 5 (sustainability) of the research-to-practice process.(12) This optimisation, which will be
562	described in detail elsewhere, is designed to enhance scalability, inclusivity and the user experience.
563	

567 **Declarations**

568 *Ethics approval and consent to participate*

- 569 The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the
- 570 Human Research Ethics Committee of The University of Queensland (protocol code #2016001743 on the
- 571 9th January 2017). All participants provided informed consent. No individual or personally identifiable
- 572 information is reported.
- 573 Consent for publication
- 574 Not applicable

575 Availability of data and materials

576 The datasets generated and analysed during the current study are available from the corresponding author

577 on reasonable request.

578 Competing interests

The BeUpstanding program does include consultancy options within the toolkit. All and any proceeds from
consultancy activities return to the research to continue to develop the research program. GNH is on the
Editorial Board of IJBNPA.

582 *Funding*

The BeUpstanding program was supported by funding from the Queensland Government "Healthier. 583 Happier. Workplaces" program, Safe Work Australia, Comcare, and the National Health and Medical 584 Research Council of Australia through a Partnership Project Grant (#1149936) conducted in partnership 585 with Comcare, Safe Work Australia, the Queensland Office of Industrial Relations, VicHealth, and 586 Healthier Workplace WA. GNH was supported by an MRFF-NHMRC Emerging Leadership Investigator 587 Grant (#1193815). NHMRC had no role in the study in terms of the design, data collection, management, 588 589 analysis and/or interpretation. SKM's salary is supported by the Health and Wellbeing Centre for Research Innovation (HWCRI), co-funded by The University of Queensland and the Queensland Government 590 591 through Health and Wellbeing Queensland.

592 *Authors contributions*

- 593 GNH, ADG, AA, DWD, EGE, NDG, LG, ADL, MM, NO, TS, LS: conception, funding, recruitment,
- 594 interpretation.
- 595 LS: recruitment
- ADG; SKM, JJ, LU: recruitment; data collection; project management; partner engagement
- 597 EAHW: data management and data analysis
- 598 GNH and EAHW wrote the paper with input from all authors.
- 599 Acknowledgements
- 600 We acknowledge and thank the partner investigators and their associated support staff, academic
- 601 investigators, project staff, and external consultants for their contribution to the BeUpstanding program of
- 602 research. We also acknowledge and thank all the participants who have taken part in the program,
- 603 particularly our workplace champions.

604

Table 1: Characteristics of champions, teams and workplaces/organisations participating in the trial

_		All t	All trial participants		Champions leading teams only	
Characteristic	Category / (units or range)	n	M(SD) or n(%)		M(SD) or n(%)	
Champion characteristics						
Location (State)	New South Wales	118	20 (16.9%)	94	17 (18.1%)	
	Australian Capital Territory		5 (4.2%)		5 (5.3%)	
	Victoria		20 (16.9%)		14 (14.9%)	
	Queensland		56 (47.5%)		43 (45.7%)	
	South Australia		8 (6.8%)		7 (7.4%)	
	Western Australia		7 (5.9%)		6 (6.4%)	
	Tasmania		1 (0.8%)		1 (1.1%)	
	Northern Territory		1 (0.8%)		1 (1.1%)	
Postcode SES ^a	Australian Percentile (1-100)	118	68.2 (27.5)	94	69.9 (27.0)	
	State Percentile (1-100)	118	67.6 (26.7)	94	69.3 (26.3)	
Age	(years)	116	42.9 (11.0)	94	43.4 (11.1)	
Female	Yes	116	88 (75.9%)	94	73 (77.7%)	
Job classification	Employee	116	53 (45.7%)	94	40 (42.6%)	
	Middle management	116	45 (38.8%)	94	38 (40.4%)	
	Upper management	116	18 (15.5%)	94	16 (17.0%)	
Occupational Health and Sa	fety Role (Yes)	116	76 (65.5%)	94	60 (63.8%)	
Prior Workplace Health Pro	motion Training (Yes)	116	66 (56.9%)	94	55 (58.5%)	
Prior Workplace Health Pro	motion Experience (Yes)	116	42 (36.2%)	94	32 (34.0%)	
Referred to BeUpstanding b	by (multiples apply)					
	Article / Publication / Newsletter	116	16 (13.8%)	94	13 (13.8%)	
	Internet / Website	116	28 (24.1%)	94	20 (21.3%)	
S	eminar / Presentation / Conference	116	15 (12.9%)	94	14 (14.9%)	
	Social Media / TV / Radio	116	4 (3.4%)	94	3 (3.2%)	
	Colleague	116	38 (32.8%)	94	31 (33.0%)	
	Other word of mouth	116	13 (11.2%)	94	12 (12.8%)	
	General Workplace	116	1 (0.9%)	94	1 (1.1%)	
	Health & Safety Organisation	116	2 (1.7%)	94	2 (2.1%)	
	Email	116	1 (0.9%)	94	1 (1.1%)	
	Other source	116	5 (4.3%)	94	4 (4.3%)	
Team characteristics						
Team Size ⁶		94	29.4 (32.1)	94	29.4 (32.1)	
Team Size Category	1-10	94	30 (31.9%)	94	30 (31.9%)	
	11-20	94	27 (28.7%)	94	27 (28.7%)	
~	>20	94	37 (39.4%)	94	37 (39.4%)	
Current participation in othe program (Yes)	er workplace health promotion	116	31 (26.7%)	94	23 (24.5%)	
Interest in Health ^c	(1-5)	116	3.4 (0.7)	94	3.4 (0.7)	
Motivation to Sit Less ^c	(1-5)	116	3.1 (0.7)	94	3.2 (0.7)	
Stress ^c	(1-5)	116	3.3 (0.6)	94	3.2 (0.5)	
Regional/remote staff include	ded (Yes) ^d	116	41 (35.3%)	94	32 (34.0%)	
Call-centre staff included (Y	(es) ^e	118	6 (5.1%)	94	6 (6.4%)	
Main Location (State) ^f	New South Wales	103	20 (19.4%)	94	17 (18.1%)	
× /	Australian Capital Territory		5 (4,9%)		5 (5.3%)	
	Victoria		17 (16.5%)		14 (14.9%)	
	Queensland		46 (44.7%)		43 (45.7%)	

	South Australia		7 (6.8%)		7 (7.4%)
	Western Australia		6 (5.8%)		6 (6.4%)
	Tasmania		1 (1.0%)		1 (1.1%)
	Northern Territory		1 (1.0%)		1 (1.1%)
Workplace or organisation	r I		· · · ·		
Organisation Size	Small (<20)	118	11 (9.3%)	94	11 (11.7%)
	Medium (20-199)		17 (14.4%)		13 (13.8%)
	Large (200-1999)		34 (28.8%)		24 (25.5%)
	Very large (2000+)		56 (47.5%)		46 (48.9%)
Small-medium enterprise	Yes	118	28 (23.7%)	94	24 (25.5%)
Sector	Public	118	62 (52.5%)	94	48 (51.1%)
	Non-profit		19 (16.1%)		14 (14.9%)
	Private		37 (31.4%)		32 (34.0%)
Blue-collar workplace ^g	Yes	118	40 (33.9%)	94	33 (35.1%)
COVID-19 impact h	None	118	6 (5.1%)	94	6 (6.4%)
*	General		66 (55.9%)		53 (56.4%)
	High		46 (39.0%)		35 (37.2%)
Organisational readiness ⁱ	8		3.8 (0.6)		3.8 (0.6)
Context	(1-5)	99	38(0.6)	94	3.8 (0.6)
Change effort	(0-1)	100	0.4 (0.4)	94	0.4 (0.4)
Change efficacy	(1-5)	116	41(05)	94	4.1 (0.5)
Main industry of	Administrative and Support		(0.0)		
organisation ^j	Services	118	6 (5.1%)	94	6 (6.4%)
-	Agriculture / Forestry and				
	Fishing		4 (3.4%)		2 (2.1%)
	Construction		6 (5.1%)		6 (6.4%)
	Education and Training		6 (5.1%)		6 (6.4%)
	Electricity / Gas / Water and				
	Waste Services		8 (6.8%)		7 (7.4%)
	Financial and Insurance Services Health Care and Social		4 (3.4%)		3 (3.2%)
	Assistance		29 (24.6%)		20 (21.3%)
	Manufacturing		9 (7.6%)		8 (8.5%)
	Other Services		8 (6.8%)		8 (8.5%)
	Professional / Scientific and				
	Technical Services		22 (18.6%)		16 (17.0%)
	Public Administration and				
	Safety		/ (5.9%)		6 (6.4%)
	Retail Trade		2 (1.7%)		2 (2.1%)
	Transport / Postal and		7 (5.00/)		1 (1 20/)
	w arenousing		1 (3.9%)		4 (4.3%)

606 Characteristics of organisations, workplaces, and teams were collected from as many of the champions as possible and linked to each team 607 mostly by using data from the team's primary champion, or if unavailable, from another co-champion for the team, an overseer with

608 knowledge of all the teams, or any other suitable respondent from that workplace.

609 ^a SES = Socioeconomic Status (postal area level Index of Relative Economic Advantage and Disadvantage in national and state

610 percentiles).(50)

 \overline{b} estimated by champions, and periodically updated by project staff during implementation checks as champions evolved their concept of who their team would be

613 ^c champion perception when signing up of team's current level (1=worst response to 5=best response)

^d some or all of the team were in regional or remote work locations as reported by champions or ascertained from staff or champion workplace
 postcodes

616 ^e some or all of the team were call-centre staff according to the champion

^f the primary state in which the team is located, calculated from champion and staff-reported workplace postcodes

618 ^g classified from the reported industry, or by champion report of a blue-collar workplace where the industry was mixed

^h Those participating prior to COVID-19 were treated as no impact. General impact (e.g., working from home) was assigned where there was

620 no evidence of a high impact (decided subjectively). Severe impacts (e.g., job losses, shutdown of core business) reported by the champion, an

621 impact strong enough to prevent participation, being located in Victoria where extended lockdowns occurred, or being part of sectors hard hit

delivering healthcare or COVID-19 response schemes all qualified as high impact.

- 623 ⁱ Higher scores are beneficial(51)
- ^j main industry of the organisation according to the 19 standard ANZIC categories(35)

625

626 **Table 2:** Characteristics of participating staff (n=2228)

Characteristic	(Unit or range) / category	n	Mean (SD) or n(%) ^a
Age	(years)	2004	42.9 (11.4)
Work hours	(hours per week)	2043	38.2 (9.0)
Pre-program exercise ^b	(days per week)	1768	3.1 (2.2)
Meeting recommendations ^b	No <5 days per week	1768	1286 (72.7%)
	Yes, ≥5 days per week		482 (27.3%)
Post-program exercise ^b	(days per week)	709	3.5 (2.1)
Meeting recommendations ^b	No, <5 days per week	709	469 (66.1%)
	Yes, ≥5 days per week		240 (33.9%)
Body Mass Index (BMI) ^c	$<25 \text{ kg/m}^2$	1705	771 (45.2%)
	25-<30 kg/m ²		561 (32.9%)
	$\geq 30 \text{ kg/m}^2$		373 (21.9%)
Female	No	2131	745 (35.0%)
	Yes		1386 (65.0%)
Non-English speaking background	No	2015	1674 (83.1%)
	Yes		341 (16.9%)
Full-time employment	No	2043	365 (17.9%)
	Yes		1678 (82.1%)
Education	High school or less	2024	191 (9.4%)
	TAFE/Trade certificate/ Diploma		530 (26.2%)
	University or other tertiary		1303 (64.4%)
Job classification	Employee	2043	1399 (68.5%)
	Middle management		167 (8.2%)
	Upper management		477 (23.3%)
Job category skill level ^d	1 (most skill / training)	2043	1028 (50.3%)
	2		266 (13.0%)
	3		505 (24.7%)
	4		56 (2.7%)
	5 (least skill / training)		188 (9.2%)
Shift worker	No Yes	2043	2001 (97.9%) 42 (2.1%)

627 ^a Mean SD corrected for clustering using linearized variance (STATA survey commands)

^b number of days per week exercised for at least 30 minutes (0-7)(52) and meeting national guidelines ($\geq 5 v < 5 days$)

^c Excludes invalid responses (truncated to plausible range of 15 to 50)(53)

^d Australian Standard Classification of Occupations (ASCO: 1 most to 5 least skilled)(54)

- 631
- 632

Table 3: Completion of core steps for all participating teams (n=94) and completing teams (n=89) by one or more

634 champion(s) for that team

Core program steps	All teams	Completing teams		
	n (%)	n (%)		
- Pre-program staff survey	94 (100.0%)	89 (100.0%)		
- Workplace audit ^a	71 (75.5%)	71 (79.8%)		
- Staff information and consultation session (e.g., workshop)	94 (100.0%)	89 (100.0%)		
- Strategy selection	73 (77.7%)	78 (87.6%)		
- Program promotion via posters and/or emails	94 (100.0%)	89 (100.0%)		
- Post-program staff survey	94 (100.0%)	89 (100.0%)		
- Program completion survey	80 (85.1%)	79 (88.8%)		
Number of core steps completed (out of 7)				
7	68 (72.3%)	68 (76.4%)		
6	17 (18.1%)	16 (18.0%)		
5	9 (9.5%)	5 (5.6%)		
<5	0 (0.0%)	0 (0.0%)		

^a The workplace audit mostly asked about the workplace in general and was considered completed if it was
 completed by any champion for the workplace

- 000
- 637
- 638

Table 4: Post-program costs reported by the champion (n=72 champions)

Total co	osts reported	Costs per team member				
Cost n (%)		Cost	n (%)			
\$0	52 (72.2%)	\$0	52 (72.2%)			
<\$100	7 (9.7%)	>\$0-\$5	7 (9.7%)			
\$100-\$499	1 (1.4%)	>\$5- \$25	4 (5.6%)			
\$500-\$999	1 (1.4%)	>\$25-\$50	2 (2.8%)			
\$1000-\$1499	4 (5.6%)	>\$50-\$100	2 (2.8%)			
\$1500-\$1999	1 (1.4%)	\$133	1 (1.4%)			
\$2000-\$2499	3 (4.2%)	\$250	1 (1.4%)			
\$7000	1 (1.4%)	\$318	1 (1.4%)			
\$10,500	1 (1.4%)	\$1500	1 (1.4%)			

640

Table 5: Changes in staff-reported main outcomes pre to post BeUpstanding program (k=82 workplaces) ^a

Outcome	Pre-									
	program Unadjusted		Adjusted ^b			Adjusted ^b & multiply imputed ^c				
	M (SD)	n	Change (95% CI)	р	n	Change (95% CI)	р	n	Change (95% CI)	р
Outcomes targeted by the intervent	tion									
Activity as % of workday ^g										
Sitting ^{d,e}	78.3 (17.9)	2131	-7.6 (-9.1, -6.1)	< 0.001	2043	-8.1 (-9.7, -6.5)	< 0.001	2248	-7.7 (-9.6, -5.8)	< 0.001
Standing ^{d,f}	10.7 (11.1)	2131	5.0 (3.3, 6.7)	< 0.001	2043	5.9 (3.8, 7.9)	< 0.001	2248	6.3 (4.3, 8.3)	< 0.001
Moving ^{d,f}	11.0 (10.5)	2131	0.8 (-0.2, 1.8)	0.109	2043	1.3 (0.1, 2.4)	0.032	2248	3.1 (2.1, 4.1)	< 0.001
% of sitting in prolonged bouts ^{d, h}	65.8 (24.5)	2130	-9.1 (-11.0, -7.1)	< 0.001	2042	-9.3 (-11.2, -7.3)	< 0.001	2248	-9.0 (-11.2, -6.7)	< 0.001
Activity preference alignment ⁱ										
Sitting	153.1 (84.0)	2091	-34.3 (-40.7, -27.8)	< 0.001	2043	-33.7 (-40.2, -27.1)	< 0.001	2248	-30.7 (-37.6, -23.9)	< 0.001
Standing	90.3 (62.6)	2091	-17.6 (-22.5, -12.6)	< 0.001	2043	-17.3 (-22.2, -12.3)	< 0.001	2248	-14.8 (-21.8, -7.8)	< 0.001
Moving	70.8 (60.6)	2091	-10.7 (-15.4, -6.0)	< 0.001	2043	-10.3 (-15.1, -5.6)	< 0.001	2248	-6.9 (-12.7, -1.0)	0.023
Perceptions of culture (0-4) ^j										
Overall culture score	2.6 (0.8)	2092	0.37 (0.31, 0.43)	< 0.001	2043	0.37 (0.30, 0.43)	< 0.001	2248	0.34 (0.28, 0.40)	< 0.001
Perceived control over sitting ^e	2.6 (1.1)	2092	0.23 (0.16, 0.30)	< 0.001	2043	0.23 (0.15, 0.30)	< 0.001	2248	0.06 (-0.02, 0.13)	0.136
Organisation supports choices ^e	2.7 (1.0)	2092	0.36 (0.30, 0.43)	< 0.001	2043	0.36 (0.30, 0.43)	< 0.001	2248	0.15 (0.09, 0.21)	< 0.001
Organisation supports moving	2.9 (0.9)	2092	0.28 (0.21, 0.35)	< 0.001	2043	0.27 (0.20, 0.34)	< 0.001	2248	0.05 (-0.02, 0.13)	0.159
Role modelling	2.1 (1.0)	2092	0.47 (0.38, 0.55)	< 0.001	2043	0.46 (0.38, 0.55)	< 0.001	2248	0.33 (0.25, 0.42)	< 0.001
Supportive culture	2.5 (1.0)	2092	0.47 (0.39, 0.55)	< 0.001	2043	0.47 (0.38, 0.55)	< 0.001	2248	0.28 (0.20, 0.36)	< 0.001
Other outcomes										
Feeling part of a team (0-4) ^k	2.6 (1.0)	2043	0.02 (-0.06, 0.10)	0.633	2043	0.01 (-0.07, 0.09)	0.869	2248	-0.05 (-0.13, 0.04)	0.296
Feeling engaged (0-4) ^k	2.5 (0.9)	2043	0.05 (-0.02, 0.12)	0.173	2043	0.03 (-0.04, 0.11)	0.368	2248	-0.04 (-0.13, 0.05)	0.383
Productivity and Health		2043	0.10 (0.02, 0.17)	0.012						
Self-rated job performance (1-7)										
e,1	5.5 (1.0)	2043	0.09 (0.01, 0.18)	0.027	2043	0.09 (0.01, 0.16)	0.019	2248	-0.07 (-0.16, 0.03)	0.148
Job satisfaction (1-7) ^{e,m}	5.1 (1.2)	2043	0.00 (0.00, 0.00)	0.923	2043	0.09 (0.00, 0.17)	0.047	2248	-0.01 (-0.10, 0.07)	0.733
Sick Days past 28 days (0-28) ^{f,n}	0.6 (1.9)	2043	0.03 (-0.06, 0.12)	0.573	2043	0.00 (0.00, 0.00)	0.923	2248	-0.14 (-0.51, 0.23)	0.441
Stress (0-4) °	1.7 (1.1)	2043	0.13 (0.08, 0.18)	< 0.001	2043	0.03 (-0.06, 0.12)	0.507	2248	0.06 (-0.02, 0.14)	0.169
Energy score (0-4) °	2.1 (0.7)	2043	0.02 (-0.05, 0.10)	0.558	2043	0.12 (0.07, 0.18)	< 0.001	2248	0.10 (0.06, 0.14)	< 0.001
Creativity (0-4) °	1.6 (1.0)	2043	0.02 (-0.06, 0.10)	0.633	2043	0.01 (-0.07, 0.08)	0.864	2248	0.05 (-0.01, 0.12)	0.093
Self-rated health (0-100) ^p										

Psychological	2.2 (1.0) 2043	0.07 (0.00, 0.14)	0.038	2043	0.07 (0.00, 0.14)	0.054	2248	0.04 (-0.03, 0.11)	0.222
Physical	2.0 (1.0) 2043	0.12 (0.05, 0.18)	< 0.001	2043	0.12 (0.05, 0.18)	< 0.001	2248	0.07 (0.00, 0.14)	0.044
Musculoskeletal discomfort (0-100) ^q									
Upper body	4.7 (3.0) 2043	-0.55 (-0.78, -0.33)	< 0.001	2043	-0.54 (-0.76, -0.31)	< 0.001	2248	-0.35 (-0.58, -0.13)	0.003
Lower back	3.9 (3.1) 2043	-0.60 (-0.84, -0.37)	< 0.001	2043	-0.58 (-0.82, -0.34)	< 0.001	2248	-0.37 (-0.68, -0.06)	0.020
Lower body	3.8 (3.1) 2043	-0.37 (-0.60, -0.13)	0.002	2043	-0.36 (-0.59, -0.12)	0.003	2248	-0.12 (-0.37, 0.13)	0.323

^a Estimated from linear mixed models, with random intercepts for workplace, and staff (nested within workplace). Reports M and SD with linearized variance to correct for workplace clustering, and contrasts of marginal means. Expressions used for transformed outcomes.

^b Adjusts for variables that may be imbalanced across pre- and post-program respondents: p<0.2 predictors of team participation in post-program surveys (Additional File 6); p<0.2 predictors of staff participation in post-program only versus pre-program only surveys (Additional File 7) using only one variable for concepts measured in multiple ways (organisational readiness, location, and postcode SES). Models adjusted for: small-medium enterprise (yes/no); regional staff (yes/no); inclusion of call-centre staff (yes/no); COVID-19 impact level (none/general/high); public sector (yes/no); organisational readiness (context score); Champion age (years); full-time employment (yes/no); Job category skill (1-5); team health interest level (0-4); Postcode SES.

^c Missing data multiply imputed by chained equations (STATA) with m=20 imputations.

^d Model also includes random slopes for workplace, as likelihood ratio test supported random slopes over random intercept models at p<0.05 (Additional File 8)

^e Inverse log transformed as ln(1 + maximum value – variable).

 $^{\rm f}$ log transformed as ln(variable) for positive or ln (variable + 0.001) for non-negative variables

^g Occupational Sitting and Physical Activity Questionnaire.(34)

^h percentage of their sitting that staff estimate is accrued by sitting for 30+ minutes continuously: 0-100;(55)

ⁱ absolute difference between desired and self-reported percentage (0= perfect alignment; 100=complete misalignment)(22)

^j 0-4;(19) higher=better.

^k single item (0-4; higher=better)

¹ single item;(56) 1-7; higher=better;

^m single item;(57) 1-7; higher=better;

ⁿ sick days in last four weeks (0-28 days; lower=better);(58)

^o 0-4 with higher=better; note: energy is mean of 3 items

^p self-rated physical and psychological health (0=poor; 4=excellent); higher=better.(59, 60)

^q musculoskeletal discomfort experienced over the last week (0=no discomfort; 10=severe discomfort);(61, 62)

Figure 1: Participant flow diagram



Figure 2: Change in staff-reported outcomes from pre to post BeUpstanding intervention

Change (95%CI) in outcomes from pre to post program

Change (95% CI) in units, p



5 **Figure 3:** Staff perceptions of the impact of the BeUpstanding program (n=654)



■ Positive □ Negligible ■ Negative

7 **References**

8	1.	Saunders TJ, McIsaac T, Douillette K, Gaulton N, Hunter S, Rhodes RE, et al. Sedentary behaviour
9		and health in adults: an overview of systematic reviews. Appl Physiol Nutr Metab. 2020;45(10
10		(Suppl. 2)):S197-S217.
11	2.	Ekelund U, Tarp J, Steene-Johannessen J, Hansen BH, Jefferis B, Fagerland MW, et al. Dose-
12		response associations between accelerometry measured physical activity and sedentary time and all
13		cause mortality: systematic review and harmonised meta-analysis. BMJ. 2019;366:14570.
14	3.	Dempsey PC, Biddle SJH, Buman MP, Chastin S, Ekelund U, Friedenreich CM, et al. New global
15		guidelines on sedentary behaviour and health for adults: broadening the behavioural targets. Int J
16		Behav Nutr Phys Act. 2020;17(1):151.
17	4.	Healy GN, Lawler SP, Thorp A, Neuhaus M, Robson EL, Owen N, et al. Reducing prolonged sitting
18		in the workplace (An evidence review: full report). Available at:
19		http://www.vichealth.vic.gov.au/Publications/Economic-
20		participation/Creating_Healthy_Workplaces.aspx. Melbourne, Australia: Victorian Health Promotion
21		Foundation; 2012.
22	5.	Chu AH, Ng SH, Tan CS, Win AM, Koh D, Muller-Riemenschneider F. A systematic review and
23		meta-analysis of workplace intervention strategies to reduce sedentary time in white-collar workers.
24		Obes Rev. 2016.
25	6.	MacEwen BT, MacDonald DJ, Burr JF. A systematic review of standing and treadmill desks in the
26		workplace. Prev Med. 2015;70:50-8.
27	7.	Neuhaus M, Eakin EG, Straker L, Owen N, Dunstan DW, Reid N, et al. Reducing occupational
28		sedentary time: a systematic review and meta-analysis of evidence on activity-permissive
29		workstations. Obes Rev. 2014;15(10):822-38.
30	8.	Tew GA, Posso MC, Arundel CE, McDaid CM. Systematic review: height-adjustable workstations
31		to reduce sedentary behaviour in office-based workers. Occup Med (Lond). 2015;65(5):357-66.

32	9.	Shrestha N, Kukkonen-Harjula KT, Verbeek JH, Ijaz S, Hermans V, Bhaumik S. Workplace
33		interventions for reducing sitting at work. Cochrane Database Syst Rev. 2016;3:CD010912.
34	10.	Nguyen P, Ananthapavan J, Tan EJ, Crosland P, Bowe SJ, Gao L, et al. Modelling the potential
35		health and economic benefits of reducing population sitting time in Australia. Int J Behav Nutr Phys
36		Act. 2022;19(1):28.
37	11.	Straker L, Coenen P, Dunstan D, Gilson N, Healy GN. Sedentary Work – Evidence on an Emergent
38		Work Health and Safety Issue – Final Report. Canberra: Safe Work Australia; 2016.
39	12.	Healy GN, Goode A, Schultz D, Lee D, Leahy B, Dunstan DW, et al. The BeUpstanding Program:
40		Scaling up the Stand Up Australia Workplace Intervention for Translation into Practice. AIMS
41		Public Health. 2016;3(2):341-7.
42	13.	The University of Queensland. BeUpstanding [Available from: <u>www.beupstanding.com.au</u> .
43	14.	Healy GN, Eakin EG, Owen N, Lamontagne AD, Moodie M, Winkler EA, et al. A Cluster
44		Randomized Controlled Trial to Reduce Office Workers' Sitting Time: Effect on Activity Outcomes.
45		Med Sci Sports Exerc. 2016;48(9):1787-97.
46	15.	LaMontagne AD, Noblet AJ, Landsbergis PA. Intervention development and implementation:
47		Understanding and addressing barriers to organisational-level interventions. In: Biron C, Karanika-
48		Murray M, Cooper CL, editors. Improving organisational interventions for stress and well-being:
49		Routledge; 2012.
50	16.	Bandura A. Health promotion by social cognitive means. Health Educ Behav. 2004;31(2):143-64.
51	17.	McLeroy KR, Bibeau D, Steckler A, Glanz K. An ecological perspective on health promotion
52		programs. Health Educ Q. 1988;15(4):351-77.
53	18.	Owen N, Sugiyama T, Eakin EE, Gardiner PA, Tremblay MS, Sallis JF. Adults' sedentary behavior
54		determinants and interventions. Am J Prev Med. 2011;41(2):189-96.
55	19.	Healy GN, Goode AD, Abbott A, Burzic J, Clark BK, Dunstan DW, et al. Supporting Workers to Sit
56		Less and Move More Through the Web-Based BeUpstanding Program: Protocol for a Single-Arm,
57		Repeated Measures Implementation Study. JMIR Res Protoc. 2020;9(5):e15756.

- Healy GN, Winkler EAH, Goode AD. A RE-AIM evaluation in early adopters to iteratively improve
 the online BeUpstanding program supporting workers to sit less and move more. BMC Public
 Health. 2021;21(1):1916.
- Goode AD, Frith M, Hyne SA, Burzic J, Healy GN. Applying a User Centred Design Approach to
 Optimise a Workplace Initiative for Wide-Scale Implementation. Int J Environ Res Public Health.
 2022;19(13).
- Healy GN, Eakin EG, Winkler EA, Hadgraft N, Dunstan DW, Gilson ND, et al. Assessing the
 Feasibility and Pre-Post Impact Evaluation of the Beta (Test) Version of the BeUpstanding
 Champion Toolkit in Reducing Workplace Sitting: Pilot Study. JMIR Form Res. 2018;2(2):e17.
- Goode AD, Hadgraft NT, Neuhaus M, Healy GN. Perceptions of an online 'train-the-champion'
 approach to increase workplace movement. Health Promot Int. 2019;34(6):1179-90.
- Glasgow RE, Harden SM, Gaglio B, Rabin B, Smith ML, Porter GC, et al. RE-AIM Planning and
 Evaluation Framework: Adapting to New Science and Practice With a 20-Year Review. Front Public
 Health. 2019;7:64.
- 72 25. Glasgow RE, Battaglia C, McCreight M, Ayele RA, Rabin BA. Making Implementation Science
- 73 More Rapid: Use of the RE-AIM Framework for Mid-Course Adaptations Across Five Health
- 74 Services Research Projects in the Veterans Health Administration. Front Public Health. 2020;8:194.
- Australian Institute of Health and Welfare. Australian bushfires 2019–20: Exploring the short-term
 health impacts. . Canberra; 2020. Contract No.: Cat. no. PHE 276.
- Stobart A, Duckett S. Australia's Response to COVID-19. Health Econ Policy Law. 2022;17(1):95106.
- Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D, et al. Better reporting of
 interventions: template for intervention description and replication (TIDieR) checklist and guide.
 Bmj. 2014;348:g1687.
- B2 29. Des Jarlais DC, Lyles C, Crepaz N. Improving the reporting quality of nonrandomized evaluations of
 behavioral and public health interventions: the TREND statement. Am J Public Health.
- 84 2004;94(3):361-6.

- 85 30. Buckley JP, Hedge A, Yates T, Copeland RJ, Loosemore M, Hamer M, et al. The sedentary office:
- 86 an expert statement on the growing case for change towards better health and productivity. Br J

87 Sports Med. 2015;49(21):1357-62.

- 88 31. Human Factors and Ergonomics Society of Australia (HFESA) Inc. Sedentary Behaviour: HFESA
- 89 Position on Prolonged Unbroken Sitting Time 2015 [Available from:
- 90 <u>https://www.ergonomics.org.au/documents/item/184</u>. Accessed: 2017-11-06. (Archived by
- 91 WebCite® at <u>http://www.webcitation.org/6ulUGJfFW</u>).
- 92 32. Colberg SR, Sigal RJ, Yardley JE, Riddell MC, Dunstan DW, Dempsey PC, et al. Physical
- 93 Activity/Exercise and Diabetes: A Position Statement of the American Diabetes Association.
- 94 Diabetes Care. 2016;39(11):2065-79.
- 95 33. Australian Institute of Health and Welfare. Changing patterns of work. 2023 7th September.
- 96 34. Chau JY, Van Der Ploeg HP, Dunn S, Kurko J, Bauman AE. Validity of the occupational sitting and
 97 physical activity questionnaire. Med Sci Sports Exerc. 2012;44(1):118-25.
- 98 35. Australian Bureau of Statistics. Australian and New Zealand Standard Industrial Classification
 99 (ANZSIC) Canberra: ABS; 2006 [Available from:
- 100 https://www.abs.gov.au/statistics/classifications/australian-and-new-zealand-standard-industrial-
- 101 <u>classification-anzsic/latest-release</u>.
- Thorp AA, Healy GN, Winkler E, Clark BK, Gardiner PA, Owen N, et al. Prolonged sedentary time
 and physical activity in workplace and non-work contexts: a cross-sectional study of office, customer
 service and call centre employees. Int J Behav Nutr Phys Act. 2012;9:128.
- 105 37. Morris A, Murphy R, Shepherd S, Graves L. Multi-Stakeholder Perspectives of Factors That
- Influence Contact Centre Call Agents' Workplace Physical Activity and Sedentary Behaviour. Int J
 Environ Res Public Health. 2018;15(7).
- 10838.Morris AS, Murphy RC, Shepherd SO, Healy GN, Edwardson CL, Graves LEF. A multi-component
- intervention to sit less and move more in a contact centre setting: a feasibility study. BMC Public
- Health. 2019;19(1):292.

- 111 39. Wang C, Lu EY, Sun W, Chang JR, Tsang HWH. Effectiveness of interventions on sedentary
- behaviors in office workers: a systematic review and meta-analysis. Public Health. 2024;230:45-51.
- 40. Healy GN, Eakin EG, Lamontagne AD, Owen N, Winkler EA, Wiesner G, et al. Reducing sitting
- time in office workers: short-term efficacy of a multicomponent intervention. Prev Med.
- 115 2013;57(1):43-8.
- 116 41. Neuhaus M, Healy GN, Dunstan DW, Owen N, Eakin EG. Workplace sitting and height-adjustable
 117 workstations: a randomized controlled trial. Am J Prev Med. 2014;46(1):30-40.
- Wilms P, Schroder J, Reer R, Scheit L. The Impact of "Home Office" Work on Physical Activity and
 Sedentary Behavior during the COVID-19 Pandemic: A Systematic Review. Int J Environ Res
 Public Health. 2022;19(19).
- 43. Sui W, Smith ST, Fagan MJ, Rollo S, Prapavessis H. The effects of sedentary behaviour
- interventions on work-related productivity and performance outcomes in real and simulated office
 work: A systematic review. Appl Ergon. 2019;75:27-73.
- 44. Oakman J, Clune S, Stuckey R. Work-related musculoskeletal disorders in Australia. Canberra;
 2019.
- 45. Bell Z, Porcellato L, Holland P, Morris A, Smith C, Haines C, et al. A systematic scoping review of
 health-promoting interventions for contact centre employees examined through a behaviour change
 wheel lens. PLoS One. 2024;19(3):e0298150.
- 129 46. Edwardson CL, Biddle SJH, Clarke-Cornwell A, Clemes S, Davies MJ, Dunstan DW, et al. A three
- arm cluster randomised controlled trial to test the effectiveness and cost-effectiveness of the SMART
- Work & Life intervention for reducing daily sitting time in office workers: study protocol. BMC
 Public Health. 2018;18(1):1120.
- 133 47. Shrestha N, Kukkonen-Harjula KT, Verbeek JH, Ijaz S, Hermans V, Pedisic Z. Workplace
- interventions for reducing sitting at work. Cochrane Database Syst Rev. 2018;12(12):CD010912.
- 135 48. Taylor WC, Suminski RR, Das BM, Paxton RJ, Craig DW. Organizational Culture and Implications
- 136 for Workplace Interventions to Reduce Sitting Time Among Office-Based Workers: A Systematic
- 137 Review. Front Public Health. 2018;6:263.

- 49. Schell SF, Luke DA, Schooley MW, Elliott MB, Herbers SH, Mueller NB, et al. Public health
- program capacity for sustainability: a new framework. Implementation Science. 2013;8(1):15.
- 140 50. Australian Bureau of Statistics. Socio-Economic Indexes for Areas (SEIFA) Australia: ABS; 2021
- [Available from: <u>https://www.abs.gov.au/statistics/people/people-and-communities/socio-economic-</u>
 indexes-areas-seifa-australia/latest-release#cite-window1.
- 143 51. Hannon PA, Helfrich CD, Chan KG, Allen CL, Hammerback K, Kohn MJ, et al. Development and
- 144 Pilot Test of the Workplace Readiness Questionnaire, a Theory-Based Instrument to Measure Small
- 145 Workplaces' Readiness to Implement Wellness Programs. Am J Health Promot. 2017;31(1):67-75.
- 146 52. Milton K, Bull FC, Bauman A. Reliability and validity testing of a single-item physical activity
- 147 measure. Br J Sports Med. 2011;45(3):203-8.
- 148 53. Miller Y. Recommendations for the truncation of body mass index in population data. . Liverpool,
- 149 NSW; 2003. Report No.: CPAH03-0005.
- 54. Australian Bureau of Statistics. ANZSCO Australian and New Zealand Standard Classification of
 Occupations.: ABS; 2022 [Available from: <u>https://www.abs.gov.au/statistics/classifications/anzsco-</u>

152 <u>australian-and-new-zealand-standard-classification-occupations/latest-release</u>.

- 153 55. Clark BK, Stephens SK, Goode AD, Healy GN, Winkler EAH. Alternatives for Measuring Sitting
 154 Accumulation in Workplace Surveys. J Occup Environ Med. 2021;63(12):e853-e60.
- 155 56. Bond FW, Bunce D. Job control mediates change in a work reorganization intervention for stress
 156 reduction. Journal of occupational health psychology. 2001;6(4):290-302.
- 57. Dolbier CL, Webster JA, McCalister KT, Mallon MW, Steinhardt MA. Reliability and validity of a
 single-item measure of job satisfaction. Am J Health Promot. 2005;19(3):194-8.
- 159 58. Kessler RC, Barber C, Beck A, Berglund P, Cleary PD, McKenas D, et al. The World Health
- 160 Organization Health and Work Performance Questionnaire (HPQ). J Occup Environ Med.
- 161 2003;45(2):156-74.
- Jurges H, Avendano M, Mackenbach JP. Are different measures of self-rated health comparable? An
 assessment in five European countries. Eur J Epidemiol. 2008;23(12):773-81.

- 164 60. Ahmad F, Jhajj AK, Stewart DE, Burghardt M, Bierman AS. Single item measures of self-rated
- 165 mental health: a scoping review. BMC Health Serv Res. 2014;14:398.
- 166 61. Dickinson CE, Campion K, Foster AF, Newman SJ, O'Rourke AM, Thomas PG. Questionnaire
- 167 development: an examination of the Nordic Musculoskeletal questionnaire. Applied ergonomics.

168 1992;23(3):197-201.

- 169 62. Brakenridge CL, Chong YY, Winkler EAH, Hadgraft NT, Fjeldsoe BS, Johnston V, et al. Evaluating
- 170 Short-Term Musculoskeletal Pain Changes in Desk-Based Workers Receiving a Workplace Sitting-
- 171 Reduction Intervention. Int J Environ Res Public Health. 2018;15(9).

172

173