

# 1 National Implementation Trial of BeUpstanding™: An Online Initiative for Workers to Sit Less and Move

2 More

3

## 4 Authors

Name	Institutional Address(es) and Affiliations
Genevieve N. Healy	The University of Queensland, School of Human Movement and Nutrition Sciences, Brisbane, Queensland, Australia
Ana D. Goode	The University of Queensland, School of Human Movement and Nutrition Sciences, Brisbane, Queensland, Australia
Lisa Ulyate	The University of Queensland, School of Human Movement and Nutrition Sciences, Brisbane, Queensland, Australia
Alison Abbott	Work Health Design (retired), Workplace Health and Safety Qld, Office of Industrial Relations
David Dunstan	1. Physical Activity Laboratory, Baker Heart and Diabetes Institute. 75 Commercial Road, Melbourne, VIC Australia 2. Institute for Physical Activity and Nutrition, Deakin University, Melbourne, VIC Australia
Elizabeth G. Eakin	The University of Queensland, School of Public Health, Brisbane, Queensland, Australia
Nicholas D. Gilson	The University of Queensland, School of Human Movement and Nutrition Science, Brisbane, Queensland, Australia
Lynn Gunning	Comcare, GPO Box 9905, Canberra ACT 2601
Jodie Jetann	The University of Queensland, School of Human Movement and Nutrition Sciences, Brisbane, Queensland, Australia
Anthony D. LaMontagne	Deakin University, Institute for Health Transformation, Determinants of Health Unit, Geelong, VIC, Australia

NOTES: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Marj Moodie	Deakin University, Institute for Health Transformation, Deakin Health Economics, Melbourne, VIC, Australia
Samantha Mulcahy	The University of Queensland, Health and Wellbeing Centre for Research Innovation, School of Human Movement and Nutrition Science, Brisbane, Queensland, Australia
Neville Owen	1. Centre for Urban Transitions, Swinburne University of Technology, Hawthorn, Victoria, Australia 2. Physical Activity Laboratory, Baker Heart & Diabetes Institute, 75 Commercial Road, Melbourne 3004, Victoria, Australia
Trevor Shilton	1. Curtin University, School of Public Health, Kent Street, Bentley, Western Australia 2. World Heart Federation
Leanne Sweeny	Work Health Design, Workplace Health and Safety Qld, Office of Industrial Relations
Leon Straker	School of Allied Health, Curtin University, Perth, Australia
Elisabeth A.H. Winkler	The University of Queensland, School of Human Movement and Nutrition Sciences, Brisbane, Queensland, Australia

5

## 6 Corresponding author

7 Genevieve N. Healy

8 The University of Queensland, School of Human Movement and Nutrition Sciences

9 St Lucia, Brisbane, Queensland, 4072

10 Email: [g.healy@uq.edu.au](mailto:g.healy@uq.edu.au)

11

## 12 Abstract

13 **Background:** The online BeUpstanding<sup>TM</sup> program is an eight-week workplace-delivered intervention for  
14 desk-based workers to raise awareness of the benefits of sitting less and moving more and build a  
15 supportive culture for change. A workplace representative (the “champion”) delivers the program, which  
16 includes a workshop where teams collectively choose their sit less/move more strategies. A toolkit  
17 provides the champion with a step-by-step guide and associated resources to support program uptake,  
18 delivery, and evaluation. Here we report on the main findings from the Australian national implementation  
19 trial of BeUpstanding.

20 **Methods:** Recruitment (12/06/2019 to 30/09/2021) was supported by five policy and practice partners,  
21 with desk-based work teams from across Australia targeted. Effectiveness was measured via a single arm,  
22 repeated-measures trial. Data were collected via online surveys, toolkit analytics, and telephone calls with  
23 champions. The RE-AIM framework guided evaluation, with adoption/reach (number and characteristics);  
24 effectiveness (primary: self-reported workplace sitting time); implementation (completion of core  
25 components; costs); and, maintenance intentions reported here. Linear mixed models, correcting for  
26 cluster, were used for effectiveness, with reach, adoption, implementation, and maintenance outcomes  
27 described.

28 **Results:** Of the 1640 website users who signed-up to BeUpstanding during the recruitment period, 233  
29 were eligible, 198 (85%) provided preliminary consent, and 118 (50.6%) champions consented and started  
30 the trial, with 94% (n=111 champions) completing. Trial participation was from across Australia and  
31 across industries, and reached 2,761 staff, with 2,248 participating in the staff survey(s): 65% female, 64%  
32 university educated; 16.9% from non-English speaking background. The program effectively changed  
33 workplace sitting (-38.5 [95%CI -46.0 to -28.7] minutes/8-hour workday) and all outcomes targeted by  
34 BeUpstanding (behaviours and culture), with small-to-moderate statistically-significant effects observed.  
35 All participating teams (n=94) completed at least 5/7 core steps; 72.4% completed all seven. Most  
36 champions spent \$0 (72%) or >\$0-\$5 (10%) per team member; most (67/70 96%) intended to continue or  
37 repeat the program.

38 **Conclusions:** BeUpstanding can be adopted and successfully implemented by a range of workplaces, reach  
39 a diversity of staff, and be effective at creating a supportive culture for teams of desk-based workers to sit  
40 less and move more. Learnings will inform optimisation of the program for longer-term sustainability.

41  
42 **Trial registration:** ACTRN12617000682347.

43 The trial was prospectively registered on the 12<sup>th</sup> May, 2017 (ACTRN12617000682347), prior to the soft  
44 launch of the program, with the last update on the 11<sup>th</sup> June, 2019 prior to the commencement of  
45 recruitment to the trial on the 12<sup>th</sup> June, 2019.

## 46 47 **Keywords**

48 Implementation trial; workplace; sitting; health promotion; activity; health and safety; public health;  
49 occupational; web-based; RE-AIM

50

51

## 52 Introduction

53 High levels of sedentary time can be associated with poor health, wellbeing and productivity outcomes and  
 54 indicators (1) with detrimental impacts particularly pronounced in those who are also physically  
 55 inactive.(2, 3) The desk-based workplace has been identified as a key target setting for intervening on  
 56 prolonged sedentary time,(4) with systematic review evidence concluding that interventions addressing  
 57 prolonged workplace sedentary time can be both effective and acceptable.(5-9) Recent economic analyses  
 58 have also indicated that reductions in sedentary time, to the levels achieved through workplace  
 59 interventions, could result in Australian healthcare cost savings of \$39 million per year.(10) Given this  
 60 evidence, and the associated identification of workplace sedentary time as an emergent workplace health  
 61 and safety issue,(11) there is demand from occupational policy and practice partners for an evidence-based,  
 62 low-cost/no-cost, scalable solution to support workers and organisations to reduce prolonged sedentary  
 63 time.(12)

64 The online BeUpstanding™ program (13) is based on a strong foundation of research evidence, including  
 65 from randomised controlled trials (14) and best-practice approaches to workplace health promotion.(15)  
 66 The 8-week program, targeted at desk-based work teams and using a champion-led approach, has a  
 67 theoretical basis in social cognitive theory (16) and a social-ecologic model.(17, 18) The program is  
 68 designed to raise awareness of the benefits of sitting less and moving more and create a culture of  
 69 sustainable change through teams collectively identifying and promoting strategies to support these  
 70 behaviours that are suitable for their work team and context.(19)

71 BeUpstanding has been iteratively developed and tested across multiple phases in collaboration with end  
 72 users and key policy and practice stakeholders,(12, 20-23) and guided by the RE-AIM (reach,  
 73 effectiveness, adoption, implementation, maintenance) framework.(24) Use of this framework is relevant  
 74 for informing broader dissemination as it guides design and evaluation of programs in applied  
 75 contexts.(25) This study reports on the adoption, reach, effectiveness, selected implementation, and  
 76 maintenance intent outcomes from the Australian national implementation trial (Phase 4) of BeUpstanding.  
 77 Cost-benefit analyses, and detailed implementation, maintenance, engagement and process analyses, will

be reported separately. Importantly, the implementation trial, which started in mid-June 2019, occurred in the context of major events in Australia and globally, namely the Australian Black Summer bushfires (July 2019 to March 2020) and the COVID-19 pandemic (March 2020--), which caused major disruptions to ways of working.(26, 27) Findings from this evaluation are intended to inform further optimisation of the BeUpstanding program prior to Phase 5 (dissemination).

## Methods

### Study design

Detailed methods for this implementation trial have been previously reported.(19) In brief, a single-arm design was used for evaluation of effectiveness, with repeated cross-sectional evaluations at pre-program (0 weeks), end-of-program ( $\approx$ 8 weeks; primary endpoint), and at 9 months post-program ( $\approx$ 12 months post sign-up). Only the pre-program and end-of-program data are reported herein, with data reporting according to the TIDieR (28) and TREND (29) checklists (Additional Files 1 and 2). The implementation trial was funded by a National Health and Medical Research Council (NHMRC) of Australia Partnership Project Grant (#1149936), which included cash and/or in-kind support from the five partners (Safe Work Australia, Comcare, Queensland Office of Industrial Relations, The Victorian Health Promotion Foundation (VicHealth), and Healthier Workplace Western Australia). These organisations are responsible for developing, implementing and/or promoting workplace health, safety and wellbeing in Australian workplaces. Ethical approval was from The University of Queensland Human Research Ethics Committee (approval #2016001743). Recruitment into the trial ceased on the 30<sup>th</sup> September, 2021, noting the online program remained available for anyone to access and use. Only data from users who signed-up during the recruitment period are reported here.

### Participants and recruitment

Promotional efforts, co-ordinated by a detailed marketing and promotional plan developed by the research team and external marketing and communication experts, were made by the five partners to direct potential

users to the BeUpstanding website (13) and the implementation trial. Multiple promotional channels were used, including social media, web links, email listservers, newsletters, workplace health promotion and occupational health networks, conferences, and workshops. Recruitment efforts were aimed at champions and decision makers of desk-based employees from a wide cross-section of industries, especially including the five sectors identified as priorities by the partners (regional, call centre, small business, blue-collar, and government).

Recruitment into the trial was a multi-step process. The BeUpstanding online program was freely available online for the entire trial period and beyond, without any requirement to be eligible or participate in the trial. A user could register (by completing preliminary details of themselves and their organisation) and fully sign-up to access the toolkit and associated resources by completing a champion profile survey and providing informed consent. Preliminary eligibility for the trial was ascertained from these two data sources (initial registration; champion profile survey), with eligibility being: not having previously run the BeUpstanding program; team size of at least five staff; and, most of the team do predominantly desk-based work. Those potentially eligible were contacted by telephone (call 1) by the research team to confirm criteria and ascertain the final eligibility requirements. Namely, that a workplace representative/employee signing up to the toolkit could perform the duties of a workplace champion and planned to run the program within the recruitment window. Potential champions were made aware that the main feature of taking part in the trial (as distinct from using the program independently) was they were committing to the evaluation steps in the BeUpstanding program and would receive expert health coaching support to deliver the program via email and phone calls.

All confirmed eligible champion participants were invited to participate in the trial, except where recruitment quota had been reached for a particular priority sector (see sample size) or a limit was reached for coaching availability to be allocated to a single organisation. Consent for the trial was in addition to that provided as part of the website sign-up process. The program was intended to be delivered to all staff

in participating teams, with no exclusion criteria at the staff level. Staff provided informed consent as a step for gaining entry to the staff surveys.

### **BeUpstanding program – national implementation trial version**

A detailed description of the BeUpstanding program is reported elsewhere.<sup>(19)</sup> In brief, the program was designed to be run by the workplace champion across three phases (plan, do, review), each with associated tasks for the champion to complete. The accompanying toolkit provided information and training for the champion on the purpose of each phase and associated resources to support implementation, with the seven core components highlighted (starred) to indicate their importance. The most critical was the staff information and consultation workshop (or equivalent), where staff were provided information about the benefits of sitting less/moving more and collectively chose three (or more) team strategies they would use to sit less/move more as a team. Champions were encouraged to promote these strategies over eight weeks via the posters and emails provided in the toolkit. This participative approach meant each intervention program was unique for each team. In line with public, occupational, and clinical guidelines,<sup>(30-32)</sup> behavioural targets were for workers to achieve a 50:50 split between sitting and upright activities during work hours and to alternate sitting/upright posture at least every 30 minutes. Increased incidental movement throughout the day was also encouraged through the move more messaging. No incentives were provided as part of the trial. The main distinguishing feature of the implementation trial (as compared with the program in general) is that the research team both supported (and collected further data on) program implementation, via email support and telephone calls at five key timepoints: 1) recruitment; 2) obtaining consent; 3) at program initiation, following the staff consultation; 4) at end of program; and, 5) after approximately 12 months (maintenance). Project staff providing this support all had a minimal Masters level qualification and were experienced in motivational interviewing for behaviour change.

Due to the major shift to hybrid work resulting from COVID-19,<sup>(33)</sup> the toolkit and resources were audited and modified, and new resources were developed (e.g., see Additional File 3) during the trial to ensure the program was suitable for delivery for all desk-workers, no matter where they were working. These new



resources went online in July 2020. Additional measures to capture work-from-home arrangements (staff completed) and the impact of COVID-19 on the workplace (champion completed) were also added (May 2020).

## **Data collection**

Detailed description of the data collection process is provided in the protocol paper.(19) Data were primarily collected via the dedicated, stand-alone BeUpstanding website, using surveys and toolkit analytics, with the project management team collecting and confirming implementation data via the telephone check-ins. Each toolkit user was required to complete the registration survey and champion profile survey as part of the sign-up process. Each champion was requested to complete a workplace audit before delivering the program and then a program completion survey after program delivery. Champions were also responsible for sending their bespoke links for the staff surveys (pre-program, post-program) to all staff in their team. For each survey, an anonymous staff identifier was constructed based on responses to three questions (month of birth; first initial of mothers first name; and, last three digits of mobile phone). This identifier, coupled with the cluster identification number, was used to ensure, per combined work team (cluster), each staff member responded only once and to match pre- and post-program responses. There was no blinding.

## **Outcomes and Measures**

The outcomes according to the RE-AIM framework are outlined below.

### ***Adoption***

Adoption was chiefly described in terms of uptake of the toolkit (n and % unlocking the toolkit), trial participation (n and %), the organisational, workplace, team and champion characteristics of those who participated in the trial, and how they had heard of BeUpstanding (in order to adopt it). Organisational characteristics were reported by champions, reconciled across champions from the same organisation, and checked against public records. Workplace characteristics were permitted to diverge between different

champions from the same workplace. Champions were classified by project staff according to their role delivering the program to a team (overseer and primary champion / primary champion / co-champion) or supporting other aspects of BeUpstanding (overseer only / decision maker / other role). Further detailed information relevant to understanding adoption were the predictors of non-adoption (comparison of participants with non-participants), reasons given for not participating or withdrawing, and reasons for adopting the program, focusing specifically on what the champions said that they aimed to achieve by running BeUpstanding (open ended).

### ***Reach***

Reach outcomes included: the number of staff in participating teams (as estimated by the champion, and as verified through staff surveys), and the characteristics of staff who took part in the evaluation (by responding to one or both staff surveys).

### ***Implementation***

Implementation outcomes reported are completion rates (yes/no) of the seven core program steps, and champion-reported costs they incurred (reported in the program completion survey) or expected to incur (reported in the planning stage implementation check, shortly after selecting team strategies) through running BeUpstanding.

### ***Effectiveness***

Program effectiveness was assessed by the difference between pre-and post-program staff survey responses regarding 13 indicators of the program's impact on workplace behaviour and workplace culture targeted directly by the program and 13 measures of productivity and health and wellbeing that were expected might improve as a result of the program. The primary outcome for effectiveness was workplace sitting, measured by the Occupational Sitting and Physical Activity Questionnaire (OSPAQ).(34) The post-staff survey also captured staff perceptions of the program and its impact, and adverse events. When described, adverse events were classified by two researchers as to whether they constituted adverse events and

whether they could reasonably have resulted from the program. Program satisfaction was collected from staff in post-program surveys and from team champions in program completion surveys.

### ***Maintenance intentions***

The maintenance intentions of the champions of completing teams were ascertained by coding and reviewing the responses to an open text question of “What are your plans now in relation to BeUpstanding?” asked during program completion calls. Their detailed actual maintenance of the program and longer-term outcomes will be addressed in detail in subsequent papers.

### **Sample size (primary effectiveness outcome)**

As described elsewhere,(19) using assumptions informed by earlier iterations of the program (SD 90,  $r=0.5$ ,  $ICC=0.1$ , post-attrition  $n/workplace = 5$ ), it was determined that 47 to 62 workplaces (235 to 310 staff) were required to provide 80% to 90% power to detect a change in workplace sitting of at least 20 minutes per 8 hours of work (i.e., 4% of the workday) with 5% two-tailed significance. Consequently, the target sample size was at least 50 workplaces per priority sector to permit subgroup analysis.

### **Statistical Analysis**

Most of the reach, adoption, implementation and maintenance outcomes are described using descriptive statistics. Additionally, logistic regression was used to compare the organisational, workplace, team, and personal characteristics of those who included in the trial (trial adopters) versus non-participants (non-adopters), as well as those who completed the post-program evaluation of their team versus those who did not. Effectiveness outcomes (continuous) were compared between pre- and post-program staff surveys using linear mixed models, correcting for cluster (workplace) as a random intercept, or as random slopes, where likelihood ratio tests indicated the random slopes model was superior. Unadjusted models were reported in addition to the main findings, which adjusted for potential confounding due to different staff and teams providing data at each survey. Specifically, these were characteristics with a  $p<0.2$  association

either with teams being part of the post-program evaluation or not (tested by logistic regression models), or with staff members providing outcome data at only the pre-program survey (n=1481) versus only the post-program survey (n=397) (tested by mixed logistic regression models, correcting for workplace cluster as a random intercept). To test the sensitivity of conclusions to missing data handling, confounder-adjusted models were also reported with missing data imputed, using multiple imputation by chained equations. In addition to the analytic variables, imputation models contained auxiliary variables (additional variables predicting outcomes at either pre- or post- program surveys at  $p < 0.2$  in backwards eliminations) to help improve prediction of missing outcomes. A further sensitivity analysis excluded staff survey responses that had been collected after the program had already started. Analyses were performed in STATA version 18, with the final date of data extraction 15 May 2024. Significance was set at  $p < 0.05$  (two-tailed).

## Results

For ease of reporting, RE-AIM outcomes are presented these in the order of Adoption, Reach, Implementation, Effectiveness, and Maintenance intentions.

---

**INSERT FIGURE 1 ABOUT HERE**

---

### Adoption

The participant flow diagram (Figure 1) shows that out of the 1640 website users who signed up during the trial recruitment period, 847 fully registered and unlocked the toolkit, and 233 were determined to be eligible for the trial, with ineligibility mostly due to not being contactable or not planning to run the program either at all or in the requisite timeframe. In total, 82 potentially would have been eligible if the recruitment quota for their sector(s) was considered not already met. Out of the 233 eligible users, the preliminary consent rate was 85.0% (n=198), which dropped to 50.6% (n=118) after excluding those who preliminarily seemed eligible and consented but withdrew before starting the program. Completion rates for the 118 who were included in the trial (who commenced the program) were high (94.1%, n=111). The

most common reasons for non-participation prior to preliminary consent were reasons related to not being able to run the program, or lack of interest in the trial. Reasons for not participating after preliminarily consenting were primarily due to changing circumstances at either the organisation or champion level. Withdrawing after having started the program was rare and most often due to it being a difficult time in the organisation (n=5/7).

Almost all trial respondents (n=116) indicated what they aimed to achieve by adopting the program, with results categorised under 22 themes (Additional File 4). The most common aim was to improve either physical health and/or wellbeing (50.9%), with changing sitting / physical activity levels (37.9%) also often mentioned. There was some evidence that aims were associated with trial participation (Additional File 5). Reporting an aim to improve culture (OR=1.95, 95%CI: 1.09, 3.48), connection (OR=1.85, 95%CI: 1.02, 3.35), and/or productivity (OR=2.78, 95%CI: 1.47, 5.24) were associated with significantly higher odds of participation. Reporting a general aim to improve health was associated with a significantly lower odds of participation (OR=0.30, 95%CI: 0.09, 0.98).

Table 1 shows the characteristics of teams participating in the trial and their champions. The sample size target was met in terms of total numbers but not in any of the subgroups, due in part to the COVID-19 pandemic. Teams were recruited from every priority sector: 48 teams (51.1%) from the public sector; 24 teams (25.5%) from small-medium enterprises (11 (11.75%) small business); 33 teams (35.1%) from blue-collar workplaces; 32 teams (34.0%) with regional staff; and, six teams with call-centre staff (6.4%). Team size was highly varied and averaged 29.4 (SD=32.1) members. Overall, 13 of the 19 standard industry classifications in Australia (35) had some representation in the trial. The most common industry of the organisations of participating teams was Health Care and Social Assistance (n=16, 17.0%) and Professional / Technical and Scientific Services (n=16, 17.0%). Teams were recruited from every state and territory of Australia and were predominantly located in Queensland (n=43, 45.7%). Champions' work locations varied widely in socioeconomic status, with a mean (SD) state percentile of 67.6 (26.7) in terms of socioeconomic index of advantage and disadvantage. Champions had a mean (SD) age of 42.9 (11.0)

years. Many were female (75.9%), in middle (38.8%) or senior (15.5%) management, had a health and safety role (65.5%), and workplace health promotion training (56.9%). Less than half (36.2%) had prior experience delivering workplace health promotion programs.

---

### INSERT TABLE 1 ABOUT HERE

---

Additional File 5 shows the odds of participation (adopting the program) by these characteristics. Significant predictors included location ( $p < 0.001$ , with highest participation in Queensland, and lowest participation in WA/NT/TAS), sector ( $p = 0.001$ , with highest participation by public followed by non-profit then private sector), and COVID-19 impact level ( $p < 0.001$ , with highest participation in high followed by general followed by no impact). Predictors significantly associated with lower odds of participation were: having an occupational health and safety role (0.60, 95% CI: 0.39, 0.91); inclusion of call centre staff (0.38, 95% CI: 0.16, 0.89); small-medium enterprise (0.50, 95% CI: 0.32, 0.79); and, hearing of the program through an article/publication/newsletter (0.56, 95% CI: 0.32, 0.98). Predictors significantly associated with higher odds of participation were: hearing of the program through a seminar/presentation/conference (1.97, 95% CI: 1.07, 3.64) or a colleague (2.29, 95% CI: 1.48, 3.52); and, higher organisational readiness (1.52, 95% CI: 1.09, 2.12 per unit change efficacy).

### Reach outcomes

Based on estimates from participating champions, the program reached 2761 staff in the trial, with 2248 able to be verified via responding to at least the identifier questions in one or both staff surveys. Staff were on average (mean (SD)) aged 42.9 (11.4) years, worked 38.2 (9.0) hours per week, and exercised for 3.1 (2.2) days per week prior to BeUpstanding, with 27.3% of staff meeting the recommended five or more days of exercise per week (Table 2). More than half of staff were female (65.0%), in full-time employment (82.1%), university educated (64.4%), non-managerial employees (68.5%), and in the highest occupational skill category (50.3%). A very small minority were shift-workers (2.1%), and 16.9% of staff were from a non-English speaking background.

---

**INSERT TABLE 2 ABOUT HERE**

---

**Implementation outcomes**

The champions had completed all core steps for most of the participating teams (72.4%) and most of the completing teams (76.4%). The steps with the lowest completion rates were the workplace audit, strategy selection, and the program completion survey (Table 3). During their planning stage implementation check-in calls, shortly after selecting their team strategies, 80 champions reported on any costs they had incurred or expected to incur in order to support or enact their team strategies. In total, 35 cost items were reported, the most common of which was sit-stand desks or similar (n=14). Other costs included: accessories to assist with standing at the desk (Bluetooth headsets (n=3); anti-fatigue mats (n=1)), aids for prompting (reminder/lockout software (n=4); timer (n=1)), enhancements to communal areas (standing benches/tables (n=3); communal bins (n=1)), equipment (n=2) and subscriptions (n=1) for physical activity at work; and, miscellaneous other (pedometers (n=1); undescribed equipment (n=2); sporting goods gift cards (n=1)). In the program completion survey, 72 champions reported what was spent in total to run the program for their team, with the majority reporting spending \$0 (n=52, 72.2%, Table 4). Sit-stand desks had been purchased for every team where the amount spent was >\$1000.

---

**INSERT TABLES 3 AND 4 ABOUT HERE**

---

**Effectiveness outcomes**

***Outcomes targeted by the intervention***

Baseline levels of the staff-reported program outcomes and their changes over the program are shown in Table 5. The main models adjusted for characteristics that might differ between those providing data at post- and pre-program surveys, based on participation in the post-program evaluation by the team (Additional File 6) or staff member (Additional File 7). These models (Figure 2) indicated significant

improvements in all outcomes directly targeted by the program: the primary outcome (self-reported work sitting); work standing and moving; the extent to which sitting was accumulated in prolonged bouts; the alignment between preferred and actual sitting, standing and moving at work; and, all measures of workplace culture. The magnitude of these effects on average were mostly small-to-moderate, with the smallest effects seen for moving, moving alignment, and perceived control over sitting. Workplaces varied significantly in their changes in activity at work and sitting in prolonged bouts, with tests for random slopes reaching statistical significance (Additional File 8).

---

**INSERT FIGURE 2 AND TABLE 5 ABOUT HERE**

---

Conclusions regarding workplace behaviours from the main adjusted models were mostly robust to modelling choices, except that moving results strengthened slightly and became significant in adjusted models and then further strengthened with imputation. Results concerning culture were unaffected by statistical adjustment but were sensitive to missing data handling, whereas results from imputation models were often weaker and sometimes no longer statistically significant. Excluding the pre-program surveys that had been collected after the program commenced had no meaningful impact on findings (Additional File 9).

***Other outcomes***

Of the 13 outcomes concerning work productivity and health, significant improvements were seen in self-rated job performance, job satisfaction, energy, self-rated physical health, and musculoskeletal discomfort in the upper body, lower back, and lower body, with improvements ranging in magnitude from very small to small (Figure 2; Table 5). Other changes were consistently all very small, and not statistically significant. Random slopes tests were not statistically significant for any of these outcomes (Additional File 8). Findings were similar to those from unadjusted models, except that adjustment brought results for engagement even closer to the null and the previously significant change in self-rated psychological health became borderline significant. There were no meaningful changes in outcomes excluding late pre-program



surveys (Additional File 9). Many findings, especially for work-related and productivity outcomes, were highly sensitive to missing data, where multiple imputation results only indicated significant changes (all improvements) in energy, self-rated physical health, and musculoskeletal discomfort in the upper body and lower back, all of which were somewhat smaller than they had been in the evaluable case analyses. In imputed models, effects for lower body discomfort were somewhat attenuated in magnitude and no longer statistically significant, while effects for estimates of changes in self-rated job performance and job satisfaction were completely attenuated. Effect sizes became further away from the null for sick days, stress, and creativity, but remained very small and did not reach statistical significance.

### ***Priority sectors***

The changes in the primary outcome (work sitting) overall amounted on average to -38.5 (-46.0, -31.0) min/8 h for the trial in general. Changes were mostly also present within each of the priority sectors: -28.2 (-39.7 to -16.6) min/8-h in small-medium enterprises (based on 24 workplaces and 496 staff); -35.8 (-49.4 to -22.1) min/8-h in blue-collar workplaces (based on 26 workplaces and 772 staff), -41.7 (-52.5 to -31.0) min/8-h in public-sector workplaces (based on 40 workplaces and 954 staff); and, -45.2 (-60.0 to -30.3) min/8-h in teams that included regional staff (based on 26 workplaces and 626 staff). Changes in the teams that included call-centre staff were -46.3 (-126.4 to 33.9) min/8-h and did not reach statistical significance with the small sample of 3 workplaces and 97 staff.

### ***Adverse events and staff perceptions of the program***

Most (n=604, 96.0%) of the 629 staff responding to the staff follow-up surveys did not experience any adverse events. Only 7 staff (1.1%) reported an adverse event that was confirmed by research staff, with six of these (1.0%) potentially being attributable to the study, and a further 18 staff (2.9%) reported that they experienced some adverse event that could not be verified as they did not describe it. The six events that could have arisen from program participation were: new or exacerbation of musculoskeletal pain or injury (n=4), specifically to the hip from standing propped on a stool, to the back from “overdoing things”,

plantar fasciitis, and lower leg and lower back pain; and, adverse work impacts (n=2), specifically feeling guilty about leaving the desk (n=1) and feeling followed by their supervisor (n=1).

The majority of staff reported positive impacts on outcomes targeted directly by BeUpstanding (culture, knowledge, attitudes, awareness), with few reporting negative impacts (Figure 3). Staff provided more mixed responses in relation to the impact of the program on activity outside work, with very few staff (n=11, 1.7%) perceiving a negative impact and the rest of the staff being almost equally divided between perceiving a negligible impact (51.1%) or a positive impact (47.2%).

---

### INSERT FIGURE 3 ABOUT HERE

---

#### ***Program satisfaction***

Staff (n=654) reported on their satisfaction with the program. Excluding those who indicated they were not aware of BeUpstanding or responded not applicable (n=35), most staff (n=477, 77.1%) indicated they enjoyed BeUpstanding, some were uncertain (n=115, 18.6%), and a minority of staff said they did not enjoy participating (n=27, 4.4%). Champions also reported on their satisfaction with the program in post-completion surveys (n=72). When asked whether the program met expectations, most said yes (n=57, 79.2%) or somewhat (n=13, 18.1%) and only two (2.8%) said no. Similarly, most said they would (n=63, 87.5%) or maybe would (n=8, 11.1%) recommend the program and only one said they would not (1.4%).

#### **Maintenance intentions**

The 8-week BeUpstanding program was completed by 89 of the 94 teams that were part of the trial. Upon completion, project staff were able to contact 70 of their team champions and ask about their planned next steps concerning BeUpstanding. Most responses to the open-ended question indicated the champions intended to continue the program (n=59, 84.3%) or repeat it (n=8, 11.4%), while very few indicated that the champion intended to stop or pause the program (n=1, 1.4%), including to restart it with different staff (n=1, 1.4%), and one champion intended to check with management (n=1, 1.4%).

423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449

# **Discussion**

This study reports the main outcomes, covering the five dimensions of the RE-AIM framework, of the first national implementation trial of a workplace health promotion initiative specifically supporting teams of desk-based workers to sit less and move more (BeUpstanding). This trial was the fourth phase in the research-to-practice translation process from an intervention that had demonstrated effectiveness in the context of a cluster randomised controlled trial.(14)

## ***Adoption and Reach***

While there was interest in the program more broadly, with over 600 organisations unlocking the toolkit during the trial period, adoption of the program as part of the research trial was more modest, with approximately half of eligible workplaces running the program. This needs to be understood in the context of the BeUpstanding program being available for free online without participation in the trial, and two significant emergencies in Australia that occurred during the trial’s recruitment period, specifically the COVID-19 pandemic and the 2019-20 bushfires. Reported reasons for not adopting the program were mostly regarding it being a difficult time in the organisation (e.g., availability of personnel) or lack of interest in the research trial. Only rarely did withdrawal occur after initial consent due to an issue with the program. As might be expected, a higher level of organisational readiness, specifically change efficacy, was significantly associated with higher odds of adoption into the trial while conversely having an occupational health and safety role was associated with lower odds of adoption. One interpretation (among many) is that champions who do not feel their workplace is well placed to run the program may require additional supports (beyond what was provided in the implementation trial) to adopt the program, while conversely those in occupational health and safety roles may have felt confident to run the program independently outside of the trial context without support. Other reasons could include that occupational health and safety staff had competing priorities especially in the COVID-19 pandemic, and unready workplaces may have wanted to avoid the oversight of a trial.

Workplaces and staff took part from every partner-identified priority sector, from every Australian state and territory (including rural/regional locations), from 13 of 19 ANZIC industries,(35) and with diverse workplace and worker characteristics. Some groups were easier to reach than others, with limited participation from shift workers, from those not in fulltime employment, and with lower-skilled occupations. While this may align with our specific focus on desk-based workers, it is important in future research to understand how this program and others like it may work in these groups, as well as other underrepresented or untested key populations, such as people with disabilities. Adoption was significantly lower for workplaces outside the public sector, of small-medium size, and especially in teams that included call-centre staff (despite deliberate recruitment efforts). Staff in call centres or contact centres generally have higher rates of sedentary time than general office workers,(36) but typically also have reduced autonomy to break up their sedentary time due to their job tasks and job demands.(37) In line with previous recommendations,(38) it will be important to work with relevant stakeholders to co-design appropriate approaches for this particular setting.

### ***Implementation***

Champions were provided automated feedback via the toolkit that was engaging and meaningful and had a streamlined online user experience that highlighted the core components. As part of the trial, they also received support from the research team at key stages: all factors that had been added and optimised based on learnings from the version used by the early adopters.(21) In the previous phase of this research-to-practice translation (Phase 3), the seven core steps were completed for only 5% of teams, due in part to retention and in part to progressing through without completing core steps.(20) These modifications appear to have resulted in a much more complete implementation of the program, with most teams (>72%) completing all seven core steps, and none completing fewer than five. Most workplaces implemented the program in a no or low-cost manner, and only a minority spent significant sums of money, usually on sit-stand workstations.

### ***Effectiveness***

The program was significantly effective in relation to the primary outcome (workplace sitting), behaviours and culture (which were the main targets of the program), and also for some indicators of productivity, health and wellbeing. The average decrease in self-reported work sitting of -38.5 (95%CI -46.0 to -31.0) minutes per 8-hour workday appeared to have co-occurred mainly with increased standing, and also a smaller amount of increased moving. Notably the extent of change strongly resembled the average effect on sedentary behaviour of 38 minutes reduction per workday (95% CI -47.3 to -28.7) achieved by sedentary behaviour interventions in office workers (across all intervention types combined) as reported by a recent systematic review and meta-analysis.(39) Given the timing of the trial, and the substantial shift to hybrid work resulting from the COVID-19 pandemic,(33) these findings also provide some evidence that BeUpstanding is suitable for work from home and hybrid work contexts. A pre-post design was required for feasibility reasons, leaving open the possibility of other explanations for changes over time, especially in light of the pandemic. Notably however, the controlled studies that informed the BeUpstanding program development had not observed any meaningful degree of changes in control groups.(14, 40, 41) It is not likely that the pandemic produced the improvements, considering COVID-19 saw an increase in sedentary time and a reduction in physical activity in desk workers(42), and the average change was -43 minutes per 8-hour workday (- 9.0%, 95% CI: - 12.0% to - 5.9%) in the pre-pandemic evaluation of the program's early adopters.(20)

Findings concerning workplace sitting, other workplace behaviours and workplace culture were robust to issues of potentially selective evaluation, being largely unaltered in sensitivity analyses. Where selective participation may have been more of an issue was in gauging the impact of the program on productivity and health, where changes were typically smaller after accounting for missing data, but notably were still present for energy, self-rated physical health, and upper body and lower back musculoskeletal discomfort. The benefits seen to at least some of the productivity and health indicators adds to the growing evidence from field-based studies that interventions aimed at reducing workplace sedentary time, particularly those where sedentary time is predominantly replaced with standing, are not detrimental to indicators of work performance.(43) The musculoskeletal findings in particular add to the currently limited evidence base in

this area.(17) Although these reductions were small on average, they are potentially important given that work-related musculoskeletal disorders are the leading work health and safety issue in Australia in terms of both frequency and costs,(44) and were not necessarily small for all participants. Future detailed evaluation will explore whether changes are possibly enhanced based on factors such as pre-existing discomfort, and nuanced aspects of the behaviour changes, such as the reallocation of sitting to standing versus to moving, the prolonged or interrupted nature of sedentary accumulation, and the specific behavioural strategies that were adopted.

A key goal of the implementation trial was to evaluate effectiveness in not just overall but also within key sectors identified *a priori*.(19) Here, significant improvements in workplace sitting were seen within almost all sectoral subgroups. The sole exception was that changes in teams with call centre staff were substantial -46.3 minutes per 8-hour workday (95%CI: -126.4 to 33.9) but inconclusive with wide confidence intervals as a consequence of only three call centres participating in the evaluation. This finding adds to a known paucity of evidence, as indicated by a recent scoping review(45) that called for further evidence on the effectiveness, acceptability and feasibility of health-promoting interventions (including for addressing sedentary behaviour) conducted in the call-centre setting. Minimal adverse events were recorded and were of mostly a similar nature (musculoskeletal) as have been reported in similar programs.(46, 47)

### ***Satisfaction and Maintenance intentions***

Satisfaction was generally high for both staff and champions and most indicated they would participate again in BeUpstanding. Although further evidence is needed,(48) shifting culture may be the key for sustainable change in workplace sedentary behaviour. The BeUpstanding program heavily targeted culture, and significantly improved many cultural aspects. Following BeUpstanding, most champions intended to continue on with the program, often also intending adaptation -- an important feature of sustainable delivery.(49) Many particularly mentioned intentions to expand to either more staff and/or embedding the program into standard practice, which is consistent with the intent of BeUpstanding. Findings from the 12-

month follow-up champions (to be reported elsewhere), will provide further insights into workplace changes following BeUpstanding.

### ***Strengths and limitations***

Strengths of the study include: the wide-ranging participation from workplaces across Australia and from a wide variety of industries and all priority sectors identified by the policy and practice partners; the data collection across the RE-AIM framework, enabling understanding of how the program is working in practice; and, the data collection on a range of measures, including those relevant to organisations (e.g., job satisfaction). Limitations include the single-group evaluation design, which was necessary for feasibility reasons, and the large workforce turnover, reflective of the broader job mobility occurring during this period.<sup>(33)</sup> To some extent this was mitigated by combining longitudinal and repeated cross-sectional designs, and using statistics that account for repeated observations but do not require them. Staff could be included in the evaluation whether they were part of the team and participating in the evaluation in either one or both evaluations. Another limitation was the measures used are self-report, or proxy-reported by champions, and both responses concerning the organisation, the program, and the sending out of staff surveys all relied on the workplace champions. While necessary for the scalable online program, this does have significant potential for measurement error and bias. Importantly, staff perspectives were collected as well as the champion perspectives, and many staff findings were robust to missing data handling. Single-item or minimal item measures were chosen where possible in order to minimise participant burden in line with our preliminary work informing the trial.<sup>(21)</sup> One possibility for better gauging the impact of this program and others like it on productivity and health is to continue the broad, brief evaluation model but add optional evaluation tools with more detailed assessments. These may have measurement properties more suitable to detecting changes.

### **Conclusion**

Findings from this Australian national implementation trial found that the workplace champion-delivered BeUpstanding program was effective in supporting teams of desk-based workers to sit less and move more and in creating a supportive culture for this change. It also demonstrated that the program can be taken up and implemented by a wide range of users from across Australia and across industries. Learnings from the trial are being used to inform the next iteration of the program (BeUpstanding 2.0), which in turn will inform Phase 5 (sustainability) of the research-to-practice process.<sup>(12)</sup> This optimisation, which will be described in detail elsewhere, is designed to enhance scalability, inclusivity and the user experience.



## 567 **Declarations**

### 568 ***Ethics approval and consent to participate***

569 The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the  
570 Human Research Ethics Committee of The University of Queensland (protocol code #2016001743 on the  
571 9th January 2017). All participants provided informed consent. No individual or personally identifiable  
572 information is reported.

### 573 ***Consent for publication***

574 Not applicable

### 575 ***Availability of data and materials***

576 The datasets generated and analysed during the current study are available from the corresponding author  
577 on reasonable request.

### 578 ***Competing interests***

579 The BeUpstanding program does include consultancy options within the toolkit. All and any proceeds from  
580 consultancy activities return to the research to continue to develop the research program. GNH is on the  
581 Editorial Board of IJB NPA.

### 582 ***Funding***

583 The BeUpstanding program was supported by funding from the Queensland Government “Healthier.  
584 Happier. Workplaces” program, Safe Work Australia, Comcare, and the National Health and Medical  
585 Research Council of Australia through a Partnership Project Grant (#1149936) conducted in partnership  
586 with Comcare, Safe Work Australia, the Queensland Office of Industrial Relations, VicHealth, and  
587 Healthier Workplace WA. GNH was supported by an MRFF-NHMRC Emerging Leadership Investigator  
588 Grant (#1193815). NHMRC had no role in the study in terms of the design, data collection, management,  
589 analysis and/or interpretation. SKM’s salary is supported by the Health and Wellbeing Centre for Research  
590 Innovation (HWCRI), co-funded by The University of Queensland and the Queensland Government  
591 through Health and Wellbeing Queensland.

### 592 ***Authors contributions***

593 GNH, ADG, AA, DWD, EGE, NDG, LG, ADL, MM, NO, TS, LS: conception, funding, recruitment,  
 594 interpretation.  
 595 LS: recruitment  
 596 ADG; SKM, JJ, LU: recruitment; data collection; project management; partner engagement  
 597 EAHW: data management and data analysis  
 598 GNH and EAHW wrote the paper with input from all authors.

# 599 *Acknowledgements*

600 We acknowledge and thank the partner investigators and their associated support staff, academic  
 601 investigators, project staff, and external consultants for their contribution to the BeUpstanding program of  
 602 research. We also acknowledge and thank all the participants who have taken part in the program,  
 603 particularly our workplace champions.

604

**Table 1:** Characteristics of champions, teams and workplaces/organisations participating in the trial

Characteristic	Category / (units or range)	All trial participants		Champions leading teams only	
		n	M(SD) or n(%)		M(SD) or n(%)
<i>Champion characteristics</i>					
Location (State)	New South Wales	118	20 (16.9%)	94	17 (18.1%)
	Australian Capital Territory		5 (4.2%)		5 (5.3%)
	Victoria		20 (16.9%)		14 (14.9%)
	Queensland		56 (47.5%)		43 (45.7%)
	South Australia		8 (6.8%)		7 (7.4%)
	Western Australia		7 (5.9%)		6 (6.4%)
	Tasmania		1 (0.8%)		1 (1.1%)
	Northern Territory		1 (0.8%)		1 (1.1%)
Postcode SES <sup>a</sup>	Australian Percentile (1-100)	118	68.2 (27.5)	94	69.9 (27.0)
	State Percentile (1-100)	118	67.6 (26.7)	94	69.3 (26.3)
Age	(years)	116	42.9 (11.0)	94	43.4 (11.1)
Female	Yes	116	88 (75.9%)	94	73 (77.7%)
Job classification	Employee	116	53 (45.7%)	94	40 (42.6%)
	Middle management	116	45 (38.8%)	94	38 (40.4%)
	Upper management	116	18 (15.5%)	94	16 (17.0%)
Occupational Health and Safety Role (Yes)		116	76 (65.5%)	94	60 (63.8%)
Prior Workplace Health Promotion Training (Yes)		116	66 (56.9%)	94	55 (58.5%)
Prior Workplace Health Promotion Experience (Yes)		116	42 (36.2%)	94	32 (34.0%)
Referred to BeUpstanding by ... (multiples apply)					
	Article / Publication / Newsletter	116	16 (13.8%)	94	13 (13.8%)
	Internet / Website	116	28 (24.1%)	94	20 (21.3%)
	Seminar / Presentation / Conference	116	15 (12.9%)	94	14 (14.9%)
	Social Media / TV / Radio	116	4 (3.4%)	94	3 (3.2%)
	Colleague	116	38 (32.8%)	94	31 (33.0%)
	Other word of mouth	116	13 (11.2%)	94	12 (12.8%)
	General Workplace	116	1 (0.9%)	94	1 (1.1%)
	Health & Safety Organisation	116	2 (1.7%)	94	2 (2.1%)
	Email	116	1 (0.9%)	94	1 (1.1%)
	Other source	116	5 (4.3%)	94	4 (4.3%)
<i>Team characteristics</i>					
Team Size <sup>b</sup>		94	29.4 (32.1)	94	29.4 (32.1)
Team Size Category	1-10	94	30 (31.9%)	94	30 (31.9%)
	11-20	94	27 (28.7%)	94	27 (28.7%)
	>20	94	37 (39.4%)	94	37 (39.4%)
Current participation in other workplace health promotion program (Yes)		116	31 (26.7%)	94	23 (24.5%)
Interest in Health <sup>c</sup>	(1-5)	116	3.4 (0.7)	94	3.4 (0.7)
Motivation to Sit Less <sup>c</sup>	(1-5)	116	3.1 (0.7)	94	3.2 (0.7)
Stress <sup>c</sup>	(1-5)	116	3.3 (0.6)	94	3.2 (0.5)
Regional/remote staff included (Yes) <sup>d</sup>		116	41 (35.3%)	94	32 (34.0%)
Call-centre staff included (Yes) <sup>e</sup>		118	6 (5.1%)	94	6 (6.4%)
Main Location (State) <sup>f</sup>	New South Wales	103	20 (19.4%)	94	17 (18.1%)
	Australian Capital Territory		5 (4.9%)		5 (5.3%)
	Victoria		17 (16.5%)		14 (14.9%)
	Queensland		46 (44.7%)		43 (45.7%)

	South Australia		7 (6.8%)		7 (7.4%)
	Western Australia		6 (5.8%)		6 (6.4%)
	Tasmania		1 (1.0%)		1 (1.1%)
	Northern Territory		1 (1.0%)		1 (1.1%)
<b>Workplace or organisation</b>					
Organisation Size	Small (<20)	118	11 (9.3%)	94	11 (11.7%)
	Medium (20-199)		17 (14.4%)		13 (13.8%)
	Large (200-1999)		34 (28.8%)		24 (25.5%)
	Very large (2000+)		56 (47.5%)		46 (48.9%)
Small-medium enterprise	Yes	118	28 (23.7%)	94	24 (25.5%)
	Public	118	62 (52.5%)	94	48 (51.1%)
	Non-profit		19 (16.1%)		14 (14.9%)
	Private		37 (31.4%)		32 (34.0%)
Blue-collar workplace <sup>g</sup>	Yes	118	40 (33.9%)	94	33 (35.1%)
	COVID-19 impact <sup>h</sup>	118	6 (5.1%)	94	6 (6.4%)
	General		66 (55.9%)		53 (56.4%)
	High		46 (39.0%)		35 (37.2%)
Organisational readiness <sup>i</sup>			3.8 (0.6)		3.8 (0.6)
	Context (1-5)	99	3.8 (0.6)	94	3.8 (0.6)
	Change effort (0-1)	100	0.4 (0.4)	94	0.4 (0.4)
	Change efficacy (1-5)	116	4.1 (0.5)	94	4.1 (0.5)
Main industry of organisation <sup>j</sup>	Administrative and Support Services	118	6 (5.1%)	94	6 (6.4%)
	Agriculture / Forestry and Fishing		4 (3.4%)		2 (2.1%)
	Construction		6 (5.1%)		6 (6.4%)
	Education and Training		6 (5.1%)		6 (6.4%)
	Electricity / Gas / Water and Waste Services		8 (6.8%)		7 (7.4%)
	Financial and Insurance Services		4 (3.4%)		3 (3.2%)
	Health Care and Social Assistance		29 (24.6%)		20 (21.3%)
	Manufacturing		9 (7.6%)		8 (8.5%)
	Other Services		8 (6.8%)		8 (8.5%)
	Professional / Scientific and Technical Services		22 (18.6%)		16 (17.0%)
	Public Administration and Safety		7 (5.9%)		6 (6.4%)
	Retail Trade		2 (1.7%)		2 (2.1%)
	Transport / Postal and Warehousing		7 (5.9%)		4 (4.3%)

Characteristics of organisations, workplaces, and teams were collected from as many of the champions as possible and linked to each team mostly by using data from the team's primary champion, or if unavailable, from another co-champion for the team, an overseer with knowledge of all the teams, or any other suitable respondent from that workplace.

<sup>a</sup> SES = Socioeconomic Status (postal area level Index of Relative Economic Advantage and Disadvantage in national and state percentiles).(50)

<sup>b</sup> estimated by champions, and periodically updated by project staff during implementation checks as champions evolved their concept of who their team would be

<sup>c</sup> champion perception when signing up of team's current level (1=worst response to 5=best response)

<sup>d</sup> some or all of the team were in regional or remote work locations as reported by champions or ascertained from staff or champion workplace postcodes

<sup>e</sup> some or all of the team were call-centre staff according to the champion

<sup>f</sup> the primary state in which the team is located, calculated from champion and staff-reported workplace postcodes

<sup>g</sup> classified from the reported industry, or by champion report of a blue-collar workplace where the industry was mixed

<sup>h</sup> Those participating prior to COVID-19 were treated as no impact. General impact (e.g., working from home) was assigned where there was no evidence of a high impact (decided subjectively). Severe impacts (e.g., job losses, shutdown of core business) reported by the champion, an impact strong enough to prevent participation, being located in Victoria where extended lockdowns occurred, or being part of sectors hard hit delivering healthcare or COVID-19 response schemes all qualified as high impact.

623 <sup>i</sup> Higher scores are beneficial(51)  
624 <sup>j</sup> main industry of the organisation according to the 19 standard ANZIC categories(35)  
625

626 **Table 2:** Characteristics of participating staff (n=2228)

Characteristic	(Unit or range) / category	n	Mean (SD) or n(%) <sup>a</sup>
Age	(years)	2004	42.9 (11.4)
Work hours	(hours per week)	2043	38.2 (9.0)
Pre-program exercise <sup>b</sup>	(days per week)	1768	3.1 (2.2)
Meeting recommendations <sup>b</sup>	No <5 days per week	1768	1286 (72.7%)
	Yes, ≥5 days per week		482 (27.3%)
Post-program exercise <sup>b</sup>	(days per week)	709	3.5 (2.1)
Meeting recommendations <sup>b</sup>	No, <5 days per week	709	469 (66.1%)
	Yes, ≥5 days per week		240 (33.9%)
Body Mass Index (BMI) <sup>c</sup>	<25 kg/m <sup>2</sup>	1705	771 (45.2%)
	25-<30 kg/m <sup>2</sup>		561 (32.9%)
	≥30 kg/m <sup>2</sup>		373 (21.9%)
Female	No	2131	745 (35.0%)
	Yes		1386 (65.0%)
Non-English speaking background	No	2015	1674 (83.1%)
	Yes		341 (16.9%)
Full-time employment	No	2043	365 (17.9%)
	Yes		1678 (82.1%)
Education	High school or less	2024	191 (9.4%)
	TAFE/Trade certificate/ Diploma		530 (26.2%)
	University or other tertiary		1303 (64.4%)
Job classification	Employee	2043	1399 (68.5%)
	Middle management		167 (8.2%)
	Upper management		477 (23.3%)
Job category skill level <sup>d</sup>	1 (most skill / training)	2043	1028 (50.3%)
	2		266 (13.0%)
	3		505 (24.7%)
	4		56 (2.7%)
	5 (least skill / training)		188 (9.2%)
Shift worker	No	2043	2001 (97.9%)
	Yes		42 (2.1%)

627 <sup>a</sup> Mean SD corrected for clustering using linearized variance (STATA survey commands)

628 <sup>b</sup> number of days per week exercised for at least 30 minutes (0-7)(52) and meeting national guidelines (≥5 v <5 days)

629 <sup>c</sup> Excludes invalid responses (truncated to plausible range of 15 to 50)(53)

630 <sup>d</sup> Australian Standard Classification of Occupations (ASCO: 1 most to 5 least skilled)(54)

631

632

**Table 3:** Completion of core steps for all participating teams (n=94) and completing teams (n=89) by one or more champion(s) for that team

Core program steps	All teams n (%)	Completing teams n (%)
- Pre-program staff survey	94 (100.0%)	89 (100.0%)
- Workplace audit <sup>a</sup>	71 (75.5%)	71 (79.8%)
- Staff information and consultation session (e.g., workshop)	94 (100.0%)	89 (100.0%)
- Strategy selection	73 (77.7%)	78 (87.6%)
- Program promotion via posters and/or emails	94 (100.0%)	89 (100.0%)
- Post-program staff survey	94 (100.0%)	89 (100.0%)
- Program completion survey	80 (85.1%)	79 (88.8%)
Number of core steps completed (out of 7)		
7	68 (72.3%)	68 (76.4%)
6	17 (18.1%)	16 (18.0%)
5	9 (9.5%)	5 (5.6%)
<5	0 (0.0%)	0 (0.0%)

<sup>a</sup> The workplace audit mostly asked about the workplace in general and was considered completed if it was completed by any champion for the workplace

**Table 4:** Post-program costs reported by the champion (n=72 champions)

Total costs reported		Costs per team member	
Cost	n (%)	Cost	n (%)
\$0	52 (72.2%)	\$0	52 (72.2%)
<\$100	7 (9.7%)	>\$0-\$5	7 (9.7%)
\$100-\$499	1 (1.4%)	>\$5- \$25	4 (5.6%)
\$500-\$999	1 (1.4%)	>\$25-\$50	2 (2.8%)
\$1000-\$1499	4 (5.6%)	>\$50-\$100	2 (2.8%)
\$1500-\$1999	1 (1.4%)	\$133	1 (1.4%)
\$2000-\$2499	3 (4.2%)	\$250	1 (1.4%)
\$7000	1 (1.4%)	\$318	1 (1.4%)
\$10,500	1 (1.4%)	\$1500	1 (1.4%)

**Table 5:** Changes in staff-reported main outcomes pre to post BeUpstanding program (k=82 workplaces) <sup>a</sup>

<b>Outcome</b>	<b>Pre-program M (SD)</b>	<b>n</b>	<b>Unadjusted Change (95% CI)</b>	<b>p</b>	<b>n</b>	<b>Adjusted <sup>b</sup> Change (95% CI)</b>	<b>p</b>	<b>n</b>	<b>Adjusted <sup>b</sup> &amp; multiply imputed <sup>c</sup> Change (95% CI)</b>	<b>p</b>
<b><i>Outcomes targeted by the intervention</i></b>										
Activity as % of workday <sup>g</sup>										
Sitting <sup>d,e</sup>	78.3 (17.9)	2131	-7.6 (-9.1, -6.1)	<0.001	2043	-8.1 (-9.7, -6.5)	<0.001	2248	-7.7 (-9.6, -5.8)	<0.001
Standing <sup>d,f</sup>	10.7 (11.1)	2131	5.0 (3.3, 6.7)	<0.001	2043	5.9 (3.8, 7.9)	<0.001	2248	6.3 (4.3, 8.3)	<0.001
Moving <sup>d,f</sup>	11.0 (10.5)	2131	0.8 (-0.2, 1.8)	0.109	2043	1.3 (0.1, 2.4)	0.032	2248	3.1 (2.1, 4.1)	<0.001
% of sitting in prolonged bouts <sup>d,h</sup>	65.8 (24.5)	2130	-9.1 (-11.0, -7.1)	<0.001	2042	-9.3 (-11.2, -7.3)	<0.001	2248	-9.0 (-11.2, -6.7)	<0.001
Activity preference alignment <sup>i</sup>										
Sitting	153.1 (84.0)	2091	-34.3 (-40.7, -27.8)	<0.001	2043	-33.7 (-40.2, -27.1)	<0.001	2248	-30.7 (-37.6, -23.9)	<0.001
Standing	90.3 (62.6)	2091	-17.6 (-22.5, -12.6)	<0.001	2043	-17.3 (-22.2, -12.3)	<0.001	2248	-14.8 (-21.8, -7.8)	<0.001
Moving	70.8 (60.6)	2091	-10.7 (-15.4, -6.0)	<0.001	2043	-10.3 (-15.1, -5.6)	<0.001	2248	-6.9 (-12.7, -1.0)	0.023
Perceptions of culture (0-4) <sup>j</sup>										
Overall culture score	2.6 (0.8)	2092	0.37 (0.31, 0.43)	<0.001	2043	0.37 (0.30, 0.43)	<0.001	2248	0.34 (0.28, 0.40)	<0.001
Perceived control over sitting <sup>e</sup>	2.6 (1.1)	2092	0.23 (0.16, 0.30)	<0.001	2043	0.23 (0.15, 0.30)	<0.001	2248	0.06 (-0.02, 0.13)	0.136
Organisation supports choices <sup>e</sup>	2.7 (1.0)	2092	0.36 (0.30, 0.43)	<0.001	2043	0.36 (0.30, 0.43)	<0.001	2248	0.15 (0.09, 0.21)	<0.001
Organisation supports moving	2.9 (0.9)	2092	0.28 (0.21, 0.35)	<0.001	2043	0.27 (0.20, 0.34)	<0.001	2248	0.05 (-0.02, 0.13)	0.159
Role modelling	2.1 (1.0)	2092	0.47 (0.38, 0.55)	<0.001	2043	0.46 (0.38, 0.55)	<0.001	2248	0.33 (0.25, 0.42)	<0.001
Supportive culture	2.5 (1.0)	2092	0.47 (0.39, 0.55)	<0.001	2043	0.47 (0.38, 0.55)	<0.001	2248	0.28 (0.20, 0.36)	<0.001
<b><i>Other outcomes</i></b>										
Feeling part of a team (0-4) <sup>k</sup>	2.6 (1.0)	2043	0.02 (-0.06, 0.10)	0.633	2043	0.01 (-0.07, 0.09)	0.869	2248	-0.05 (-0.13, 0.04)	0.296
Feeling engaged (0-4) <sup>k</sup>	2.5 (0.9)	2043	0.05 (-0.02, 0.12)	0.173	2043	0.03 (-0.04, 0.11)	0.368	2248	-0.04 (-0.13, 0.05)	0.383
<i>Productivity and Health</i>		2043	0.10 (0.02, 0.17)	0.012						
Self-rated job performance (1-7) <sup>e,l</sup>	5.5 (1.0)	2043	0.09 (0.01, 0.18)	0.027	2043	0.09 (0.01, 0.16)	0.019	2248	-0.07 (-0.16, 0.03)	0.148
Job satisfaction (1-7) <sup>e,m</sup>	5.1 (1.2)	2043	0.00 (0.00, 0.00)	0.923	2043	0.09 (0.00, 0.17)	0.047	2248	-0.01 (-0.10, 0.07)	0.733
Sick Days past 28 days (0-28) <sup>f,n</sup>	0.6 (1.9)	2043	0.03 (-0.06, 0.12)	0.573	2043	0.00 (0.00, 0.00)	0.923	2248	-0.14 (-0.51, 0.23)	0.441
Stress (0-4) <sup>o</sup>	1.7 (1.1)	2043	0.13 (0.08, 0.18)	<0.001	2043	0.03 (-0.06, 0.12)	0.507	2248	0.06 (-0.02, 0.14)	0.169
Energy score (0-4) <sup>o</sup>	2.1 (0.7)	2043	0.02 (-0.05, 0.10)	0.558	2043	0.12 (0.07, 0.18)	<0.001	2248	0.10 (0.06, 0.14)	<0.001
Creativity (0-4) <sup>o</sup>	1.6 (1.0)	2043	0.02 (-0.06, 0.10)	0.633	2043	0.01 (-0.07, 0.08)	0.864	2248	0.05 (-0.01, 0.12)	0.093
Self-rated health (0-100) <sup>p</sup>										



Psychological	2.2 (1.0)	2043	0.07 (0.00, 0.14)	0.038	2043	0.07 (0.00, 0.14)	0.054	2248	0.04 (-0.03, 0.11)	0.222
Physical	2.0 (1.0)	2043	0.12 (0.05, 0.18)	<0.001	2043	0.12 (0.05, 0.18)	<0.001	2248	0.07 (0.00, 0.14)	0.044
Musculoskeletal discomfort (0-100) <sup>a</sup>										
Upper body	4.7 (3.0)	2043	-0.55 (-0.78, -0.33)	<0.001	2043	-0.54 (-0.76, -0.31)	<0.001	2248	-0.35 (-0.58, -0.13)	0.003
Lower back	3.9 (3.1)	2043	-0.60 (-0.84, -0.37)	<0.001	2043	-0.58 (-0.82, -0.34)	<0.001	2248	-0.37 (-0.68, -0.06)	0.020
Lower body	3.8 (3.1)	2043	-0.37 (-0.60, -0.13)	0.002	2043	-0.36 (-0.59, -0.12)	0.003	2248	-0.12 (-0.37, 0.13)	0.323

<sup>a</sup> Estimated from linear mixed models, with random intercepts for workplace, and staff (nested within workplace). Reports M and SD with linearized variance to correct for workplace clustering, and contrasts of marginal means. Expressions used for transformed outcomes.

<sup>b</sup> Adjusts for variables that may be imbalanced across pre- and post-program respondents: p<0.2 predictors of team participation in post-program surveys (Additional File 6); p<0.2 predictors of staff participation in post-program only versus pre-program only surveys (Additional File 7) using only one variable for concepts measured in multiple ways (organisational readiness, location, and postcode SES). Models adjusted for: small-medium enterprise (yes/no); regional staff (yes/no); inclusion of call-centre staff (yes/no); Champion Occupational Health & Safety role (yes/no); COVID-19 impact level (none/general/high); public sector (yes/no); organisational readiness (context score); Champion age (years); full-time employment (yes/no); Job category skill (1-5); team health interest level (0-4); Postcode SES.

<sup>c</sup> Missing data multiply imputed by chained equations (STATA) with m=20 imputations.

<sup>d</sup> Model also includes random slopes for workplace, as likelihood ratio test supported random slopes over random intercept models at p<0.05 (Additional File 8)

<sup>e</sup> Inverse log transformed as  $\ln(1 + \text{maximum value} - \text{variable})$ .

<sup>f</sup> log transformed as  $\ln(\text{variable})$  for positive or  $\ln(\text{variable} + 0.001)$  for non-negative variables

<sup>g</sup> Occupational Sitting and Physical Activity Questionnaire.(34)

<sup>h</sup> percentage of their sitting that staff estimate is accrued by sitting for 30+ minutes continuously: 0-100;(55)

<sup>i</sup> absolute difference between desired and self-reported percentage (0= perfect alignment; 100=complete misalignment)(22)

<sup>j</sup> 0-4;(19) higher=better.

<sup>k</sup> single item (0-4; higher=better)

<sup>l</sup> single item;(56) 1-7; higher=better;

<sup>m</sup> single item;(57) 1-7; higher=better;


<sup>n</sup> sick days in last four weeks (0-28 days; lower=better);(58)

<sup>o</sup> 0-4 with higher=better; note: energy is mean of 3 items


<sup>p</sup> self-rated physical and psychological health (0=poor; 4=excellent); higher=better.(59, 60)

<sup>q</sup> musculoskeletal discomfort experienced over the last week (0=no discomfort; 10=severe discomfort);(61, 62)

# 1 Figure 1: Participant flow diagram




3 **Figure 2: Change in staff-reported outcomes from pre to post BeUpstanding intervention**



4

5 **Figure 3: Staff perceptions of the impact of the BeUpstanding program (n=654)**



6

## 7 References

1. Saunders TJ, McIsaac T, Douillette K, Gaulton N, Hunter S, Rhodes RE, et al. Sedentary behaviour and health in adults: an overview of systematic reviews. *Appl Physiol Nutr Metab*. 2020;45(10 (Suppl. 2)):S197-S217.
2. Ekelund U, Tarp J, Steene-Johannessen J, Hansen BH, Jefferis B, Fagerland MW, et al. Dose-response associations between accelerometry measured physical activity and sedentary time and all cause mortality: systematic review and harmonised meta-analysis. *BMJ*. 2019;366:l4570.
3. Dempsey PC, Biddle SJH, Buman MP, Chastin S, Ekelund U, Friedenreich CM, et al. New global guidelines on sedentary behaviour and health for adults: broadening the behavioural targets. *Int J Behav Nutr Phys Act*. 2020;17(1):151.
4. Healy GN, Lawler SP, Thorp A, Neuhaus M, Robson EL, Owen N, et al. Reducing prolonged sitting in the workplace (An evidence review: full report). Available at: [http://www.vichealth.vic.gov.au/Publications/Economic-participation/Creating\\_Healthy\\_Workplaces.aspx](http://www.vichealth.vic.gov.au/Publications/Economic-participation/Creating_Healthy_Workplaces.aspx). Melbourne, Australia: Victorian Health Promotion Foundation; 2012.
5. Chu AH, Ng SH, Tan CS, Win AM, Koh D, Muller-Riemenschneider F. A systematic review and meta-analysis of workplace intervention strategies to reduce sedentary time in white-collar workers. *Obes Rev*. 2016.
6. MacEwen BT, MacDonald DJ, Burr JF. A systematic review of standing and treadmill desks in the workplace. *Prev Med*. 2015;70:50-8.
7. Neuhaus M, Eakin EG, Straker L, Owen N, Dunstan DW, Reid N, et al. Reducing occupational sedentary time: a systematic review and meta-analysis of evidence on activity-permissive workstations. *Obes Rev*. 2014;15(10):822-38.
8. Tew GA, Posso MC, Arundel CE, McDaid CM. Systematic review: height-adjustable workstations to reduce sedentary behaviour in office-based workers. *Occup Med (Lond)*. 2015;65(5):357-66.

- 32 9. Shrestha N, Kukkonen-Harjula KT, Verbeek JH, Ijaz S, Hermans V, Bhaumik S. Workplace  
33 interventions for reducing sitting at work. *Cochrane Database Syst Rev.* 2016;3:CD010912.
- 34 10. Nguyen P, Ananthapavan J, Tan EJ, Crosland P, Bowe SJ, Gao L, et al. Modelling the potential  
35 health and economic benefits of reducing population sitting time in Australia. *Int J Behav Nutr Phys*  
36 *Act.* 2022;19(1):28.
- 37 11. Straker L, Coenen P, Dunstan D, Gilson N, Healy GN. Sedentary Work – Evidence on an Emergent  
38 Work Health and Safety Issue – Final Report. Canberra: Safe Work Australia; 2016.
- 39 12. Healy GN, Goode A, Schultz D, Lee D, Leahy B, Dunstan DW, et al. The BeUpstanding Program:  
40 Scaling up the Stand Up Australia Workplace Intervention for Translation into Practice. *AIMS*  
41 *Public Health.* 2016;3(2):341-7.
- 42 13. The University of Queensland. BeUpstanding [Available from: [www.beupstanding.com.au](http://www.beupstanding.com.au).
- 43 14. Healy GN, Eakin EG, Owen N, Lamontagne AD, Moodie M, Winkler EA, et al. A Cluster  
44 Randomized Controlled Trial to Reduce Office Workers' Sitting Time: Effect on Activity Outcomes.  
45 *Med Sci Sports Exerc.* 2016;48(9):1787-97.
- 46 15. LaMontagne AD, Noblet AJ, Landsbergis PA. Intervention development and implementation:  
47 Understanding and addressing barriers to organisational-level interventions. In: Biron C, Karanika-  
48 Murray M, Cooper CL, editors. *Improving organisational interventions for stress and well-being:*  
49 *Routledge;* 2012.
- 50 16. Bandura A. Health promotion by social cognitive means. *Health Educ Behav.* 2004;31(2):143-64.
- 51 17. McLeroy KR, Bibeau D, Steckler A, Glanz K. An ecological perspective on health promotion  
52 programs. *Health Educ Q.* 1988;15(4):351-77.
- 53 18. Owen N, Sugiyama T, Eakin EE, Gardiner PA, Tremblay MS, Sallis JF. Adults' sedentary behavior  
54 determinants and interventions. *Am J Prev Med.* 2011;41(2):189-96.
- 55 19. Healy GN, Goode AD, Abbott A, Burzic J, Clark BK, Dunstan DW, et al. Supporting Workers to Sit  
56 Less and Move More Through the Web-Based BeUpstanding Program: Protocol for a Single-Arm,  
57 Repeated Measures Implementation Study. *JMIR Res Protoc.* 2020;9(5):e15756.

20. Healy GN, Winkler EAH, Goode AD. A RE-AIM evaluation in early adopters to iteratively improve the online BeUpstanding program supporting workers to sit less and move more. BMC Public Health. 2021;21(1):1916.
21. Goode AD, Frith M, Hyne SA, Burzic J, Healy GN. Applying a User Centred Design Approach to Optimise a Workplace Initiative for Wide-Scale Implementation. Int J Environ Res Public Health. 2022;19(13).
22. Healy GN, Eakin EG, Winkler EA, Hadgraft N, Dunstan DW, Gilson ND, et al. Assessing the Feasibility and Pre-Post Impact Evaluation of the Beta (Test) Version of the BeUpstanding Champion Toolkit in Reducing Workplace Sitting: Pilot Study. JMIR Form Res. 2018;2(2):e17.
23. Goode AD, Hadgraft NT, Neuhaus M, Healy GN. Perceptions of an online 'train-the-champion' approach to increase workplace movement. Health Promot Int. 2019;34(6):1179-90.
24. Glasgow RE, Harden SM, Gaglio B, Rabin B, Smith ML, Porter GC, et al. RE-AIM Planning and Evaluation Framework: Adapting to New Science and Practice With a 20-Year Review. Front Public Health. 2019;7:64.
25. Glasgow RE, Battaglia C, McCreight M, Ayele RA, Rabin BA. Making Implementation Science More Rapid: Use of the RE-AIM Framework for Mid-Course Adaptations Across Five Health Services Research Projects in the Veterans Health Administration. Front Public Health. 2020;8:194.
26. Australian Institute of Health and Welfare. Australian bushfires 2019–20: Exploring the short-term health impacts. . Canberra; 2020. Contract No.: Cat. no. PHE 276.
27. Stobart A, Duckett S. Australia's Response to COVID-19. Health Econ Policy Law. 2022;17(1):95-106.
28. Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D, et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. Bmj. 2014;348:g1687.
29. Des Jarlais DC, Lyles C, Crepaz N. Improving the reporting quality of nonrandomized evaluations of behavioral and public health interventions: the TREND statement. Am J Public Health. 2004;94(3):361-6.



- 85 30. Buckley JP, Hedge A, Yates T, Copeland RJ, Loosemore M, Hamer M, et al. The sedentary office:  
86 an expert statement on the growing case for change towards better health and productivity. Br J  
87 Sports Med. 2015;49(21):1357-62.
- 88 31. Human Factors and Ergonomics Society of Australia (HFESA) Inc. Sedentary Behaviour: HFESA  
89 Position on Prolonged Unbroken Sitting Time 2015 [Available from:  
90 <https://www.ergonomics.org.au/documents/item/184>. Accessed: 2017-11-06. (Archived by  
91 WebCite® at <http://www.webcitation.org/6ulUGJfFW>).
- 92 32. Colberg SR, Sigal RJ, Yardley JE, Riddell MC, Dunstan DW, Dempsey PC, et al. Physical  
93 Activity/Exercise and Diabetes: A Position Statement of the American Diabetes Association.  
94 Diabetes Care. 2016;39(11):2065-79.
- 95 33. Australian Institute of Health and Welfare. Changing patterns of work. 2023 7th September.
- 96 34. Chau JY, Van Der Ploeg HP, Dunn S, Kurko J, Bauman AE. Validity of the occupational sitting and  
97 physical activity questionnaire. Med Sci Sports Exerc. 2012;44(1):118-25.
- 98 35. Australian Bureau of Statistics. Australian and New Zealand Standard Industrial Classification  
99 (ANZSIC) Canberra: ABS; 2006 [Available from:  
100 <https://www.abs.gov.au/statistics/classifications/australian-and-new-zealand-standard-industrial->  
101 [classification-anzsic/latest-release](https://www.abs.gov.au/statistics/classifications/australian-and-new-zealand-standard-industrial-classification-anzsic/latest-release).
- 102 36. Thorp AA, Healy GN, Winkler E, Clark BK, Gardiner PA, Owen N, et al. Prolonged sedentary time  
103 and physical activity in workplace and non-work contexts: a cross-sectional study of office, customer  
104 service and call centre employees. Int J Behav Nutr Phys Act. 2012;9:128.
- 105 37. Morris A, Murphy R, Shepherd S, Graves L. Multi-Stakeholder Perspectives of Factors That  
106 Influence Contact Centre Call Agents' Workplace Physical Activity and Sedentary Behaviour. Int J  
107 Environ Res Public Health. 2018;15(7).
- 108 38. Morris AS, Murphy RC, Shepherd SO, Healy GN, Edwardson CL, Graves LEF. A multi-component  
109 intervention to sit less and move more in a contact centre setting: a feasibility study. BMC Public  
110 Health. 2019;19(1):292.



111 39. Wang C, Lu EY, Sun W, Chang JR, Tsang HWH. Effectiveness of interventions on sedentary  
112 behaviors in office workers: a systematic review and meta-analysis. *Public Health*. 2024;230:45-51.

113 40. Healy GN, Eakin EG, Lamontagne AD, Owen N, Winkler EA, Wiesner G, et al. Reducing sitting  
114 time in office workers: short-term efficacy of a multicomponent intervention. *Prev Med*.  
115 2013;57(1):43-8.

116 41. Neuhaus M, Healy GN, Dunstan DW, Owen N, Eakin EG. Workplace sitting and height-adjustable  
117 workstations: a randomized controlled trial. *Am J Prev Med*. 2014;46(1):30-40.

118 42. Wilms P, Schroder J, Reer R, Scheit L. The Impact of "Home Office" Work on Physical Activity and  
119 Sedentary Behavior during the COVID-19 Pandemic: A Systematic Review. *Int J Environ Res*  
120 *Public Health*. 2022;19(19).

121 43. Sui W, Smith ST, Fagan MJ, Rollo S, Prapavessis H. The effects of sedentary behaviour  
122 interventions on work-related productivity and performance outcomes in real and simulated office  
123 work: A systematic review. *Appl Ergon*. 2019;75:27-73.

124 44. Oakman J, Clune S, Stuckey R. Work-related musculoskeletal disorders in Australia. Canberra;  
125 2019.

126 45. Bell Z, Porcellato L, Holland P, Morris A, Smith C, Haines C, et al. A systematic scoping review of  
127 health-promoting interventions for contact centre employees examined through a behaviour change  
128 wheel lens. *PLoS One*. 2024;19(3):e0298150.

129 46. Edwardson CL, Biddle SJH, Clarke-Cornwell A, Clemes S, Davies MJ, Dunstan DW, et al. A three  
130 arm cluster randomised controlled trial to test the effectiveness and cost-effectiveness of the SMART  
131 Work & Life intervention for reducing daily sitting time in office workers: study protocol. *BMC*  
132 *Public Health*. 2018;18(1):1120.

133 47. Shrestha N, Kukkonen-Harjula KT, Verbeek JH, Ijaz S, Hermans V, Pedisic Z. Workplace  
134 interventions for reducing sitting at work. *Cochrane Database Syst Rev*. 2018;12(12):CD010912.

135 48. Taylor WC, Suminski RR, Das BM, Paxton RJ, Craig DW. Organizational Culture and Implications  
136 for Workplace Interventions to Reduce Sitting Time Among Office-Based Workers: A Systematic  
137 Review. *Front Public Health*. 2018;6:263.

138 49. Schell SF, Luke DA, Schooley MW, Elliott MB, Herbers SH, Mueller NB, et al. Public health  
139 program capacity for sustainability: a new framework. *Implementation Science*. 2013;8(1):15.

140 50. Australian Bureau of Statistics. Socio-Economic Indexes for Areas (SEIFA) Australia: ABS; 2021  
141 [Available from: [https://www.abs.gov.au/statistics/people/people-and-communities/socio-economic-](https://www.abs.gov.au/statistics/people/people-and-communities/socio-economic-indexes-areas-seifa-australia/latest-release#cite-window1)  
142 [indexes-areas-seifa-australia/latest-release#cite-window1](https://www.abs.gov.au/statistics/people/people-and-communities/socio-economic-indexes-areas-seifa-australia/latest-release#cite-window1).

143 51. Hannon PA, Helfrich CD, Chan KG, Allen CL, Hammerback K, Kohn MJ, et al. Development and  
144 Pilot Test of the Workplace Readiness Questionnaire, a Theory-Based Instrument to Measure Small  
145 Workplaces' Readiness to Implement Wellness Programs. *Am J Health Promot*. 2017;31(1):67-75.

146 52. Milton K, Bull FC, Bauman A. Reliability and validity testing of a single-item physical activity  
147 measure. *Br J Sports Med*. 2011;45(3):203-8.

148 53. Miller Y. Recommendations for the truncation of body mass index in population data. . Liverpool,  
149 NSW; 2003. Report No.: CPAH03-0005.

150 54. Australian Bureau of Statistics. ANZSCO - Australian and New Zealand Standard Classification of  
151 Occupations.: ABS; 2022 [Available from: [https://www.abs.gov.au/statistics/classifications/anzsco-](https://www.abs.gov.au/statistics/classifications/anzsco-australian-and-new-zealand-standard-classification-occupations/latest-release)  
152 [australian-and-new-zealand-standard-classification-occupations/latest-release](https://www.abs.gov.au/statistics/classifications/anzsco-australian-and-new-zealand-standard-classification-occupations/latest-release).

153 55. Clark BK, Stephens SK, Goode AD, Healy GN, Winkler EAH. Alternatives for Measuring Sitting  
154 Accumulation in Workplace Surveys. *J Occup Environ Med*. 2021;63(12):e853-e60.

155 56. Bond FW, Bunce D. Job control mediates change in a work reorganization intervention for stress  
156 reduction. *Journal of occupational health psychology*. 2001;6(4):290-302.

157 57. Dolbier CL, Webster JA, McCalister KT, Mallon MW, Steinhardt MA. Reliability and validity of a  
158 single-item measure of job satisfaction. *Am J Health Promot*. 2005;19(3):194-8.

159 58. Kessler RC, Barber C, Beck A, Berglund P, Cleary PD, McKenas D, et al. The World Health  
160 Organization Health and Work Performance Questionnaire (HPQ). *J Occup Environ Med*.  
161 2003;45(2):156-74.

162 59. Jurges H, Avendano M, Mackenbach JP. Are different measures of self-rated health comparable? An  
163 assessment in five European countries. *Eur J Epidemiol*. 2008;23(12):773-81.

60. Ahmad F, Jhajj AK, Stewart DE, Burghardt M, Bierman AS. Single item measures of self-rated mental health: a scoping review. BMC Health Serv Res. 2014;14:398.
61. Dickinson CE, Campion K, Foster AF, Newman SJ, O'Rourke AM, Thomas PG. Questionnaire development: an examination of the Nordic Musculoskeletal questionnaire. Applied ergonomics. 1992;23(3):197-201.
62. Brakenridge CL, Chong YY, Winkler EAH, Hadgraft NT, Fjeldsoe BS, Johnston V, et al. Evaluating Short-Term Musculoskeletal Pain Changes in Desk-Based Workers Receiving a Workplace Sitting-Reduction Intervention. Int J Environ Res Public Health. 2018;15(9).

