

1 Infant- and child-level predictors of mortality in low-resource settings: the WHO Child
2 Mortality Risk Stratification Multi-Country Pooled Cohort

3

4 **The WHO Risk Stratification Working Group (WHO-RSWG)**

5 WHO-RSWG authors, affiliations and roles are listed at the end of this manuscript.

6

7

8 **Corresponding author:** Catherine Schwinger c.schwinger@uib.no

9

10

11 **Keywords:** mortality, low- and middle income, prioritisation, preterm birth, low birth weight,
12 breastfeeding, guidelines, anthropometry, community, hospital, wasting.

13

14 **Funding**

15 WHO gratefully acknowledges funding support from USAID to establish the pooled database and
16 conduct the main analyses.

17

18 **Disclaimer:**

19 The authors alone are responsible for the views expressed in this paper and they do not necessarily
20 represent the views, decisions, or policies of the institutions with which they are affiliated.

21 **SUMMARY (235 words)**

22 **Background**

23 Despite impressive reductions in overall global child mortality, rates of decline have slowed during
24 the last decade. Current guidelines for the care of at-risk children in low-resource settings mostly
25 focus on broad clinical syndromes or undernutrition rather than children's individual contextualized
26 risk. We aimed to identify readily assessable child-level characteristics that can predict child mortality
27 risk in a range of community and healthcare settings in high-burden settings.

28

29 **Methods**

30 We analysed pooled data from 33 cohorts including 75,287 children under five years of age living in
31 low resource settings to estimate the absolute risks of death associated with risk exposures
32 separately and combined with anthropometry. Children were grouped according to population types
33 studied: general population (GP), selected on anthropometric criteria (A-S), and selected on the
34 presence of illness (I-S).

35

36 **Findings**

37 During a total of 69,085 child-years of follow-up in the pooled dataset, 2,805 (3.7%) children died.
38 Age <24 months, low anthropometry, preterm birth, low birthweight, and absence of breastfeeding
39 were each associated with increased mortality: risks were additive and declined with increasing age.
40 However, overall mortality and the association between child-level characteristics and mortality
41 differed according to the type of study population and child age.

42

43 **Interpretation**

44 Risk assessments combining individual child-level characteristics including anthropometry can enable
45 programmes to identify children at high and lower risk of mortality and, thereafter, differentiate care
46 accordingly. Such a strategy may reduce mortality and optimise health system efficiency and
47 effectiveness.

48

49 **INTRODUCTION**

50 Global rates of under-5 child mortality (U5M) have declined from 93 to 38 deaths per 1,000 live
51 births between 1990 and 2021.¹ However, since 2010, U5M reductions have slowed; 53 countries will
52 not meet the 2030 UN Sustainable Development Goals (SDG) U5M target.² More than 80% of global
53 U5M occurs in sub-Saharan Africa and South Asia, with significant sub-national heterogeneity.^{3,4}
54 Mortality also varies markedly by age, with a third of U5M in some settings occurring within the first
55 three days of life⁵ and almost half occurring in the first month.²

56

57 Low birth weight (LBW), including from preterm birth (PTB), is associated with half of neonatal
58 deaths and confers life-long health risks.² Additionally, undernutrition in all its forms contributes to
59 45% of U5M.⁶ Chronic disease and disability, maternal illiteracy, poor housing conditions, and limited
60 access to healthcare are less often assessed, but may all directly or indirectly be associated with child
61 mortality.⁷ Focusing on interventions that target underlying pathways associated with these risks may
62 offer opportunity to reduce mortality. Importantly, however, while a focus on higher-risk children
63 may avert some mortality, identifying lower risk children may provide opportunities for de-escalation
64 of care and to optimize available resources while improving care for those at highest risk. While
65 prognostic scores have been devised among hospitalised children to identify those at high risk, they
66 are rarely used in practice in LMICs.^{8,9}

67

68 We estimated absolute and relative mortality risks associated with separate and combined
69 assessable, child-level characteristics among children aged 0-59 months through an individual data
70 pooled analysis of the WHO Child Mortality Risk Stratification Multi-Country Pooled Cohort (WHO
71 CMRS).

72 **METHODS**

73 Study design

74 We pooled data from individual children enrolled in observational or randomized controlled trials in
75 LMICs (Table S1). The criteria for inclusion of a dataset were 1) documentation of age, weight, vital
76 status, date of death, and 2) at least two observations per participant under the age of 60 months.
77 Details of the study selection procedure are reported fully in the cohort profile.¹⁰

78

79 Data management

80 Thirty-three datasets were included and pooled into a single database with consistent variable
81 names and structures. Data cleaning included checking for missing data, duplicates, and outliers¹⁰.
82 Anthropometric measurements were defined as outliers if mid-upper arm circumference (MUAC)
83 >300mm or <55mm, weight-for-age (WAZ) >10 or <-10, length <35cm, length- or height-for-age
84 (HAZ) >10 or <-12, weight-for-length or -height (WHZ) >20 or <-10 and values set to missing. As we
85 used anthropometric measurements as categorical variables in analyses, and an extreme value
86 would likely be in the severe deficit category (Z-score <-3), we did not apply more strict cleaning
87 criteria.

88

89 Indicator assessment and definition

90 In all analyses, the outcome was defined as death within the respective observation period. If the
91 vital status of a child was recorded but age at death was missing, we estimated the age at death by
92 adding half of the median observation period of the study to the age at the last study contact point
93 for that child.

94

95 Child-level characteristics were chosen as risk exposure variables for analysis based on availability
96 across datasets. Weight and height measurements were converted to Z-scores using WHO Child
97 Growth Standards.¹¹ We defined a moderate anthropometric deficit as Z-score in WAZ, HAZ, or WHZ
98 <-2 and \geq -3, and a severe deficit as Z-score <-3. MUAC was included as an absolute measure (mm).
99 The WHO guidelines recommend the cut-off for moderate and severe deficits for MUAC at 125mm
100 and 115mm for children aged 6-59 months.¹² For infants <6 months of age, we also applied cut-offs
101 of 120mm and 110mm for moderate and severe deficits.¹²

102

103 All other child-level characteristics were dichotomized. Breastfeeding was categorized as yes if the
104 mother reported any breastfeeding in the 30-day period before assessment.¹⁰ LBW was defined as
105 <2500g, PTB as gestational age <37 weeks. Due to limitations in gestational assessments, we did not

106 include a variable for small for gestational age (SGA). We captured illness by including any reported
107 episode of diarrhoea within the prior 14 days, and study physician diagnoses of pneumonia or
108 caretaker reports of WHO symptom classifications in the last 14 days for lower respiratory tract
109 infections (LRTI). Severe illness was defined as being admitted to hospital or having any WHO
110 Integrated Management of Childhood Illness (IMCI) danger sign, i.e., not able to drink or breastfeed,
111 convulsions, lethargic or unconscious¹³. The IMCI danger sign of vomiting everything was not
112 consistently available for inclusion in the pooled dataset.

113

114 Each study was classified into one of three population types based on the assumption that
115 associations between risk exposures and mortality would differ between subsets of children: (1)
116 “General population” (GP) if enrolment to the original study was not based on an anthropometric
117 deficit or the presence of an illness, (2) Anthropometry-Selected (A-S) if children were enrolled based
118 on an anthropometric deficit, or (3) Illness-Selected (I-S) if children were enrolled on the basis of
119 signs or symptoms of an illness (potentially in addition to other eligibility criteria)¹⁰.

120

121 Statistical analysis

122 All statistical analyses were done using R version 4.3.1 (R Foundation for Statistical Computing,
123 Vienna, Austria, 2023). To estimate the probability of dying within an observation period, we used
124 multiple generalized linear mixed effects regression (glmer) models of the binomial distribution
125 family and logit link function. The dependent variable of these models was whether a child died
126 during the observation period, and all models included child and study as random intercepts. As the
127 time intervals between observations affect the possibility of capturing death, we also included the
128 log transformed time in months between each visit as an offset.

129

130 We built separate models for each exposure; all models included age at time of visit. From these
131 models, we calculated odds ratios (see supplementary material) and the predicted absolute risk of
132 death per child-month. We report these according to age groups 0-5, 6-11, 12-23, and 24-59 months.
133 In addition, we graphically display the predicted absolute mortality risk over the age range of 0-24
134 months. All models included an interaction term between age at the time of visit and the other
135 independent variable. To account for differences in the completeness of data, missing observations
136 from independent variables were included as a distinct category, such that the risk of dying was
137 separately calculated for children with missing values.

138

139 To estimate the association of not being breastfed, LBW, or PTB, combined with an anthropometric
140 deficit, we estimated the absolute risk using the same statistical approach noted above. We estimate
141 the risk for none, only one of the exposures (e.g., underweight but breastfed) and both together
142 (e.g., underweight AND not breastfed). Only children with information on both exposures were used
143 in the analyses. Data availability for each study is detailed in the cohort profile.¹⁰

144

145 All analyses were run separately for the three population categories: GP, A-S, or I-S. In additional
146 analyses, we report absolute risks of death only for children with death age available (Table S2). We
147 decided a priori to report only those estimates where there were more than 20 observations within
148 each age-specific risk exposure group.

149

150 Ethics

151 The funders of the studies had no role in study design, data collection, data analysis, data
152 interpretation, or writing of the report. All principal investigators contributing data signed an
153 agreement with WHO for pooling and use of anonymous study data for pre-defined analyses. All
154 study investigators confirmed that each included study had appropriate ethical review approval and
155 that informed consent was obtained from all participants. The WHO Ethics Review Committee
156 granted an exemption in March 2022 (ERC.0003745). The pooled data are kept on a secure server at
157 the University of Bergen, Norway with restricted access.

158 **RESULTS**

159 Pooled dataset

160 Of 35 datasets received, two did not meet the inclusion criteria (Figure 1), and data from 33 studies
161 were included and pooled. In total, 4.5% observations were excluded from analyses (i.e., 949
162 duplicate observations, 22,342 observations without age at a single observation timepoint, 160
163 observations occurring at age >60 months, and 1,047 observations without information on sex).¹⁰
164 The sample sizes of included studies ranged from 247 to 7,956 children. The final pooled dataset
165 included individual-level data from 75,287 children aged 0-5 years and 546,459 observations. There
166 were 23 studies classified as GP (49,740 children), six studies classified as A-S (14,759 children), and
167 four studies classified as I-S (10,788 children) (Figure 1).

168

169 **[Figure 1: Study flow]**

170

171 Baseline characteristics of children differed between the three population types (Table 1). Children in
172 GP studies were younger at enrolment (median age of one month as compared to 13.6 months (A-S)
173 and 10.7 months (I-S)). In studies in which birthweight and gestational age were collected, the
174 proportion of children without information on birthweight was 6% among GP studies (1,659/28,326
175 children), 14% among A-S studies (174/1,286), and 10% among I-S studies (956/9,515). The
176 proportion of missing data on PTB was 2% in GP studies (574/28,274 children) and 0.7% in I-S studies
177 (47/6,414 children). None of the A-S studies had information on PTB. The proportion of missing data
178 increased with age; among children aged 12-23 months, this information was not available for 24-
179 39% of the children (Supplementary Table S3). The anthropometric values among children in A-S
180 studies at time of first measurement were lower than in GP and I-S studies. Symptoms and signs of
181 illness were more common among children in I-S studies. Median maternal HIV prevalence also
182 differed significantly by study type (Table 1).

183

184 Follow-up duration and frequency varied between studies: median (IQR) follow-up time per child
185 was 17 (11,23), 3 (1,6), and 6 (5.9,6.2) months, in the GP, A-S and I-S populations, respectively. The
186 median (IQR) interval between visits was 1 (0.9,1.3) month in the GP population, 0.5 (0.4,0.9)
187 months in the A-S population, and 3 (1.6,6) months in the I-S population. Fourteen of the 23 GP
188 studies were birth cohorts (enrolling up to 17 days after birth) including 25,992 of 49,739 children
189 (52%); only one of six A-S studies (2,079/14,760 children; 14%) and none of four I-S studies enrolled
190 from birth. See Table S1 and the cohort description.¹⁰

191

192 There were 2,805 deaths during a total of 69,085 child-years of follow up. Death occurred in 2.5% of
193 children in the GP population (1,238/49,740), 2.5% in the A-S population (366/14,759) and 11% in
194 the I-S population (1,201/10,788). Age at time of death was documented for 2,660 (95%) children.

195

196 **[Table 1]**

197

198 Mortality risks in the GP population

199 Age was a strong predictor of death, with the highest absolute risk of death within the first months
200 after birth, declining with increasing age (see Figure 2). The risk of death increased with lower
201 anthropometric status at all ages and for all anthropometric indices (Figure 2). The mortality risks in
202 the age groups <6, 6-11, 12-23, and 24-59 months can be found in Supplementary Tables S3 and S4.
203 The highest absolute mortality risks were among those with very low WAZ and youngest age.
204 However, the age gradient was not apparent for MUAC, where the absolute mortality risk was
205 relatively stable between birth and 24 months (Figure 2). Applying modified cut-offs for moderate
206 and severe deficits among children aged 0-5 months (i.e., 120 and 110mm) did not substantially
207 change the risk estimates (Supplementary Table S3). The mortality risk associated with a deficit in
208 WAZ by stunting status changed with age (Supplementary Table S5). Among children with low WAZ
209 aged less than 12 months, concurrent stunting increased the mortality risk; among those with low
210 WAZ and aged 12 months or more, concurrent stunting was associated with lower mortality risk.

211

212 **[Figure 2: The absolute predicted risk of death in GP children 0-24 months of age according to**
213 **A) age; B) weight-for-age Z-score (WAZ), C) length/height-for-age Z-score (HAZ), D) weight-for-**
214 **length/height Z-score (WHZ), E) mid-upper arm circumference in mm (MUAC), F)**
215 **breastfeeding practices, G) birthweight, and H) age of gestation at birth]**

216

217 Reported symptoms of diarrhea or LRTI did not further discriminate mortality risks (Supplementary
218 Table S3). Very few children were reported to have any severe illness as defined. Infants who were
219 reported as not breastfeeding had a higher absolute mortality risk as compared to those who were
220 receiving any breastfeeding (Figure 2). The mortality risk associated with not breastfeeding was
221 highest in younger children and decreased gradually with increasing age. In the age group 24-59
222 months, there were few children still being breastfed with no differences in absolute mortality risk by
223 breastfeeding status (Supplementary Table S3). The absolute mortality risk was increased among
224 LBW or PTB infants and remained elevated until about one year of age (Figure 2).

225

226 In analyses that combined risk exposures, the absolute risk of mortality in children with even a
227 moderate anthropometric deficit (<-2 Z-scores) was further increased when there was a history of
228 LBW, PTB, or no reported breastfeeding (Table 2 and Figure 3).

229

230 **[Figure 3. The absolute risk of death in GP children 0-24 months of age with combined risk**
231 **exposures: either not breastfeeding, low birthweight, or being born preterm, combined with**
232 **low weight-for-age (WAZ), low height-for-age (HAZ), low weight-for-height (WHZ) or low mid-**
233 **upper arm circumference (MUAC). Analyses are restricted to children where data on both**
234 **indicators are available (see also Table 2)]**

235

236 **[Table 2]**

237

238 Mortality risks in the anthropometry-selected (A-S) and illness-selected (I-S) populations

239 Although children in the A-S population were older than those in the GP population, mortality was
240 similar in both groups (2.5% in both). However, in the I-S population, mortality was significantly
241 higher (11%) and did not decrease as clearly with increasing age (Supplementary Table S6).

242

243 Anthropometric deficits in the A-S and I-S populations tended to be more severe, but these deficits
244 did not identify mortality risk as clearly as in GP studies. (Tables S6 and S7) There was only one A-S
245 study with information on birthweight and LBW did not increase the risk of death. In I-S studies,
246 there was an increased mortality risk in LBW and PTB (although estimates for PTB were imprecise
247 due to low number of preterm children (Tables S8 and S9).

248

249 Among infants less than six months of age in I-S studies, the mortality risk of those with an IMCI
250 danger sign was six times higher compared to those without. The absolute mortality risk decreased
251 with age but remained higher than in children without danger signs in each age group. The risk
252 increased if there was more than one danger sign present – though confidence intervals were
253 overlapping. There were too few children in A-S studies with multiple dangers signs to enable
254 analyses.

255

256 In both the A-S and the I-S child populations, the mortality risk of not being breastfed added to the
257 risks associated with anthropometric deficits for all measures, although sample sizes were limited
258 with overlapping confidence intervals (Tables 3 and 4).

259

260 **[Table 3]**

261

262 [Table 4]

263 **DISCUSSION**

264 Among 75,287 children under five years old in 33 pooled cohorts across a range of low resource
265 settings, four readily assessable child-level characteristics – age less than 24 months, low weight-for-
266 age, history of low birthweight or preterm birth, and not breastfeeding in the immediate period
267 preceding – identified children at high risk of mortality. The highest mortality risk (GP studies) was
268 seen at the youngest ages, with a relatively low risk of death beyond two years of age. Young children
269 with even a moderate anthropometric deficit and an additional risk exposure (LBW, PTB or not
270 breastfeeding) were at very high risk.

271

272 Our analyses are consistent with other reports that most child deaths occur in the first days of life^{2,5}
273 and that mortality declines rapidly during the first year. Despite these data, infants under twelve
274 months of age are not consistently prioritized in health and nutrition programmes¹⁴, though some
275 efforts are trying to address this.^{15,16} In contrast to WAZ, WHZ identified mortality risk better with
276 increasing age, but showed poor discrimination among infants as reported elsewhere.^{17,18} An
277 important finding was that concurrent stunting and underweight among infants was associated with
278 increased mortality; whereas, after 12 months of age, concurrent stunting was associated with a
279 lower mortality risk. This possibly reflects specific vulnerabilities associated with small size at birth
280 that are largely lost by one year and points to the need for sustained interventions throughout the
281 first 12 months of life for LBW infants. The mortality risk associated with low MUAC was relatively
282 stable between 1-24 months of age reflecting the balance between older children meeting MUAC
283 criteria being relatively more wasted while their mortality risks tending to decline with increasing
284 age. In our analyses, using cut-offs of <110mm and <120mm for severe and moderate wasting in
285 infants <6 months of age (mentioned in WHO 2023 Wasting guidelines) did not substantially change
286 mortality risks compared to 115mm and 125mm. Nevertheless, it might have implications for
287 caseload and choice of intervention.¹⁹ Further analyses of the pooled dataset will examine the
288 performance of anthropometric criteria.

289

290 As noted by others, children born preterm or with LBW had a higher mortality risk²⁰⁻²³. In our
291 analyses, mortality was high throughout the first year of life and remained increased until 24 months
292 of age. The addition of information about LBW, PTB, or not breastfeeding to anthropometry added
293 considerably to the identification of infants at high and lower mortality risk. Between birth and 24
294 months, children in the GP population who were even moderately underweight but had a history of
295 LBW had increased mortality risk. Notably however, children who were born LBW but who were not
296 underweight, i.e., who had caught up, were not at increased risk. Infants who were moderately

297 underweight and not breastfed were similarly at increased mortality risk. In these infants, recovering
298 or maintaining WAZ probably reflects general nurturing care and health, in addition to appropriate
299 breastfeeding. In A-S and I-S studies, only the combination of any low anthropometric status and not
300 being breastfed carried an additional risk. These analyses demonstrate how the context and inclusion
301 criteria of individual studies needs to be considered when interpreting the performance of risk
302 indicators.

303

304 Reported symptoms of diarrhea and LRTI did not add to the predicted mortality risk in the GP and A-
305 S populations. These conditions clearly contribute to child mortality, but the value of self-reported
306 illness is limited. Diarrhea episodes were likely mild and thus did not pose a measurable mortality
307 risk. Pneumonia is similarly difficult to diagnose, and commonly used syndromic definitions have low
308 specificity. There were also few deaths in these children leading to imprecision and probably an
309 underestimation of the risks associated with pneumonia²⁴. In I-S studies, the hospitalization period
310 accounted for approximately 50% deaths occurring during 6-months of observation; about 50% post-
311 discharge deaths also occurred within one month of discharge, highlighting both the risks of serious
312 conditions requiring admission and the continued mortality risk in children that persists after
313 discharge from acute care.^{25,26}

314

315 Strengths of the analyses include a large sample size originating from 33 relatively recent studies
316 conducted in different regions. The analytical techniques were robust and, different from previous
317 analyses^{27,28}, accounted for repeated measures, clustering within studies as well as differences
318 between measurement intervals. Importantly, we report predicted absolute risks by age groups
319 within the first five years of age. Several analyses have previously reported relative risk measures,
320 such as a hazard ratio, and have not generally considered the effects of age^{29,30}. While a relative risk
321 informs about the magnitude of association between an exposure and an outcome, it neglects the
322 magnitude of the problem, i.e. the absolute risk of dying in different populations and age groups.

323 Limitations of the analyses include that studies in the pooled dataset were not necessarily
324 representative of countries or specific populations. Eligibility criteria of original studies may also have
325 skewed the age and exposure distribution which should be considered when interpreting the
326 findings. A paucity of data in the first week of life limited our ability to examine early life risk
327 exposures in more detail.

328

329 The analyses demonstrate that assessable, child-level characteristics can be used to identify young
330 children at high and lower risk of mortality. Individual child-centred risk assessment could improve

331 care and outcomes of high-risk children while improving management and efficiencies for lower-risk
332 children. To mitigate risks, low birth weight and preterm infants can be identified at birth and
333 immediately followed-up and supported over a 12-month period to optimise care and nutrition. Low
334 weight-for-age routinely identified at facility or community health visits should prompt the
335 identification of other concurrent risks such as LBW, PTB or not breastfeeding. This would identify
336 children needing closer follow-up and ensure, at minimum, continued monitoring and that
337 vaccinations are received, and mothers are well and able to breastfeed. Young children being
338 discharged from hospital could be actively followed-up in communities and facilities through the first
339 month post-discharge. We need to learn how treatment of specific conditions such as pneumonia or
340 diarrhoea may differ by risk status.

341

342 Finally, it is important to recognise that treatment or support for a high-risk child may not primarily
343 target the exposure that identifies the child as being vulnerable. For example, it is not possible to
344 change the birth weight or gestational age. However, it is possible to implement interventions to
345 modify their risk such as improved support and follow-up care. A strength of the findings is their
346 relative simplicity that point to an opportunity and paradigm shift toward more child-centred and
347 targeted care. Avoiding the temptation to over-complicate risk assessments will be essential for
348 programmes.

349

350 More effective targeting of interventions to those children at highest risk, while de-escalating care
351 for those at significantly lower risk, offers enormous opportunity. In settings where resources are
352 constrained, it is imperative that we optimize the efficiency of programmes while reducing mortality
353 and morbidity as effectively as possible. These data suggest that simple risk assessment tools that
354 are predicated on individual child characteristics, in addition to the severity of illnesses, may allow
355 for differentiation in care and approaches that further improve child survival, health and
356 development.

357

358 [Authors' contributions](#)

359 Conceptualization: Catherine Schwinger, Nigel Rollins, Rajiv Bahl, Tor Strand; Design: all
360 WHO-RSWG members; Statistical analysis: Catherine Schwinger, Tor Strand, Ayushi, James A.
361 Berkley; Data acquisition and interpretation: all WHO-RSWG members; Drafting the
362 manuscript: Catherine Schwinger, Nigel Rollins, James A. Berkley, Tor Strand, Judd L. Walson;
363 Reviewing the manuscript: all WHO-RSWG members.

364

365 [Declaration of interest](#)

366 All authors confirm no conflict of interest

367

368 [Role of the funding source](#)

369 Funding sources had no role in study design; in the collection, analysis, and interpretation of data; in
370 the writing of the report; and in the decision to submit the paper for publication.

371

372 [Acknowledgments](#)

373 We thank all study participants and their families for their valuable time in the studies; the teams of
374 each individual study for conducting the study; Tejshri Shah for initial coordination of the working
375 group; and Håkon Gjessing for helpful discussions.

376

377 References

- 378 1. UNICEF. Under-five mortality. 2023. [https://data.unicef.org/topic/child-survival/under-](https://data.unicef.org/topic/child-survival/under-five-mortality)
379 five-mortality (accessed 28th August 2023).
- 380 2. Sharrow D, Hug L, You D, et al. Global, regional, and national trends in under-5 mortality
381 between 1990 and 2019 with scenario-based projections until 2030: a systematic
382 analysis by the UN Inter-agency Group for Child Mortality Estimation. *Lancet Glob*
383 *Health* 2022; **10**(2): e195-e206.
- 384 3. India State-Level Disease Burden Initiative Child Mortality C. Subnational mapping of
385 under-5 and neonatal mortality trends in India: the Global Burden of Disease Study
386 2000-17. *Lancet* 2020; **395**(10237): 1640-58.
- 387 4. Macharia PM, Giorgi E, Thurania PN, et al. Sub national variation and inequalities in
388 under-five mortality in Kenya since 1965. *BMC Public Health* 2019; **19**(1): 146.
- 389 5. Sankar MJ, Natarajan CK, Das RR, Agarwal R, Chandrasekaran A, Paul VK. When do
390 newborns die? A systematic review of timing of overall and cause-specific neonatal
391 deaths in developing countries. *J Perinatol* 2016; **36 Suppl 1**(Suppl 1): S1-S11.
- 392 6. Black RE, Victora CG, Walker SP, et al. Maternal and child undernutrition and overweight
393 in low-income and middle-income countries. *Lancet* 2013; **382**(9890): 427-51.
- 394 7. Childhood Acute Illness & Nutrition Network. Childhood mortality during and after
395 acute illness in Africa and south Asia: a prospective cohort study. *Lancet Glob Health*
396 2022; **10**(5): e673-e84.
- 397 8. Wen B, Brals D, Bourdon C, et al. Predicting the risk of mortality during hospitalization in
398 sick severely malnourished children using daily evaluation of key clinical warning signs.
399 *BMC medicine* 2021; **19**(1): 222.
- 400 9. George EC, Walker AS, Kiguli S, et al. Predicting mortality in sick African children: the
401 FEAST Paediatric Emergency Triage (PET) Score. *BMC Med* 2015; **13**: 174.
- 402 10. Schwinger C, Kaldenbach S, Berkley JA, et al. Cohort profile: The WHO Child Mortality
403 Risk Stratification Multi-Country Pooled Cohort (WHO-CMRS) to identify predictors of
404 mortality through early childhood. *medRxiv* 2024: 2024.03.06.24303859.
- 405 11. WHO. WHO Child Growth Standards: Methods and development: Length/height-for-age,
406 weight-for-age, weight-for-length, weight-for-height and body mass index-for-age.
407 Geneva: World Health Organization Multicentre Growth Reference Study; 2006.
- 408 12. WHO. WHO guideline on the prevention and management of wasting and nutritional
409 oedema (acute malnutrition) in infants and children under 5 years. Geneva: World
410 Health Organization, 2023.
- 411 13. WHO. Integrated Management of Childhood Illness (IMCI): Chart Booklet. Geneva,
412 Switzerland: WHO, 2014.
- 413 14. Grey K, Rana R, McGrath M, Kerac M. Key considerations for developing patient
414 management tools for small and nutritionally at-risk infants: A scoping review. *World*
415 *Nutrition* 2023; **14**(1): 58-75.
- 416 15. Engler R, McGrath M, Kerac M. Training Packages and Patient Management Tools for
417 Healthcare Staff Working with Small, Nutritionally At-Risk Infants Aged under 6 Months:
418 A Mixed-Methods Study. *Children (Basel)* 2023; **10**(9).
- 419 16. Grey K, Brennan E, Kerac M, McGrath M. The MAMI Care Pathway Package: A resource
420 to support the management of small and nutritionally at-risk infants under six months
421 of age and their mothers (MAMI). *South Sudan Medical Journal* 2021; **14**(3).

- 422 17. Lelijveld N, Kerac M, McGrath M, Mwangome M, Berkley JA. A review of methods to
423 detect cases of severely malnourished infants less than 6 months for their admission
424 into therapeutic care: Emergency Nutrition Network, 2017.
- 425 18. Mwangome M, Ngari M, Fegan G, et al. Diagnostic criteria for severe acute malnutrition
426 among infants aged under 6 mo. *Am J Clin Nutr* 2017; **105**(6): 1415-23.
- 427 19. Khara T, Myatt M, Sadler K, et al. Anthropometric criteria for best-identifying children at
428 high risk of mortality: a pooled analysis of twelve cohorts. *Public Health Nutr* 2023;
429 **26**(4): 1-17.
- 430 20. Lawn JE, Ohuma EO, Bradley E, et al. Small babies, big risks: global estimates of
431 prevalence and mortality for vulnerable newborns to accelerate change and improve
432 counting. *Lancet* 2023; **401**(10389): 1707-19.
- 433 21. Upadhyay RP, Martines JC, Taneja S, et al. Risk of postneonatal mortality, hospitalisation
434 and suboptimal breast feeding practices in low birthweight infants from rural Haryana,
435 India: findings from a secondary data analysis. *BMJ Open* 2018; **8**(6): e020384.
- 436 22. Zoleko-Manego R, Mischlinger J, Dejon-Agobé JC, et al. Birth weight, growth, nutritional
437 status and mortality of infants from Lambaréné and Fougamou in Gabon in their first
438 year of life. *PLoS One* 2021; **16**(2): e0246694.
- 439 23. O'Leary M, Edmond K, Floyd S, Newton S, Thomas G, Thomas SL. A cohort study of low
440 birth weight and health outcomes in the first year of life, Ghana. *Bull World Health
441 Organ* 2017; **95**(8): 574-83.
- 442 24. Martin H, Falconer J, Addo-Yobo E, et al. Assembling a global database of child
443 pneumonia studies to inform WHO pneumonia management algorithm: Methodology
444 and applications. *J Glob Health* 2022; **12**: 04075.
- 445 25. CHAIN Network. Childhood mortality during and after acute illness in Africa and south
446 Asia: a prospective cohort study. *Lancet Glob Health* 2022; **10**(5): e673-e84.
- 447 26. Wiens MO, Bone JN, Kumbakumba E, et al. Mortality after hospital discharge among
448 children younger than 5 years admitted with suspected sepsis in Uganda: a prospective,
449 multisite, observational cohort study. *Lancet Child Adolesc Health* 2023; **7**(8): 555-66.
- 450 27. Thurstans S, Wrottesley SV, Fenn B, et al. Anthropometric deficits and the associated risk
451 of death by age and sex in children aged 6-59 months: A meta-analysis. *Matern Child
452 Nutr* 2023; **19**(1): e13431.
- 453 28. Schwinger C, Golden MH, Grellety E, Roberfroid D, Guesdon B. Severe acute
454 malnutrition and mortality in children in the community: Comparison of indicators in a
455 multi-country pooled analysis. *PLoS One* 2019; **14**(8): e0219745.
- 456 29. Olofin I, McDonald CM, Ezzati M, et al. Associations of suboptimal growth with all-cause
457 and cause-specific mortality in children under five years: a pooled analysis of ten
458 prospective studies. *PLoS One* 2013; **8**(5): e64636.
- 459 30. McDonald CM, Olofin I, Flaxman S, et al. The effect of multiple anthropometric deficits
460 on child mortality: meta-analysis of individual data in 10 prospective studies from
461 developing countries. *The American journal of clinical nutrition* 2013; **97**(4): 896-901.
462

1 **Table 1: Characteristics of children and their mothers in the pooled dataset according to enrolment criteria groups (general populations**
 2 **(GP), anthropometry-selected (A-S) and illness-selected (I-S) populations)**

Characteristic (where data available)	GP ¹		A-S ¹		I-S ¹	
	N ²	% / median (IQR)	N ²	% / median (IQR)	N ²	% / median (IQR)
Maternal characteristics at enrolment						
Mother alive	19,702	99.8%	12,467	98.2%	10,721	99.0%
Mother primary caretaker	15,052	98.1%	12,866	94.6%	9,444	93.9%
Maternal age, years	42,549	24 (20, 29)	2,079	23 (21, 25)	10,78	26 (22, 30)
Maternal education	34,622		5,941		7,472	
no education		35%		37%		17%
primary education		25%		10%		25%
secondary or higher education		40%		53%		58%
Mother HIV positive	10,872	0.2%	4,418	14.6%	1,517	18.5%
Child's characteristics at enrolment						
	N ²		N ²		N ²	
Child's sex, male	49,740	50.5%	14,759	42.8%	10,788	55.6%
Child age, months	49,740	1.0 (0, 7.8)	14,759	13.6 (8.3, 22.6)	10,788	10.7 (4.7, 18.7)
Low birthweight, <2500g	26,667	13.3%	1,112	5.0%	8,559	7.8%
Birthweight, g	26,667	3000 (2700, 3400)	1,112	3000 (2900, 3400)	2,145	3000 (2500, 3500)
Preterm birth, <37 weeks of gestation	27,700	15.7%	N/A	N/A	6,367	4.3%
WAZ	42,636	-0.9 (-1.7, -0.1)	14,754	-2.9 (-3.6, -2.3)	10,767	-1.4 (-2.8, -0.4)
HAZ	41,251	-1.1 (-1.9, -0.3)	14,747	-2.5 (-3.5, -1.7)	10,721	-1.2 (-2.4, 0.1)
WHZ	39,853	-0.3 (-1.2, 0.7)	14,168	-2.2 (-2.8, -1.6)	10,665	-1.3 (-2.6, -0.1)

MUAC, mm	33,423	120 (104, 135)	14,717	118 (110, 123)	10,762	128 (114, 141)
Reported illness or feeding practice at any visit	Obs²		Obs²		Obs²	
Acute diarrhoea ³	313,381	7.6%	53,279	30.2%	18,672	43.7%
Acute lower respiratory tract infection ⁴	150,874	3.2%	N/A	N/A	18,671	40.1%
Severe illness ⁵	242,980	1.2%	36,476	9.6%	17,461	28.4%
Any breastfeeding ⁶						
at age 0-5 months	92,024	98.9%	7,947	98.8%	3,622	91.2%
at age 6-23 months	202,644	87.0%	25,314	81.6%	12,673	70.6%
at age 24-59 months	21,306	18.8%	7,521	16.0%	2,423	17.1%

¹ We categorized studies on the basis of their enrolment criteria, where GP = study cohorts where enrolment was not based on anthropometry or the presence of illness, A-S = study cohorts where children were enrolled on the basis of an anthropometric deficit, I-S = study cohorts where children were enrolled on the basis of the presence of illness using study specific definitions.

² N = total number of children with data available for this variable, Obs = total number of observations available for this variable

³ Any diarrheal episode reported by a clinician or caregiver within the last 24 hours, 3, 7 or 14 days.

⁴ Pneumonia or acute lower respiratory tract infection reported by a clinician or caregiver within the last 14 days.

⁵ Any of the following: being admitted to hospital, not being able to drink or breastfeed, having had convulsions, being lethargic or unconscious.

⁶ Any breastfeeding within the past 14 days reported by caregiver at the study visit.

Abbreviations: HAZ = height-for-age Z-score (includes length-for-age Z-scores); IQR = interquartile range; MUAC = mid-upper arm circumference; N = number of children; Obs = number of observations, WAZ = weight-for-age Z-score; WHZ = weight-for-height Z-score (includes weight-for-length Z-scores).

15 **Table 2: Absolute risk of death per 1,000 child months by age group and by combinations of exposures in General Population (GP)**
 16 **cohorts, i.e., enrolment not on the basis of anthropometric deficit or the presence of illness**

Indicator	<6 months				6-11 months				12-23 months				24-59 months			
	N	Death	Risk ¹	95% CI	N	Death	Risk ¹	95% CI	N	Death	Risk ¹	95% CI	N	Death	Risk ¹	95% CI
BF & WAZ ≥-2	78,448	72	0.4	0.2, 0.8	70,186	99	0.4	0.2, 0.8	50,120	37	0.4	0.2, 0.8	1,608	1	0.4	0.2, 0.9
No BF & WAZ ≥-2	706	3	1.4	0.5, 4.3	2,110	10	1.0	0.4, 2.5	12,040	9	0.7	0.3, 1.8	8,897	1	0.5	0.2, 1.7
BF & WAZ <-2	9,564	58	2.0	0.9, 4.6	17,841	76	1.5	0.7, 3.3	18,938	44	1.1	0.5, 2.5	981	4	0.8	0.3, 2.1
No BF & WAZ <-2	192	5	5.3	1.5, 18.4	648	4	3.7	1.5, 9.2	5,466	7	2.6	1.0, 6.3	5,454	4	1.8	0.5, 5.9
NBW & WAZ ≥-2	75,955	241	0.9	0.4, 2.0	57,350	94	0.6	0.2, 1.3	35,137	34	0.4	0.1, 0.8	14,458	1	0.2	0.1, 0.6
LBW & WAZ ≥-2	4,962	31	1.6	0.7, 4.0	4,055	10	0.8	0.3, 2.0	2,524	3	0.4	0.1, 1.3	1,159	0	0.2	0.0, 0.9
NBW & WAZ <-2	5,032	30	2.3	0.9, 5.5	8,697	50	1.6	0.7, 3.8	7,419	15	1.1	0.5, 2.8	4,681	1	0.8	0.3, 2.3
LBW & WAZ <-2	5,808	140	6.6	2.8, 15.3	3,689	19	2.5	1.0, 6.1	2,693	10	1.0	0.4, 2.7	1,352	0	0.4	0.1, 1.3
Term & WAZ ≥-2	73,293	245	1.7	1.0, 3.1	55,343	94	1.0	0.6, 1.9	32,746	33	0.6	0.3, 1.1	9,632	1	0.4	0.2, 0.7
PTB & WAZ ≥-2	9,723	31	2.4	1.2, 4.7	8,232	19	1.7	0.9, 3.2	4,622	7	1.2	0.5, 2.6	784	0	0.8	0.3, 2.3
Term & WAZ <-2	5,598	75	6.2	3.3, 11.4	7,662	54	3.6	2.0, 6.6	5,305	20	2.1	1.1, 4.1	1,192	1	1.2	0.6, 2.7
PTB & WAZ <-2	2,496	97	22.3	12.2, 40.7	1,802	18	8.9	4.7, 17.0	1,126	9	3.6	1.6, 8.1	163	0	1.4	0.5, 4.1
BF & WAZ ≥-3	85,076	87	0.4	0.2, 1.0	82,904	140	0.5	0.2, 1.0	63,805	60	0.5	0.2, 1.1	2,331	4	0.5	0.2, 1.2
No BF & WAZ ≥-3	796	3	1.3	0.4, 4.0	2,537	11	1.1	0.5, 2.6	15,948	14	0.9	0.4, 2.1	12,969	2	0.7	0.3, 2.2
BF & WAZ <-3	2,936	43	4.8	2.1, 10.8	5,123	35	2.8	1.3, 6.0	5,253	21	1.6	0.7, 3.8	258	1	0.9	0.3, 2.6
No BF & WAZ <-3	102	5	13.2	3.7, 45.7	221	3	7.8	3.0, 19.9	1,558	2	4.6	1.6, 12.9	1,382	3	2.7	0.6, 11.7
NBW & WAZ ≥-3	79,918	254	0.9	0.4, 2.1	64,021	122	0.6	0.3, 1.4	41,024	41	0.4	0.2, 1.0	18,321	2	0.3	0.1, 0.7
LBW & WAZ ≥-3	8,422	69	2.1	0.9, 5.0	6,116	14	0.9	0.3, 2.1	4,114	4	0.3	0.1, 1.0	1,989	0	0.1	0.0, 0.5
NBW & WAZ <-3	1,069	17	5.0	2.0, 12.8	2,026	22	3.6	1.5, 8.6	1,532	8	2.6	0.9, 6.9	818	0	1.8	0.5, 6.3
LBW & WAZ <-3	2,348	102	13.5	5.8, 31.3	1,628	15	5.1	2.1, 12.3	1,103	8	1.9	0.7, 5.5	522	0	0.7	0.2, 2.7
Term & WAZ ≥-3	77,551	280	1.9	1.1, 3.4	61,205	120	1.2	0.7, 2.1	37,010	42	0.7	0.4, 1.3	10,644	2	0.4	0.2, 0.8
PTB & WAZ ≥-3	11,066	49	3.2	1.7, 6.0	9,423	25	2.1	1.1, 3.8	5,441	10	1.3	0.6, 2.8	913	0	0.8	0.3, 2.2
Term & WAZ <-3	1,340	40	13.4	7.0, 25.3	1,800	28	8.5	4.6, 15.7	1,041	11	5.4	2.5, 11.3	180	0	3.4	1.3, 9.0
PTB & WAZ <-3	1,153	79	40.1	22.0, 72.1	611	12	17.9	9.1, 34.9	307	6	7.9	3.1, 20.2	34	0	3.5	0.9, 12.6
BF & HAZ ≥-2	73,307	71	0.4	0.2, 1.0	68,367	113	0.4	0.2, 0.9	41,912	38	0.4	0.2, 1.0	1,310	2	0.4	0.2, 1.0
No BF & HAZ ≥-2	632	3	1.6	0.5, 4.8	1,983	10	1.3	0.6, 3.1	8,186	11	1.1	0.5, 2.6	6,453	1	0.9	0.3, 2.9
BF & HAZ <-2	13,276	58	1.6	0.7, 3.4	19,417	62	0.8	0.5, 2.4	27,001	42	0.8	0.4, 1.8	1,279	3	0.6	0.2, 1.4
No BF & HAZ <-2	261	5	3.3	1.0, 11.2	776	4	1.3	0.8, 5.1	9,123	4	1.3	0.5, 3.2	7,783	4	0.8	0.2, 2.7
NBW & HAZ ≥-2	71,054	221	0.9	0.4, 2.0	55,216	103	0.6	0.3, 1.4	29,039	32	0.4	0.2, 1.0	10,737	0	0.3	0.1, 0.8
LBW & HAZ ≥-2	5,116	36	1.7	0.7, 4.2	4,002	14	1.2	0.5, 2.9	1,903	5	0.8	0.3, 2.3	811	0	0.9	0.1, 2.1
NBW & HAZ <-2	8,591	45	1.9	0.8, 4.6	10,650	41	1.2	0.5, 2.7	13,251	17	0.7	0.3, 1.7	8,269	2	0.4	0.2, 1.1
LBW & HAZ <-2	5,540	133	6.7	2.9, 15.6	3,708	15	2.0	0.8, 4.8	3,299	8	0.6	0.2, 1.7	1,689	0	0.2	0.0, 0.6
Term & HAZ ≥-2	68,724	240	1.8	0.9, 3.4	52,045	101	1.1	0.6, 2.1	25,958	28	0.7	0.4, 1.4	7,046	0	0.4	0.2, 0.9
PTB & HAZ ≥-2	8,451	29	2.5	1.2, 5.1	7,188	18	2.3	1.1, 4.5	3,191	9	2.0	0.9, 4.7	467	0	1.8	0.6, 5.4
Term & HAZ <-2	9,234	75	4.4	2.3, 8.5	10,841	47	2.3	1.2, 4.5	11,886	24	1.2	0.6, 2.5	3,653	2	0.7	0.3, 1.5
PTB & HAZ <-2	3,567	97	17.4	9.0, 33.2	2,801	19	5.3	2.7, 10.7	2,533	7	1.6	0.7, 3.9	457	0	0.5	0.2, 1.5
BF & WHZ ≥-2	81,063	100	0.5	0.2, 1.1	78,155	123	0.5	0.2, 1.0	60,701	58	0.5	0.2, 1.0	2,311	3	0.4	0.2, 1.0
No BF & WHZ ≥-2	789	6	2.2	0.8, 6.0	2,439	12	1.4	0.6, 3.1	15,545	11	0.9	0.4, 2.0	12,899	3	0.6	0.2, 1.6

BF & WHZ <-2	5,194	25	1-6	0-7, 3-6	9,529	52	1-5	0-7, 3-1	8,182	22	1-4	0-6, 3-1	277	2	1-3	0-5, 3-5
No BF & WHZ <-2	101	2	3-0	0-6, 15-3	317	2	3-3	1-1, 9-8	1,760	4	3-7	1-4, 10-1	1,325	2	4-2	1-0, 18-3
NBW & WHZ ≥-2	74,596	241	1-0	0-5, 2-2	60,139	109	0-7	0-3, 1-4	38,869	36	0-4	0-2, 0-9	17,957	2	0-3	0-1, 0-6
LBW & WHZ ≥-2	7,719	65	2-3	1-0, 5-2	5,826	15	0-9	0-4, 2-1	3,931	4	0-3	0-1, 1-0	2,086	0	0-1	0-0, 0-5
NBW & WHZ <-2	4,447	18	1-4	0-6, 3-3	5,634	35	1-7	0-8, 3-8	3,380	13	2-1	0-9, 4-9	1,037	0	2-5	0-9, 7-0
LBW & WHZ <-2	1,896	32	4-0	1-7, 9-3	1,879	14	3-3	1-4, 7-4	1,269	9	2-7	1-0, 7-2	414	0	2-2	0-6, 7-7
Term & WHZ ≥-2	73,105	268	2-0	1-2, 3-4	57,925	110	1-2	0-7, 2-0	35,400	37	0-7	0-4, 1-2	10,328	2	0-4 0-8	0-2, 0-7
PTB & WHZ ≥-2	10,801	49	3-4	1-9, 6-0	9,121	27	2-1	1-2, 3-7	5,272	9	1-3	0-6, 2-6	891	0	3-8	0-3, 1-9
Term & WHZ <-2	4,249	35	3-9	2-2, 7-1	4,878	38	3-9	2-2, 6-8	2,413	15	3-8	2-0, 7-3	362	0	10-3	1-6, 8-7
PTB & WHZ <-2	729	14	9-7	4-7, 20-0	851	10	9-9	5-3, 18-6	441	7	10-1	4-3, 23-9	32	0	-	2-9, 35-7
BF & MUAC ≥125	34,445	42	1-0	0-6, 1-9	63,418	110	1-0	0-6, 1-8	58,937	59	1-1	0-6, 1-9	936	4	1-1	0-5, 2-1
No BF & MUAC ≥125	452	3	3-2	1-1, 9-1	1,316	11	2-8	1-4, 5-6	10,574	12	2-4	1-2, 4-8	3,468	2	2-1	0-7, 5-9
BF & MUAC <125	13,304	35	2-1	1-1, 3-9	8,251	47	2-8	1-6, 4-9	5,104	21	3-7	2-0, 7-1	64	1	5-0	2-2, 11-4
No BF & MUAC <125	141	3	5-4	1-2, 23-1	94	1	6-6	2-6, 16-6	1,108	2	8-0	2-9, 22-0	264	2	9-7	1-9, 48-2
NBW & MUAC ≥125	38,552	99	1-4	0-8, 2-6	48,363	108	1-1	0-6, 2-0	32,485	41	0-9	0-3, 1-7	8,944	2	0-7	0-3, 1-4
NBW & MUAC <125	3,225	16	2-2	1-1, 4-7	4,865	16	1-7	0-9, 3-4	3,107	8	1-4	0-3, 3-1	745	0	1-1	0-3, 3-2
LBW & MUAC <125	22,849	153	2-8	1-5, 5-1	3,792	22	3-2	1-7, 6-2	1,291	8	3-7	1-7, 8-3	47	0	4-3	1-5, 12-0
LBW & MUAC <125	5,611	142	11-5	6-3, 21-1	1,425	10	5-5	2-7, 11-2	560	5	2-6	0-9, 7-2	24	0	1-2	0-3, 5-0
Term & MUAC ≥125	35,504	98	1-5	0-7, 3-1	44,378	100	1-1	0-5, 2-3	30,080	36	0-8	0-4, 1-8	9,456	2	0-6	0-3, 1-4
PTB & MUAC ≥125	3,818	17	2-9	1-2, 6-5	5,722	19	2-5	1-2, 5-4	3,704	11	2-2	0-9, 5-1	559	0	1-9	0-6, 5-6
Term & MUAC <125	21,433	182	3-3	1-6, 6-7	3,664	19	3-2	1-5, 6-9	1,283	10	3-2	1-3, 7-8	59	0	3-1	1-0, 9-4
PTB & MUAC <125	3,578	86	13-9	6-7, 28-6	689	7	8-8	3-8, 20-4	264	4	5-6	1-7, 18-1	12	0	-	-

¹ Risk = predicted absolute risk of dying within 1 month per 1,000 child months modelled using generalized linear mixed effects models accounting for repeated measures in each child, clustering within studies, and time at risk

Abbreviations: BF = breastfeeding; HAZ = height-for-age Z-score; LBW= low birthweight (<2500g); MUAC = mid-upper arm circumference; NBW = normal birthweight (≥2500g); PTB = preterm birth (<37 weeks of gestation at birth); WAZ = weight-for-age Z-score; WHZ= weight-for-height Z-score

17
18
19
20
21

22 **Table 3: Absolute risk of death per 1000 child months by age group and by combinations of exposures in Anthropometry-Selected (A-S)**
 23 **cohorts, i.e., enrolment on the basis of anthropometric deficit**

Indicator	<6 months				6-11 months				12-23 months				24-59 months			
	N	Deaths	Risk ²	95% CI	N	Deaths	Risk ²	95% CI	N	Deaths	Risk ²	95% CI	N	Deaths	Risk ²	95% CI
BF & WAZ ≥-2	4,123	0	1.3	0.2, 8.9	2,439	8	1.6	0.3, 7.4	1,944	3	2.0	0.4, 9.5	113	1	2.5	0.3, 18.2
No BF & WAZ ≥-2	40	0	0.8	0.0, 20.4	142	0	1.3	0.1, 13.7	983	4	2.3	0.4, 11.5	1,314	7	3.8	0.8, 17.4
BF & WAZ <-2	3,706	16	13.3	3.3, 51.7	6,997	79	9.6	2.5, 36.3	9,283	74	7.0	1.8, 26.8	1,206	10	5.0	1.2, 20.7
No BF & WAZ <-2	52	1	33.9	7.7, 136	390	15	19.4	4.8, 74.8	3,129	48	11.0	2.8, 41.8	5,041	49	6.2	1.6, 24.3
NBW & WAZ ≥-2	1	0	570	5	7.5	3.2, 17.6	507	2	3.6	1.0, 13.4	64	0	1.8	0.1, 25.7
LBW & WAZ ≥-2	0	0	10	0	17	0	4	0
NBW & WAZ <-2	0	0	1,564	15	10.1	6.4, 16.2	2,141	19	8.5	5.9, 12.2	528	3	7.1	3.4, 15.0
LBW & WAZ <-2	0	0	90	1	7.8	1.1, 54.7	117	1	12.5	4.0, 38.4	37	1	20.1	2.7, 134
Term & WAZ ≥-2	0	0	0	0	0	0	0	0
PTB & WAZ ≥-2	0	0	0	0	0	0	0	0
Term & WAZ <-2	0	0	0	0	0	0	0	0
PTB & WAZ <-2	0	0	0	0	0	0	0	0
BF & WAZ ≥-3	6,814	3	2.7	0.6, 11.3	5,911	23	2.8	0.8, 10.3	6,122	17	3.0	0.8, 11.0	480	1	3.1	0.7, 13.7
No BF & WAZ ≥-3	65	0	6.0	1.0, 36.4	266	3	5.8	1.3, 25.6	2,259	14	5.6	1.5, 20.5	3,202	21	5.4	1.4, 19.9
BF & WAZ <-3	1,015	13	22.9	6.3, 79.8	3,525	64	15.4	4.4, 52.1	5,105	60	10.3	2.9, 35.7	839	10	6.9	1.8, 25.8
No BF & WAZ <-3	27	1	48.4	11.8, 177	266	12	26.4	7.2, 92.5	1,853	38	14.2	4.0, 49.0	3,153	35	7.6	2.1, 27.2
NBW & WAZ ≥-3	1	0	1,590	12	7.7	4.6, 13.1	1,661	12	6.1	3.6, 10.4	219	0	4.9	1.6, 14.7
LBW & WAZ ≥-3	0	0	34	0	6.9	0.2, 17.4	60	1	7.7	1.1, 52.7	25	0	8.5	0.3, 209
NBW & WAZ <-3	0	0	544	8	14.4	7.4, 27.7	987	9	10.2	6.4, 16.3	373	3	7.2	2.8, 18.4
LBW & WAZ <-3	0	0	66	1	6.5	0.5, 73.4	74	0	14.8	3.7, 57.9	16	1
Term & WAZ ≥-3	0	0	0	0	0	0	0	0
PTB & WAZ ≥-3	0	0	0	0	0	0	0	0
Term & WAZ <-3	0	0	0	0	0	0	0	0
PTB & WAZ <-3	0	0	0	0	0	0	0	0
BF & HAZ ≥-2	4,099	2	5.1	1.0, 24.8	3,775	19	3.4	0.8, 14.1	2,357	5	2.3	0.5, 10.8	158	0	1.6	0.2, 10.5
No BF & HAZ ≥-2	41	0	2.6	0.2, 32.8	193	1	3.2	0.5, 21.4	823	5	3.9	0.9, 17.7	1,115	6	4.8	1.0, 22.4
BF & HAZ <-2	3,643	14	14.8	3.6, 58.4	5,646	68	10.1	2.6, 38.8	8,865	72	6.9	1.7, 26.9	1,161	11	4.7	1.1, 19.6
No BF & HAZ <-2	50	1	36.3	8.1, 148	339	14	19.6	4.8, 76.9	3,288	47	10.5	2.7, 40.5	5,237	50	5.6	1.4, 22.2
NBW & HAZ ≥-2	1	0	765	6	7.4	3.4, 16.1	481	3	6.0	2.1, 17.1	52	0	4.9	0.6, 41.1
LBW & HAZ ≥-2	0	0	10	0	15	0	13	0
NBW & HAZ <-2	0	0	1,367	14	10.4	6.4, 16.9	2,162	18	8.1	5.6, 11.7	539	3	6.3	2.9, 13.5
LBW & HAZ <-2	0	0	90	1	7.3	1.0, 52.7	119	1	12.9	4.1, 39.6	28	1	22.8	3.0, 153
Term & HAZ ≥-2	0	0	0	0	0	0	0	0
PTB & HAZ ≥-2	0	0	0	0	0	0	0	0
Term & HAZ <-2	0	0	0	0	0	0	0	0
PTB & HAZ <-2	0	0	0	0	0	0	0	0
BF & WHZ ≥-2	6,966	8	3.8	0.9, 15.9	5,381	22	3.5	0.9, 13.1	5,917	15	3.2	0.8, 12.4	601	4	3.0	0.7, 13.4
No BF & WHZ ≥-2	75	0	5.9	0.9, 37.0	259	2	5.5	1.2, 25.3	2,457	16	5.1	1.3, 19.6	3,854	24	4.7	1.2, 18.2

BF & WHZ <-2	771	8	17.9	4-6, 66.4	4,038	65	13.1	3-6, 46.9	5,305	62	9.6	2-6, 35.0	717	7	7.0	1-8, 27.6
No BF & WHZ <-2	16	1	273	13	24.2	6.2, 89.1	1,652	36	14.7	4.0, 52.7	2,497	32	8.8	2.3, 33.0
NBW & WHZ ≥-2	1	0	1,645	15	8.5	5.2, 13.9	1,754	11	6.6	4.1, 10.5	338	2	5.1	1.9, 13.3
LBW & WHZ ≥-2	0	0	81	1	12.6	2.1, 73.1	91	1	7.7	1.3, 45.0	20	0	4.6	0.1, 162
NBW & WHZ <-2	0	0	486	5	12.1	5.7, 25.5	888	10	10.0	5.9, 16.8	253	1	8.2	2.8, 23.8
LBW & WHZ <-2	0	0	19	0	43	0	0.0	0, 1000	21	1	57.9	7.7, 327
Term & WHZ ≥-2	0	0	0	0	0	0	0	0
PTB & WHZ ≥-2	0	0	0	0	0	0	0	0
Term & WHZ <-2	0	0	0	0	0	0	0	0
PTB & WHZ <-2	0	0	0	0	0	0	0	0
BF & MUAC ≥125	3,226	1	2.5	0.4, 15.2	2,201	6	2.1	0.4, 9.9	3,903	8	1.8	0.4, 8.3	602	1	1.5	0.2, 9.0
No BF & MUAC ≥125	42	0	7.9	1.0, 60.8	157	2	5.9	1.0, 32.8	1,737	11	4.5	1.0, 20.1	4,072	21	3.4	0.8, 15.0
BF & MUAC <125	4,573	14	10.3	2.4, 44.2	7,207	81	9.0	2.1, 37.3	7,301	69	7.9	1.9, 33.1	708	10	6.9	1.5, 30.9
No BF & MUAC <125	49	1	21.8	4.4, 100	372	13	15.8	3.6, 67.8	2,368	41	11.5	2.7, 47.7	2,264	35	8.3	1.9, 35.6
NBW & MUAC ≥125	0	0	438	2	4.9	1.8, 13.2	949	8	5.9	3.3, 10.8	279	1	7.2	2.3, 22.1
NBW & MUAC <125	0	0	15	0	47	0	0.0	0.0, 1000	18	0
LBW & MUAC <125	1	0	1693	18	11.2	7.1, 17.5	1,700	13	8.4	5.4, 13.1	313	2	6.4	2.5, 15.8
LBW & MUAC <125	0	0	85	1	8.8	1.2, 61.6	87	1	19.0	6.1, 58.0	23	1	40.5	5.5, 244
Term & MUAC ≥125	0	0	0	0	0	0	0	0
PTB & MUAC ≥125	0	0	0	0	0	0	0	0
Term & MUAC <125	0	0	0	0	0	0	0	0
PTB & MUAC <125	0	0	0	0	0	0	0	0

24 ¹ Risk = predicted absolute risk of dying within 1 month per 1,000 child months modelled using generalized linear mixed effects models accounting for repeated measures in
 25 each child, clustering within studies, and time at risk

26 Abbreviations: BF = breastfeeding; HAZ = height-for-age Z-score; LBW= low birthweight (<2500g); MUAC = mid-upper arm circumference; NBW = normal birthweight
 27 (≥2500g); PTB = preterm birth (<37 weeks of gestation at birth); WAZ = weight-for-age Z-score; WHZ= weight-for-height Z-score

28 **Table 4: Absolute risk of death per 1000 child months by age group and by combinations of exposures in Illness-Selected (I-S) cohorts,**
 29 **i.e., enrolment on the basis of the presence of illness**

Indicator	<6 months				6-11 months				12-23 months				24-59 months			
	N	Deaths	Risk ¹	95% CI	N	Deaths	Risk ¹	95% CI	N	Deaths	Risk ¹	95% CI	N	Deaths	Risk ¹	95% CI
BF & WAZ \geq -2	2,235	181	3.6	1.3, 9.8	2,255	63	3.0	1.2, 7.5	2,533	39	2.6	0.9, 7.4	190	0	2.2	0.5, 8.8
No BF & WAZ \geq -2	88	7	6.8	1.5, 29.4	384	19	6.0	1.9, 18.7	1,353	55	5.3	2.0, 13.8	1,371	85	4.7	1.7, 13.0
BF & WAZ <-2	1,045	230	30.7	11.6, 78.9	1,560	120	20.5	8.4, 49.5	2,107	86	13.7	5.0, 36.5	192	2	9.1	2.6, 30.7
No BF & WAZ <-2	226	37	59.6	18.6, 174	563	74	37.3	14.1, 95.2	1,249	119	23.2	9.2, 57.5	598	55	14.3	5.0, 40.4
NBW & WAZ \geq -2	2,164	178	5.0	3.4, 7.3	1,818	56	3.9	3.0, 5.1	2,617	57	3.1	2.2, 4.4	1,056	59	2.5	1.4, 4.2
LBW & WAZ \geq -2	46	5	15.1	1.8, 115.6	154	2	8.2	2.2, 29.8	289	3	4.4	0.9, 22.0	51	2	2.4	0.2, 34.6
NBW & WAZ <-2	835	192	41.6	26.2, 65.4	1,023	102	24.4	17.9, 33.2	1,902	126	14.2	9.8, 20.6	512	43	8.3	4.6, 14.8
LBW & WAZ <-2	239	46	19.4	7.2, 51.1	375	24	18.0	9.0, 35.4	528	19	16.7	6.4, 42.9	54	3	15.5	3.3, 68.6
Term & WAZ \geq -2	1,919	171	3.3	0.6, 19.1	800	43	3.9	0.7, 21.2	936	50	4.6	0.8, 26.4	813	60	5.4	0.8, 36.2
PTB & WAZ \geq -2	34	5	5.1	0.2, 139	20	0	3.5	0.2, 65.8	15	0	15	1
Term & WAZ <-2	553	172	46.2	7.7, 232	278	46	31.0	5.6, 155	428	76	20.7	3.5, 114	357	40	13.7	1.9, 92.6
PTB & WAZ <-2	132	34	28.5	3.2, 210	28	5	28.0	3.2, 203	18	5	12	1
BF & WAZ \geq -3	2,671	261	4.8	1.9, 12.2	2,968	94	3.7	1.6, 8.4	3,619	61	2.8	1.1, 7.3	299	1	2.1	0.6, 7.5
No BF & WAZ \geq -3	145	13	8.6	2.3, 31.1	549	25	7.0	2.5, 19.3	1,871	86	5.8	2.4, 13.7	1,711	109	4.7	1.8, 12.0
BF & WAZ <-3	609	150	45.3	17.0, 115	847	89	34.3	14.4, 79.6	1,021	64	26.0	9.4, 69.5	83	1	19.6	5.1, 72.1
No BF & WAZ <-3	169	31	69.4	20.7, 208	398	68	52.0	20.0, 129	731	88	38.7	15.5, 93.7	258	31	28.8	9.3, 85.3
NBW & WAZ \geq -3	2,537	250	6.3	4.5, 8.9	2,278	83	4.8	3.8, 6.2	3,539	95	3.7	2.7, 5.0	1,366	81	2.8	1.7, 4.5
LBW & WAZ \geq -3	101	12	7.7	1.3, 43.1	293	3	4.2	1.3, 13.0	500	4	2.2	0.4, 11.1	68	2	1.2	0.1, 16.3
NBW & WAZ <-3	462	120	59.8	32.4, 108	563	75	38.2	25.5, 56.9	880	88	24.2	14.6, 39.9	202	21	15.3	6.7, 34.3
LBW & WAZ <-3	184	39	28.5	9.3, 83.9	236	23	32.7	15.0, 69.6	317	18	37.5	13.0, 103	37	3	42.9	8.0, 200
Term & WAZ \geq -3	2,183	235	4.5	0.7, 26.9	934	59	5.0	0.9, 28.5	1,154	78	5.6	0.9, 33.2	1,027	80	6.3	0.9, 42.2
PTB & WAZ \geq -3	61	10	7.6	0.5, 112	24	0	3.7	0.2, 55.9	22	0	1.8	0.0, 96.9	20	1	0.9	0.0, 245
Term & WAZ <-3	289	108	79.7	12.2, 379	144	30	52.9	0.2, 55.9	210	48	34.8	5.3, 197	143	20	22.7	2.6, 172
PTB & WAZ <-3	105	29	33.6	3.4, 263	24	5	48.8	8.9, 259	11	5	-	-	7	1	-	-
BF & HAZ \geq -2	2,357	227	5.5	2.6, 11.5	2,474	91	4.0	2.1, 7.5	2,334	45	2.9	1.3, 6.8	179	0	2.1	0.6, 7.1
No BF & HAZ \geq -2	128	16	15.1	4.7, 47.1	433	35	9.5	4.1, 22.1	1,157	71	6.0	3.0, 12.0	1,305	77	3.8	1.7, 8.4
BF & HAZ <-2	915	182	26.9	12.3, 57.7	1,335	91	16.0	8.3, 30.6	2,300	77	9.4	4.3, 20.7	203	2	5.6	1.9, 16.5
No BF & HAZ <-2	183	28	47.9	16.3, 132	507	57	31.3	14.1, 67.9	1,435	98	20.3	10.2, 40.2	657	61	13.1	5.6, 30.4
NBW & HAZ \geq -2	2,236	219	6.6	4.7, 9.5	2,023	83	4.9	3.8, 6.3	2,360	73	3.6	2.5, 5.1	1,011	53	2.6	1.5, 4.5
LBW & HAZ \geq -2	77	10	9.1	1.4, 57.6	161	4	6.1	1.8, 20.8	228	3	4.1	0.7, 22.5	38	3	2.8	0.2, 43.9
NBW & HAZ <-2	729	149	31.0	18.5, 51.6	813	74	18.8	13.3, 26.4	2,048	104	11.3	7.6, 16.7	551	45	6.8	3.7, 12.5
LBW & HAZ <-2	208	41	25.0	8.6, 70.7	367	22	20.2	9.8, 41.0	589	19	16.2	6.2, 41.6	67	2	13.1	2.8, 58.1
Term & HAZ \geq -2	1,960	206	4.9	1.3, 18.8	885	57	4.7	1.3, 16.9	917	58	4.5	1.1, 17.3	830	53	4.3	0.9, 20.1
PTB & HAZ \geq -2	48	10	12.3	0.9, 146	24	1	9.0	1.0, 74.2	18	1	19	2
Term & HAZ <-2	508	136	33.2	7.9, 128	188	31	23.7	6.3, 84.8	438	63	16.9	4.2, 65.3	334	43	12.0	2.4, 58.3

PTB & HAZ <-2	118	29	29-7	4-3, 177	23	4	23-1	3-1, 152	15	4	8	0
BF & WHZ \geq -2	2,270	228	7-0	3-8, 13-0	2,668	77	4-2	2-5, 7-1	3,267	48	2-6	1-3, 5-2	290	1	1-6	0-5, 4-4
No BF & WHZ \geq -2	165	14	12-7	4-4, 36-0	526	26	9-1	4-4, 18-9	1,736	65	6-5	3-7, 11-5	1,462	88	4-7	2-4, 9-2
BF & WHZ <-2	956	165	16-2	8-3, 31-0	1,140	105	14-6	8-6, 24-8	1,367	74	13-2	6-5, 26-6	91	1	12-0	4-2, 33-4
No BF & WHZ <-2	146	30	52-2	20-5, 126	414	66	33-0	17-0, 63-1	855	104	20-7	11-6, 36-8	493	46	12-9	6-0, 27-6
NBW & WHZ \geq -2	2,114	212	7-6	5-3, 10-7	2,006	67	5-4	4-2, 6-9	3,211	78	3-8	2-8, 5-3	1,180	64	2-7	1-6, 4-5
LBW & WHZ \geq -2	143	20	21-6	5-7, 77-9	321	7	9-3	3-9, 22-0	498	5	3-9	1-1, 14-4	75	2	1-7	0-2, 14-2
NBW & WHZ <-2	849	149	20-0	12-5, 31-8	830	90	15-8	11-5, 21-7	1,197	99	12-4	8-3, 18-6	380	33	9-8	5-2, 18-4
LBW & WHZ <-2	125	22	10-6	3-0, 37-2	207	19	17-2	7-6, 38-3	319	17	27-5	9-8, 75-0	30	3	43-9	8-3, 202
Term & WHZ \geq -2	1,781	199	5-1	1-1, 23-4	749	46	5-1	1-2, 22-2	963	61	5-1	1-1, 23-7	848	64	5-2	0-9, 28-0
PTB & WHZ \geq -2	72	15	16-5	1-5, 155	24	2	10-6	1-2, 88-4	19	2	16	1
Term & WHZ <-2	667	133	16-9	3-4, 78-7	324	42	15-2	3-4, 65-5	392	60	13-6	2-8, 63-6	314	31	12-2	2-0, 70-3
PTB & WHZ <-2	70	19	31-1	3-1, 251	14	3	14	3	11	1
BF & MUAC \geq 125	969	49	3-2	1-4, 7-4	2,428	59	2-7	1-4, 5-0	3,374	51	2-2	1-0, 4-9	322	0	1-9	0-6, 5-9
No BF & MUAC \geq 125	50	3	6-4	1-6, 24-4	435	22	5-5	2-1, 14-0	1,810	70	4-7	2-4, 9-1	1,837	118	4-0	2-0, 8-2
BF & MUAC <125	2,308	361	17-0	8-6, 33-1	1,390	126	19-2	10-6, 34-5	1,268	74	21-7	10-0, 46-4	60	2	24-5	8-2, 71-0
No BF & MUAC <125	264	41	35-6	13-2, 92-2	511	71	40-7	20-1, 80-8	794	105	46-6	22-9, 92-4	131	25	53-2	19-8, 136
NBW & MUAC \geq 125	886	46	2-8	1-5, 5-0	1,875	52	3-0	2-1, 4-3	3,309	80	3-2	2-4, 4-4	1,463	84	3-5	2-2, 5-6
NBW & MUAC \geq 125	45	4	7-7	0-7, 77-2	228	3	5-7	1-5, 21-6	514	4	4-3	1-2, 15-5	82	3	3-2	0-3, 31-1
LBW & MUAC <125	2,110	323	16-4	11-7, 22-9	966	107	20-4	15-3, 27-1	1,110	103	25-4	15-7, 40-8	105	18	31-6	15-0, 65-2
LBW & MUAC <125	240	47	20-9	7-8, 54-7	303	23	23-9	11-8, 47-9	304	18	27-4	9-5, 76-3	23	2	31-3	5-8, 152
Term & MUAC \geq 125	706	49	2-8	1-4, 5-5	827	40	3-0	2-0, 4-6	1,117	70	3-2	2-3, 4-7	1,109	84	3-5	2-0, 6-0
PTB & MUAC \geq 125	26	0	0-4	0-0, 153	24	1	0-9	0-0, 43-2	23	1	2-1	0-1, 29-5	25	2	4-9	0-1, 160
Term & MUAC <125	1,763	293	15-7	10-6, 23-3	250	49	22-2	14-5, 33-9	247	56	31-4	14-8, 65-6	61	16	44-2	14-0, 131
PTB & MUAC <125	140	39	49-1	13-5, 162	24	4	38-5	6-1, 207	10	4	2	0

¹ Risk = predicted absolute risk of dying within 1 month per 1,000 child months modelled using generalized linear mixed effects models accounting for repeated measures in each child, clustering within studies, and time at risk

Abbreviations: BF = breastfeeding; HAZ = height-for-age Z-score; LBW= low birthweight (<2500g); MUAC = mid-upper arm circumference; NBW = normal birthweight (\geq 2500g); PTB = preterm birth (<37 weeks of gestation at birth); WAZ = weight-for-age Z-score; WHZ= weight-for-height Z-score

30
31
32
33
34

Signed data pooling agreement
95 datasets

Excluded using inclusion criteria

- 1 study: no weight data
- 1 study: cross-sectional study

Pooled dataset
33 studies
572 491 observation timepoints
76 375 children

Data cleaning

Rows excluded from dataset

- Re-structuring: 1 228*
- Duplicates: 949
- Sex not reported: 1 047
- Missing age at assessment: 22 342
- Reported age at assessment outside of range 0-60 months: 176

Single values set to missing

- Anthropometry: 54
- Diarrhea: 1
- Fever: 1

Analysis pooled dataset
33 studies
546 459 observation timepoints
75 287 children

GP
23 studies /
49 740 children

Observations per age group (months):

- <6: 142 059
- 6-11: 135 464
- 12-23: 150 232
- 24-59: 35 450

A-S
6 studies /
14 759 children

Observations per age group (months):

- <6: 10 767
- 6-11: 16 734
- 12-23: 23 090
- 24-59: 13 929


I-S
4 studies /
10 788 children

Observations per age group (months):


- <6: 3 625
- 6-11: 4 999
- 12-23: 7 676
- 24-59: 2 434

*Restructuring the format of the dataset from wide to long, and from balanced to unbalanced


A) All children




B) WAZ




C) HAZ




D) WHZ




E) MUAC




F) Breastfeeding



G) Birthweight



H) Preterm birth



medRxiv preprint doi: <https://doi.org/10.1101/2024.07.06.24309988>; this version posted July 7, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

