

## **Stress from Cadaver Dissection Linked to Learning Conditions: Evidence from Sub-Saharan Africa**

**Authors:** Jude Amechi Nnaka<sup>1</sup>, Victor Udochukwu Ezeike<sup>2</sup>, Kristjan Thompson<sup>3</sup>, Izuchukwu Azuka Okafor<sup>1\*</sup>

### **Affiliations**

<sup>1</sup>Department of Anatomy, Faculty of Basic Medical Sciences, College of Health Sciences, Nnamdi Azikiwe University, Nnewi Campus, PMB 5001, Nnewi, Anambra State, Nigeria

<sup>2</sup>Department of Mental Health, University of Abuja Teaching Hospital, Gwagwalada-Zuba, 902101, Gwagwalada, P.M.B. 228, Abuja, Federal Capital Territory, Nigeria

<sup>3</sup>Department of Biomedical Sciences, Mercer University School of Medicine, Savannah, GA, USA

**\*Corresponding author:** Dr. Izuchukwu Azuka Okafor, Department of Anatomy, Faculty of Basic Medical Sciences, College of Health Sciences, Nnamdi Azikiwe University, Nnewi Campus, PMB 5001, Nnewi, Nigeria. E-mail: [okaforizuchukwu33@gmail.com](mailto:okaforizuchukwu33@gmail.com); [iza.okafor@unizik.edu.ng](mailto:iza.okafor@unizik.edu.ng)

## **Abstract**

### **Objective**

This study examined the relationship between students' perceptions of cadaver dissection (CD) and some learning conditions during CD.

### **Methods**

This is a cross-sectional study involving 2968 medical students and graduates exposed to CD at nine countries systematically and proportionally selected from Sub-Saharan African countries based on their ranking on the four World Bank indices of development and education: population, literacy, human capital index (HCI), and human development index (HDI). A self-administered questionnaire was used to collect relevant data on learning conditions during CD experience of the participants, using online channels, including email, Whatsapp, Facebook, Instagram, and Twitter.

### **Results**

About half (48.76%) of the participants perceived CD as stressful, while 51.24% thought otherwise. However, 57% of participants from institutions where a CD session lasts three hours described their experience as stressful, whereas 69.67% of participants whose institution spent one hour in a single CD session agreed that CD is not stressful. Similarly, 60.63% of participants from institutions with a student-donor ratio between 5 to 10 students per cadaver described their experience as 'not stressful'. In comparison, 57.51% of participants from institutions with a student-donor ratio of 10–20 students per donor and 53.80% of participants from institutions with over 20 students per donor described their experience as stressful.

### **Conclusions**

Students from institutions with CD instructors, shorter CD sessions, and a smaller student-donor ratio are more likely to perceive CD as stress-free. Providing more support for students during CD and reducing time for CD may be an antidote to CD-related stress.

**Keywords:** anatomy education; cadaver dissection; learning conditions; stress; teaching method

## **Introduction**

Cadaver dissection (CD) is an age-long anatomy-teaching method that has stood the test of time.<sup>1-</sup>  
<sup>2</sup> Since its first use as a pedagogical tool in the thirteenth century<sup>3</sup>, it has undergone modifications such as the introduction of prosections and plastination<sup>4-5</sup>, with recent technology-driven integration like digital imaging technology, virtual dissection, and 3D printing.<sup>6-7</sup> While these new methods offer advantages such as accessibility, convenience, and the ability to learn without the need for a donor<sup>8-9</sup>, CD is still considered by many modern scholars and educators to be an indispensable tool in medical education, as it offers a thorough understanding of the human body's complexities through an immersive and practical learning experience that is impossible to achieve through virtual simulations alone.<sup>10-11</sup>

CD, despite its benefits and importance in anatomy education, is not without its unique challenges. Among these challenges is the issue of anxiety, mental stress, and emotional distress<sup>12-14</sup> which is considered to be a side effect of working closely with deceased human bodies.<sup>14-15</sup> These challenges are often compounded by the pressure to pass exams and very high academic workloads. Additional stressors due to the scarcity of donors have been shown to contribute to these challenges, particularly in low-income countries, where there may be reduced opportunities for every student to have ample hands-on experience during CD.<sup>14,16</sup>

Medical and allied health science students encounter numerous stressors throughout their training, including those associated with CD. While many students can navigate CD-related stressors effectively, a subset may find the experience highly challenging and overwhelming.<sup>15,17</sup> These variations in students' responses to CD-related stress are influenced by diverse factors, including past traumatic experiences, prior medical exposure, and individual resilience levels.<sup>12,17-20</sup> Studies targeting the reduction and management of CD-related stress have emerged, with authors presenting various recommendations such as providing humanizing information about the donor<sup>1</sup>, carrying out desensitizing pre-CD sessions<sup>19,21-23</sup> and the use of background music during CD.<sup>24</sup>

Considering these known strategies, questions persist on the impact of specific learning conditions on CD-related stress and management. While previous studies have found that the perception of one's learning environment is associated with the academic performance and well-being of medical students<sup>25</sup>, no study has specifically examined the relationship between learning conditions during CD and students' stress levels. Understanding this relationship is crucial to improving student's CD experience. Medical educators can implement strategies to mitigate these stressors and create a more supportive learning environment by identifying the factors that contribute to elevated stress levels during CD. This study investigated the poor learning conditions during CD and examines their relationship with students' perceptions of CD.

## **Methods**

### **Study plan, participant recruitment, and data collection**

This cross-sectional study assessed the relationship between CD learning conditions and students' perception of CD. The study systematically and proportionally targeted students and graduates from Sub-Saharan African countries based on their ranking on the four World Bank indices of

development and education - population, literacy, human capital index (HCI), and human development index (HDI) - to avoid selection bias and to ensure that the countries that have more medical schools, CD-based programs, students, and medical graduates are reached. The country scoring and selection based on the above rankings have been earlier published.<sup>26</sup> After ranking, the countries that scored the highest were selected for inclusion till 20% of countries were included, with at least one country from each Sub-Saharan region. A total of 9 countries were selected – Cameroon, Ghana, Kenya Malawi, Nigeria, South Africa, Uganda, Zambia, and Zimbabwe - with a total of 2968 respondents with a sample size ranging from 150 to 788 in each country. The data for all the countries has been published in an open repository.<sup>26</sup> Only students or graduates who had dissection experience in any institution within the selected countries were included in this study. Students without CD experience were excluded to reduce bias and ensure reliability. This study considered a wide range of students and graduates from different medical science and health fields who completed an online self-administered questionnaire shared through different platforms (e-mails, texts, WhatsApp, Facebook, and Twitter).

### **Study questionnaire and variables**

This study utilized an online self-administered questionnaire (<https://forms.app/judennaka/BM-in-dissection>), with questions focusing on the key objectives of the study. A pilot survey was done with a heterogeneous population of 15 participants, and modifications were made to the questionnaire following the outcome. The final questionnaire draft was validated using internal consistency (Cronbach's alpha of 0.7), test-retest reliability, and inter-rater reliability (Kappa statistic,  $K = 0.9$ ) before commencement of data collection.

### **Statistical Analysis**

The study data were automatically generated from the site where the questionnaire was hosted and were prechecked for errors using Openrefine version 3.6 before transfer to IBM SPSS version 25 for analysis. Descriptive analysis, Chi-squared test, and one-sample non-parametric test were used to determine proportions, association, and one-sample differences respectively. Data analysis was conducted with significance set at  $p < 0.05$ .

## **Results**

### **Stress Perception during Cadaver Dissection**

The perception of stress during CD remains uncertain, as 51% of respondents assert that it is not stressful, while 48.36% agree that it induces stress (Figure 1).

### **Cadaver Dissection Learning Conditions**

The duration of CD sessions in most institutions is predominantly between 2 hours (46.77%) and 3 hours (29.41%), with less than 2% of institutions reporting CD sessions lasting less than 1 hour (Figure 2). Student-donor ratios vary, with 39.12% having 5-10 students, and 29.57% having 10-20 students per donor. About 21% of students reported a ratio exceeding 20 students per donor, while only 8.13% reported a ratio of less than five students per donor (Figure 3).

Regarding supervision, 35.71% of respondents reported always having tutors present during CD, while 24.29% and 24.43% reported having tutors often and sometimes, respectively. Less than 2% reported never having a tutor during CD sessions (Figure 5).

Audiovisual aids were not utilized in the majority (71%) of institutions, with only 26.26% confirming their use during CD (Figure 4). In terms of safety measures, 66.86% of respondents

reported consistently using personal protective equipment (PPE) during dissection, while 9.20% admitted to never using PPE during the process (Figure 6).

### **Relationship Between Learning Conditions and Stress Perception in Cadaver Dissection**

There was also a significant association ( $p < 0.001$ ) between student-donor ratio and stress perception in CD. Respondents in CD sessions with a student-donor ratio of less than 5 students predominantly perceived CD as not stressful, while those with ratios of 10 to 20 students and more than 20 students showed a higher proportion perceiving it as stressful.

The duration of CD sessions showed a statistically significant relationship with the participants' stress perception ( $p < 0.001$ ). Sessions lasting less than 1 hour were predominantly associated with low-stress responses, whereas sessions exceeding 3 hours had a higher proportion of respondents perceiving CD as stressful ( $p < 0.001$ ).

Stress perception in CD exhibited a significant association ( $p < 0.001$ ) with the presence or frequency of tutors. Participants who reported always having tutors during CD were more likely to perceive it as not stressful, whereas those who reported “seldom” or “sometimes” having tutors showed a higher proportion of perceiving it as stressful. The association between stress perception of CD and the utilization of audio-visual aids was found to be significant ( $p < 0.001$ ). Respondents using audio-visual aids during CD were more likely to perceive it as not stressful, while those who did not use such aids showed a higher proportion of perceiving CD as stressful.

The association between stress perception in CD and the frequency of personal protective equipment (PPE) use was significant ( $p < 0.001$ ). Participants reporting always using PPE during dissection were more likely to perceive it as not stressful, while those reporting never or sometimes using PPE showed a higher proportion of perceiving it as stressful.

## Discussion

This study explores students' perceptions of stress during CD and some of the learning factors that could contribute to their stress levels. The result from this study shows that students' views on CD were evenly split between perceiving it as stressful or not stressful. This suggests that stress levels during CD are likely subjective and influenced by both internal and external factors (citation needed). This finding is consistent with prior research, which suggests that responses to stress during CD are influenced by factors such as gender, previous traumatic experiences, and prior medical experiences.<sup>12,17</sup>

Most of the surveyed students reported that CD sessions last between 2 to 3 hours, a duration comparable to findings from studies conducted in Australia<sup>27</sup> and South Korea.<sup>28</sup> This suggests a commonality in the timeframe of CD sessions across different geographical regions, indicating standardized approaches to teaching gross anatomy in Sub-Saharan Africa. However, student-donor ratios in this presented study are high, with nearly 30% having 10-20 students per donor and 21% of students reporting a concerning ratio exceeding 20 students per donor. This distribution is significantly higher than the student-donor ratio of 5 students per donor reported in US and Canadian institutions.<sup>29-31</sup> This reflects the challenges faced by institutions in resource-constrained settings, where large class sizes may be necessitated by limited donor availability and funding constraints. This issue is further compounded by the very low utilization of audiovisual aids during CD in sub-Saharan African institutions in this study (less than 2%). These disparities in access to technology and educational resources, which are more pronounced in Sub-Saharan Africa compared to more economically developed regions<sup>31</sup>, could hinder the enhancement of learning experiences and the development of a visual understanding of anatomical structures among students.



In examining the relationship between learning conditions and students' perceptions of stress during CD, this study found a significant correlation between the duration of CD sessions and students' stress perception. Specifically, nearly 70% students who experienced CD sessions lasting for only one hour perceived it positively (Table 1). On the other hand, 57% of participants from institutions where a CD session lasts three hours described their experience as stressful. This suggests that there may be a threshold for how long students can effectively engage with CD before it becomes overwhelming. Previous studies have shown that not only do long working hours increase the likelihood of occupational stress, but they also have a direct positive effect on mental disturbances like depression.<sup>32-33</sup> Longer CD sessions, combined with the emotional and psychological challenges they present, may contribute to physical and mental fatigue, resulting in increased actual and perceived stress levels among students.<sup>34-35</sup> It could be beneficial for educators to consider breaking up CD sessions into shorter intervals to help students manage their stress levels and maintain focus.

The level of guidance and support offered during CD sessions significantly influences student's perceptions of CD. Those who reported receiving more guidance and support from instructors were less likely to perceive CD as stressful. This highlights the crucial role of active and engaged educators during CD to support students. A study by Manyama and colleagues<sup>36</sup> found that engaged and reciprocal teaching in CD not only improves students' performance but also boosts their confidence and helps them to communicate better. Also, Trigueros and coworkers<sup>37</sup> demonstrated that encouraging autonomy and providing support for students significantly reduces academic stress. These findings suggest that the role of educators in CD goes beyond simply teaching anatomy, but also includes providing emotional support and creating a supportive learning environment for students.

Access to visual aids and videos during CD also reduces the likelihood of perceiving CD as stressful. It is known that visual aids and videos provide a more comprehensive understanding of the CD process, allowing students to visualize and better comprehend the task at hand.<sup>38</sup> Also, the visual aids and videos may serve as a source of reassurance and guidance, helping to alleviate any anxiety or uncertainty that students may experience during CD.<sup>39</sup>

Results from this study show a significant relationship between class size and stress levels during CD. Students in classes with 10 - 20 students or more per donor were more likely to perceive CD as stressful compared to students in classes with less than 10 student-donor ratios. This suggests that larger class sizes may contribute to heightened stress levels during CD, potentially due to factors such as increased competition for access to resources, reduced individualized attention from instructors, and a higher likelihood of feeling overwhelmed in larger groups.<sup>36</sup> While there is yet to be any consensus on the ideal student-donor ratio for CD, it has been shown elsewhere that a team of three to four students per donor is adequate to carry out dissection tasks.<sup>36,40</sup>

Taken together, this study demonstrates that teaching duration, class size, instructional support, and educational resources can shape students' stress experiences during CD. As such, medical educators must continually review their learning conditions and strategies to ensure that CD-related stress is reduced to the barest minimum. This review could include ensuring that the time allocated for a CD session is within 2 to 3 hours, increasing access to tutors during CD, and reducing to student-donor ratio to not more than 10 per donor. By addressing these elements, students can feel more confident and supported throughout the dissection process, ultimately enhancing their educational experience and overall well-being.<sup>30</sup>

This study provides valuable insights into the relationship between learning conditions and students' stress perception during CD; however, there exists some limitations. Firstly, the survey

relied on self-reporting, which introduces the possibility of response bias and inaccuracies in data collection. Additionally, the cross-sectional design of this study restricts the ability to establish causality or temporal relationships between the study variables. Therefore, it is important to interpret the results of this study with caution and consider the possibility of confounding variables such as individual differences in stress coping strategies or the influence of participant's previous experiences in CD on the study outcomes. Future research employing longitudinal designs and diverse participant samples could provide a more comprehensive understanding of the factors influencing students' stress perception during CD.

## **Conclusion**

This study shows that students from institutions with CD instructors, shorter CD sessions, and a smaller student-donor ratio are more likely to perceive CD as stress-free. This suggests that institutions that prioritize creating a comfortable and inclusive learning environment during CD sessions are more likely to have reduced levels of stress among students. Providing more support for students during CD and reducing time for CD may be an antidote to CD-related stress.

## **Source of Funding**

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## **Conflict of Interest**

Authors have no conflict of interest to declare

## **Ethical Approval**

This study obtained ethical clearance from the Nnamdi Azikiwe University Faculty of Basic Medical Science Ethical Committee with the approval number NAU/CHS/NC/FMBS/663. The study also complied with all the ethical guidelines of Helsinki Declaration and guidelines for questionnaire and surveys of the National Health Research Ethics Committee of Nigeria (NHREC) guidelines.

### **Consent**

The online survey used for the study was anonymized and was only accessed by the participants upon submitting an informed consent.

### **Author Contributions**

IAO conceived and designed the study. JAN and IAO collected and analyzed the data. All the authors participated in the data interpretation, writing and revision of the manuscript. All authors have critically reviewed and approved the final draft and are responsible for the content and similarity index of the manuscript.

### **References**

1. Magee R. Art macabre: Resurrectionists and anatomists. *ANZ Journal of Surgery/ANZ Journal of Surgery* 2001;71:377–80. <https://doi.org/10.1046/j.1440-1622.2001.02127.x>.
2. Ghosh, SK. Human cadaveric dissection: a historical account from ancient Greece to the modern era. *Anatomy & cell biology* 2015;48:153–169. <https://doi.org/10.5115/acb.2015.48.3.153>
3. Daynes-Diallo S. A touch of France: Theory and practice - European Renaissance medicine. *Medicographia*, 2011;33:334–343.

4. Riederer BM. Plastination and its importance in teaching anatomy. Critical points for long-term preservation of human tissue. *Journal of Anatomy* 2013;224:309–15. <https://doi.org/10.1111/joa.12056>.
5. Lackey-Cornelison WL, Bauler LD, Smith J. A comparison of the effectiveness of dissection and prosection on short-term anatomic knowledge retention in a reciprocal peer-teaching program. *Advances in physiology education* 2020;44: 239–246. <https://doi.org/10.1152/advan.00031.2020>
6. Azam M, Raja B, Chandra A, Das S, Agarwal A. Anatomage - the virtual dissection tool and its uses: A narrative review. *Journal of Postgraduate Medicine* 2022;68:156. [https://doi.org/10.4103/jpgm.jpgm\\_1210\\_21](https://doi.org/10.4103/jpgm.jpgm_1210_21).
7. Brumpt E, Bertin E, Tatu L, Louvrier A. 3D printing as a pedagogical tool for teaching normal human anatomy: a systematic review. *BMC Medical Education* 2023;23. <https://doi.org/10.1186/s12909-023-04744-w>
8. Kazoka D, Pilmane M, Edelmers E. Facilitating student understanding through incorporating digital images and 3D-printed models in a human anatomy course. *Education Sciences* 2021;11:380. <https://doi.org/10.3390/educsci11080380>
9. Anjankar V, Chavan GN, Wankhede KP, Hajare S. The Scope of Virtual Dissection Modalities in Today’s Technological Era over the Conventional Anatomical Teaching. *Journal of Datta Meghe Institute of Medical Sciences University* 2023;18:559–62. [https://doi.org/10.4103/jdmimsu.jdmimsu\\_280\\_23](https://doi.org/10.4103/jdmimsu.jdmimsu_280_23).
10. Korf H-W, Wicht H, Snipes RL, Timmermans J-P, Paulsen F, Rune G, et al. The dissection course – necessary and indispensable for teaching anatomy to medical students. *Annals of Anatomy* 2008;190:16–22. <https://doi.org/10.1016/j.aanat.2007.10.001>.

11. Bond G, Franchi T. Resuming cadaver dissection during a pandemic. *Medical Education Online* 2020;26. <https://doi.org/10.1080/10872981.2020.1842661>.
12. Chang HJ, Kim HJ, Rhyu IJ, Lee YM, Uhm CS. Emotional experiences of medical students during cadaver dissection and the role of memorial ceremonies: a qualitative study. *BMC Medical Education* 2018;18. <https://doi.org/10.1186/s12909-018-1358-0>.
13. Wisenden PA, Budke KJ, Klemetson CJ, Kurtti TR, Patel CM, Schwantz TL, et al. Emotional response of undergraduates to cadaver dissection. *Clinical Anatomy* 2017;31:224–30. <https://doi.org/10.1002/ca.22992>.
14. Chia TI, Oyeniran OI, Ajagbe AO, Onigbinde OA, Oraebosi MI. The symptoms and stress experienced by medical students in anatomy dissection halls. *Journal of Taibah University Medical Sciences* 2020;15:8–13. <https://doi.org/10.1016/j.jtumed.2020.01.001>.
15. Zubair A, Waheed S, Shuja F. Psychological impact of cadaveric dissection on first-year medical students. *The journal of the Royal College of Physicians of Edinburgh* 2021;51:173–176. <https://doi.org/10.4997/JRCPE.2021.419>
16. Chen D, Zhang Q, Deng J, Cai Y, Huang J, Li F, et al. A shortage of cadavers: The predicament of regional anatomy education in mainland China. *Anatomical Sciences Education* 2018;11:397–402. <https://doi.org/10.1002/ase.1788>.
17. Greene SJ, Rosen L. Tracking medical student emotionality in relation to whole body dissection and donation. *Clinical Anatomy* 2020;34:128–42. <https://doi.org/10.1002/ca.23676>.
18. Dempster M, Black A, McCorry N, Wilson D. Appraisal and Consequences of Cadaver Dissection. *Med Educ Online* 2006;11:4592. <https://doi.org/10.3402/meo.v11i.4592>.

19. Boeckers A, Brinkmann A, Jerg-Bretzke L, Lamp C, Traue HC, Boeckers TM. How can we deal with mental distress in the dissection room?—An evaluation of the need for psychological support. *Annals of Anatomy* 2010;192:366–72. <https://doi.org/10.1016/j.aanat.2010.08.002>.
20. Quince TA, Barclay SIG, Spear M, Parker RA, Wood DF. Student attitudes toward cadaveric dissection at a UK medical school. *Anatomical Sciences Education* 2011;4:200–7. <https://doi.org/10.1002/ase.237>.
21. Arráez-Aybar L, Casado-Morales MI, Castaño-Collado G. Anxiety and dissection of the human cadaver: An unsolvable relationship? *Anatomical Record Part B, New Anatomist* 2004;279B:16–23. <https://doi.org/10.1002/ar.b.20022>.
22. Albabish W, Newton G, Jadeski L. Using a Dissection-based Introductory Laboratory Video to Reduce the Anxiety State of Dissection- and Prosection-based Anatomy Students Prior to their First Cadaver-based Laboratory Experience. *The FASEB Journal* 2017;31. [https://doi.org/10.1096/fasebj.31.1\\_supplement.582.25](https://doi.org/10.1096/fasebj.31.1_supplement.582.25).
23. Biswas R, Bandyopadhyay R. Attitude of first year medical students towards cadaveric dissection: a cross sectional study in a medical college of West Bengal, India. *IMSEAR* 2019. <https://pesquisa.bvsalud.org/portal/resource/pt/sea-201347>.
24. Okafor I, Nnaka J. Background Music: The Missing Stress Antidote in Cadaver Dissection Laboratories. *The Ulutaş Medical Journal* 2020;6:193. <https://doi.org/10.5455/umj.20201004115944>.
25. Chew QH, Holmboe E, Sim K. Learning environment, stress and coping in psychiatry residents within a national training program: a longitudinal study. *Perspectives on Medical Education* 2019;8:160–6. <https://doi.org/10.1007/s40037-019-0512-7>.

26. Okafor I, Nnaka J. Use of background music and cadaver dissection-associated stress in Sub-Saharan Africa. *Figshare* 2023. <https://doi.org/10.6084/m9.figshare.24271675>.
27. Dissabandara LO, Nirthanan SN, Khoo TK, Tedman R. Role of cadaveric dissections in modern medical curricula: a study on student perceptions. *Anatomy & cell biology* 2015; 48, 205–212. <https://doi.org/10.5115/acb.2015.48.3.205>
28. Chung E-K, Yun H, Nam K-I, Cho Y-S, Han E-R. The application of flipped learning to a gross anatomy dissection course. *PloS One* 2024;19:e0298067. <https://doi.org/10.1371/journal.pone.0298067>.
29. Fitzharris TP. Survey of gross anatomy courses in the United States and Canada. *The Anatomical Record* 1998;253:162–6. [https://doi.org/10.1002/\(sici\)1097-0185\(199812\)253:6](https://doi.org/10.1002/(sici)1097-0185(199812)253:6).
30. Precht MC, Bennett CC, Cope JM. Quantifying Cadaver Use in Physician Assistant Anatomy Education. *Journal of Physician Assistant Education* 2017;28:182–8. <https://doi.org/10.1097/jpa.0000000000000171>.
31. Shin M, Prasad A, Sabo G, Macnow ASR, Sheth NP, Cross MB, et al. Anatomy education in US Medical Schools: before, during, and beyond COVID-19. *BMC Medical Education* 2022;22. <https://doi.org/10.1186/s12909-022-03177-1>.
32. Hong Y, Zhang Y, Xue P, Fang X, Zhou L, Wei F, Lou X, Zou H. The Influence of Long Working Hours, Occupational Stress, and Well-Being on Depression Among Couriers in Zhejiang, China. *Front. Psychol* 2022;13:928928. <https://doi.org/10.3389/fpsyg.2022.928928>



33. Bondagji D, Fakeerh M, Alwafi H, Khan AA. The Effects of Long Working Hours on Mental Health Among Resident Physicians in Saudi Arabia. *Psychology Research and Behavior Management* 2022;15:1545–57. <https://doi.org/10.2147/prbm.s370642>.
34. Sándor I, Birkás E, Gyórfy Z. The effects of dissection-room experiences and related coping strategies among Hungarian medical students. *BMC Medical Education* 2015;15. <https://doi.org/10.1186/s12909-015-0355-9>.
35. Romo-Barrientos C, Criado-Álvarez JJ, González-González J, Ubeda-Bañon I, Flores-Cuadrado A, Saiz-Sánchez D, et al. Anxiety levels among health sciences students during their first visit to the dissection room. *BMC Medical Education* 2020;20. <https://doi.org/10.1186/s12909-020-02027-2>.
36. Manyama M, Stafford R, Mazyala E, Lukanima A, Magele N, Kidenya BR, et al. Improving gross anatomy learning using reciprocal peer teaching. *BMC Medical Education* 2016;16. <https://doi.org/10.1186/s12909-016-0617-1>.
37. Trigueros R, Padilla A, Aguilar-Parra JM, Lirola MJ, García-Luengo AV, Rocamora-Pérez P, et al. The Influence of Teachers on Motivation and Academic Stress and Their Effect on the Learning Strategies of University Students. *International Journal of Environmental Research and Public Health* 2020;17:9089. <https://doi.org/10.3390/ijerph17239089>.
38. Nabirye P, Paul LB, Mwaka ES. Attitudes and usage of visual-aids in graduate student learning of gross anatomy at Makerere University. *African Health Sciences* 2023;23:631–7. <https://doi.org/10.4314/ahs.v23i1.67>.

39. Vangala N. Student's preference of various audiovisual aids used in teaching pre- and para-clinical areas of medicine. *Medical Journal of Dr D Y Patil University* 2015;8:739.  
<https://doi.org/10.4103/0975-2870.169912>.
40. Bentley BS, Hill RV. Objective and subjective assessment of reciprocal peer teaching in medical gross anatomy laboratory. *Anatomical Sciences Education* 2009;2:143–9.  
<https://doi.org/10.1002/ase.96>.

## **Figure Legend**

**Figure 1:** Student's stress perception during CD.

**Figure 2:** CD Duration in sub-Saharan African medical schools.

**Figure 3:** Student-donor ratio in sub-Saharan African medical schools.

**Figure 4:** Prevalence of the use of audio-visual aids during CD in sub-Saharan medical schools.

**Figure 5:** Availability of tutors/instructors during CD in sub-Saharan African medical schools.

**Figure 6:** The Use of PPEs by students during CD in sub-Saharan African medical schools.





Figure 1





Figure 2











Figure 3



**Figure 4**



**Figure 5**



**Figure 6**

### Table Legend

**Table 1:** The relationship Between Learning Conditions and Stress Perception in Cadaver Dissection

		<b>Do you think Cadaver dissection is a stressful exercise</b>		<b>Total</b>	<b>P value</b>
		<b>No</b>	<b>Yes</b>		
What is the student-donor ratio based on your experience?	10 to 20 students	397	538	935	
	5 to 10 students	750	487	1237	P< 0.001*
	<5 students	146	111	257	
	>20 students	303	352	655	
	Total	1596	1488	3084	

How long does a single dissection period last in your University/College?	<1 hour	39	21	60	
	>3 hours	168	172	340	
	1 hour	246	107	353	P< 0.001*
	2 hours	780	699	1479	
	3 hours	400	530	930	
	Total	1633	1529	3162	
How often do you have tutors/prosectors/teachers during cadaver dissection?	Always	681	448	1129	
	Never	21	19	40	P< 0.001*
	Often	391	377	768	
	Seldom	141	231	372	
	Sometimes	381	423	804	
	Total	1615	1498	3113	
Do you make use of any audio-visual aid during dissection?	No	1103	1159	2262	P< 0.001*
	Yes	503	328	831	
	Total	1606	1487	3093	
How often do you use personal protective equipment (PPE) during dissection?	Always	1154	960	2114	
	Never	137	154	291	P< 0.001*
	Often	170	227	397	
	Seldom	25	18	43	
	Sometimes	137	143	280	

	Total	1623	1502	3125	
--	-------	------	------	------	--

Data was analyzed using student's Chi-squared test and values are considered significant at

$P < 0.05^*$