

Abstract

The global burden of invasive fungal disease (IFD) is substantial and escalating. Combination antifungal therapy (CAF) may improve patient outcomes by reducing development of resistance, improving drug penetration and rate of fungal clearance, and allowing for lower, less toxic antifungal drug doses; yet, increased cost, antagonism, drug-drug interactions, and toxicity are concerns. Clinical practice guidelines recommend antifungal monotherapy, rather than CAF, for most IFDs due to a lack of comparative randomized clinical trials. An examination of the existing body of CAF research should frame new hypotheses and determine priorities for future CAF clinical trials. We performed a systematic review of CAF clinical studies for invasive candidiasis, cryptococcosis, invasive aspergillosis, and mucormycosis. Additionally, we summarize findings from animal models of CAF and assess laboratory methods available to evaluate CAF efficacy. Future CAF trials should be prioritized according to animal models showing improved survival and observational clinical data supporting efficacy and safety.

Introduction

Continued dismal outcomes for patients with serious fungal infections

The global burden of invasive fungal disease (IFD) is substantial, causing

approximately 1.5 million deaths per year, and escalating due to a growing population of

immunosuppressed individuals, global health events such as the COVID-19 pandemic,

and improvements in molecular diagnostics leading to increased reporting of previously

unrecognized or undiagnosed IFD.(1-6)

IFDs have considerable economic impact in terms of direct medical costs,

productivity loss, and pre-mature deaths.(7) In 2019, the estimated economic burden of

fungal disease in the U.S. was \$11.5 billion, and the mean cost of an inpatient visit

associated with fungal disease diagnosis was more than twice that of the average

inpatient stay. Within the healthcare system, IFDs lead to increased length of stay and

mortality, and inappropriate empirical antimicrobial therapy.(8-10) Disparate mortality

rates are observed across continents and countries. For example, 30-day all-cause

mortality in patients with candidemia is approximately 30% in Europe and the U.S. and

60-72% in South Africa and Brazil.(11)

Potential and challenges of combination antifungal therapy

Given the considerable individual and public health burden of IFDs, emphasis on research pertaining to antifungal therapeutics is essential. Except for cryptococcal meningitis, current clinical practice guidelines recommend antifungal monotherapy for most IFDs; yet, morbidity and mortality remains unacceptably high.(12-19) Combination antifungal therapy (CAF) is one putative strategy to reduce risk and improve outcomes. Potential reasons to pursue CAF for IFDs are: (1) To broaden the spectrum of activity

for empirical therapy, (2) to enhance the rate of killing through antifungal synergy/ increased potency, (3) to delay or prevent development of resistance (4) to reduce antifungal toxicity by using lower drug dosages (5) or to enhance the effectiveness of older antifungals, and (6) to treat breakthrough infections.(20-23)

CAF may be particularly useful in difficult clinical situations, e.g., infections due to highly resistant pathogens or biofilm formation, or in tissues with compromised drug penetration. However, some potentially harmful effects of CAF include reduced fungal killing (antagonism), drug-drug interactions, increased toxicity, and increased cost.

The concept of combining drugs with different mechanisms of action to treat a disease is a longstanding practice supported by rigorous randomized controlled trials (RCTs) for other complex and severe infections including HIV and tuberculosis, as well as oncologic diseases.(24-27) Unfortunately, multiple challenges to conducting large-scale RCTs exist for patients with IFDs including cost, the significant time to conduct the study, particularly for rare fungi (e.g., Mucorales), and difficulty enrolling medically complex and/or critically ill patients.(28) Due to a lack of RCTs comparing CAF to monotherapy, CAF is reserved for salvage therapy or specific severe disease 83 manifestations, e.g., *Candida* endocarditis, in U.S. and international guidelines. (13) To frame new hypotheses and carefully inform clinical trial design, it is important to examine and build upon the existing body of preclinical research and human studies. We perform a systematic review of CAF clinical studies for the most prevalent and 87 dangerous fungal pathogens, Candida spp., Cryptococcus neoformans, Aspergillus spp., and the Mucorales. We further summarize findings from experimental research

- and assess laboratory methods available to evaluate CAF efficacy. Table 1 provides a
- "bird's eye view" qualitative summary of the evidence for CAF.
- Clinical studies of combination antifungal therapy
- Systematic review methods

A systematic review to identify clinical studies of CAF for invasive candidiasis (IC), cryptococcosis, invasive aspergillosis (IA), and invasive mucormycosis was conducted in accordance with the PRISMA guidelines.(29) PubMed was searched for English-language publications using the following terms "antifungal combination therapy" or "combination antifungal therapy" and "candida" or "candidiasis", "cryptococcus" or "cryptococcosis" or "cryptococcal", "aspergillus" or "aspergillosis", and "mucorales" or "mucormycosis" or "zygomycetes" or "zygomycosis" (Appendix). We limited our search to manuscripts published between January 1, 2003 (coinciding with the year of U.S. FDA approval of voriconazole) and February 1, 2023. Reference lists from the included studies were additionally screened to identify potentially relevant evidence. We included studies whose primary aim was to evaluate the CAF efficacy for proven or probable IFD according to consensus definitions; fungal speciation was not required.(30, 31) Papers were excluded if they were literature reviews or meta-analyses, if they did not specify antifungal drug dosages, or if they evaluated combinations including non-systemic antifungal agents, e.g., topical or aerosolized formulations, or non-antifungal chemical compounds.

Systematic review results

Only one RCT of CAF for IC was identified in the systematic review, and the trial was conducted prior to the availability of echinocandins. The trial included 219 non-neutropenic patients who received fluconazole (FLC) 800 mg/d plus placebo vs. FLC plus deoxycholate amphotericin B (dAmB) 0.7 mg/kg/d for candidemia (C. albicans in 62%).(32) There was no difference between groups in the primary analysis of time-to-failure, nor in mortality, although overall success rates were higher in the combination 125 vs. monotherapy arm (69% vs. 56%, respectively, P=0.043).

Further clinical data pertaining to CAF for IC was limited to 20 case reports and small case series, most often describing azole-echinocandin or polyene-echinocandin combinations used as salvage therapy. Four case reports and a series of 13 patients 129 with peritoneal dialysis-associated Candida peritonitis described favorable outcomes in patients treated with combination dAmB or liposomal AmB (L-AmB) plus flucytosine (5- 131 FC), a regimen still recommended for treatment of deep-seated Candida infections, e.g., meningitis and endocarditis, based on limited experimental and clinical data.(13, 33-37)

133 This combination has shown synergy in vitro and is widely used for cryptococcal meningitis (CM).(38) Moreover, excellent 5-FC levels are achieved in cerebrospinal fluid (CSF).(39)

Cryptococcosis

The systematic review identified 12 RCTs and one prospective, non-randomized trial of induction therapy for CM, all in individuals with HIV/AIDS, as well as 3 retrospective studies and a case report in persons with other immunocompromising conditions. There are no prospective trials in persons without HIV, nor has CAF been evaluated for other manifestations of cryptococcosis, e.g., pulmonary disease.

Several trials established that the combinations of AmB plus 5-FC and AmB plus FLC both achieve improved clinical outcomes as compared to AmB alone.(40-42) However, fungicidal activity, as measured by the rate of reduction in CSF yeast CFU from serial quantitative cultures, is greater with AmB plus 5-FC as compared to AmB plus FLC.(40, 41)

Combination 5-FC and FLC is appealing due to its oral availability and low rates of nephrotoxicity, although prolonged 5-FC is associated with leukopenia.(43) In a pivotal trial in Africa (ACTA) including 721 patients, the all-oral induction regimen of FLC 1200 mg/d plus 5-FC 100 mg/kg/d was non-inferior to dAmB plus FLC or 5-FC for 7 or 14 days for the primary endpoint of 2-week mortality.(44) However, faster CSF clearance was observed with AmB plus 5-FC compared to FLC plus 5-FC.

L-AmB was investigated in a landmark trial, Ambition, in 844 individuals with CM in sub-Saharan Africa. A single 10 mg/kg dose of L-AmB plus 14 days of high-dose FLC

and 5-FC was non-inferior to the WHO-recommended 1-week regimen of dAmB plus 5- FC and was associated with fewer adverse events.(45) Following the results of the ACTA and Ambition trials, as well as advocacy efforts, generic 5-FC is more widely available at a reduced cost and utilized in routine care.(46) Three-drug combinations of AmB, 5-FC, and FLC have also been studied. While the triple therapy had the most potent fungicidal effect in a murine model of CM, it was not more efficacious than AmB plus 5-FC in humans.(40, 47) Other triazole antifungal agents have excellent in vitro activity against Cryptococcus spp., and the combination of AmB and voriconazole (VOR) had similar early fungicidal activity as compared to AmB plus 5-FC and AmB plus FLC in a small randomized trial.(48, 49) However, VOR, isavuconazole (ISA), and posaconazole (POS) have more side effects and drug-drug interactions, and higher costs than FLC, thus limiting their utility for a disease that is most prevalent in resource-limited settings. Invasive Aspergillosis Two randomized trials, 10 observational studies, and 33 case reports/series met inclusion criteria in the systematic review of CAF for IA. Prior to widespread use of VOR as standard of care, prospective studies primarily investigated polyene-echinocandin combination. In the randomized, open-label Combistrat trial including 30 patients with hematologic malignancy (HM) and IA, those receiving L-AmB 3 mg/kg/d plus caspofungin (CAS) had higher favorable response rate (67%) and less acute kidney injury than those receiving high-dose L-AmB 10 mg/kg/d alone (27%, P=0.028 for comparison) for primary therapy.(50) However, L-

AmB ≥ 7.5 mg/kg/d plus CAS combination led to poorer clinical outcomes compared to POS suspension for salvage therapy in a single center compassionate use trial.(51) Further clinical trials with polyene-based combination therapy have not been pursued in the past two decades given a lack of compelling preclinical data, high rates of AmB-associated nephrotoxicity, and studies confirming improved IA-related mortality with VOR.(52, 53)

Observational studies of echinocandin-triazole combination therapy for IA have had inconsistent results depending on the indication and study population. Marr and colleagues compared VOR plus CAS to VOR alone in 47 patients with HM and hematopoietic cell transplant (HCT) recipients with IA who had failed initial therapy with AmB. Improved 3-month survival was observed in the combination group compared to monotherapy (HR, 0.42; 95% CI, 0.17–1.1; P=0.048), independent of other variables associated with prognosis.(54) However, in a retrospective study spanning 12 years (1998-2010), VOR plus CAS did not achieve better outcomes than VOR alone, as primary or salvage therapy.(55) In solid organ transplant recipients, compared to a historical cohort receiving L-AmB alone, VOR plus CAS for primary therapy of IA led to similar overall survival at Day 90 (67.5% combination vs. 51% L-AmB monotherapy controls; P=0.117). (56)

To date, one RCT has evaluated echinocandin-triazole combination for IA. This multicenter, multinational study randomized patients with HM or HCT recipients to VOR plus anidulafungin (AFG) versus VOR alone for primary therapy of proven or probable IA.(57) Amongst 277 patients, there was no difference in 6-week mortality (combination: 19.4%, monotherapy: 27.5%; P=0.087). However, in a post-hoc

analysis, only receipt of combination therapy was significantly associated with improved outcome (OR, 10.9; P=0.02).

Kyvernitakis and colleagues performed the largest analysis to date in a retrospective study including 106 HM and HCT patients with mucormycosis at a single U.S. cancer center between 1994-2014.(61) For initial therapy, 44% of patients received monotherapy (L-AmB 87%, POS 13%) and 56% of patients received CAF (L-AmB + echinocandin, 46%; L-AmB + POS, 27%; triple therapy, 27%), highlighting the frequency with which CAF is employed in clinical practice. Using a propensity-score adjusted analysis, they found a similar 6-week survival between patients receiving initial monotherapy vs. combination therapy (56% vs. 60%; P=0.71).

An important feature of the abovementioned studies is the relatively long study period spanning over 10 years, which may introduce biases pertaining to changes in fungal diagnostics, supportive care practices, and availability of specific antifungal agents and formulations. An additional limitation to observational studies of IFD treatment is the use of different dosages of antifungal agents, which may affect efficacy and toxicity.

Animal studies of combination antifungal therapy

In the absence of large-scale RCTs, animal models of IFDs have several strengths. They allow for pharmacokinetics of the administered drugs, tissue burden, and rate of clearance to be assessed. Researchers can integrate host factors and study a range of doses and dose combinations. Finally, resistant species may be studied; whereas, enrolling patients with rare and highly resistant fungal pathogen into

clinical trials is quite challenging. With the implementation of ARRIVE (Animal Research: Reporting In Vivo Experiments) and ARRIVE 2.0 guidelines on the design, conduct, and reporting of animal studies, it is reasonable to conclude that animal studies with IFD and CAF will be able to bridge the quality gap with RCTs.(62-64) Candida spp. Studies of CAF in experimental models of infections due to Candida spp. have yielded mixed results depending on the species, drugs, and methodology. For example, 250 the combination of FLC plus CAS for murine candidemia (C. albicans) yielded no change in tissue fungal burden as compared to FLC monotherapy; whereas POS plus CAS led to improved survival compared to either drug alone.(65, 66)

An in vivo study by Louie and colleagues further illustrates the complexity of antifungal interactions.(67, 68) In a rabbit model of Candida albicans endocarditis and pyelonephritis, both AmB monotherapy and sequential treatment with AmB alone followed by AmB plus FLC rapidly sterilized kidneys and cardiac vegetations. In contrast, simultaneous AmB plus FLC, FLC monotherapy, and FLC followed by AmB were all slower to clear fungi from infected tissues.(68)

Polyene-echinocandin combination is the most frequently studied regimen in the 260 experimental models of IC, particularly due to C. glabrata. Compared to monotherapy, AmB plus CAS or L-AmB plus CAS or micafungin (MFG) led to reduced tissue fungal 262 burden in a murine model of systemic infection due to azole-resistant C. albicans and C. 263 glabrata and in immunosuppressed mice with C. glabrata infection. (69-71)

Cryptococcus spp.

Echinocandins do not have in vitro activity against Mucorales in standard 303 susceptibility tests, however, some agents of mucormycosis, including Rhizopus oryzae, express (1→3)-β-D-glucan synthase, which is the target enzyme for echinocandins.(88) Experimental studies of murine mucormycosis due to R. oryzae have found improved outcomes when combining lipid formulations of Amb plus echinocandins as compared to either agent alone.(89, 90) The benefit noted with the addition of echinocandins may

be in part due to the class' immunomodulatory effect on the activity of phagocytic cells.(91) However, no evidence of synergy nor antagonism was noted for the 310 combination of ISA and MFG in neutropenic mice with pulmonary disease due to R. delemar.(92)

Laboratory methods

There are no standard CLSI or EUCAST methods for in vitro testing of antifungal agents in combinations (93). Our literature analysis revealed numerous publications on laboratory testing of antifungal drugs in combinations against yeasts and molds (Supplementary Fig 1). Only a small selection of published studies included details consistent with the inoculum, inclusion of quality control strains, end-point reading, and interpretations of test results per CLSI or EUCAST methods for testing of molds and yeasts (94-112). Even publications describing combination testing with standard methods originated from single institutions and the reproducibility of the described method at other institutions remains unknown. The development of a consensus method with standardized parameters is imperative for the meaningful testing of antifungal combinations in the clinical laboratory. Towards this end, one of us conducted a series of multi-laboratory studies to define the parameters that would allow 325 reproducible antifungal combination testing. We identified a Candida krusei QC strain, inoculum size, and 100% inhibition end-point as highly reproducible features for 327 combination testing of Candida species and Aspergillus fumigatus among six laboratories (113-115). The result interpretations were consistent among laboratories using summation fractional inhibitory concentration indices (ΣFICI) (113, 116). Recent re-analysis of the data revealed significant pharmacodynamic interactions with Loewe

Future prospects

Table 2 summarizes ongoing clinical trials of CAF, including those incorporating investigational and non-systemic (e.g., aerosolized) antifungal agents. Promising drugs in clinical phases of development with novel mechanisms of action include fosmanogepix, olorofim, and ibrexafungerp.(117) These agents offer hope to expand our antifungal armamentarium, particularly against resistant yeasts and molds, yet will face the same environmental and host pressures for emergence of resistance over time. As such, drug development research should investigate these compounds in combinations and even look towards an antifungal "polypill" that might improve patient adherence and lessen healthcare costs.(23, 118)

While our review focused on a combinatorial strategy involving antifungal drugs, a particularly novel and emerging area of research is the role of cancer immunotherapies as adjunctive therapy for IFDs. Evidence that fungi induce activation of checkpoint pathways has led to preclinical data demonstrating that blockade of the Programmed Cell Death Protein 1 (PD-1) and Cytotoxic T-Lymphocyte Antigen 4 pathways augments antifungal immunity and improves outcomes.(119) Thus far,

clinical data are limited to case reports for invasive mold diseases.(120, 121) Besides providing support towards future interventional trials, such investigations of immunomodulatory therapies to combat IFDs highlight the need to understand the role of host immune responses in the outcomes of IFDs when designing and interpreting experimental and clinical studies of CAF. Conclusion Our systematic review of the literature highlights the paucity of rigorous clinical investigations of CAF for invasive candidiasis and mucormycosis; whereas, multiple RCTs establish the superiority of polyene-5-FC CAF for cryptococcosis and one large RCT of azole-echinocandin CAF for IA suggests a benefit in patients with hematologic malignancy and probable IA. Given the challenges to conducting large prospective RCTs for most IFDs, it is reasonable to prioritize future trials of CAF based upon animal models demonstrating improved survival and observational clinical data supporting efficacy and safety. Recent multi-laboratory investigations demonstrated the feasibility 368 of a standardized method of in vitro testing of antifungal combinations for Candida species that should be further investigated with other pathogen and drug combinations.

- Author contributions. S.J. and V.C: Conceptualization, systematic literature review,
- writing of original draft, and review and editing.
- Conflict of interest: The authors have no conflicts of interest to declare.
- Funding: None.
- Patient consent statement: Not applicable.

383 1. Brown GD, Denning DW, Gow NAR, Levitz SM, Netea MG, White TC. Hidden killers: Human

381 References

382

384 fungal infections. Science Translational Medicine. 2012;4(165):1-10. 385 2. Seagle EE, Williams SL, Chiller TM. Recent Trends in the Epidemiology of Fungal Infections. 386 Infectious Disease Clinics of North America. 2021;35(2):237-60. 387 3. Chowdhary A, Gupta N, Wurster S, Kumar R, Mohabir JT, Tatavarthy S, et al. Multimodal analysis 388 of the COVID-19-associated mucormycosis outbreak in Delhi, India indicates the convergence of clinical 389 and environmental risk factors. Mycoses. 2023;66(6):515-26. 390 4. Steinbrink JM, Hong DK, Bergin SP, Al-Rohil RN, Perfect JR, Maziarz EK. The robust and rapid role 391 of molecular testing in precision fungal diagnostics: A case report. Med Mycol Case Rep. 2020;27:77-80. 392 5. Consortium O, Gabaldon T. Recent trends in molecular diagnostics of yeast infections: from PCR 393 to NGS. FEMS Microbiol Rev. 2019;43(5):517-47. 394 6. White PL, Alanio A, Brown L, Cruciani M, Hagen F, Gorton R, et al. An overview of using fungal 395 DNA for the diagnosis of invasive mycoses. Expert Rev Mol Diagn. 2022;22(2):169-84. 396 7. Benedict K, Whitham HK, Jackson BR. Economic Burden of Fungal Diseases in the United States. 397 Open Forum Infectious Diseases. 2022;9(4):6-9. 398 8. Rayens E, Norris KA. Prevalence and Healthcare Burden of Fungal Infections in the United States, 399 2018. Open Forum Infectious Diseases. 2022;9(1). 400 9. Valerio M, Rodriguez-Gonzalez CG, Munoz P, Caliz B, Sanjurjo M, Bouza E, et al. Evaluation of 401 antifungal use in a tertiary care institution: antifungal stewardship urgently needed. J Antimicrob 402 Chemother. 2014;69(7):1993-9. 403 10. Kara E, Metan G, Bayraktar-Ekincioglu A, Gulmez D, Arikan-Akdagli S, Demirkazik F, et al. 404 Implementation of Pharmacist-Driven Antifungal Stewardship Program in a Tertiary Care Hospital. 405 Antimicrob Agents Chemother. 2021;65(9):e0062921. 406 11. Lamoth F, Lockhart SR, Berkow EL, Calandra T. Changes in the epidemiological landscape of 407 invasive candidiasis. Journal of Antimicrobial Chemotherapy. 2018;73:i4-i13. 408 12. Cornely OA, Bassetti M, Calandra T, Garbino J, Kullberg BJ, Lortholary O, et al. ESCMID guideline 409 for the diagnosis and management of Candida diseases 2012: Non-neutropenic adult patients. Clinical 410 Microbiology and Infection. 2012;18(SUPPL.7):19-37. 411 13. Pappas PG, Kauffman CA, Andes DR, Clancy CJ, Marr KA, Ostrosky-Zeichner L, et al. Clinical 412 Practice Guideline for the Management of Candidiasis: 2016 Update by the Infectious Diseases Society 413 of America. Clinical Infectious Diseases. 2015;62(4):e1-e50. 414 14. Patterson TF, Thompson GR, 3rd, Denning DW, Fishman JA, Hadley S, Herbrecht R, et al. Practice 415 Guidelines for the Diagnosis and Management of Aspergillosis : 2016 Update by the Infectious Diseases 416 Society of America. Clinical Infectious Diseases. 2016;63:1-60. 417 15. Ullmann AJ, Aguado JM, Arikan-Akdagli S, Denning DW, Groll AH, Lagrou K, et al. Diagnosis and 418 management of Aspergillus diseases: executive summary of the 2017 ESCMID-ECMM-ERS guideline. 419 Clinical Microbiology and Infection. 2018;24:e1-e38. 420 16. Cornely OA, Alastruey-Izquierdo A, Arenz D, Chen SCA, Dannaoui E, Hochhegger B, et al. Global 421 guideline for the diagnosis and management of mucormycosis: an initiative of the European 422 Confederation of Medical Mycology in cooperation with the Mycoses Study Group Education and 423 Research Consortium. The Lancet Infectious Diseases. 2019;19(12):e405-e21. 424 17. Organization WH. Guidelines for diagnosing, preventing and managing cryptococcal disease 425 among adults, adolescents and children living with HIV Geneva2022 [updated 27 June 2022. Available 426 from: https://www.who.int/publications/i/item/9789240052178.

427 18. Guidelines for the Prevention and Treatment of Opportunistic Infections in Adults and

428 Adolescents with HIV 2021 [Available from: https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-429 guidelines-adult-and-adolescent-opportunistic-infections/cryptococcosis?view=full.

430 19. Chang CC, Harrison TS, Bicanic TA, Chayakulkeeree M, Sorrell TC, Warris A, et al. Global guideline

431 for the diagnosis and management of cryptococcosis: an initiative of the ECMM and ISHAM in 432 cooperation with the ASM. Lancet Infect Dis. 2024.

433 20. Johnson MD, Perfect JR. Use of Antifungal Combination Therapy: Agents, Order, and Timing. 434 Current Fungal Infection Reports. 2010;4(2):87-95.

435 21. Tyers M, Wright GD. Drug combinations: a strategy to extend the life of antibiotics in the 21st 436 century. Nat Rev Microbiol. 2019;17(3):141-55.

437 22. Coates ARM, Hu Y, Holt J, Yeh P. Antibiotic combination therapy against resistant bacterial 438 infections: synergy, rejuvenation and resistance reduction. Expert Rev Anti Infect Ther. 2020;18(1):5-15.

439 23. Wake RM, Allebone-Salt PE, John LLH, Caswall BA, Govender NP, Ben-Ami R, et al. Optimising

440 the Treatment of Invasive Candidiasis – A Case for Combination Therapy. Open Forum Infectious 441 Diseases. 2024.

442 24. Torres RA, Barr M. Impact of combination therapy for HIV infection on inpatient census. N Engl J 443 Med. 1997;336(21):1531-2.

444 25. Collaboration H-C, Ray M, Logan R, Sterne JA, Hernandez-Diaz S, Robins JM, et al. The effect of 445 combined antiretroviral therapy on the overall mortality of HIV-infected individuals. AIDS.

446 2010;24(1):123-37.

447 26. Kerantzas CA, Jacobs WR, Jr. Origins of Combination Therapy for Tuberculosis: Lessons for 448 Future Antimicrobial Development and Application. mBio. 2017;8(2).

449 27. Marshall HT, Djamgoz MBA. Immuno-Oncology: Emerging Targets and Combination Therapies. 450 Front Oncol. 2018;8:315.

451 28. Spellberg B, Ibrahim A, Roilides E, Lewis RE, Lortholary O, Petrikkos G, et al. Combination

452 therapy for mucormycosis: Why, what, and how? Clinical Infectious Diseases. 2012;54(SUPPL. 1):73-8.

453 29. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020

454 statement: An updated guideline for reporting systematic reviews. J Clin Epidemiol. 2021;134:178-89. 455 30. Donnelly JP, Chen SC, Kauffman CA, Steinbach WJ, Baddley JW, Verweij PE, et al. Revision and

456 Update of the Consensus Definitions of Invasive Fungal Disease From the European Organization for

457 Research and Treatment of Cancer and the Mycoses Study Group Education and Research Consortium. 458 Clin Infect Dis. 2020;71(6):1367-76.

459 31. De Pauw B, Walsh TJ, Donnelly JP, Stevens DA, Edwards JE, Calandra T, et al. Revised definitions 460 of invasive fungal disease from the European Organization for Research and Treatment of

461 Cancer/Invasive Fungal Infections Cooperative Group and the National Institute of Allergy and Infectious

462 Diseases Mycoses Study Group (EORTC/MSG) Consensus Group. Clin Infect Dis. 2008;46(12):1813-21. 463 32. Rex JH, Pappas PG, Karchmer AW, Sobel J, Edwards JE, Hadley S, et al. A randomized and blinded

464 multicenter trial of high-dose fluconazole plus placebo versus fluconazole plus amphotericin B as 465 therapy for candidemia and its consequences in nonneutropenic subjects. Clin Infect Dis.

466 2003;36(10):1221-8.

467 33. Abele-Horn M, Kopp A, Sternberg U, Ohly A, Dauber A, Russwurm W, et al. A randomized study 468 comparing fluconazole with amphotericin B/5-flucytosine for the treatment of systemic Candida 469 infections in intensive care patients. Infection. 1996;24(6):426-32.

470 34. Pelletier R, Alarie I, Lagace R, Walsh TJ. Emergence of disseminated candidiasis caused by

471 Candida krusei during treatment with caspofungin: case report and review of literature. Med Mycol.

472 2005;43(6):559-64.

473 35. Wong PN, Lo KY, Tong GM, Chan SF, Lo MW, Mak SK, et al. Treatment of fungal peritonitis with a 474 combination of intravenous amphotericin B and oral flucytosine, and delayed catheter replacement in 475 continuous ambulatory peritoneal dialysis. Perit Dial Int. 2008;28(2):155-62. 476 36. Ahuja T, Fong K, Louie E. Combination antifungal therapy for treatment of Candida parapsilosis 477 prosthetic valve endocarditis and utility of T2Candida Panel(R): A case series. IDCases. 2019;15:e00525. 478 37. Wee LE, Wong CSL, Tan AL, Oh HM. Negative cerebrospinal fluid beta-d-glucan levels as an 479 indicator for treatment cessation ahead of biochemical resolution: A case report of Candida glabrata 480 meningitis. Med Mycol Case Rep. 2021;32:47-9. 481 38. O'Brien B, Chaturvedi S, Chaturvedi V. In Vitro Evaluation of Antifungal Drug Combinations 482 against Multidrug-Resistant Candida auris Isolates from New York Outbreak. Antimicrobial Agents and 483 Chemotherapy. 2020;64(4). 484 39. Kethireddy S, Andes D. CNS pharmacokinetics of antifungal agents. Expert Opin Drug Metab 485 Toxicol. 2007;3(4):573-81. 486 40. Brouwer AE, Rajanuwong A, Chierakul W, Griffin GE, Larsen RA, White NJ, et al. Combination 487 antifungal therapies for HIV-associated cryptococcal meningitis: A randomised trial. Lancet. 488 2004;363(9423):1764-7. 489 41. Day JN, Chau TTH, Wolbers M, Mai PP, Dung NT, Mai NH, et al. Combination Antifungal Therapy 490 for Cryptococcal Meningitis. New England Journal of Medicine. 2013;368(14):1291-302. 491 42. Pappas PG, Chetchotisakd P, Larsen RA, Manosuthi W, Morris MI, Anekthananon T, et al. A 492 phase II randomized trial of amphotericin B alone or combined with fluconazole in the treatment of HIV-493 associated cryptococcal meningitis. Clin Infect Dis. 2009;48(12):1775-83. 494 43. Stamm AM, Diasio RB, Dismukes WE, Shadomy S, Cloud GA, Bowles CA, et al. Toxicity of 495 amphotericin B plus flucytosine in 194 patients with cryptococcal meningitis. Am J Med. 1987;83(2):236- 496 42. 497 44. Molloy SF, Kanyama C, Heyderman RS, Loyse A, Kouanfack C, Chanda D, et al. Antifungal 498 Combinations for Treatment of Cryptococcal Meningitis in Africa. N Engl J Med. 2018;378(11):1004-17. 499 45. Jarvis JN, Lawrence DS, Meya DB, Kagimu E, Kasibante J, Mpoza E, et al. Single-Dose Liposomal 500 Amphotericin B Treatment for Cryptococcal Meningitis. N Engl J Med. 2022;386(12):1109-20. 501 46. Hurt WJ, Harrison TS, Molloy SF, Bicanic TA. Combination Therapy for HIV-Associated 502 Cryptococcal Meningitis-A Success Story. J Fungi (Basel). 2021;7(12). 503 47. Diamond DM, Bauer M, Daniel BE, Leal MA, Johnson D, Williams BK, et al. Amphotericin B 504 colloidal dispersion combined with flucytosine with or without fluconazole for treatment of murine 505 cryptococcal meningitis. Antimicrob Agents Chemother. 1998;42(3):528-33. 506 48. Thompson GR, 3rd, Wiederhold NP, Fothergill AW, Vallor AC, Wickes BL, Patterson TF. 507 Antifungal susceptibilities among different serotypes of Cryptococcus gattii and Cryptococcus 508 neoformans. Antimicrob Agents Chemother. 2009;53(1):309-11. 509 49. Loyse A, Wilson D, Meintjes G, Jarvis JN, Bicanic T, Bishop L, et al. Comparison of the early 510 fungicidal activity of high-dose fluconazole, voriconazole, and flucytosine as second-line drugs given in 511 combination with amphotericin B for the treatment of HIV-associated cryptococcal meningitis. Clin 512 Infect Dis. 2012;54(1):121-8. 513 50. Caillot D, Thiébaut A, Herbrecht R, de Botton S, Pigneux A, Bernard F, et al. Liposomal 514 amphotericin B in combination with caspofungin for invasive aspergillosis in patients with hematologic 515 malignancies: a randomized pilot study (Combistrat trial). Cancer. 2007;110(12):2740-6. 516 51. Raad, II, Hanna HA, Boktour M, Jiang Y, Torres HA, Afif C, et al. Novel antifungal agents as 517 salvage therapy for invasive aspergillosis in patients with hematologic malignancies: posaconazole 518 compared with high-dose lipid formulations of amphotericin B alone or in combination with 519 caspofungin. Leukemia. 2008;22(3):496-503.

520 52. Upton A, Kirby KA, Carpenter P, Boeckh M, Marr KA. Invasive aspergillosis following 521 hematopoietic cell transplantation: outcomes and prognostic factors associated with mortality. Clin 522 Infect Dis. 2007;44(4):531-40. 523 53. Herbrecht R, Denning DW, Patterson TF, Bennett JE, Greene RE, Oestmann JW, et al. 524 Voriconazole versus amphotericin B for primary therapy of invasive aspergillosis. N Engl J Med. 525 2002;347(6):408-15. 526 54. Marr KA, Boeckh M, Carter RA, Hyung WK, Corey L. Combination antifungal therapy for invasive 527 aspergillosis. Clinical Infectious Diseases. 2004;39(6):797-802. 528 55. Raad, II, Zakhem AE, Helou GE, Jiang Y, Kontoyiannis DP, Hachem R. Clinical experience of the 529 use of voriconazole, caspofungin or the combination in primary and salvage therapy of invasive 530 aspergillosis in haematological malignancies. Int J Antimicrob Agents. 2015;45(3):283-8. 531 56. Singh N, Limaye AP, Forrest G, Safdar N, Muñoz P, Pursell K, et al. Combination of voriconazole 532 and caspofungin as primary therapy for invasive aspergillosis in solid organ transplant recipients: a 533 prospective, multicenter, observational study. Transplantation. 2006;81(3):320-6. 534 57. Marr KA, Schlamm HT, Herbrecht R, Rottinghaus ST, Bow EJ, Cornely OA, et al. Combination 535 antifungal therapy for invasive aspergillosis: a randomized trial. Ann Intern Med. 2015;162(2):81-9. 536 58. Kontoyiannis DP, Boktour M, Hanna H, Torres HA, Hachem R, Raad, II. Itraconazole added to a 537 lipid formulation of amphotericin B does not improve outcome of primary treatment of invasive 538 aspergillosis. Cancer. 2005;103(11):2334-7. 539 59. Rojas R, Molina JR, Jarque I, Montes C, Serrano J, Sanz J, et al. Outcome of Antifungal 540 Combination Therapy for Invasive Mold Infections in Hematological Patients is Independent of the 541 Chosen Combination. Mediterr J Hematol Infect Dis. 2012;4(1):e2012011. 542 60. Reed C, Bryant R, Ibrahim AS, Edwards J, Jr., Filler SG, Goldberg R, et al. Combination polyene-543 caspofungin treatment of rhino-orbital-cerebral mucormycosis. Clin Infect Dis. 2008;47(3):364-71. 544 61. Kyvernitakis A, Torres HA, Jiang Y, Chamilos G, Lewis RE, Kontoyiannis DP. Initial use of 545 combination treatment does not impact survival of 106 patients with haematologic malignancies and 546 mucormycosis: a propensity score analysis. Clinical Microbiology and Infection. 2016;22(9):811.e1-.e8. 547 62. Kilkenny C, Browne WJ, Cuthill IC, Emerson M, Altman DG. Improving bioscience research 548 reporting: the ARRIVE guidelines for reporting animal research. PLoS Biol. 2010;8(6):e1000412. 549 63. Percie du Sert N, Ahluwalia A, Alam S, Avey MT, Baker M, Browne WJ, et al. Reporting animal 550 research: Explanation and elaboration for the ARRIVE guidelines 2.0. PLoS Biol. 2020;18(7):e3000411. 551 64. Percie du Sert N, Hurst V, Ahluwalia A, Alam S, Avey MT, Baker M, et al. The ARRIVE guidelines 552 2.0: Updated guidelines for reporting animal research. J Cereb Blood Flow Metab. 2020;40(9):1769-77. 553 65. Graybill JR, Bocanegra R, Najvar LK, Hernandez S, Larsen RA. Addition of caspofungin to 554 fluconazole does not improve outcome in murine candidiasis. Antimicrob Agents Chemother. 555 2003;47(7):2373-5. 556 66. Chen YL, Lehman VN, Averette AF, Perfect JR, Heitman J. Posaconazole exhibits in vitro and in 557 vivo synergistic antifungal activity with caspofungin or FK506 against Candida albicans. PLoS One. 558 2013;8(3):e57672. 559 67. Fioriti S, Brescini L, Pallotta F, Canovari B, Morroni G, Barchiesi F. Antifungal Combinations 560 against Candida Species: From Bench to Bedside. J Fungi (Basel). 2022;8(10). 561 68. Louie A, Kaw P, Banerjee P, Liu W, Chen G, Miller MH. Impact of the order of initiation of 562 fluconazole and amphotericin B in sequential or combination therapy on killing of Candida albicans in 563 vitro and in a rabbit model of endocarditis and pyelonephritis. Antimicrob Agents Chemother. 564 2001;45(2):485-94. 565 69. Hossain MA, Reyes GH, Long LA, Mukherjee PK, Ghannoum MA. Efficacy of caspofungin 566 combined with amphotericin B against azole-resistant Candida albicans. J Antimicrob Chemother. 567 2003;51(6):1427-9.

568 70. Barchiesi F, Spreghini E, Fothergill AW, Arzeni D, Greganti G, Giannini D, et al. Caspofungin in 569 combination with amphotericin B against Candida glabrata. Antimicrob Agents Chemother. 570 2005;49(6):2546-9. 571 71. Olson JA, Adler-Moore JP, Smith PJ, Proffitt RT. Treatment of Candida glabrata infection in 572 immunosuppressed mice by using a combination of liposomal amphotericin B with caspofungin or 573 micafungin. Antimicrob Agents Chemother. 2005;49(12):4895-902. 574 72. Schwarz P, Dromer F, Lortholary O, Dannaoui E. Efficacy of amphotericin B in combination with 575 flucytosine against flucytosine-susceptible or flucytosine-resistant isolates of Cryptococcus neoformans 576 during disseminated murine cryptococcosis. Antimicrob Agents Chemother. 2006;50(1):113-20. 577 73. Larsen RA, Bauer M, Thomas AM, Graybill JR. Amphotericin B and fluconazole, a potent 578 combination therapy for cryptococcal meningitis. Antimicrob Agents Chemother. 2004;48(3):985-91. 579 74. Barchiesi F, Schimizzi AM, Caselli F, Novelli A, Fallani S, Giannini D, et al. Interactions between 580 triazoles and amphotericin B against Cryptococcus neoformans. Antimicrob Agents Chemother. 581 2000;44(9):2435-41. 582 75. Hossain MA, Mukherjee PK, Reyes G, Long L, Ghannoum MA. Effects of fluconazole singly and in 583 combination with 5-fluorocytosine or amphotericin B in the treatment of cryptococcal 584 meningoencephalitis in an intracranial murine model. J Chemother. 2002;14(4):351-60. 585 76. Silva EG, Paula CR, Dias AL, Chang MR, Ruiz Lda S, Gambale V, et al. Combination efficacy of 586 voriconazole and amphotericin B in the experimental disease in immunodeficient mice caused by 587 fluconazole-resistant Cryptococcus neoformans. Mycopathologia. 2011;171(4):261-6. 588 77. Martin-Vicente A, Capilla J, Guarro J. Synergistic effect of anidulafungin combined with 589 posaconazole in experimental aspergillosis. Med Mycol. 2017;55(4):457-60. 590 78. Petraitis V, Petraitiene R, McCarthy MW, Kovanda LL, Zaw MH, Hussain K, et al. Combination 591 Therapy with Isavuconazole and Micafungin for Treatment of Experimental Invasive Pulmonary 592 Aspergillosis. Antimicrob Agents Chemother. 2017;61(9). 593 79. Petraitis V, Petraitiene R, Hope WW, Meletiadis J, Mickiene D, Hughes JE, et al. Combination 594 Therapy in Treatment of Experimental Pulmonary Aspergillosis: In Vitro and In Vivo Correlations of the 595 Concentration- and Dose- Dependent Interactions between Anidulafungin and Voriconazole by Bliss 596 Independence Drug Interaction Analysis. Antimicrobial agents and chemotherapy. 2009;53(6):2382-91. 597 80. Schaffner A, Frick PG. The effect of ketoconazole on amphotericin B in a model of disseminated 598 aspergillosis. J Infect Dis. 1985;151(5):902-10. 599 81. Martin-Vicente A, Capilla J, Guarro J. In Vivo Synergy of Amphotericin B plus Posaconazole in 600 Murine Aspergillosis. Antimicrob Agents Chemother. 2016;60(1):296-300. 601 82. Clemons KV, Parmar R, Martinez M, Stevens DA. Efficacy of Abelcet alone, or in combination 602 therapy, against experimental central nervous system aspergillosis. J Antimicrob Chemother. 603 2006;58(2):466-9. 604 83. Clemons KV, Stevens DA. Efficacy of micafungin alone or in combination against experimental 605 pulmonary aspergillosis. Med Mycol. 2006;44(1):69-73. 606 84. Dennis CG, Greco WR, Brun Y, Youn R, Slocum HK, Bernacki RJ, et al. Effect of amphotericin B 607 and micafungin combination on survival, histopathology, and fungal burden in experimental aspergillosis 608 in the p47phox-/- mouse model of chronic granulomatous disease. Antimicrob Agents Chemother. 609 2006;50(2):422-7. 610 85. Gebremariam T, Gu Y, Singh S, Kitt TM, Ibrahim AS. Combination treatment of liposomal 611 amphotericin B and isavuconazole is synergistic in treating experimental mucormycosis. J Antimicrob 612 Chemother. 2021;76(10):2636-9. 613 86. Ibrahim AS, Gebremariam T, Schwartz JA, Edwards JE, Jr., Spellberg B. Posaconazole mono- or 614 combination therapy for treatment of murine zygomycosis. Antimicrob Agents Chemother. 615 2009;53(2):772-5.

616 87. Rodríguez MM, Serena C, Mariné M, Pastor FJ, Guarro J. Posaconazole combined with 617 amphotericin B, an effective therapy for a murine disseminated infection caused by Rhizopus oryzae. 618 Antimicrob Agents Chemother. 2008;52(10):3786-8. 619 88. Ibrahim AS, Bowman JC, Avanessian V, Brown K, Spellberg B, Edwards JE, Jr., et al. Caspofungin 620 inhibits Rhizopus oryzae 1,3-beta-D-glucan synthase, lowers burden in brain measured by quantitative 621 PCR, and improves survival at a low but not a high dose during murine disseminated zygomycosis. 622 Antimicrob Agents Chemother. 2005;49(2):721-7. 623 89. Spellberg B, Fu Y, Edwards JE, Jr., Ibrahim AS. Combination therapy with amphotericin B lipid 624 complex and caspofungin acetate of disseminated zygomycosis in diabetic ketoacidotic mice. Antimicrob 625 Agents Chemother. 2005;49(2):830-2. 626 90. Ibrahim AS, Gebremariam T, Fu Y, Edwards JE, Jr., Spellberg B. Combination echinocandin-627 polyene treatment of murine mucormycosis. Antimicrob Agents Chemother. 2008;52(4):1556-8. 628 91. Lamaris GA, Lewis RE, Chamilos G, May GS, Safdar A, Walsh TJ, et al. Caspofungin-mediated 629 beta-glucan unmasking and enhancement of human polymorphonuclear neutrophil activity against 630 Aspergillus and non-Aspergillus hyphae. J Infect Dis. 2008;198(2):186-92. 631 92. Gebremariam T, Wiederhold NP, Alqarihi A, Uppuluri P, Azie N, Edwards JE, Jr., et al. 632 Monotherapy or combination therapy of isavuconazole and micafungin for treating murine 633 mucormycosis. J Antimicrob Chemother. 2017;72(2):462-6. 634 93. Bidaud A-L, Schwarz P, Herbreteau G, Dannaoui E. Techniques for the assessment of in vitro and 635 in vivo antifungal combinations. Journal of Fungi. 2021;7(2):113. 636 94. Meletiadis J, Beredaki MI, Elefanti A, Pournaras S, Muller A. In Vitro-In Vivo Correlation of 637 Posaconazole-Amphotericin B Combination against Candida albicans: In Vitro Interacting Concentrations 638 Are Associated with In Vivo Free Drug Levels. J Fungi (Basel). 2023;9(4). 639 95. John LL, Thomson DD, Bicanic T, Hoenigl M, Brown AJ, Harrison TS, et al. Heightened efficacy of 640 anidulafungin when used in combination with manogepix or 5-flucytosine against Candida auris in vitro. 641 Antimicrob Agents Ch. 2023:e01645-22. 642 96. O'Brien B, Chaturvedi S, Chaturvedi V. In vitro evaluation of antifungal drug combinations 643 against multidrug-resistant Candida auris isolates from New York outbreak. Antimicrobial agents and 644 chemotherapy. 2020;64(4):10.1128/aac. 02195-19. 645 97. Shi C, Liu CL, Liu JY, Wang Y, Li J, Xiang MJ. Anti-Candida Activity of New Azole Derivatives Alone 646 and in Combination with Fluconazole. Mycopathologia. 2015;180(3-4):203-7. 647 98. Planche V, Ducroz S, Alanio A, Bougnoux ME, Lortholary O, Dannaoui E. In vitro combination of 648 anidulafungin and voriconazole against intrinsically azole-susceptible and -resistant Aspergillus spp. 649 Antimicrob Agents Chemother. 2012;56(8):4500-3. 650 99. Venturini TP, Rossato L, Spader TB, Tronco-Alves GR, Azevedo MI, Weiler CB, et al. In vitro 651 synergisms obtained by amphotericin B and voriconazole associated with non-antifungal agents against 652 Fusarium spp. Diagn Microbiol Infect Dis. 2011;71(2):126-30. 653 100. Ben-Ami R, Lewis RE, Kontoyiannis DP. In vitro interactions among echinocandins against 654 Aspergillus fumigatus: lack of concordance among methods. Medical mycology. 2011;49(3):285-8. 655 101. Cuenca-Estrella M, Alastruey-Izquierdo A, Alcazar-Fuoli L, Bernal-Martinez L, Gomez-Lopez A, 656 Buitrago MJ, et al. In vitro activities of 35 double combinations of antifungal agents against 657 Scedosporium apiospermum and Scedosporium prolificans. Antimicrob Agents Ch. 2008;52(3):1136-9. 658 102. Arikan S, Sancak B, Alp S, Hascelik G, McNicholas P. Comparative in vitro activities of 659 posaconazole, voriconazole, itraconazole, and amphotericin B against Aspergillus and Rhizopus, and 660 synergy testing for Rhizopus. Sabouraudia. 2008;46(6):567-73. 661 103. Cuenca-Estrella M, Gomez-Lopez A, Buitrago MJ, Mellado E, Garcia-Effron G, Rodriguez-Tudela 662 JL. In vitro activities of 10 combinations of antifungal agents against the multiresistant pathogen 663 Scopulariopsis brevicaulis. Antimicrob Agents Ch. 2006;50(6):2248-50.

664 104. Philip A, Odabasi Z, Rodriguez J, Paetznick VL, Chen E, Rex JH, et al. In vitro synergy testing of 665 anidulafungin with itraconazole, voriconazole, and amphotericin B against Aspergillus spp. and Fusarium 666 spp. Antimicrob Agents Ch. 2005;49(8):3572-4.

667 105. Lewis RE, Kontoyiannis DP. Micafungin in combination with voriconazole in Aspergillus species: a 668 pharmacodynamic approach for detection of combined antifungal activity in vitro. J Antimicrob 669 Chemoth. 2005;56(5):887-92.

670 106. Cuenca-Estrella M, Gomez-Lopez A, Garcia-Effron G, Alcazar-Fuoli L, Mellado E, Buitrago MJ, et

671 al. Combined activity in vitro of caspofungin, amphotericin B, and azole agents against itraconazole-

672 resistant clinical isolates of Aspergillus fumigatus. Antimicrob Agents Ch. 2005;49(3):1232-5.

- 673 107. Srimuang S, Prariyachatigul C, Chaiprasert A, Rungsipanuratn W, Tanphaichitra D. Antifungal 674 drug combinations for Cryptococcus neoformans and Prototheca spp. J Med Assoc Thai. 2000;83(1):57- 675 60.
- 676 108. Barchiesi F, Di Francesco LF, Compagnucci P, Arzeni D, Giacometti A, Scalise G. In-vitro
- 677 interaction of terbinafine with amphotericin B, fluconazole and itraconazole against clinical isolates of 678 Candida albicans. J Antimicrob Chemother. 1998;41(1):59-65.
- 679 109. Hanson LH, Stevens DA. Comparison of antifungal activity of amphotericin B deoxycholate
- 680 suspension with that of amphotericin B cholesteryl sulfate colloidal dispersion. Antimicrob Agents 681 Chemother. 1992;36(2):486-8.
- 682 110. CLSI. Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Fifth Informational 683 Supplement. CLSI document M100-S25. Wayne, PA: Clinical and Laboratory Standards Institute; 2015. 684 2015.
- 685 111. CLSI. Performance standards for antifungal susceptibility testing of yeasts: CLSI supplement 686 M60. Wayne, PA, USA: CLSI; 2017.
- 687 112. Meletiadis J, Pournaras S, Roilides E, Walsh TJ. Defining fractional inhibitory concentration index

688 cutoffs for additive interactions based on self-drug additive combinations, Monte Carlo simulation 689 analysis, and in vitro-in vivo correlation data for antifungal drug combinations against Aspergillus 690 fumigatus. Antimicrob Agents Chemother. 2010;54(2):602-9.

691 113. Ren P, Luo M, Lin S, Ghannoum MA, Isham N, Diekema DJ, et al. Multilaboratory testing of 692 antifungal drug combinations against Candida species and Aspergillus fumigatus: utility of 100 percent

- 693 inhibition as the endpoint. Antimicrobial Agents and Chemotherapy. 2015;59(3):1759-66.
- 694 114. Chaturvedi V, Ramani R, Andes D, Diekema DJ, Pfaller MA, Ghannoum MA, et al. Multilaboratory 695 testing of two-drug combinations of antifungals against Candida albicans, Candida glabrata, and Candida 696 parapsilosis. Antimicrob Agents Chemother. 2011;55(4):1543-8.
- 697 115. Chaturvedi V, Ramani R, Ghannoum MA, Killian SB, Holliday N, Knapp C, et al. Multilaboratory 698 testing of antifungal combinations against a quality control isolate of Candida krusei. Antimicrob Agents 699 Chemother. 2008;52(4):1500-2.
- 700 116. Meletiadis J, Andes DR, Lockhart SR, Ghannoum MA, Knapp CC, Ostrosky-Zeichner L, et al. 701 Multicenter Collaborative Study of the Interaction of Antifungal Combinations against Candida Spp. by
- 702 Loewe Additivity and Bliss Independence-Based Response Surface Analysis. J Fungi (Basel). 2022;8(9).

703 117. Jacobs SE, Zagaliotis P, Walsh TJ. Novel antifungal agents in clinical trials. F1000Res. 704 2021;10:507.

705 118. Castellano JM, Pocock SJ, Bhatt DL, Quesada AJ, Owen R, Fernandez-Ortiz A, et al. Polypill 706 Strategy in Secondary Cardiovascular Prevention. N Engl J Med. 2022;387(11):967-77.

707 119. Wurster S, Watowich SS, Kontoyiannis DP. Checkpoint inhibitors as immunotherapy for fungal 708 infections: Promises, challenges, and unanswered questions. Front Immunol. 2022;13:1018202.

709 120. Serris A, Ouedrani A, Uhel F, Gazzano M, Bedarida V, Rouzaud C, et al. Case Report: Immune

- 710 Checkpoint Blockade Plus Interferon-Gamma Add-On Antifungal Therapy in the Treatment of Refractory
- 711 Covid-Associated Pulmonary Aspergillosis and Cerebral Mucormycosis. Front Immunol. 2022;13:900522.

- 712 121. Grimaldi D, Pradier O, Hotchkiss RS, Vincent JL. Nivolumab plus interferon-gamma in the
- 713 treatment of intractable mucormycosis. Lancet Infect Dis. 2017;17(1):18.

714

716 Table 1. Bird's eye view: Summary of clinical and experimental evidence supporting 717 combination antifungal therapy for the most prevalent and dangerous fungal pathogens

720 AmB, liposomal amphotericin B; POS, posaconazole; RCT, randomized clinical trial; VOR,

721 voriconazole

722 Table 2: Ongoing clinical trials of combination antifungal therapy for invasive fungal diseases

723 cAmB, encochleated amphotericin B; Inh, inhaled; SOC, standard of care