

1 **EFFECT OF HEALTH EDUCATION IN THE CONTROL OF**
2 **SCHISTOSOMIASIS IN DASS EMIRATE COUNCIL OF BAUCHI**
3 **STATE, NIGERIA: AN INTERVENTION STUDY**

4 ^aSunday Charles Adeyemo, ^bGbadebo Jimoh Oyedeji, ^cJames Atolagbe, ^cOladunni Opeyemi,
5 ^dSunday Olanrewaju, ^eCalistus Akinleye, ^eFunsho Olagunju ^fEniola Dorcas Olabode,
6 ^gAbdulwaris Salisu Maleka

7 ^aInstitut Superieur de Sante, Niamey, Niger Republic

8 charlespatho@gmail.com

9 ^bDepartment of Medical Microbiology and Parasitology, LAUTECH, Osogbo, Nigeria

10 oyedejigbadebo@gmail.com

11 ^cPublic Health Department, Adeleke University, Ede, Nigeria

12 jatolagbe@yahoo.com

13 opeyemi.oladunni@adelekeuniversity.edu.ng

14 ^dCommunity Medicine Department, Osun State University, Osogbo, Nigeria

15 sunday.olarewaju@uniosun.edu.ng

16 callistus.akinleye@uniosun.edu.ng

17

18 ^eDepartment of Paediatrics Medicine, Osun State University, Osogbo, Nigeria

19 funso.olagunju@uniosun.edu.ng

20 ^fDepartment of Community Health, Obafemi Awolowo University, Ile-Ife, Nigeria

21 oluwafemieniola.ed@gmail.com

22 ^g Disease control and immunization unit, Bauchi State Primary Health Care Development
23 Agency, Dass Emirate, Nigeria

24 abdulwarismaleka1@gmail.com

25 **Corresponding author:**

26 Sunday Charles Adeyemo

27 Institut Superieur de Sante

28 +22796579435, charlespatho@gmail.com

29

30 **Abstract**

31 **Objective:** Schistosomiasis has been recognized by WHO as major contributor to mortality
32 and morbidity particularly in Sub Sahara Africa where it is mostly prevalent. There is a lack
33 of reliable data on the effectiveness of health education interventions in reducing the
34 prevalence of Schistosomiasis in Bauchi State. Hence, the study assessed the prevalence of
35 Schistosomiasis and the knowledge, attitude and practices of community members of Dass
36 Emirate towards the prevention and control of Schistosomiasis before and after health
37 education intervention.

38 **Results:** At pre-intervention, majority of the respondents 234 (66.9%) have been diagnosed,
39 or have family members or community members who have been diagnosed with
40 Schistosomiasis. Ninety-six (27.5%) of respondents have good knowledge about
41 Schistosomiasis. 79 (22.6%) of the respondents strongly agree that they can confidently
42 recognize symptoms of schistosomiasis. Only 91 (26.0%) strongly agreed to taking
43 responsibilities for taking preventive measures. At post-intervention, the prevalence of
44 Schistosomiasis dropped to 55.1%. This was statistically significant at $p = 0.043$. Knowledge
45 about Schistosomiasis increased from 27.5% to 87.0% at post-intervention. This was
46 statistically significant at $p < 0.05$. Regarding attitude and practices, good attitude and
47 practices increased from 59.1% at pre-intervention 71.0% at post-intervention. However, this
48 was not statistically significant ($p > 0.05$).

49 **Keywords:** Schistosomiasis, infected water, eggs, health, education, knowledge, prevalence,
50 intervention

51 **Introduction**

52 Schistosomiasis, commonly known as bilharzia, is a parasitic disease caused by
53 trematode flatworms of Schistosoma. It is a major public health concern in many tropical and
54 subtropical regions, affecting millions worldwide. The disease is transmitted through contact
55 with fresh water contaminated with parasitic larvae, which penetrate human skin and mature
56 in the blood vessels, leading to various clinical manifestations. [1] Globally, schistosomiasis
57 is recognized as a significant contributor to morbidity and mortality, especially in sub-
58 Saharan Africa. The World Health Organization (WHO) estimates that over 200 million
59 people are infected with schistosomiasis worldwide, with most cases occurring in Africa. [2]
60 The disease can lead to severe health complications, including liver and spleen enlargement,
61 kidney damage, bladder cancer, and even death if left untreated. [3]

62 In a study on the challenges of controlling schistosomiasis through mass drug
63 administration (MDA) campaigns were highlighted. While MDA has shown short-term
64 benefits in reducing the prevalence and intensity of infection, the long-term impact remains
65 uncertain. [4] The study emphasized the need for comprehensive strategies, including health
66 education and improved sanitation, to achieve sustainable disease control. There is a lack of
67 reliable data on the effectiveness of health education interventions in reducing the prevalence
68 of Schistosomiasis in Bauchi State. This knowledge gap makes it difficult to design effective
69 control programs and allocate resources efficiently. Therefore, this study aims to evaluate the
70 effectiveness of health education as an intervention in controlling Schistosomiasis in Dass
71 Emirate Council of Bauchi State.

72

73 **Methods**

74 The study employed a quasi-experimental design which involved pre-intervention
75 data collection, intervention and post-intervention data collection. The intervention was a
76 health education intervention programme which was designed with training manual adapted
77 from Federal Ministry of Health manual. The training targeted major stakeholders including
78 school pupils, market people, religious leaders, farmers and herders in 10 communities of
79 Dass Emirate of Bauchi where schistosomiasis are mostly prevalent. Self-administered semi-
80 structured questionnaire was used to collect baseline data and post intervention data. The
81 Fisher's formula ($n=z^2pq/d^2$) was used to estimate a total of 450 respondents. Multistage
82 sampling technique was used to recruit respondents. The study population of this study
83 encompassed residents of the Dass Emirate Council. The data collected were analyzed with
84 SPSS software package version 25 and p - value less than 0.05 was taken as statistically
85 significant. Descriptive statistics were done for all variables. Comparison of variables at pre-
86 and post-intervention was done using chi-square as a test of association.

87 **Results**

88 At pre-intervention, Majority of the respondents 234 (66.9%) have been diagnosed, or
89 have family members or community members who have been diagnosed with
90 Schistosomiasis. Ninety-five (27.1%) reported that Schistosomiasis is very common in their
91 community, 131 (37.5%) reported that Schistosomiasis is somewhat common in their
92 community, 43 (12.3%) reported that Schistosomiasis is rare in their community while 81
93 (23.1%) are unsure about the commonness of Schistosomiasis in their community. More than
94 half, (207, 59.1%) reported that someone in their community has been diagnosed with
95 Schistosomiasis.

96 About knowledge of Schistosomiasis at pre-intervention, overall good knowledge was
97 27.5%. Majority (310, 88.6%) of the respondents have heard about Schistosomiasis. Majority
98 of the respondents 200 (64.5%) heard about Schistosomiasis from health care workers. One
99 hundred and twenty-eight (41.3%) of the respondents reported abdominal pain as a common
100 symptom of Schistosomiasis, 36 (11.6%) respondents reported blood in urine or stool as a
101 common symptom of Schistosomiasis, 71 (22.9%) reported fatigue as a common symptom of
102 Schistosomiasis, while 85 (27.4%) respondents reported weight loss as a common symptom
103 of Schistosomiasis. About transmission of Schistosomiasis, 83 (26.8%) reported that it can be
104 transmitted through infected water sources, 140 (45.2%) reported that it can be transmitted
105 through mosquito bites, 75 (24.2%) reported that it can be transmitted through contaminated
106 food while 52 (16.8%) reported that it can be transmitted through human contact. Ninety-one
107 (29.4%) reported that Schistosomiasis is preventable, while 140 (45.2%) reported that it is not
108 preventable and 79 (25.4%) do not know. On the preventive measures of Schistosomiasis, out
109 of 91 that said it can be prevented, 20 (30.0%) reported that it can be prevented by avoiding
110 contact with contaminated water, 31 (34.1%) reported that it can be prevented by boiling or
111 treating water before use, 29 (31.9%) reported that it can be prevented by proper sanitation
112 and hygiene practices while 10 (11.0%) reported that it can be prevented by use of
113 insecticide-treated nets.

114 About attitude and practices towards the prevention of schistosomiasis, at pre-
115 intervention, 79 (22.6%) of the respondents strongly agree that they can confidently
116 recognize symptoms of schistosomiasis. Fifty-five (15.7%) strongly agree that there is
117 community engagement in efforts towards prevention of Schistosomiasis. Only 91 (26.0%)
118 strongly agreed to taking responsibilities for taking preventive measures and only 23 (6.6%)
119 strongly agreed that they discuss prevention of Schistosomiasis with their family. However,
120 118 (33.7%) agreed that their actions can reduce the risk of Schistosomiasis. One hundred

121 and twenty (34.3%) avoided contact with contaminated water to prevent and control
122 Schistosomiasis, 51 (14.6%) boiled or treated water to prevent and control Schistosomiasis,
123 129 (36.8%) practiced proper hygiene and sanitation to prevent and control Schistosomiasis
124 while 50 (14.3%) used insect treated nets to prevent and control Schistosomiasis.

125 At post-intervention, the prevalence reduced to 190 (55.1%) and this was statistically
126 significant at $p=0.043$. (Figure 1) Knowledge of schistosomiasis increased to 87.0% at post-
127 intervention and this was statistically significant at $p < 0.001$. (Figure 2) About attitude and
128 practices towards schistosomiasis prevention and control, 126 (36.5%) of the respondents
129 strongly agree that they can confidently recognize symptoms of schistosomiasis. This
130 difference was not statistically significant at $p=0.045$. Eighty-eight (25.5%) strongly agree
131 that there is community engagement in efforts towards prevention of Schistosomiasis.
132 However, this difference was not statistically significant ($p>0.05$). One hundred and eighteen
133 (34.2%) strongly agreed to taking responsibilities for taking preventive measure. The
134 difference is statistically significant at $p = 0.035$. One hundred and fifty-eight (45.8%) agreed
135 that they discuss prevention of Schistosomiasis with their family. This difference was
136 statistically significant at $p=0.022$. One hundred and twenty-four (35.9%) respondents
137 strongly agreed that their actions can reduce the risk of Schistosomiasis. However, this was
138 not statistically significant ($p>0.05$). About actions taken to prevent schistosomiasis, 120
139 (34.8%) reported avoiding contact with contaminated water, 51 (14.8%) reported boiling or
140 treating water before use, 124 (35.9%) reported proper sanitation and hygiene practices and
141 50 (14.5%) reported use of insecticide-treated nets. However, this difference was not
142 statistically significant ($p>0.05$) (Table 1)

143

144

145

146 **Table 1: Comparison of Attitude and Practices towards Schistosomiasis prevention and**
 147 **control among respondents at pre-intervention and post-intervention**

Attitude and practices towards prevention and control of Schistosomiasis	Frequency (%)		Statistics
	Baseline (n=350)	Post-intervention (n=345)	
I can confidently recognize Schistosomiasis symptoms			$\chi^2 = 2.242$ df = 5 p = 0.045*
Strongly Disagree	109 (31.1)	50 (20.4)	
Disagree	71(20.3)	51 (14.8)	
Neutral	55 (15.7)	49 (14.3)	
Agree	36 (10.3)	69 (20.0)	
Strongly agree	79 (22.6)	126 (36.5)	
There is community engagement in prevention efforts			$\chi^2 = 13.698$ df = 5 p = 0.058
Strongly disagree	120 (34.3)	59 (17.1)	
Disagree	91 (26.0)	64 (18.6)	
Neutral	49 (14.0)	48 (13.9)	
Agree	35 (10.0)	86 (24.9)	
Strongly agree	55 (15.7)	88 (25.5)	
I take responsibility for taking preventive measures			$\chi^2 = 15.090$ df = 5 p = 0.035*
Strongly disagree	108 (30.9)	20 (5.8)	
Disagree	55 (15.7)	56 (16.2)	
Neutral	37 (10.5)	47 (13.6)	
Agree	59 (16.9)	104 (30.1)	
Strongly agree	91 (26.0)	118 (34.2)	
I have discussions about prevention with family			$\chi^2 = 12.781$ df = 5 p = 0.022*
Strongly disagree	141 (40.2)	54 (15.7)	
Disagree	70 (20.0)	52 (15.1)	
Neutral	65 (18.6)	17 (4.9)	
Agree	51 (14.6)	158 (45.8)	
Strongly agree	23 (6.6)	64 (18.6)	
I belief that my actions can reduce risk of Schistosomiasis			$\chi^2 = 18.889$ df = 5 p =0.055
Strongly disagree	90 (25.7)	51 (14.8)	
Disagree	45 (12.9)	53 (15.4)	
Neutral	36 (10.3)	40 (11.6)	
Agree	118(33.7)	77 (22.3)	
Strongly agree	61 (17.4)	124 (35.9)	
Actions you/ family members has taken to prevent Schistosomiasis			$\chi^2 = 18.889$ df = 4 p =0.075
Avoiding contact with contaminated water	80 (22.9)	120 (34.8)	
Boiling or treating water before use	81 (23.1)	51 (14.8)	
Proper sanitation and hygiene practices	89 (25.4)	124 (35.9)	
Use of insecticide-treated nets	100 (28.6)	50 (14.5)	

148 χ^2 = Chi-square, df= degree of freedom, p= p-value, *= statistically significant

149 ***Discussion***

150 The prevalence of Schistosomiasis was high at baseline. More than 3 out of 5
151 respondents have been diagnosed with Schistosomiasis. This can be attributed to the fact that
152 two-fifth of the respondents are adolescents and are prone to activities that expose them to the
153 disease. Also, more than four-fifth are rural dwellers which are prone to poor sanitation and
154 hygiene practices. This is similar to a study who reported that children and adolescents are
155 more prone to Schistosomiasis and poor sanitation is also linked with the disease. [5]

156 The Knowledge regarding Schistosomiasis was poor at baseline. Only about one-third
157 had good knowledge about Schistosomiasis. More than four-fifth of the respondents have
158 heard about Schistosomiasis and two-fifth of the respondents reported abdominal pain as a
159 common symptom of Schistosomiasis. More than two-fifth reported that it can be transmitted
160 through mosquito bites. One-third reported that Schistosomiasis is preventable, out of which
161 one-third reported that it can be prevented by proper sanitation and hygiene practices. The
162 poor knowledge can be attributed to low level of education among respondents. This is
163 similar to the findings on the knowledge attitude and practices regarding rural communities in
164 Kano State. [6]

165 The attitude and practices towards Schistosomiasis prevention and control was poor at
166 baseline. One-third of the respondents reported that there was no community engagement in
167 efforts towards prevention of Schistosomiasis. Less than one-third were taking
168 responsibilities for taking preventive measures and only less than one-tenth of the
169 respondents discuss prevention of Schistosomiasis with their family. However, slightly above
170 one-third of the respondents agreed that some actions can reduce the risk of Schistosomiasis.
171 In order to prevent and control schistosomiasis, slightly above one-third avoided contact with

172 contaminated water and practiced proper hygiene and sanitation. This reveals significant
173 shifts in the community's perceptions and reported experiences with the disease.

174 At post-intervention, the prevalence of Schistosomiasis decreased significantly to
175 slightly more than half, which was statistically significant at $p=0.043$. This could be
176 attributed to heightened community awareness of Schistosomiasis as a result of health
177 education, which aligns with research that suggests heightened awareness through health
178 education can increase community concern thereby reducing prevalence of Schistosomiasis.
179 [7]

180 At post-intervention, overall knowledge improved from less than one-third at baseline
181 to fourth-fifth after health education intervention. This aligns with the findings which
182 reported that improved symptom recognition was linked to effective health education,
183 potentially leading to earlier disease detection and treatment. [8] The attitude and practices
184 concerning Schistosomiasis control improved after the health education intervention. One-
185 third of the respondents strongly agreed to taking responsibilities for taking preventive
186 measures while more than two-fifth of the respondents agreed that they discuss prevention of
187 Schistosomiasis with their family. A study by [9] supports this finding, suggesting that
188 enhanced community engagement is often observed following targeted health education
189 campaigns, which empower communities to take an active role in disease prevention. The
190 role of family discussion in promoting health practices has also been well-documented, [7]
191 and the result of this study echo the potential of family-based interventions to amplify health
192 messages within a community. More than one-third of the respondents strongly agreed that
193 some actions can reduce the risk of Schistosomiasis. This is indicative of an internalization of
194 health education messages, where community members not only learn about preventive
195 measures but also believe in their efficacy. This internal belief system is essential for
196 sustained behavior change and has been cited in previous studies as a critical component of

197 health education success. [10] The increase in the items indicating Knowledge, Attitude and
198 Practices may be attributed to an enhancement in disease detection and diagnosis efforts, a
199 finding consistent with studies, that demonstrate improved identification of endemic diseases
200 following concerted health interventions. [11] Similar patterns have also been noted in other
201 literature, where increased awareness following educational interventions correlates with
202 improved health behaviors. [12]

203 An essential aspect of the study's success was its culturally sensitive approach. By
204 tailoring the intervention to fit the local cultural context and incorporating local beliefs and
205 practices, the program achieved higher engagement and effectiveness. This approach is in
206 line with the recommendations of [7], which emphasize the importance of cultural sensitivity
207 in health education interventions.

208 The findings highlight the importance of health education interventions in improving
209 knowledge, attitudes, and practices towards Schistosomiasis control. They underscore the
210 need for addressing perceived barriers and emphasize the role of community engagement in
211 Schistosomiasis prevention and control. The study recommends prioritizing formulation of
212 comprehensive health policies that address Schistosomiasis within the broader context of
213 public health and increased allocation of resources while encouraging partnership from
214 stakeholders and participation from individuals in combatting schistosomiasis.

215 ***Acknowledgements***

216 The authors wish to acknowledge the respondents and the spouses of the authors for their
217 understanding during the course of this study.

218 **References**

- 219 [1] Kokaliaris, C., Garba, A., Matuska, M., Bronzan, R. N., Colley, D. G., Dorkenoo, A. M.,
220 and Vounatsou, P. (2022). Effect of preventive chemotherapy with praziquantel on
221 schistosomiasis among school-aged children in sub-Saharan Africa: a spatiotemporal
222 modelling study. *The Lancet Infectious Diseases*, 22(1), 136-149.
- 223 [2] World Health Organization (2019, April 17). *Schistosomiasis*. Who.int; World Health
224 Organization: WHO. <https://www.who.int/news-room/fact-sheets/detail/schistosomiasis>.
225 [Accessed 25th November 2022](#)
- 226 [3] Castro VN, Rodrigues JL, Cardoso DT, Resende SD, Magalhães FC, Souza DC, Requeijo
227 MH, Negrão-Corrêa D and Geiger SM (2018) Systemic Cytokine and Chemokine Profiles in
228 Individuals with *Schistosoma mansoni* Infection and Low Parasite Burden. *Front. Immunol.*
229 9:2975. doi: 10.3389/fimmu.2018.02975
- 230 [4] Olveda, D. U., McManus, D., and Ross, A. (2016). Mass drug administration and the
231 global control of schistosomiasis: successes, limitations and clinical outcomes. *Clinical*
232 *Microbiology and Infection*, 22(6), 499-507.
- 233 [5] Karunamoorthi, K., Almalki, M.J. and Ghailan, K.Y., 2018. Schistosomiasis: A neglected
234 tropical disease of poverty: A call for intersectoral mitigation strategies for better
235 health. *Journal of Health Research and Reviews (In Developing Countries)*, 5(1), pp.1-12.
- 236 [6] Dawaki, S., Al-Mekhlafi, H. M., Ithoi, I., Ibrahim, J., Abdulsalam, A. M., Ahmed, A.,
237 Sady, H., Nasr, N. A., & Atroosh, W. M. (2015). The Menace of Schistosomiasis in Nigeria:
238 Knowledge, Attitude, and Practices Regarding Schistosomiasis among Rural Communities in
239 Kano State. *PloS One*, 10 (11), e0143667. doi:10.1371/journal.pone.0143667.
- 240 [7] Wang, M., Han, X., Fang, H., Xu, C., Lin, X., Xia, S., Yu, W., He, J., Jiang, S., and Tao,
241 H. (2018). Impact of Health Education on Knowledge and Behaviors toward Infectious

242 Diseases among Students in Gansu Province, China. *BioMed Research International*, 2018,
243 1–12. <https://doi.org/10.1155/2018/6397340>.

244 [8] Sousa Pereira J, Neves H, Lobao C, Goncalves R, Santos M. The effectiveness of
245 education on symptoms recognition in heart failure patients to manage self-care: a systematic
246 review protocol. *Prof Inferm.* 2019 Jan-Mar;72(1):50-54. English. doi:
247 10.7429/pi.2019.721050. PMID: 31162043.

248 [9] Sacolo H, Chimbari M, Kalinda C. Knowledge, attitudes and practices on Schistosomiasis
249 in sub-Saharan Africa: a systematic review. *BMC Infect Dis.* 2018 Jan 18;18(1):46. doi:
250 10.1186/s12879-017-2923-6. PMID: 29347919; PMCID: PMC5773048.

251 [10] Musuva RM, Awiti A, Omedo M, Ogutu M, Secor WE, Montgomery SP, Alaii J,
252 Mwinzi PN. Community knowledge, attitudes and practices on schistosomiasis in western
253 Kenya--the SCORE Project. *Am J Trop Med Hyg.* 2014 Apr;90(4):646-52. doi:
254 10.4269/ajtmh.13-0488. Epub 2014 Feb 17. PMID: 24534810; PMCID: PMC3973508.

255 [11] Kong, S., Qin, Z., Sun, J., Cao, Y., Wang, L., & Guan, W. (2020). The impact of a rural
256 health promotion program on knowledge and practice of preventive schistosomiasis and
257 intestinal parasitic infections in communities at high risk of snail-borne diseases in rural
258 China. *Journal of Infection and Public Health*, 13 (8), 1164–1171.

259 [12] Gazzinelli Maria Flávia, Lucas Lobato, Gisele Andrade, Leonardo Ferreira Matoso,
260 David J. Diemert, Andréa Gazzinelli. Improving the understanding of schistosomiasis among
261 adolescents in endemic areas in Brazil: A comparison of educational methods, Patient
262 Education and Counseling, Volume 99, Issue 10,2016, Pages 1657-1662, ISSN 0738-
263 3991.<https://doi.org/10.1016/j.pec.2016.04.010>.

264 **Limitation**

265 The limitation of the study is the reliance on self-reported data, which may be subject
266 to recall bias and social desirability bias. Future research could consider using objective
267 measures, such as medical records, to validate self-reported prevalence of Schistosomiasis.
268 The duration of the health education was also limited due to the nature of occupation of the
269 respondents who were farmers who needed to go to their various farms and the insecurity
270 issues peculiar with the study location.

271 **Declarations**

272 ***Ethical approval***

273 Ethical approval was obtained from Research Ethics Committee of Ministry of
274 Health, Bauchi State and Research Ethics Committee of Adeleke University, Ede. Informed
275 consent was also obtained from participants after providing them with detailed information
276 about the study's purpose, the procedures involved, the potential risks and benefits, and their
277 rights as participants. Confidentiality was strictly maintained throughout the study. All
278 participant data were anonymized and stored securely, accessible only to the research team.
279 Identifiers were removed or coded to prevent the identification of individual participants in
280 any reports or publications resulting from the research.

281 ***Consent for publication***

282 Not applicable

283 ***Availability of data and material***

284 The data for this study is available on reasonable request from the corresponding author.

285 ***Funding***

286 The study was funded by the authors

287 ***Competing interests***

288 The authors know no competing interest for this study.

289 ***Authors' contribution***

290 Sunday Charles Adeyemo- Study design, data collection and analysis, Gbadebo Jimoh


291 Oyedeji- Study design and training of health educators, James Atolagbe- Supervision,

292 Oladunni Opeyemi- Study design and supervision, Sunday Olanrewaju- Ethical compliance,

293 Calistus Akinleye- Study design, Funsho Olagunju- Data analysis, Eniola Dorcas Olabode-

294 Data Analysis and manuscript, Abdulwaris Salisu Maleka- Data collection

Comparison between pre-intervention and post-intervention knowledge about Schistosomiasis prevention and control



Comparison of the prevalence of Schistosomiasis before and after intervention

