

1 **Title:** Topical Probiotics Decrease the Severity of Atopic Dermatitis. A Systematic Review and
2 Meta-Analysis of Double-Blind, Randomized, Placebo Control Trials

3

4 **Elliot Flint¹, Nabeel Ahmad MEd¹, Kevin Rowland PhD¹, Charles Hildebolt PhD^{2*},**
5 **David Raskin PhD¹**

6 ¹**Tilman J Fertitta Family College of Medicine, University of Houston, Houston, TX, and**

7 ²**Department of Anthropology, Washington University in St. Louis, St. Louis, WI**

8

9 *Corresponding author:* David Raskin, Ph.D. Tilman Fertitta Family College of Medicine,
10 University of Houston, 5055 Medical Circle, Houston, TX 77204-6064, USA Email:
11 [draskin@central.uh.edu]

12 **ABSTRACT**

13 Atopic Dermatitis (AD) is a chronic skin disease that commonly appears during childhood
14 but can present at any age. There are many reports showing that probiotics relieve AD
15 symptoms in children. This systematic review and meta-analysis sought to determine the
16 efficacy of topical probiotic treatment for AD in adult populations. A database search was
17 conducted of peer-reviewed, double-blind clinical trials, and studies underwent a systematic
18 exclusion and inclusion process, yielding four that met the criteria. Disease severity, as
19 measured by a standardized scoring tool (SCORAD), was collected, and compared to placebo
20 at two-week and four-week time points. All studies showed improvement in SCORAD in the
21 treatment groups compared to baseline at all time points. Two showed significant decreases
22 in SCORAD after two weeks of treatment, and three studies showed long-lasting
23 improvement after four weeks of treatment. Interestingly, while each study showed a
24 reduction in severity of AD at the two- and four-week time points, a pooled meta-analysis did
25 not show a statistically significant difference between treatment and control at four weeks of
26 treatment. Clinically, there may be benefits to topical probiotic usage as evidenced by the
27 individual studies, more studies need to be performed including adults to show statistical
28 significance.

29

30 INTRODUCTION

31 Atopic dermatitis (AD) is an inflammatory skin disorder that often appears in early
32 childhood, affecting approximately 8 - 12% of children¹. AD can become a chronic
33 condition, extending into adulthood, with 6 - 9% of U.S. adults affected¹. An isolated cause
34 has not been identified; however, a combination of family history, loss of function in the
35 filaggrin (FLG) protein, which helps maintain the integrity of the skin, and exogenous
36 environmental triggers, such as dust mites, heat, dry or humid climate, may play a role in its
37 development². The skin of AD lesions is characterized by microbial dysbiosis, with a
38 reduction of diversity and overrepresentation of *Staphylococcus aureus*, correlating with
39 increased lesion infection and flare-ups³. The normal microbiomes of the gut and the skin
40 may produce molecules that inhibit the growth of *S. aureus*, and they interact with the
41 immune system to downregulate inflammatory responses^{4,5}. Given the relationship between
42 the skin microbiome and AD, investigations into the use of adjuvant probiotics are warranted.
43 Probiotics are living organisms that provide health benefits when consumed or applied to the
44 body. Several systematic reviews have studied the effects of oral probiotics and their effect
45 on decreasing AD severity in adult and pediatric populations. Such reviews explored various
46 oral probiotic strains of Bifidobacterium, Lactobacillus, and/or Streptococcus with most trials
47 concluding a reduction in the Scoring Atopic Dermatitis (SCORAD) index⁶⁻¹².

48 There is a growing body of research investigating the effectiveness of probiotics on reducing
49 the severity of AD. Most of these studies use oral probiotics and many test effectiveness only
50 in children. There have been fewer studies using topical probiotics and testing their
51 effectiveness in adults. Thus, a systematic review of randomized control trials of topical
52 probiotics is needed to establish the validity of existing research regarding treating AD. To
53 date, there has been no systematic review of topical probiotics for AD. In this systematic
54 review, we explored the literature on topical probiotics' role in decreasing AD in adult and
55 pediatric populations. We performed a meta-analysis of the reported data to evaluate the
56 effectiveness of the topical probiotics on reducing AD severity.

57 MATERIALS AND METHODS

58 *Study Design*

59

60 Web of Science and PubMed were searched using the terms (atopic dermatitis OR eczema)
61 AND (probiotic* OR synbiotic*). After removing duplicates, the titles and abstracts of 979
62 articles were screened by two reviewers (EF and NA). Reviewers voted to exclude articles or
63 advance them to the next stage of the review using the inclusion and exclusion criteria listed
64 in Table 1. Reviewers' decisions were blinded to one another and organized by the systematic
65 review software Rayyan (Qatar Computing Research Institute). Any article that received that
66 did not receive two votes to "exclude" was advanced to the next stage of the screening
67 process (n=80). Article retrieval and screening of the full text of studies was done by four
68 reviewers (EF, NA, DR, and KR). Each article was reviewed by two reviewers and voted to
69 include or exclude based on the eligibility criteria. Conflicts between reviewer decisions were
70 resolved by group consensus of the four reviewers. At the end of the screening process, only
71 one article met the eligibility criteria for inclusion in the review. To expand the pool of
72 eligible studies, the reviewers eliminated "Study subjects of 18 years or older" as part of the
73 eligibility criteria. Articles were screened using the same process as above, and three
74 additional articles were identified for inclusion in the review. Fig 1 highlights the preferred
75 reporting items for systematic reviews and meta-analysis (PRISMA).

76

77 *Data Extraction*

78 Data was extracted from the included articles by one reviewer (EF). Data extracted from each
79 article included the probiotic(s) used, the study's duration, the days at which treatment effects
80 were measured, and the age of the study subjects. Raw data of SCORAD values were not
81 included in the Axt-Gadermann study, so SCORAD values were obtained using the software
82 NIH Image J to estimate values reliably based on graphs provided by the paper¹³.

83

84 *Statistical Analysis*

85 Meta-analyses were performed for three time periods: (1) baseline, (2) two weeks, and four
86 weeks. The intervals at which outcomes were measured varied slightly between studies. For
87 our analysis, measurements taken on days 0 and 1 were grouped as "baseline." Days 14 and
88 15 were grouped as "two weeks," and days 28, 29, or 30 were grouped as "four weeks." Four
89 studies were used for the baseline assessments, two for the two-week assessments, and three
90 for the four-week assessments. The standardized difference in means was used as the effect
91 size index. Because we assumed that the studies in each analysis represented a random
92 sample from the universe of potential studies, we employed a random-effects model for each
93 analysis, with each analysis being used to make an inference to the universe of potential

94 studies. In support of our using a random-effects model, we provide the following two
95 statements from a book on meta-analysis. “When studies are pulled from the literature, a
96 random-effects model should be used because common sense indicates that the true effect
97 size varies across studies”¹⁴; and “This model assumes that the studies in the analysis are
98 representative of a universe of comparable studies, and that the results of the analysis will be
99 generalized to that universe¹⁴.” In addition, one of our goals was to create a prediction
100 interval for each time period, and the creation of a prediction interval requires the use of a
101 random-effects model. As indicated in another recently published book¹⁵, a prediction
102 interval includes the true effect size for 95% of all populations in the universe.” In this book,
103 the importance of prediction intervals in meta-analyses is stressed,¹⁵ and an article¹⁶ is cited
104 that makes a plea for routinely presenting prediction intervals in meta-analyses; therefore, to
105 help interpret the results of our meta-analysis, a prediction interval (which is based upon the
106 random-effects model) was calculated for each time. “The prediction interval reflects the
107 variation in treatment effects over different settings, including what effect is to be expected in
108 future patients...”¹⁶ “(More accurately, in 95% of all meta-analyses the mean effect size will
109 fall within the confidence interval.)”¹⁴ As explained in this cited book: The 95% confidence
110 interval for a fixed-effect model tells us that the mean prevalence in the set of studies falls
111 within this range, and the 95% confidence interval for a random-effects model tells us that the
112 mean in the universe of comparable populations falls within this range; whereas, the 95%
113 prediction interval tells us that the prevalence in any single population can be as low or as
114 high as the lower limit and the upper limit of the interval¹⁴.

115

116 For our meta-analysis, the standardized mean difference (SMD) for a fixed-effect model was
117 calculated with Hedges g statistic,¹⁷ and the heterogeneity statistic was used to calculate the
118 summary standardized mean difference for a random effects model¹⁸. “If the value 0 is not
119 within the 95% confidence interval (CI), the SMD is statistically significant at the 5% level
120 ($P < 0.05$)¹⁹.” “Cohen's rule of thumb for interpretation of the SMD statistic is: a value of 0.2
121 indicates a small effect, a value of 0.5 indicates a medium effect and a value of 0.8 or larger
122 indicates a large effect¹⁹.”

123

124 Alpha was set at 0.05; however, Amrhein et al. emphasized that the clinical importance of
125 findings, not their statistical significance, be emphasized,²⁰ and for the interpretation of
126 results of our meta-analysis, the clinical importance of findings are emphasized. For our
127 meta-analysis, the standardized mean difference (SMD) for a fixed-effect model was

128 calculated with Hedges g statistic,¹⁷ and the heterogeneity statistic was used to calculate the
129 summary standardized mean difference for a random effects model¹⁸. “If the value 0 is not
130 within the 95% confidence interval (CI), the SMD is statistically significant at the 5% level
131 ($P < 0.05$)²⁰.” “Cohen's rule of thumb for interpretation of the SMD statistic is a value of 0.2
132 indicates a small effect, a value of 0.5 indicates a medium effect and a value of 0.8 or larger
133 indicates a large effect¹⁹.” We performed our meta-analysis with Comprehensive Meta-
134 Analysis Version 4²¹.

135

136

137 **RESULTS**

138 Our search strategy yielded four articles that met the inclusion criteria²²⁻²⁵ (Table 2). Articles
139 tested the use of probiotics in treating atopic dermatitis, although different timelines were
140 used for testing.

141 **SCORAD**

142 SCORAD is a standardized scoring tool to rate the severity of atopic dermatitis by physician
143 and patient ratings of pruritis, dryness, and redness. Higher SCORAD values indicate more
144 severe symptoms, whereas a low SCORAD indicates milder disease severity. In the articles
145 that were included in our analysis, the percent change in SCORAD demonstrated reduced
146 eczema severity with the use of probiotics in all four studies (Fig 2a), whereas placebos only
147 reduced eczema severity in two of the four studies (Fig 2b). The percent change in SCORAD
148 from the probiotic groups was subtracted from the percent change in placebo at each
149 respective time point and graphed; positive values indicate that the probiotics outperformed
150 the placebo at that charted point (Fig 2c). This figure demonstrates how probiotics
151 outperformed placebos across all studies, except for the Butler et al. study at 28 days. It
152 should be noted, however, that the Butler study showed a benefit from the probiotics
153 compared to placebo later in the study at 56 days.

154 *Meta-analysis*

155 Meta-analyses were performed for the placebo and probiotic treatment groups. For the
156 baseline data, the mean effect size is 0.372 with a 95% confidence interval of -0.021 to 0.765
157 (Fig 3a). The mean effect size in the universe of comparable studies could fall anywhere in

158 this interval. The Z-value tests the null hypothesis that the mean effect size is zero. The Z-
159 value is 1.854 with $p = 0.064$. Using a criterion alpha of 0.050, we cannot reject this null
160 hypothesis of no effect. If the true effects are normally distributed (in g units), an estimate of
161 the prediction interval is -0.889 to 1.633 (Fig 3a). The true effect size in 95% of all
162 comparable populations falls in this interval.

163

164 These calculations for baseline data included only four studies. It has been suggested that ten
165 studies are a useful minimum for a meta-analysis¹⁵. Therefore, estimates of heterogeneity
166 based on less than ten studies may not be reliable. A prediction interval and other functions
167 could not be performed for the two-week data because a minimum sample size of 3 is
168 required, and for the two-week assessments, there were only two studies. For the two-week
169 data, the mean effect size is -3.700 with a 95% confidence interval of -10.481 to 3.0815 (Fig
170 3b). The mean effect size in the universe of comparable studies could fall anywhere in this
171 interval. The Z-value is -1.069 with $p = 0.285$, which does not reject the null hypothesis of no
172 effect. For the four-week data, the mean effect size for the three studies is -0.292 with a 95%
173 confidence interval of -0.904 to 0.321 (Fig 3c). The mean effect size in the universe of
174 comparable studies could fall anywhere in this interval. The Z-value is -0.934 with $p = 0.350$.
175 Using a criterion alpha of 0.050, we cannot reject this null hypothesis of no effect. An
176 estimate of the prediction interval is -7.221 to 6.637 (Fig 4b). The true effect size in 95% of
177 all comparable populations falls in this interval. These calculations for four-week data
178 included only three studies; therefore, heterogeneity estimates may not be reliable.

179

180 To summarize the results of our assessments, none of our meta-analyses rejected the null
181 hypothesis of no effect ($p \geq 0.064$). The prediction interval at baseline indicated that with
182 treatment, many future patients would experience benefits; however, at four weeks, the
183 prediction interval indicated that it would be essentially a coin toss as to whether a future
184 patient would experience a benefit. A limitation of our assessments is that our sample sizes
185 for baseline, two weeks, and four weeks were small—respectively, four studies, two studies,
186 and three studies.

187 **DISCUSSION**

188 AD lesions show an increased abundance of *S. aureus* and reduced diversity of the skin
189 microbiome^{3,5}. Due to a lack of long-term, effective AD treatments, there has been interest
190 in modifying the skin microbiome using probiotics. Probiotics are easily assembled,
191 relatively inexpensive, and can be purchased without a prescription, representing a practical
192 avenue of treatment for individuals afflicted with AD. We performed a systematic review and
193 meta-analysis of studies that investigated the use of topically applied probiotics for AD
194 treatment. Overall, only a few studies met our inclusion criteria. The included studies
195 demonstrate a benefit of probiotic application in treating atopic dermatitis as measured by a
196 standard scoring system (SCORAD); however, that benefit varied at the different time points
197 measured.

198 This systematic review had multiple challenges. Initially, we tried to perform a review of AD
199 in adults, but there were not enough studies to carry that out. We modified the criteria to
200 accept studies that included both children and adults. Even so, only four studies met the
201 criteria. These studies used different probiotic formulations and different study periods. The
202 meta-analysis showed a beneficial effect at two weeks but not at four weeks. The one study
203 with enough participants to show the non-effect at four weeks showed a beneficial effect at
204 eight weeks, suggesting that topical probiotics are effective, but effects may be better
205 appreciated in studies of longer durations. Until more clinical trials are performed using
206 standardized treatments and treatment lengths, it will be challenging to determine the
207 effectiveness of topical probiotics. The one study that showed little effect on SCORAD for up
208 to four weeks used a probiotic with only a single organism, *Lactobacillus reuteri*²³. Two of
209 the other studies used *Vitreoscilla filiformis*, and one study used a symbiotic, a mix of several
210 bacterial species, along with a prebiotic (Table 1). Future studies should focus on probiotics
211 that show effectiveness, and while there are only two studies, *V. filiformis* is promising as a
212 probiotic treatment for AD^{24,25}.

213

214 Atopic dermatitis commonly affects infants and young children, but adults are affected as
215 well, with as many as 10% of adults reporting eczema^{26,27}. The gut microbiome has been
216 linked with skin health and may be important in preventing inflammatory processes
217 associated with AD^{28,29}. Probiotics may interfere with *S. aureus* growth through directly
218 killing *S. aureus*, or occupying niches in the skin preventing *S. aureus* from colonizing.
219 Decreasing *S. aureus* growth on the skin could increase diversity and reduce inflammation.
220 Clinical trials using oral probiotics to modify the gut microbiome have been performed, but
221 in adults, oral probiotics have shown mixed effects^{30,31}. Using oral probiotics on those with
222 mature gut microbiomes might not be as effective compared to using them for infants or
223 young children. Our systematic review evaluated topical probiotics to determine whether
224 there was an effective treatment for adults.

225 A recent systematic review analyzed many AD treatments, including oral and topical
226 probiotics, topical emollients, biologics, pharmaceuticals, and many other therapies³⁰. They
227 found that treatments across the spectrum had positive effects on AD outcomes. Oral
228 probiotics generally improved AD severity in children and adults, although some studies in
229 children showed no effect. They found that topical probiotics generally improved AD, but
230 they also included non-clinical trials in their results. They also found that there were
231 significant benefits from biologics. Our systematic review and meta-analysis used only
232 double-blind, randomized clinical trials, and all studies we analyzed included adults. We
233 found that while there was a general reduction in SCORAD from topical probiotic use in
234 three out of four studies, we could not reject the null hypothesis, that the treatment had no
235 effect at four weeks. This is consistent with other research. Greenzaid et al. found that most
236 of the reported studies show a positive effect on SCORAD or other indicators of AD, but
237 there were also studies showing no effect. Many treatment studies show a benefit to AD
238 patients, but the mixed results show that there is a need to determine which treatments work,
239 why they work, and the appropriate length of treatment. For probiotics, there are a variety of
240 different bacteria that are administered, and there are a variety of vehicles to carry the
241 bacteria. Future studies need to determine the most effective treatments.

242 An advantage of topical probiotic treatment is that it is low-cost, easy to administer, and has
243 few adverse effects. Using biologics or pharmaceuticals will be more costly. Oral
244 administration of probiotics carries a greater risk of non-adherence to treatment. All four
245 clinical trials we identified that used topical probiotics in randomized, double-blind studies
246 show positive effects over the length of the trial. Our meta-analysis showed that this
247 treatment is only effective over the first two weeks of treatment. Because endpoints varied
248 between studies, the longest common endpoint, four weeks, was analyzed and showed no
249 significant effect. We recommend that additional clinical trials be performed using the
250 treatments that look successful (break out which probiotics were successful vs not), using
251 more patients over a longer period to determine treatments that will be effective enough for
252 widespread use.

253 **ACKNOWLEDGEMENTS**

254 **Funding sources:** n/a

255 **Conflict of Interest Disclosures:** the authors declare no conflicts interest

256 **IRB approval status:** n/a

257

REFERENCES

1. Chiesa Fuxench ZC, Block JK, Boguniewicz M, Boyle J, Fonacier L, Gelfand JM, et al. Atopic Dermatitis in America Study: A Cross-Sectional Study Examining the Prevalence and Disease Burden of Atopic Dermatitis in the US Adult Population. *J Invest Dermatol* 2019; 139: 583-590.
2. Eichenfield LF, Tom WL, Chamlin SL, Feldman SR, Hanifin JM, Simpson EL, et al. Guidelines of care for the management of atopic dermatitis: section 1. Diagnosis and assessment of atopic dermatitis. *J Am Acad Dermatol* 2014; 70: 338-351.
3. Notay M, Foolad N, Vaughn AR, Sivamani RK. Probiotics, Prebiotics, and Synbiotics for the Treatment and Prevention of Adult Dermatological Diseases. *Am J Clin Dermatol* 2017; 18: 721-732.
4. Nakatsuji T, Chen TH, Narala S, Chun KA, Two AM, Yun T, et al. Antimicrobials from human skin commensal bacteria protect against. *Sci Transl Med* 2017; 9.
5. Totté JE, van der Feltz WT, Hennekam M, van Belkum A, van Zuuren EJ, Pasmans SG. Prevalence and odds of *Staphylococcus aureus* carriage in atopic dermatitis: a systematic review and meta-analysis. *Br J Dermatol* 2016; 175: 687-695.
6. Matsumoto M, Aranami A, Ishige A, Watanabe K, Benno Y. LKM512 yogurt consumption improves the intestinal environment and induces the T-helper type 1 cytokine in adult patients with intractable atopic dermatitis. *Clin Exp Allergy* 2007; 37: 358-370.
7. Matsumoto M, Ebata T, Hirooka J, Hosoya R, Inoue N, Itami S, et al. Antipruritic effects of the probiotic strain LKM512 in adults with atopic dermatitis. *Ann Allergy Asthma Immunol* 2014; 113: 209-216.e207.
8. Drago L, Iemoli E, Rodighiero V, Nicola L, De Vecchi E, Piconi S. Effects of *Lactobacillus salivarius* LS01 (DSM 22775) treatment on adult atopic dermatitis: a randomized placebo-controlled study. *Int J Immunopathol Pharmacol* 2011; 24: 1037-1048.
9. Drago L, De Vecchi E, Toscano M, Vassena C, Altomare G, Pigatto P. Treatment of atopic dermatitis eczema with a high concentration of *Lactobacillus salivarius* LS01 associated with an innovative gelling complex: a pilot study on adults. *J Clin Gastroenterol* 2014; 48 Suppl 1: S47-51.

10. Moroi M, Uchi S, Nakamura K, Sato S, Shimizu N, Fujii M, et al. Beneficial effect of a diet containing heat-killed *Lactobacillus paracasei* K71 on adult type atopic dermatitis. *J Dermatol* 2011; 38: 131-139.
11. Iemoli E, Trabattoni D, Parisotto S, Borgonovo L, Toscano M, Rizzardini G, et al. Probiotics reduce gut microbial translocation and improve adult atopic dermatitis. *J Clin Gastroenterol* 2012; 46 Suppl: S33-40.
12. Roessler A, Friedrich U, Vogelsang H, Bauer A, Kaatz M, Hipler UC, et al. The immune system in healthy adults and patients with atopic dermatitis seems to be affected differently by a probiotic intervention. *Clin Exp Allergy* 2008; 38: 93-102.
13. Rueden CT, Schindelin J, Hiner MC, DeZonia BE, Walter AE, Arena ET, et al. ImageJ2: ImageJ for the next generation of scientific image data. *BMC Bioinformatics* 2017; 18: 529.
14. Borenstein M. *Common Mistakes in Meta-Analysis and How to Avoid Them*: Biostat Inc; 2019.
15. Borenstein M, Hedges V. L, Higgins P.T. J, Rothstein R. H. *Introduction to Meta-Analysis*, 2nd Edition . . 2nd ed. Wiley, 2021: p. 544.
16. IntHout J, Ioannidis JP, Rovers MM, Goeman JJ. Plea for routinely presenting prediction intervals in meta-analysis. *BMJ Open* 2016; 6: e010247.
17. Hedges LV, Olkin I. *Statistical methods for meta-analysis*. Orlando, Florida ; London, England: Academic Press, Inc; 1985.
18. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986; 7: 177-188.
19. Schoonjans F. *Medcalc Manual: Easy-to-Use Statistical Software* 2017.
20. Amrhein V, Greenland S, McShane B. Scientists rise up against statistical significance. *Nature* 2019; 567: 305-307.
21. Borenstein M, Hedges L, Higgins JPT, Rothstein H. *Comprehensive Meta-Analysis Version 4*. Englewood, NJ, 2022.
22. Noll M, Jäger M, Lux L, Buettner C, Axt-Gademmann M. Improvement of Atopic Dermatitis by Synbiotic Baths. *Microorganisms* 2021; 9:527.
23. Butler É, Lundqvist C, Axelsson J. DSM 17938 as a Novel Topical Cosmetic Ingredient: A Proof of Concept Clinical Study in Adults with Atopic Dermatitis. *Microorganisms* 2020; 8:1026.

24. Seité S, Zelenkova H, Martin R. Clinical efficacy of emollients in atopic dermatitis patients - relationship with the skin microbiota modification. *Clin Cosmet Investig Dermatol* 2017; 10: 25-33.
25. Gueniche A, Knaudt B, Schuck E, Volz T, Bastien P, Martin R, et al. Effects of nonpathogenic gram-negative bacterium *Vitreoscilla filiformis* lysate on atopic dermatitis: a prospective, randomized, double-blind, placebo-controlled clinical study. *Br J Dermatol* 2008; 159: 1357-1363.
26. Shaw TE, Currie GP, Koudelka CW, Simpson EL. Eczema prevalence in the United States: data from the 2003 National Survey of Children's Health. *J Invest Dermatol* 2011; 131: 67-73.
27. Silverberg JI, Hanifin JM. Adult eczema prevalence and associations with asthma and other health and demographic factors: a US population-based study. *J Allergy Clin Immunol* 2013; 132: 1132-1138.
28. Lee SY, Lee E, Park YM, Hong SJ. Microbiome in the Gut-Skin Axis in Atopic Dermatitis. *Allergy Asthma Immunol Res* 2018; 10: 354-362.
29. Polkowska-Pruszyńska B, Gerkowicz A, Krasowska D. The gut microbiome alterations in allergic and inflammatory skin diseases - an update. *J Eur Acad Dermatol Venereol* 2020; 34: 455-464.
30. Greenzaid JD, Chan LJ, Chandani BM, Kiritsis NR, Feldman SR. Microbiome modulators for atopic eczema: a systematic review of experimental and investigational therapeutics. *Expert Opin Investig Drugs* 2024; 33: 415-430.
31. Umborowati MA, Damayanti D, Anggraeni S, Endaryanto A, Surono IS, Effendy I, et al. The role of probiotics in the treatment of adult atopic dermatitis: a meta-analysis of randomized controlled trials. *J Health Popul Nutr* 2022; 41: 37.

FIGURE LEGENDS

Fig. 1. PRISMA flow diagram




Fig. 2. Percent Change from baseline in SCORAD vs time. A) Percent change in SCORAD vs time for probiotics for all studies. B) Percent change in SCORAD vs time for placebo for all studies. C) Difference in percent change in SCORAD between groups vs time for all studies.




Fig. 3. Summary values for dermatitis, (A) Baseline values for four studies and the pooled values for the random effects model. (B) Two-week values for two studies and the pooled values for the random effects model. (C) Four-week values for four studies and the pooled values for the random effects model.





Fig. 4. Prediction intervals for dermatitis at baseline and at four weeks—a two-week prediction model could not be created because a minimum sample size of 3 is required, (A) Prediction interval for baseline values. (B) Prediction interval for four-week values.

A)



B)




TABLE LEGENDS

Table I. Inclusion and Exclusion Criteria

Inclusion	Exclusion
Subject age of 18 years or older ¹	Systematic Reviews/Review Articles
Randomized Control Trial	Published Before 2007
Placebo	Receiving Other Treatments Targeted at Symptoms of Atopic Dermatitis
Use of SCORAD	
Human Subjects	
Topical Probiotic(s)	
English Language	

¹This criterion was removed for the second pass of the literature review

Table II. Summary of all included articles

Study	Probiotic	Duration of Study/ Days SCORAD Measured	n / Age Range of Subjects in years (placebo, treatment)	Percent Change from Baseline in SCORAD	Additional Findings
Noll et al, 2021	Mixture (<i>Bifidobacterium breve</i> , <i>Bifidobacterium animalis lactis</i> , <i>Lactobacillus casei</i> , <i>L. gasseri</i> , <i>L. planatarum</i> , <i>L. rhamnosus</i>)	2 weeks/ 0,9,11,14	7,7 / 5+	Synbiotic baths: -40.7%, -49.7% and -56.8% on days 9, 11, and 14, respectively. Placebo: -2.3%, +1.3%, and +7.1% on days 9, 11, and 14, respectively.	Analysis of skin flora showed successful colonization with bacteria from the probiotic treatment, but the abundance of <i>Staphylococcus aureus</i> did not significantly change. Subjects in probiotic group subjectively reported less pruritis and less dryness at the end of study compared to control group.
Butler et al, 2020	<i>Lactobacillus reuteri</i> DSM17938	8 weeks/ 0,28,56	17,17 / 19-66, 19-57	Probiotic ointment: -27.7% and -45.7% on days 28 and 56, respectively. Placebo: -31.9% and -41.0% on days 28 and 56, respectively.	Improvement of skin dryness in the probiotic group was the most pronounced change during the study.
Gueniche et al, 2008	<i>Vitreoscilla filiformis</i>	4 weeks/ 0,15,29	25,26 / 6-70, 6-70	Probiotic cream: -29.7% and -56.1% on days 15 and 29, respectively. Placebo: -7.8% and -16.3% on days 15 and 29, respectively.	Probiotic group showed significantly greater reduction in SCORAD when compared to control at days 15 and 29. Skin barrier function was measured using the biomarker TEWL. There was a significant reduction in TEWL in both groups from their baseline values at the end of the study, but there was no significant difference between the probiotic vs control. The group treated with probiotic had decreased colonization of <i>Staphylococcus aureus</i> and <i>Escherichia coli</i> compared to control at the end of the study, but the finding was not statistically significant.
Seite et al, 2017	<i>Vitreoscilla filiformis</i> (LRP-VFB)	4 weeks/ 1,28	22,24 / 0-63, 0-63	Probiotic cream: -15.5% on day 28. Placebo: +32.0% on day 28.	The probiotic group had significantly increased levels of <i>Xanthomonas</i> at the end of the study compared to control. Colonization with <i>Staphylococcus aureus</i> increased in the control group, but not the treatment group. The difference was not statistically significant.