

Study on the Situation of Growth and Development of Kindergarten Children and the Relationship between Vitamin D and Body Composition, Chengdu, Sichuan Province, China

Wang Yanli¹, Ren Ying¹, Xing Fenglin², Liu Dan¹, Hu Mengjuan¹, Luo Tongyong^{3*}

1: Jinniu maternity and child health hospital of Chengdu

2: Qingyang maternity and child health hospital of Chengdu

3: Sichuan Provincial Women's and Children's Hospital, Affiliated Women's and Children's Hospital of Chengdu Medical College

Wang Yanli, Clinical Nutrition Department, Chengdu Jinniu District Maternal and Child Health Hospital, Chengdu, Sichuan, 610081, China, 404585602@qq.com

Ren Ying, Clinical Nutrition Department, Chengdu Jinniu District Maternal and Child Health Hospital, Chengdu, Sichuan, 610081, China, 923073864@qq.com

Xing Fenglin, Health Care Department, Chengdu Qingyang District Maternal and Child Health Hospital, Chengdu, Sichuan, 610017, China, xingfenglinvv@163.com

Liu Dan, Medical Check-up Center, Chengdu Jinniu District Maternal and Child Health Hospital, Chengdu, Sichuan, China, 120948849@qq.com

Hu Mengjuan, Health Care Department, Chengdu Jinniu District Maternal and Child Health Hospital, Chengdu, Sichuan, 610081, China, 850794839@qq.com

Luo Tongyong, Clinical Nutrition Department, Sichuan Provincial Women's and Children's Hospital & Affiliated Women's and Children's Hospital of Chengdu Medical College, whnic@msn.com

Corresponding author: Luo Tongyong

Vitamin D plays a critical role in bone health, mineral metabolism, and overall well-being, including immune function and muscle strength. Its potential impact on body composition in children, particularly in relation to obesity and undernutrition, is an area of growing interest. However, existing literature presents inconsistent findings regarding the correlation between Vitamin D levels and body composition parameters in pediatric populations. The aim of this study was to explore the relationship between Vitamin D levels and body composition among kindergarten-aged children in Chengdu, Sichuan Province, China, providing insights that could contribute to health promotion strategies for this demographic. A cross-sectional study was conducted among children aged 3 to 6 years across 47 kindergartens in Chengdu. Participants were selected via stratified random sampling to ensure representativeness. Data collection took place during April to June to minimize seasonal effects on Vitamin D levels. Body composition was assessed using bio-electrical impedance analysis (BIA). Vitamin D status was determined through blood tests, and the distribution of Vitamin D levels by sex and age groups was analyzed. The study included 1239 children, with 75.54% having insufficient Vitamin D levels and 24.46% having sufficient levels. There was no statistically significant difference in Vitamin D status between sexes ($\chi^2=0.676$, $p=0.411$) or age groups ($\chi^2=4.700$, $p=0.095$). Correlation analysis revealed that BMI had a weak negative correlation with Vitamin D (-0.028 , $p=0.320$) but no significant correlation with Vitamin A (-0.007 , $p=0.799$). Similar results were observed for other body composition components. This study suggests that a majority of kindergarten children in Chengdu have insufficient Vitamin D levels, regardless of sex or age. While some correlations between Vitamin D levels and body composition were observed, they were weak and not clinically significant. Further research is needed to elucidate the complex interplay between Vitamin D, body composition, and other influencing factors in this age group. Our findings contribute to the ongoing debate about the role of Vitamin D in pediatric health and may guide future interventions aimed at improving Vitamin D status and overall body composition in children.

Keywords: Vitamin D, Body Composition, Kindergarten Children, China, Chengdu, Bio-Electrical Impedance Analysis (BIA)

1. Introduction

Vitamin D is essential for bone health and mineral metabolism, with its roles extending into various aspects of human health, including immune function, muscle strength, and the prevention of chronic diseases. Recent research has highlighted its potential influence on body composition, including body fat percentage, muscle mass, and overall growth in children [1-5]. These attributes of Vitamin D suggest that it could be a pivotal factor in addressing pediatric health issues, such as obesity and under-nutrition, which have become significant public health concerns globally.

Despite these insights, the literature presents mixed findings on the relationship between Vitamin

D levels and body composition in children. Some studies have indicated that sufficient Vitamin D levels in childhood are associated with a lower body fat percentage and higher muscle mass, potentially contributing to healthier bone growth and a reduced risk of developing obesity^[6-8]. Conversely, other studies have not found significant associations, suggesting that the relationship might be influenced by various factors including genetic predispositions, dietary intake, and physical activity levels^[9-10] .

The current study aims to contribute to this ongoing debate by exploring the relationship between Vitamin D levels and body composition among kindergarten children in Chengdu, Sichuan Province, China. By focusing on a demographically specific population, this study seeks to provide insights that could inform both global and local health promotion strategies for children.

2. Materials and Methods

A cross-sectional study design was employed in this study. The retrospective data were collected from the archived physical examination of kindergartens in Jinniu, Chengdu, Sichuan Province, China, targeting kindergarten children aged 3 to 6 years old. The research aimed to investigate the relationship between Vitamin D levels and body composition among this population.

The study received the ethics approval from the ethics committee of Jinniu maternity and child health hospital of Chengdu (2023NT01). Authors cannot access to information that could identify individual participants during or after data collection.

The kindergartens varied in size, socio-economic status of the attending families, and urban or suburban locations. Data collection occurred in three months, from April to June, to minimize the seasonal effect on Vitamin D levels.

Height and weight were measured using standardized procedures. Body composition was assessed using a bio-electrical impedance analysis (BIA) machine. Measurements of Body Mass Index(BMI), Total Body Water (TBW), Intracellular Water (ICW), Extracellular Water (ECW), Protein, Minerals, Soft Lean Mass (SLM), SLM Index (SLMI), Fat Free Mass (FFM), SMM (Skeletal Muscle Mass), PBF (Percent Body Fat), Visceral Fat Area (VFA), Body Cell Mass (BCM), Fat Free Mass Index (FFMI), Fat Mass Index(FMI), Measured Circumference of Abdomen (MCA), Skeletal Muscle Index (SMI), ECW/TBW, TBW/FFM were listed and analyzed.

Blood samples were collected by trained medical professionals to measure serum Vitamin 25-hydroxyvitamin D and Vitamin A concentrations, using a electrochemiluminescence method. Vitamin D and A assessing standards based on serum concentrations listed as below. Vitamin A Concentration: Deficiency < 0.70 $\mu\text{mol/L}$, Marginal deficiency 0.70~1.05 $\mu\text{mol/L}$, Sufficiency $\geq 1.05 \mu\text{mol/L}$. Vitamin D Concentration: Deficiency, < 30nmol/L, Insufficiency 30~50nmol/L, Sufficiency $\geq 50\text{nmol/L}$ ^[11].

Children's height, weight, and BMI, adjusted for age and sex, were categorized into eight hierarchies based on the Chinese Growth Standard for Children under 7 Years of Age (WS/T

423-2022), namely $\geq P97$, $\geq P90$, $\geq P75$, $\geq P50$, $\geq P25$, $\geq P10$, $\geq P3$, and $< P3$. A height $\geq P3$ indicates stunted growth, while $< P3$ indicates severe stunting; a weight $\geq P3$ indicates underweight, while $< P3$ indicates severe underweight; a BMI $\geq P97$ indicates severe obesity, $P90 \leq \text{BMI} < P97$ indicates obesity, $P75 \leq \text{BMI} < P90$ indicates overweight, $P3 \leq \text{BMI} < P10$ indicates thinness, and $< P3$ indicates severe thinness.

Statistical analysis was performed using SPSS version 25.0. Descriptive statistics summarized the demographic characteristics of the sample. Pearson correlation analyses were conducted to explore the relationships between Vitamin D levels and various body composition measures. Independent t-tests were used to compare differences in Vitamin D levels, body composition, height, and weight across age groups and between genders. A p-value < 0.05 was considered statistically significant. Rank sum tests are used for rank data. Hierarchy analysis of height, weight, and BMI by sex and age, gender differences were assessed using the Mann-Whitney U test, while age group (3-, 4-, 5-) differences were analyzed using the Kruskal-Wallis H test.

3. Results

A total of 2035 children participated, with 925 girls and 1110 boys. The age distribution was 329 children in the 3-year-old group, 675 in the 4-year-old group, and 1030 in the 5-year-old group. Vitamin D and A levels were measured in a sub-sample of 1239 children.

For height, 3.20% of children were severe stunted growth, 5.31% stunted growth. For weight, 2.26% of children were severe underweight, while 5.01% were underweight. For BMI, 2.02% were severe thinness, while 5.90% thinness, whereas 5.90% were severe obesity, 6.49% obesity, and 15.0% overweight. Boys displayed significantly higher hierarchy values for height, weight, and BMI (< 0.001). In age groups distribution, the overall hierarchy values of height and weight were statistical differences ($p < 0.001$), however no statistical difference of the overall BMI was shown ($P=0.122$). (Table 1).

No significant differences in Vitamin D and Vitamin A levels were observed between boys and girls across all age groups ($P=0.095$ and $P=0.437$ respectively). Although boys exhibited a lower Vitamin D insufficiency rate (74.63%) than girls (76.65%), the difference was not statistically significant ($p = 0.411$). The prevalence of Vitamin A marginal deficiency was slightly higher in girls (23.35%) compared to boys (21.53%), although this was not statistically significant ($p = 0.445$). Vitamin D deficiency and vitamin A deficiency were not found in our study. (Table 2 and 3).

Correlation between the serum Vitamin D and Vitamin A is statistically significant at the 5% significance level ($t=0.038$), however the level of linear association is weak ($r= 0.186$). (Figure 1)

Except statistical association between BW/FFM, MCA and Vitamin D, there were no significant correlations between Vitamin D levels and BMI, TBW, ICW, ECW, Protein, Minerals, BFM, SLM, FFM, SMM, PBF, and other measured indices. This lack of significant association extends to Vitamin A levels across similar parameters, highlighting the complex interaction of micronutrient status with body composition in children. Table 4.

Table 1 Distribution of eight hierarchies of Height, Weight and BMI by sex and age groups in 2034 children

	sum		Female (n=925)		Male (n=1109)		<i>Z/P</i>	3~ (n=329)		4~ (n=675)		5~ (n=1030)		<i>H/P</i>	
	n	%	n	%	n	%		n	%	n	%	n	%		
	Height	<P3	65	3.20	45	4.86		20	1.80		3	0.91	14		2.07
≥P3		108	5.31	55	5.95	53	4.78		3	0.91	40	5.93	65	6.31	
≥P10		266	13.08	143	15.46	123	11.09		24	7.29	86	12.74	156	15.15	
≥P25		355	17.45	176	19.03	179	16.14	-6.269/	43	13.07	124	18.37	188	18.25	129.619/
≥P50		440	21.63	193	20.86	247	22.27	<0.001	44	13.37	156	23.11	240	23.30	<0.001
≥P75		335	16.47	145	15.68	190	17.13		60	18.24	114	16.89	161	15.63	
≥P90		233	11.46	96	10.38	137	12.35		61	18.54	72	10.67	100	9.71	
≥P97		232	11.41	72	7.78	160	14.43		91	27.66	69	10.22	72	6.99	
Weight	<P3	46	2.26	39	4.22	7	0.63		0	0.00	15	2.22	31	3.01	
	≥P3	102	5.01	57	6.16	45	4.06	-6.275/	5	1.52	33	4.89	64	6.21	55.905/
	≥P10	260	12.78	141	15.24	119	10.73	<0.001	26	7.90	98	14.52	136	13.20	<0.001

	≥P25	422	20.75	203	21.95	219	19.75		52	15.81	135	20.00	235	22.82	
	≥P50	495	24.34	210	22.70	285	25.70		81	24.62	176	26.07	238	23.11	
	≥P75	330	16.22	129	13.95	201	18.12		70	21.28	114	16.89	146	14.17	
	≥P90	195	9.59	74	8.00	121	10.91		52	15.81	55	8.15	88	8.54	
	≥P97	184	9.05	72	7.78	112	10.10		43	13.07	49	7.26	92	8.93	
	<P3	41	2.02	26	2.81	15	1.35		6	1.82	12	1.78	23	2.23	
	≥P3	120	5.90	68	7.35	52	4.69		20	6.08	54	8.00	46	4.47	
	≥P10	320	15.73	160	17.30	160	14.43		54	16.41	103	15.26	163	15.83	
BMI	≥P25	473	23.25	217	23.46	256	23.08	-3.905/	74	22.49	152	22.52	247	23.98	4.213/
	≥P50	523	25.71	227	24.54	296	26.69	<0.001	99	30.09	181	26.81	243	23.59	0.122
	≥P75	305	15.00	116	12.54	189	17.04		43	13.07	103	15.26	159	15.44	
	≥P90	132	6.49	66	7.14	66	5.95		16	4.86	44	6.52	72	6.99	
	≥P97	120	5.90	45	4.86	75	6.76		17	5.17	26	3.85	77	7.48	

Table 2 Distribution of Vitamin D status by Sex and Age groups in 1239 children

		insufficient		sufficient		Sum	χ^2	<i>P</i>
		n	%	n	%			
Sex	girl	430	76.65	131	23.35	561	0.676	0.411
	boy	506	74.63	172	25.37	678		
Sum		936	75.54	303	24.46	1239		
AgeC	3~	140	71.79	55	28.21	195	4.700	0.095
	4~	348	78.91	93	21.09	441		
	5~	448	74.30	155	25.70	603		
Sum		936	75.54	303	24.46	1239		

		marginal deficient		sufficient		Sum	χ^2	<i>P</i>
		n	%	n	%			
Sex	girl	131	23.35	430	76.65	561	0.584	0.445
	boy	146	21.53	532	78.47	678		
Sum		277	22.36	962	77.64	1239		
AgeC	3~	50	25.64	145	74.36	195	1.657	0.437
	4~	99	22	342	78	441		
	5~	128	21.23	475	78.77	603		
Sum		277	22.36	962	77.64	1239		

Table 3 Distribution of Vitamin A status by Sex and Age groups in 1239 children




Figure 1 Correlation of Vitamin A and Vitamin D in 1239 children

Table 4 Correlation analysis of body components and Vitamin D and Vitamin A in 1239 children

	Vitamin D		VitA	
	r	P	r	P
BMI (Body Mass Index)	-0.028	0.320	-0.007	0.799
TBW (Total Body Water)	0.031	0.280	-0.012	0.680
ICW (Intracellular Water)	0.034	0.232	-0.010	0.718
ECW (Extracellular Water)	0.025	0.384	-0.014	0.620
Protein	0.034	0.230	-0.009	0.749
Minerals	-0.026	0.357	-0.027	0.342
SLM (Soft Lean Mass)	0.031	0.269	-0.011	0.689
SLM Index	0.006	0.826	-0.007	0.817
FFM (Fat Free Mass)	0.027	0.346	-0.013	0.644
SMM (Skeletal Muscle Mass)	0.033	0.248	-0.010	0.726
PBF (Percent Body Fat)	-0.029	0.313	0.005	0.847
ECW/TBW	-0.052	0.067	-0.033	0.252
VFA (Visceral Fat Area)	-0.032	0.255	-0.040	0.155
BCM (Body Cell Mass)	0.033	0.242	-0.009	0.747
TBW/FFM*	0.071	0.012	0.029	0.315
FFMI (Fat Free Mass Index)	-0.019	0.501	-0.008	0.782
FMI (Fat Mass Index)	-0.032	0.266	-0.001	0.984
MCA (Measured Circumference of Abdomen)**	0.105	<0.001	0.024	0.391
SMI (Skeletal Muscle Index)	0.004	0.898	-0.009	0.755

4. Discussion

The findings of this study provide a comprehensive examination of the relationship between Vitamin D levels and body composition in a significant sample of kindergarten children from Chengdu, Sichuan Province, China. We found that 75.54% of the preschoolers were Vitamin D insufficiency and 22.36% were marginal Vitamin A deficiency, but none Vitamin D deficiency and vitamin A deficiency were observed. The prevalence of Vitamin D insufficiency was lower in boys (74.63%) than in girls (76.65%) ,and the marginal deficiency rates of Vitamin A in boys and girls was consistent with Vitamin D (21.53% and 23.35%), but neither was statistically significant. The prevalence of vitamin D insufficiency has been reported to vary around the world, with prevalence rates of 20% in Australia, 23.3% in the United States, 34.2% in Africa, 40.4% in Europe, and 53.6% in Japan [12-16]. A systematic review and meta-analysis in 2021 showed that the overall prevalence of vitamin D deficiency and insufficiency of chinese children and adolescents was 23.0% and 46.8% respectively, which means 69.8% of them were vitamin D undernourished[17]. It is generally consistent with the results of this study. Song P et al. reported that the prevalence of both Vitamin A deficiency and marginal Vitamin A deficiency was the highest in children aged 0-5 years at 19.53% and 28.22%, respectively, with both decreasing to 14.48% and 24.28% in children aged 6-12years in low- and middle-income countries[18]. The results of a systematic evaluation by the School of Public Health of Peking University in 2017 showed that the vitamin A deficiency rate among children aged 0 to 12 years in China was 5.16%, and the marginal deficiency rate was 24.29%[19]. This indicates that the nutritional status of vitamin A in preschool children in Chengdu is better than the national level. The prevalence estimates also declined with increasing age, from 31.53% in 0-4 years' children to 31.68% in children aged 5-9 years, which is generally consistent with this study[19].

Our research suggested gender differences in anthropometric measurements, with boys showing higher mean values for height, weight, and BMI. A cross-sectional study in Jiangsu, China showed that boys had lower rates of growth retardation, underweight and wasting than girls (1.45% vs. 1.78%, 0.54% vs. 0.67%, 1.36% vs. 2.00%) [20], This is consistent with our study. Wang Fuman et al. reported that the obesity rate of preschool children in seven cities in China was 12.4%[21], which was higher than this study; and the rates of overweight and obesity in boys were higher than those in girls, with the rates increasing with age, which was consistent with the present study. This finding contributes to the growing evidence on gender disparities in growth patterns and body composition during early childhood. However, the similar rates of Vitamin D sufficiency between genders suggest that Vitamin D's role in these differences is minimal, if existent.

Consistent with previous research[22-24], our study found no significant correlation between Vitamin D levels and various measures of body composition, including BMI, BFM, SLM, and SMM. Moreover, many researchers have studied the effects of vitamin D supplementation on normal weight or overweight/obese children deficient in vitamin D, including body weight, body composition, growth and development the effects remain controversial[25-27]. This aligns with the body of evidence suggesting that while Vitamin D is crucial for bone health and metabolism, its impact on body fat and muscle mass in children may not be as significant as previously thought.

The lack of a significant association between Vitamin D levels and body composition in this population might be due to several factors. Firstly, genetic variations, dietary patterns, sun exposure habits, and overall health status can all influence Vitamin D metabolism and its effects on the body. Additionally, the narrow range of Vitamin D levels observed in our study population could have limited our ability to detect significant associations.

Our study found a correlation between vitamin A and vitamin D, but the correlation was weak ($r=0.186$, $P=0.038$). Multiple studies show interaction between vitamin A and vitamin D. A study in Brazilian covering 468 children suggested that serum retinol concentrations were significantly and positively correlated with vitamin D ($P=0.001$), and vitamin D adequacy increased up to 1.38-fold for each 1- μ mol/L increase in vitamin A concentration (95% CI:1.18-1.61)^[28]. Li XY reported that vitamin D₃ only can induced catalase (CAT) activity by about 100%,when together with 9-cis retinoic acid, the CAT activity was increased by 230%^[29]. Given the prevalence of vitamin A and vitamin D deficiencies in children and the synergistic effects of vitamin A and vitamin D, combined supplementation of the two may help to improve the nutritional status of both vitamin A and vitamin D.

Future research should aim to address these limitations by incorporating longitudinal designs, examining genetic factors that might influence Vitamin D metabolism, and considering the role of dietary Vitamin D intake and sun exposure. Understanding the complex interactions between these factors and body composition will be crucial for developing targeted interventions to improve child health outcomes.

5. Conclusion

This study contributes to the existing literature by exploring the relationship between Vitamin D levels and body composition among kindergarten children in Chengdu, Sichuan Province, China. While it found no significant correlation between Vitamin D levels and body composition, it highlighted important gender differences in growth patterns. These findings underscore the complexity of the relationship between Vitamin D and body composition and point to the need for further research in this area.

References

1. Montenegro KR, Cruzat V, Carlessi R, Newsholme P. Mechanisms of vitamin D action in skeletal muscle. *Nutr Res Rev.* 2019 Dec;32(2):192-204. doi: 10.1017/S0954422419000064. Epub 2019 Jun 17. PMID: 31203824.
2. Sonne J, Kofoed-Enevoldsen P, Andersen TB, et al. Association between 25-hydroxyvitamin D level and body fat percentage in preschool children. *Pediatr Obes.* 2015;10(8):600-6.

doi:10.1111/pedob.12225.

3. Healthier bone growth and overall growth: Aksoy N, Alagol R, Kose N, et al. The relationship between serum 25-hydroxyvitamin D levels and linear growth in healthy schoolchildren. *J Int Med Res.* 2011;39(5):849-54. doi:10.1177/0300926811413731.
4. Bezrati I, Hammami R, Ben Fradj MK, Martone D, Padulo J, Feki M, Chaouachi A, Kaabachi N. Association of plasma 25-hydroxyvitamin D with physical performance in physically active children. *Appl Physiol Nutr Metab.* 2016 Nov;41(11):1124-1128. doi: 10.1139/apnm-2016-0097. Epub 2016 Jul 13. PMID: 27764544.
5. Cediel, G., Corvalán, C., Aguirre, C. et al. Serum 25-Hydroxyvitamin D associated with indicators of body fat and insulin resistance in prepubertal Chilean children. *Int J Obes* 40, 147–152 (2016). <https://doi.org/10.1038/ijo.2015.148>
6. Doaei S, Jarrahi S, Torki S, Haghshenas R, Jamshidi Z, Rezaei S, Moslem A, Ghorat F, Khodabakhshi A, Gholamalizadeh M. Serum vitamin D level may be associated with body weight and body composition in male adolescents; a longitudinal study. *Pediatr Endocrinol Diabetes Metab.* 2020;26(3):125-131. English. doi: 10.5114/pedm.2020.97466. PMID: 32901469.
7. Kouda K, Nakamura H, Fujita Y, Ohara K, Iki M. Vitamin D status and body fat measured by dual-energy X-ray absorptiometry in a general population of Japanese children. *Nutrition.* 2013 Oct;29(10):1204-8. doi: 10.1016/j.nut.2013.03.010. Epub 2013 Jun 22. PMID: 23800567.
8. Li H, Huang T, Xiao P, Zhao X, Liu J, Cheng H, Dong H, Morris HA, Mi J; China Child and Adolescent Cardiovascular Health (CCACH) Collaboration Group. Widespread vitamin D deficiency and its sex-specific association with adiposity in Chinese children and adolescents. *Nutrition.* 2020 Mar;71:110646. doi: 10.1016/j.nut.2019.110646. Epub 2019 Nov 9. PMID: 31896064
9. Kasvis P, Cohen TR, Loiselle SÈ, Hazell TJ, Vanstone CA, Weiler HA. Associations between Body Composition and Vitamin D Status in Children with Overweight and Obesity Participating in a 1-Year Lifestyle Intervention. *Nutrients.* 2022 Jul 30;14(15):3153. doi: 10.3390/nu14153153. PMID: 35956333; PMCID: PMC9370728.
10. White Z, White S, Dalvie T, Kruger MC, Van Zyl A, Becker P. Bone Health, Body Composition, and Vitamin D Status of Black Preadolescent Children in South Africa. *Nutrients.* 2019 May 31;11(6):1243. doi: 10.3390/nu11061243. PMID: 31159206; PMCID: PMC6627122
11. Chinese Preventive Medicine Association.Child Health Branch. Expert consensus on the clinical application of vitamin A and vitamin D in Chinese children (2024)[J]. *Chinese Journal of Child Health*,2024, 32(4):349-358.DOI:10.11852/zgetbjzz2024-0279.

12. Malacova E, Cheang PR, Dunlop E, Sherriff JL, Lucas RM, Daly RM, et al. Prevalence and predictors of vitamin D deficiency in a nationally representative sample of adults participating in the 2011-2013 Australian Health Survey. *Br J Nutr* 2019;121(8):894 – 904. doi: 10.1017/s0007114519000151.
13. Herrick KA, Storandt RJ, Afful J, Pfeiffer CM, Schleicher RL, Gahche JJ, et al. Vitamin D status in the United States, 2011-2014. *Am J Clin Nutr* 2019;110(1):150 – 7. doi: 10.1093/ajcn/nqz037.
14. Mogire RM, Mutua A, Kimita W, Kamau A, Bejon P, Pettifor JM, et al. Prevalence of vitamin D deficiency in Africa: a systematic review and meta-analysis. *Lancet Glob Health* 2020;8(1):e134 – e42. doi: 10.1016/s2214-109x(19)30457-7.
15. Cashman KD, Dowling KG, Skrabakova Z, Gonzalez-Gross M, Valtuena J, De Henauw S, et al. Vitamin D deficiency in Europe: pandemic? *Am J Clin Nutr* 2016;103(4):1033 – 44. doi: 10.3945/ajcn.115.120873.
16. Nakamura K, Kitamura K, Takachi R, Saito T, Kobayashi R, Oshiki R, et al. Impact of demographic, environmental, and lifestyle factors on vitamin D sufficiency in 9084 Japanese adults. *Bone* 2015;74:10 – 7. doi: 10.1016/j.bone.2014.12.064.
17. Liu W, Hu J, Fang Y, Wang P, Lu Y, Shen N. Vitamin D status in Mainland of China: A systematic review and meta-analysis. *EClinicalMedicine*. 2021 Jul 14;38:101017. doi: 10.1016/j.eclinm.2021.101017. PMID: 34308318; PMCID: PMC8283334.
18. Song P, Adeloye D, Li S, Zhao D, Ye X, Pan Q, Qiu Y, Zhang R, Rudan I; Global Health Epidemiology Research Group (GHERG). The prevalence of vitamin A deficiency and its public health significance in children in low- and middle-income countries: A systematic review and modelling analysis. *J Glob Health*. 2023 Aug 11;13:04084. doi: 10.7189/jogh.13.04084. PMID: 37565390; PMCID: PMC10416138.
19. Song P, Wang J, Wei W, Chang X, Wang M, An L. The Prevalence of Vitamin A Deficiency in Chinese Children: A Systematic Review and Bayesian Meta-Analysis. *Nutrients*. 2017 Nov 25;9(12):1285. doi: 10.3390/nu9121285. PMID: 29186832; PMCID: PMC5748736.
20. Liu Y, Wang YY, Cheng Y, et al. Growth and development of children and related influencing factors: a cross-sectional study of the families with children aged 0-6 years in Jiangsu Province. *Zhongguo Dang Dai Er Ke Za Zhi*. 2022 Jun 15;24(6):693-698. Chinese. doi: 10.7499/j.issn.1008-8830.2202072. PMID: 35652424; PMCID: PMC9250398.
21. Wang Fuman, Yao Yi, Yang Qi. A cohort study of wasting, overweight and obesity among preschool children in seven cities in China[J]. *Chinese Journal of Disease Control & Prevention*, 2019, 23(5): 522-526. DOI: 10.16462/j.cnki.zhjbkz.2019.05.05.006.
22. Pruszkowska-Przybylska P, Sitek A, Rosset I, Sobalska-Kwapis M, Słomka M, Strapagiel D, Żądzińska E, Morling N. Associations between second to fourth digit ratio, cortisol, vitamin

- D, and body composition among Polish children. *Sci Rep.* 2021 Mar 29;11(1):7029. doi: 10.1038/s41598-021-86521-7. PMID: 33782473; PMCID: PMC8007693.
23. Correa-Rodríguez M, Schmidt-RioValle J, Ramírez-Vélez R, Correa-Bautista JE, González-Jiménez E, Rueda-Medina B. Influence of Calcium and Vitamin D Intakes on Body Composition in Children and Adolescents. *Clin Nurs Res.* 2020 May;29(4):243-248. doi: 10.1177/1054773818797878. Epub 2018 Aug 31. PMID: 30168345.
 24. White Z, White S, Dalvie T, Kruger MC, Van Zyl A, Becker P. Bone Health, Body Composition, and Vitamin D Status of Black Preadolescent Children in South Africa. *Nutrients.* 2019 May 31;11(6):1243. doi: 10.3390/nu11061243. PMID: 31159206; PMCID: PMC6627122.
 25. Ganmaa D, Bromage S, Khudyakov P, Erdenenbaatar S, Delgererekh B, Martineau AR. Influence of Vitamin D Supplementation on Growth, Body Composition, and Pubertal Development Among School-aged Children in an Area With a High Prevalence of Vitamin D Deficiency: A Randomized Clinical Trial. *JAMA Pediatr.* 2023 Jan 1;177(1):32-41. doi: 10.1001/jamapediatrics.2022.4581. PMID: 36441522; PMCID: PMC9706398.
 26. Corsello A, Macchi M, D'Oria V, Pigazzi C, Alberti I, Treglia G, De Cosmi V, Mazzocchi A, Agostoni C, Milani GP. Effects of vitamin D supplementation in obese and overweight children and adolescents: A systematic review and meta-analysis. *Pharmacol Res.* 2023 Jun;192:106793. doi: 10.1016/j.phrs.2023.106793. Epub 2023 May 11. PMID: 37178775.
 27. Brzeziński M, Jankowska A, Słomińska-Frączek M, Metelska P, Wiśniewski P, Socha P, Szlagatys-Sidorkiewicz A. Long-Term Effects of Vitamin D Supplementation in Obese Children During Integrated Weight-Loss Programme-A Double Blind Randomized Placebo-Controlled Trial. *Nutrients.* 2020 Apr 15;12(4):1093. doi: 10.3390/nu12041093. PMID: 32326621; PMCID: PMC7230345.
 28. Lourenço BH, Silva LL, Fawzi WW, Cardoso MA; ENFAC Working Group. Vitamin D sufficiency in young Brazilian children: associated factors and relationship with vitamin A corrected for inflammatory status. *Public Health Nutr.* 2020 May;23(7):1226-1235. doi: 10.1017/S1368980019002283. Epub 2019 Aug 23. PMID: 31439064; PMCID: PMC10200414.
 29. Li XY, Xiao JH, Feng X, Qin L, Voorhees JJ. Retinoid X receptor-specific ligands synergistically upregulate 1, 25-dihydroxyvitamin D3-dependent transcription in epidermal keratinocytes in vitro and in vivo. *J Invest Dermatol.* 1997 Apr;108(4):506-12. doi: 10.1111/1523-1747.ep12289733. PMID: 9077482.