

1 **PREVALENCE OF ABNORMAL SEMINAL FLUID AND ASSOCIATED FACTORS**
2 **AMONG PATIENTS ATTENDING FERTILITY CLINICS IN OSOGBO: A CROSS**
3 **SECTIONAL STUDY**

4 ^aKehinde Awodele, ^bSunday Charles Adeyemo, ^cEniola Dorcas Olabode, ^dAdeniyi
5 Olaonipekun Fasanu, ^eLanre Olaitan, ^dAkintunde Rasaq Akindele, ^fOlufunso Abidemi
6 Olagunju, ^gAdeola Dorcas Aderinwale

7

8 ^aObstetrics and Gynecology Department, Uniosun Teaching Hospital, Osogbo, Nigeria.

9 Kawodele@uth-osogbo.org.ng

10 ^bInstitut Supérieur de Santé, Niamey, Niger Republic

11 charlespatho@gmail.com

12 ^cDepartment of Community Health, Obafemi Awolowo University, Ile-Ife, Nigeria

13 oluwafemieniola.ed@gmail.com

14 ^dDepartment of Obstetrics and Gynecology, Osun State University, Osogbo, Nigeria.

15 adeniyi.fasanu@uniosun.edu.ng

16 drakindele@yahoo.com

17 ^eDepartment of public health/fb. University of Ilorin.

18 Lanreolive@gmail.com

19 ^fDepartment of Pediatrics, Osun State University, Osogbo, Osun State, Nigeria

20 funso.olagunju@uniosun.edu.ng

21 ^gPrimary Health Care Unit, Adeleke University, Ede, Nigeria

22 aderinwaleadeola@gmail.com

23 **Corresponding author:**

24 Kehinde Awodele

25 +2348033992462, Kawodele@uth-osogbo.org.ng

26

27

28

29 **ABSTRACT**

30 **Background:** Infertility is a global health issue that affects millions of couples worldwide, In
31 Nigeria, the prevalence of infertility is particularly high underscoring the need for a better
32 understanding of the factors contributing to male infertility at which Seminal Fluid Analysis
33 (SFA) is the most important characteristics. This study aimed to assess the prevalence of
34 abnormal seminal fluid and its associated factors among patients attending fertility clinics in
35 Osogbo, Nigeria.

36 **Methods and Findings:** A descriptive cross-sectional study design was used among male
37 patients attending fertility centers in Osogbo, Osun State. Fisher's formula ($n = z^2pq/d^2$) was
38 used to determine the sample size. 305 respondents were selected using multistage
39 sampling technique. Pre-tested questionnaire was used to collect data from respondents.
40 Patients who consented to the study were also made to undergo seminal fluid analysis. The
41 results were analysed using IBM Statistical Product for Service Solution (SPSS) version
42 27. Descriptive statistics was used for all variables. Bivariate and multivariate analysis
43 were done at $p < 0.05$ as level of significance. Majority of the respondents, 257 (84.3%) had
44 at least one abnormality of Seminal fluid. Multiple regression analysis revealed that
45 respondents who were habitual drunkard were about five times more likely to have at least
46 one abnormality in their seminal fluid (OR: 4.990, p : 0.004, C.I.: 1.688-14.749) and
47 smokers were three times more likely to have at least one abnormality in their seminal fluid
48 ($p=0.005$, OR=3.300 and C.I.=1.396-4.273). Also, respondents with history of sexually
49 transmitted infection were 3.5 times more likely to have at least one abnormality in their
50 seminal fluid ($p=0.039$, OR=3.595 and C.I.=1.072-14.146).

51 **Conclusions:** The study observed high prevalence of abnormal seminal fluid which was
52 significantly associated with lifestyle habits such as smoking, alcohol as well as previous
53 history of sexually transmitted infection. The study recommended that advocacy program for

54 healthy lifestyle, early screening and public health education will further reduce the burden of
55 abnormal seminal fluid and by implication, infertility among couples.

56 **Keywords:** Male Infertility, Implications, Seminal fluid Analysis, Sperm count.

57

58

59 **INTRODUCTION**

60 Infertility is a global health issue that affects millions of couples worldwide. The inability to
61 conceive a child after a year of regular, unprotected intercourse is a distressing experience for
62 many couples, leading to emotional, psychological, and social challenges. [1] In Nigeria, the
63 prevalence of infertility is particularly high, with studies showing that around 20-30% of
64 couples face fertility challenges. [2]

65 One of the most important aspects of male infertility is the quality and characteristics of the
66 seminal fluid. Seminal fluid, also known as semen, is a complex mixture of spermatozoa and
67 secretions from the male reproductive glands, including the seminal vesicles, prostate, and
68 bulbourethral glands. Seminal fluid plays a crucial role in the fertilisation process, providing
69 a supportive environment for sperm transport, nourishment, and protection against the female
70 reproductive tract's hostile conditions. [3]

71 Infertility is a significant public health concern affecting millions of couples worldwide, with
72 male infertility accounting for approximately half of these cases.[4] In Southwest Nigeria,
73 particularly, there is a noticeable gap in research regarding the seminal fluid analysis (SFA)
74 and its implications on male infertility among patients attending fertility clinics [5] Therefore,
75 this study aims to assess the prevalence of abnormal seminal fluid among male patients
76 attending fertility centers in Osogbo, Osun State, Nigeria.

77 **METHODS**

78 A descriptive cross-sectional study design was used in the study. The study population
79 were male patients attending fertility centers in Osogbo, Osun State. Fisher's formula

80 $(n=z^2pq/d^2)$ was used to determine the sample size. 305 respondents were selected using
81 multistage sampling technique. Out of 10 major fertility centers in Osogbo, Osun State, 5
82 centers were randomly selected. Proportional allocation was used to determine the number
83 of respondents from each center. Male patients aged 18-50 were randomly selected in each
84 center. Pre-tested questionnaire was used to collect data from respondents. Patients who
85 consented to the study were also made to undergo seminal fluid analysis. The semen was
86 collected aseptically after an abstinence period of 2–7 days using sterile universal bottles in
87 a private room near the laboratory to limit the exposure of the semen to fluctuations in
88 temperature and to control the time between collection and analysis. The semen was then
89 analysed in the laboratory for sperm count, motility and morphology of the semen. The
90 data from the questionnaire were analysed using IBM Statistical Product for Service
91 Solution (SPSS) version 27. Descriptive statistics was used for all variables. Bivariate and
92 multivariate analysis were done at $p<0.05$ as level of significance.

93 **RESULTS**


94 **Sociodemographic characteristics of respondents**

95 All the respondents were married male attending the infertility clinic in Osogbo. More than
96 half 164 (53.8%) of the respondents were with the ages of 31-40 years, while the majority,
97 267(87.5%) were Yoruba. 164 (53.8%) had tertiary level of education followed by
98 secondary 83 (27.2%). About half 140 (45.9%) were skilled workers while 110 (36.1%)
99 were unskilled.

100 **Prevalence of abnormal seminal fluid**

101 Majority of the respondents, 84.3% had at least one abnormality while 15.7% have normal
102 seminal fluid. About two-thirds 217 (71.1%) had normal sperm count (>32 million per
103 ejaculation) while 88 (28.9%) had abnormal sperm count. Only 97 (31.8%) had normal
104 progressive motility (>32 percent) while 208 (68.2%) had abnormal (Athenospermia)

105 progressive motility and 178 (58.4%) were found to have abnormal morphology
106 (Teratospermia i.e., <4%). (Figure 1)



107

108 **Figure 1: Seminal fluid Analysis of Respondents**

109

110 **Lifestyle factors of respondents**

111 A little above one-third were known smokers (35.1), out of which 78.5% smokes cigarette
112 and 21.5% smokes marijuana. Less than half (47.5%) were known drunkard while only
113 8.5% currently used drugs. The majority (68.2%) eats any type of food available while
114 23.6% regularly consumed carbohydrate/starch food and only 1.0% consumed vegetables.
115 (Table 1)

116

117

118 **Table 1: Lifestyle behaviours among the respondents**

Variables	Frequency (n=305)	Percentage
Known smoker		
Yes	107	35.1
No	198	64.9
Type of smoke used (n=107)		
Cigarette	84	78.5
Marijuana	23	21.5
Known Drunkard		
Yes	145	47.5
No	160	52.5
Currently using any drug		
Yes	26	8.5
No	279	91.5
Type of food consumed regularly		
Any food	208	68.2
Carbohydrate/Starch	72	23.6
Protein	22	7.2
Vegetables	3	1.0

119

120

121 **Association between medical history of respondents and abnormal seminal fluid among**
122 **respondents**

123 Majority of the respondents 65 (91.5%) who were diabetic had at least one abnormality in
124 seminal fluid. Majority of the respondents 29 (96.7%) who were hypertensive had at least
125 one abnormality in seminal fluid. Majority of the respondents 49 (90.7%) who were
126 overweight had at least one abnormality in seminal fluid. However, there was no
127 statistically significant association between these factors and abnormal seminal fluid.
128 Majority of the respondents 84 (96.6%) who had history of sexually transmitted infection
129 had at least one abnormality in seminal fluid and this was statistically significant
130 (p<0.001). (Table 2)

131

132

133 **Table 2: Association between medical history of respondents and abnormal seminal**
 134 **fluid among respondents**

Variables	Seminal Fluid Analysis Category		Statistics
	At least One No Abnormality (n=257) %	Abnormality (n=48) %	
Diabetic			$x^2 = 3.706$ $p = 0.054$
Yes	65 (91.5)	6 (8.5)	
No	192 (82.1)	42 (17.9)	
Hypertensive			$x^2 = 2.893^*$ $p = 0.089$
Yes	29 (96.7)	1 (3.3)	
No	228 (82.9)	47 (17.1)	
History of Overweight			$x^2 = 2.077$ $p = 0.150$
Yes	49 (90.7)	5 (9.3)	
No	208 (82.9)	43 (17.1)	
History of STI			$x^2 = 13.863$ $p < 0.001^{**}$
Yes	84 (96.6)	3 (3.4)	
No	173 (79.4)	45 (20.6)	

135 χ^2 = chi-square, p = p-value, * Continuity Correction, ** Statistically Significant

136

137 **Predictors of abnormal seminal fluid among respondents**

138 Multiple regression analysis revealed that respondents who were habitual drunkard were
 139 about five times more likely to have at least one abnormality in their seminal fluid (OR:
 140 4.990, p: 0.004, C.I.: 1.688-14.749) and smokers were three times more likely to have at
 141 least one abnormality in their seminal fluid (p=0.005, OR=3.300 and C.I.=1.396-4.273).
 142 Also, respondents with history of sexually transmitted infection were 3.5 times more likely

143 to have at least one abnormality in their seminal fluid ($p=0.039$, $OR=3.595$ and $C.I=1.072-$
144 14.146).

145

146 **Discussion**

147 This study assessed the prevalence of abnormal seminal fluid and its associated factors
148 among patients attending fertility clinics in Osogbo, Osun State, Nigeria. More than half of
149 the respondents were between the ages of 31 and 40 years old while more than two-fifth
150 were aged 21-30 years. More than half of the respondents have tertiary as their highest
151 educational attainment. This is similar to another study conducted among male partners
152 attending infertility clinic in a tertiary Nigerian hospital [6]

153 More than four-fifth of the respondents had at least one abnormality. About one-third had
154 abnormal sperm count, which is below the WHO 2010 reference for semen analysis. [7]

155 This proportion is also lower compared to the study of [8] where more than four-fifths were
156 found to have abnormal sperm count. Almost 7 out of 10 respondents had abnormal
157 (Athenospermia) progressive motility) which is more than the WHO 2010 reference of
158 $<32\%$. [7] Furthermore, this proportion is higher compared to another study carried out
159 among male partners of infertile couples seeking care at the Lagos University Teaching
160 Hospital where 55.8% had abnormal progressive motility. [9] Also, about three-fifth had
161 abnormal morphology (Teratospermia i.e., $<4\%$). This proportion is much higher compared
162 to 60.8% of abnormal morphology reported in the study of [6] among male patients in a
163 tertiary Nigerian hospital.

164 The predictors of abnormal seminal fluid among respondents included smoking, drinking,
165 and previous history of sexual transmitted infection such that respondents who were
166 habitual drunkard were about five times more likely to have at least one abnormality in

167 their seminal fluid (OR: 4.990, p: 0.004, C.I.: 1.688-14.749) and smokers were three times
168 more likely to had at least one abnormality in their seminal fluid (p=0.005, OR=3.300 and
169 C.I.=1.396-4.273). Also, respondents with history of sexually transmitted infection were 3.5
170 times more likely to have at least one abnormality in their seminal fluid (p=0.039,
171 OR=3.595 and C.I.=1.072-14.146). This was also evident in the study of [6] where age,
172 smoking and alcohol consumption were reported as the possible risk factors for abnormal
173 sexually transmitted infection. This suggests that substance abuse, habitual smoker,
174 drinking habit have detrimental impact on reproductive health and fertility as it has been
175 associated with reduced sperm count, motility, and normal morphology. It is also linked to
176 increased levels of sperm aneuploidy, lower seminal plasma antioxidant levels, and
177 increased oxidative damage to sperm deoxyribonucleic acid (DNA). [10]

178 **Conclusions:** The findings indicate a prevalence of abnormal seminal fluid parameters (low
179 sperm count, motility, and morphology) among the participants in which motility has the
180 highest prevalence followed by morphology and sperm count respectively. Lifestyle factors
181 such as smoking, alcohol consumption and sexually transmitted infections were found to be
182 significantly associated with abnormal seminal fluid. The implications of abnormal seminal
183 fluid parameters on infertile males are substantial, extending beyond physical health to
184 encompass psychological, emotional, and social well-being. The findings from this study
185 indicates the need for increased awareness about male infertility and accessibility to male
186 fertility testing in order to combat the issue of infertility among couples. Further studies can
187 assess the effect on seminal fluid analysis on male infertility.

188 **Acknowledgements**

189 The authors wish to acknowledge the respondents who gave their consent to be part of this
190 study as well as the spouses of the authors for their understanding during the course of this
191 study.

192 **REFERENCES**

- 193 [1] Zegers-Hochschild, F., Adamson, G. ., De Mouzon, J., Ishihara, O., Mansour, R., and
194 Nygren, K. (2009). The International Committee for Monitoring Assisted Reproductive
195 Technology (ICMART) and the World Health Organization (WHO) Revised Glossary on
196 ART Terminology,. *Human Reproduction*, *24*, 2683–2687.
- 197 [2] Okonofua, F., Ntoimo, L., Ogungbangbe, J., Anjorin, S., Imongan, W., and Yaya, S.
198 (2018). Predictors of women’s utilization of primary health care for skilled pregnancy care
199 in rural Nigeria. *BMC Pregnancy and Childbirth*, *18*(1), 1–15.
200 <https://doi.org/10.1186/s12884-018-1730-4>
- 201 [3] Robertson, S. A., and Sharkey, D. J. (2016). Seminal fluid and fertility in women.
202 *Fertility and Sterility*, *106*(3), 511–519. <https://doi.org/10.1016/j.fertnstert.2016.07.1101>
- 203 [4] Agarwal, A., Mulgund, A., Hamada, A., & Chyatte, M. (2015). A unique view on male
204 infertility around the globe. *Reproductive Biology and Endocrinology*, *13*(1), 37.
- 205 [5] Ibitoye, B. O., Fasasi, A. O., Imosemi, I. O., Alabi, O., Olaniyan, O. T., Ibitoye, F. O.,
206 Faduola, P., Bodun, D. S., and Wike, N. Y. (2023). The spermiogram and correlation of
207 seminal fluid parameters in patient attending fertility centre in Lagos, Southwest Nigeria.
208 *Morphologie*, *107*(359), 100606. <https://doi.org/10.1016/j.morpho.2023.05.004>
- 209 [6] Umar, A. ., Panti, A. ., Mbakwe, M., Ahmed, Y., Garba, J. ., & Nnadi, D. C. (2020).
210 The Pattern of Seminal Fluid Analysis among Male Partners Attending an Infertility
211 Clinicin a Nigerian Tertiary Health Institution. *Open Journal of Obstetrics and*
212 *Gynecology*, *10*, 957–967.
- 213 [7] WHO. (2021a). WHO laboratory manual for the examination and processing of human
214 semen. In *WHO laboratory manual for the examination and processing of human semen*
215 (5th ed.) (5th ed.).

216 [8] Tilahun, T., Oljira, R., and Getahun, A. (2022). Pattern of semen analysis in male
217 partners of infertile couples in Western Ethiopia: Retrospective cross-sectional study.
218 *SAGE Open Medicine*, 10, 1–7. <https://doi.org/10.1177/20503121221088100>

219 [9] Makwe, C. C., Ugwu, A. O., Ojewola, R. W., and Onyeze, C. I. (2021). Seminal fluid
220 parameters of male partners of infertile couples seeking care at the Lagos University
221 Teaching Hospital. *International Journal of Reproduction, Contraception, Obstetrics and*
222 *Gynecology*, 10(4), 1347. <https://doi.org/10.18203/2320-1770.ijrcog20211109>

223 [10] Joo, K., Kwon, Y., Myung, S., & Kim, T. (2012). The Effects of Smoking and
224 Alcohol Intake on Sperm Quality: Light and Transmission Electron Microscopy Findings.
225 *Journal of International Medical Research*, 40(6), 2327–2335.
226 <https://doi.org/10.1177/030006051204000631>

227 **Limitation**

228 The study is limited to patients attending fertility centers in Osogbo, which can have effect
229 on the generalization of the study. Also, due to the nature of the study as a cross-sectional
230 study, causal relationship may not be established.

231 **Ethical approval and consent to participate**

232 Ethical approvals for this study were obtained from the Ethics and Research Committee of
233 Adeleke University, Ede, and Uniosun Teaching Hospital. The Protocol number
234 UTH/EC/2024/06/957. All ethical principles guiding the conduct of research such as
235 informed consent, beneficence, non- maleficence, confidentiality, justice, autonomy etc
236 were strictly taken into consideration. All information gathered were confidential and
237 privacy and confidentiality of the respondents were guaranteed. All respondents were made
238 to sign the informed consent before filling the questionnaire.

239 **Consent for publication**

240 Not applicable

241 **Availability of data and material**

242 The data for this study is available on reasonable request from the corresponding author.

243 **Competing interest**

244 The authors know no competing interest for the study

245 **Funding**

246 The study was funded by the authors.

247 **Authors' contribution**

248 Kehinde Awodele-Research design, ethical compliance and data collection, Sunday Charles
249 Adeyemo-Data Analysis and manuscripting, Eniola Dorcas Olabode- Data analysis and
250 manuscripting, Adeniyi Olaonipekun Fasanu- Data collection and ethical compliance, Lanre
251 Olaitan- Supervision and ethical compliance, Akintunde Rasaq Akindele- Data collection and
252 ethical compliance, Olufunso Abidemi Olagunju- Data collection and Analysis, Adeola
253 Dorcas Aderinwale-Data collection

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

Seminal Fluid Analysis of Respondents




Figure 1: Seminal fluid Analysis of Respondents