

1 Global Stroke Burden from Metabolic Risks Across Demographics: Findings from the
2 2021 Global Burden of Disease Study

3

4 Song Xue^{1,2}, Guoqing Wu^{2*}

5

6 ¹ School of Minerals Processing and Bioengineering, Central South University,
7 Changsha, Hunan 410011, China.

8 ² Affiliated Hospital of Jiangxi University of Traditional Chinese Medicine, Nanchang,
9 Jiangxi 330006, China

10

11 * Corresponding author: 13970986129@163.com

12 Song Xue: Ph. D. Candidates, xuesong2022@csu.edu.cn, School of Minerals
13 Processing and Bioengineering, Central South University, Changsha, Hunan 410011,
14 China.

15 Guoqing Wu: Master's degree, 13970986129@163.com, Affiliated Hospital of Jiangxi
16 University of Traditional Chinese Medicine, Nanchang, Jiangxi 330006, China

17

18

Abstract

19 **Background:** Stroke is the second leading cause of death and the primary cause of
20 disability worldwide. Metabolic risks are major contributors to stroke. The global
21 trends in metabolic risk-related stroke from 1990 to 2019, and the differences in
22 mortality and DALYs across various demographic factors, remain unclear.

23 **Methods:** All analyses were based on rates derived from the GBD2021 results
24 (<https://vizhub.healthdata.org/gbd-results/>). Data were stratified by gender, region,
25 and age. Joinpoint software was used to perform regression analysis of the average
26 annual percent change (AAPC) and its 95% confidence interval to analyze trends
27 from 1990 to 2019. Excel, PowerPoint, and R software were used for plotting and
28 analysis, with $p < 0.05$ considered statistically significant.

29 **Results:** From 1990 to 2019, the average annual percent change (AAPC) for
30 age-standardized rates (ASR) of DALYs was -1.70% (-1.81%, -1.58%), and for
31 mortality, the AAPC was -1.57% (-1.68%, -1.46%). As the SDI increased, both the
32 ASR of DALYs and mortality in 2019 showed a significant decline. The AAPC from
33 1990 to 2019 also exhibited a downward trend with increasing SDI levels. The
34 DALYs and mortality rates of metabolic risk-related stroke predominantly affected
35 individuals aged 75 and above, with a lesser impact on those under 55. For both
36 genders, the 10-55 and 50-74 age groups had the highest DALYs and mortality rates
37 due to metabolic-related intracerebral hemorrhage. For those aged 75-84 and over 85,
38 ischemic stroke was the leading subtype of metabolic-related stroke contributing to
39 DALYs and mortality rates.

40 **Conclusion:** This is the first retrospective study on metabolic risk-related stroke on a
41 global scale, summarizing its temporal trends and demographic distribution
42 characteristics. Effective public health strategies are needed to address these
43 disparities and continue reducing the global burden of metabolic risk-related strokes.

44

45 **Introduction:**

46 Stroke is the second leading cause of death and the primary cause of disability
47 worldwide.¹ Over the past three decades, the global burden of stroke has seemingly
48 decreased, primarily reflecting declines in high-income countries.² In contrast, the
49 burden of stroke has rapidly increased in low- and middle-income countries,³ with
50 variations in age and gender compositions contributing to differences in stroke burden
51 across countries.⁴

52 Metabolic factors are key contributors to stroke risk.⁵ Elevated fasting blood
53 glucose levels,⁶ high low-density lipoprotein cholesterol,⁷ high systolic blood
54 pressure,⁸ elevated body mass index,⁹ low bone mineral density,¹⁰ and impaired
55 renal function¹¹ significantly increase stroke risk. However, comprehensive
56 epidemiological information on stroke burden related to metabolic risk factors
57 remains limited.

58 In this study, we extracted epidemiological data on stroke related to metabolic
59 risks from the Global Burden of Disease (GBD) 2021 study. The GBD study is a
60 collaborative research project between the World Health Organization (WHO) and
61 many academic institutions that aims to assess the burden of disease and injury at the
62 global, regional and national levels.¹² The data of GBD 2021 study were stratified by
63 location, gender, age, and Socio-Demographic Index (SDI) status, and we further
64 analyzed the trends in disease burden over the demographics and time dimensions.
65 This study represents the first global epidemiological analysis specifically addressing
66 stroke related to metabolic factors, providing valuable insights for developing
67 prevention strategies tailored to the metabolic risk profiles of different countries and
68 regions.

69

70 **Methods**

71 2.1 Data Sources

72 The GBD 2021 study provided a systematic and comprehensive assessment of
73 diseases and injuries worldwide from 1990 to 2021.¹³ Data sources included field
74 surveys, censuses, vital registration data, and other health-related data sources.¹² The
75 GBD study followed the Guidelines for Accurate and Transparent Health Estimates
76 Reporting (GATHER).¹⁴

77 The results were available from the GBD online results tool
78 (<https://vizhub.healthdata.org/gbd-results/>) and could be viewed interactively via the
79 GBD compare tool (<https://vizhub.healthdata.org/gbd-compare/>).).

80 2.2 Exposure and Outcomes

81 There were three primary risk factors in the GBD study (Supplemental Table 1).
82 Metabolic risk was a primary risk factor, including high fasting plasma glucose, high
83 LDL cholesterol, high systolic blood pressure, high body mass index, low bone
84 mineral density, and impaired kidney function. (Supplemental Table 1) Stroke was a
85 third-level disease in the GBD study, which included three fourth-level diseases. The
86 definition of each disease could be found in previous research.¹⁵

87 SDI, a composite indicator of social and economic factors influencing health
88 outcomes in each region, was calculated by GBD 2021 for each country.¹⁶ SDI was
89 categorized into five levels: low, low-middle, middle, high-middle, and high based on
90 GBD result Website (<https://vizhub.healthdata.org/gbd-results/>). The 204 countries
91 were classified into different SDI regions based on their SDI scores: low SDI (0–
92 0.455), low-middle SDI (0.456–0.608), middle SDI (0.609–0.690), high-middle SDI
93 (0.690–0.805), and high SDI (0.806–1).¹⁷

94 2.3 Data Analysis

95 To avoid the potential impact of the COVID-19 pandemic^{18,19} on the results, we
96 limited our analysis to data from 1990 to 2019. Age-standardized rates (ASR) of
97 DALYs and deaths were used to compare the effect of metabolic-related stroke in
98 different years, sexes, regions, and age groups.²⁰ Unless otherwise specified, DALYs

99 and mortality rates were age-standardized and reported per 100,000 persons.

100 The calculation of time trend was based on the Average Annual Percent Change
101 (AAPC) indicator,²¹ AAPCs were calculated using the Joinpoint software (Version
102 5.2.0).²² For calculating the AAPC from 1990 to 2019, we selected 0 joinpoint index.

103 To determine the proportion of strokes attributable to various risk factors, we
104 utilized the GBD results to obtain DALYs and mortality rates for each risk factor and
105 calculated the proportion by dividing the value for each specific risk factor by the
106 total. Statistical analysis and graphical mapping were conducted using Excel 2020,
107 PowerPoint 2020, and R software (version 4.1.1), with packages ggplot2,²³ dplyr,²⁴
108 countrycode,²⁵ and scatterpie.²⁶ Confidence intervals were two-tailed at 95%, and
109 p-values < 0.05 were considered statistically significant.

110 **Results:**

111 3.1 Global Burden of Stroke Attributable to Metabolic Risks

112 In 2019, the age-standardized DALYs rate due to metabolic risk-related stroke
113 globally was 1321.94 (1104.76, 1507.9) and the mortality rate was 62.73 (52.26, 71.7),
114 both showing significant declines compared to 1990 (Table 1). From 1990 to 2019,
115 the AAPC for age-standardized DALYs was -1.70% (-1.81%, -1.58%), and for
116 mortality rate, the AAPC was -1.57% (-1.68%, -1.46%) (Table 1). Geographically, the
117 world was divided into 33 regions (Table 1). In 2019, Southeast Asia had the highest
118 age-standardized mortality rate (ASMR) of 116.21 (98.02, 133.41) and the highest
119 age-standardized DALYs rate of 2500.54 (2104.75, 2894.53) (Table 1). Except for
120 Southern Africa and Southern Sub-Saharan Africa, the other 31 regions showed
121 significant declines in both rates from 1990 to 2019, with AAPCs less than 0 (Table
122 1).

123 At the country level (Supplemental Table 2, Figure 1), the Republic of Nauru had
124 the highest rates in 1990 with an ASR of DALYs 4759.53 (3678.31, 6011.55) and the
125 Republic of Serbia had an ASMR of 252.39 (212.12, 290.61). In 2019, the highest
126 rates were again in the Republic of Nauru with a DALYs rate of 4527.26 (3412.25,
127 5837.06) and North Macedonia with a mortality rate of 240.36 (199.66, 277.11).
128 Singapore had the lowest ASMR in 2019 at 9.48 (7.39, 11.45), and the Swiss
129 Confederation had the lowest DALYs ASR at 234.72 (186.91, 277.78) (Supplemental
130 Table 2, Figure 2). A total of 170 countries showed significant declines in DALYs
131 rates, and 165 countries showed significant declines in mortality rates (Supplemental
132 Table 2, Figure 2). The Republic of Korea had the most pronounced decrease in
133 DALYs rate with an AAPC of -6.55% (-6.92%, -6.28%), and the Republic of Estonia
134 had the most pronounced decrease in mortality rate with an AAPC of -6.66% (-7.31%,
135 -6.01%). Conversely, the Kingdom of Lesotho exhibited the most significant increase
136 in both DALYs and mortality rates (Supplemental Table 2, Figure 2).

137 3.2 Impact of SDI on Stroke Burden Attributable to Metabolic Risks

138 We found that in 2019, regions with high SDI had significantly lower ASMR

139 (23.15 [18.51, 26.85]) and DALYs ASR (504.31 [417.8, 582.61]) compared to other
140 SDI regions (Table 1, Figure 2a). The AAPC was also the lowest in five SDI level
141 regions, with AAPC of mortality at -3.47% (-3.61%, -3.32%) and DALYs at -3.03%
142 (-3.17%, -2.89%) (Table 1). However, regardless of whether it was in 1990 or 2019,
143 the regions with the highest ASR of DALYs or mortality rates were not the low SDI
144 regions (Table 1). The regions with high-medium SDI exhibited a significant upward
145 trend in both DALYs and mortality rates from 1990 to 1994 (Figure 2 a, f).

146 Among 204 countries, we found that as SDI increased, both ASR of DALYs
147 (Figure 2d) and mortality rates (Figure 2i) in 2019 showed a significant decline. The
148 AAPC from 1990 to 2019 also displayed a downward trend with the SDI upward
149 (Figure 2e, j), and this decline was exponential; the higher the SDI, the more
150 pronounced the decrease (Figure 2d, e, i, j).

151 3.3 Gender and Age Heterogeneity in Stroke Burden Due to Metabolic Risks

152 Globally, the impact of metabolic risks on males was greater than on females
153 both on DALYs ASR and ASMR, and the trend of change was consistent for both
154 genders, with AAPCs being less than 0 (Table 1). Except for the low-middle SDI
155 region, where the ASMR of male was lower than that of female during 1990-1999,
156 male consistently had higher age-standardized mortality rates and DALYs rates in all
157 other SDI level regions and time periods during 1990-2019. (Figure 2b, c, g, h).

158 In different age groups, the stroke-related DALYs and deaths rates
159 predominantly affected individuals aged 75 and above, with a lesser impact on those
160 under 55. (Figure 3a). For the 20-24 age group, the AAPC from 1990 to 2019 was
161 greater than 0 for both DALYs 5.18 (4.59, 5.78) and mortality rates 5.84 (5.07, 6.61)
162 (Figure 3b), but the 2019 rates of DALYs 7.40 (0.52, 16.74) and deaths 0.08 (0.003,
163 0.18) was visible lower than other age groups (Figure 2a). All other age groups
164 showed significant declines (Figure 3b). Additionally, as the SDI level increased, the
165 percentage of DALYs or mortality rates among the population over 85 years old rose
166 compared to other age groups. In North America, high-income Asia Pacific, Western
167 Europe, and Australasia, individuals over 85 accounted for more than 70% of the

168 2019 mortality rate from metabolic-related strokes (Figure 3c).

169 3.4 Proportional Contribution of Risk Factors to Stroke

170 Comparing age-standardized mortality and DALYs due to the three primary risk
171 factors identified in the GBD study (Environmental/Occupational risks, Behavioral
172 risks, and Metabolic risks), we found that metabolic risks contributed the highest
173 percentage to stroke-related ASMR (50.82%) and DALYs (48.80%) globally in 2019
174 (Figure 4a), followed by Environmental/Occupational risks. The 1990 composition
175 ratio ranking was also similar to 2019 (Figure 4a). As SDI increased, the composition
176 ratio of DALYs ASR and ASMR due to Environmental/Occupational risks decreased,
177 while Behavioral and Metabolic risks increased (Figure 4a). In high SDI regions,
178 metabolic risks accounted for 58.59% of stroke-related deaths in 2019 (Figure 4a).
179 This trend was also observed across different genders (Figure 4b, c). For males, the
180 proportion of behavioral risks was higher than for females regardless of the data
181 collection year or outcome measure, while the proportion of metabolic and
182 environmental risks was lower for females (Figure 4d).

183 3.5 Burden Proportions of Stroke Types Attributable to Metabolic Risks

184 Based on the GBD study's four-level causes, stroke was categorized into
185 intracerebral hemorrhage, ischemic stroke, and subarachnoid hemorrhage.¹⁵ From
186 1990 to 2019, age-standardized mortality and DALYs rates for these stroke subtypes
187 attributable to metabolic factors showed a declining trend globally and across all SDI
188 regions (Figure 5a, b). The most significant decline in DALYs for ischemic stroke was
189 observed in high-middle SDI regions, dropping from 1478.48 (1276.71, 1648.34) in
190 1990 to 869.25 (727.48, 981.48) in 2019 (Figure 5a, b).

191 Globally in 2019, ischemic stroke due to metabolic risks had the highest ASMR
192 (34.67 [28.44, 39.67]) and DALYs rate (661.26 [554.38, 749.68]) among the three
193 subtypes, while subarachnoid hemorrhage had the lowest ASMR (2.41 [1.79, 3.03])
194 and DALYs rate (67.61 [49.87, 85.50]) (Figure 5a, b). With decreasing SDI levels
195 (from High to Low), the ASMR for intracerebral hemorrhage increased (Figure 5b). In
196 high SDI regions, the ASMR for ischemic stroke was 15.11 (12.20, 17.67) compared

197 to 6.44 (5.20, 7.66) for intracerebral hemorrhage, and the rates were 37.20 (29.70,
198 45.89) for ischemic stroke and 38.02 (29.19, 46.39) for intracerebral hemorrhage in
199 the low SDI regions (Figure 5b). Except for the slightly lower age-standardized
200 mortality rate for subarachnoid hemorrhage in high SDI regions for males (1.55 [1.13,
201 1.89]) compared to females (1.61 [1.16, 2.00]), age-standardized DALYs and
202 mortality rates were higher in males for all subtypes globally and across SDI regions
203 (Figure 5c, d). For both genders, the 10-55 and 50-74 age groups had the highest
204 DALYs and mortality rates due to intracerebral hemorrhage among the three subtypes
205 (Figure 5e, f). For those aged 75-84 and over 85, ischemic stroke was the leading
206 subtype contributing to DALYs and mortality rates (Figure 5c, d). Across all age
207 groups, subarachnoid hemorrhage had the lowest DALYs and mortality rates among
208 the three subtypes (Figure 5c, d).

209

210 **Discussion:**

211 Despite the similarities in global distribution patterns between metabolic risk
212 related stroke and stroke caused by all factors¹⁵—such as overall global levels and the
213 decreasing ASMR and DALY rates across the five SDI levels—there are notable
214 differences in how metabolic risk-related stroke is distributed by region and SDI
215 compared to stroke from all causes.¹⁵ For example, in 2019, the highest stroke
216 mortality rate from all causes was observed in the low SDI category, while the lowest
217 incidence of metabolic stroke occurred in the low-middle SDI category. Oceania had
218 the highest all-cause stroke-related DALYs in 2019, with a rate of 2,998.97 (2,432.79,
219 3,617.12). In contrast, Southeast Asia had the highest metabolic stroke-related DALYs,
220 with a rate of 2,500.54 (2,104.75, 2,894.53).

221 Southeast Asia had the highest ASMR for metabolic stroke among the 33 regions,
222 while Singapore, also located in Southeast Asia, had the lowest ASMR for metabolic
223 stroke in 2019. (Supplemental Table 2) This suggests that ethnicity and geographic
224 location may not be the primary factors influencing metabolic stroke mortality or
225 DALY rates.^{2 27} Instead, public health and healthcare systems, socioeconomic factors,
226 and lifestyle choices likely play a more significant role.^{28 29} From 1990 to 2019,
227 South Korea saw the most significant decline in the ASR of metabolic stroke DALYs,
228 while Estonia experienced the most notable decrease in mortality rates.³⁰ During this
229 period, South Korea's Gross Domestic Product (GDP) per capita rose from \$6,000 in
230 1990 to \$31,500 in 2019 from World Bank website. Similarly, Estonia underwent
231 significant socioeconomic improvements after gaining independence from the Soviet
232 Union in 1991, transforming into a high-income country with an efficient economic
233 and social system.³⁰

234 In contrast, sub-Saharan Africa and Southern Africa did not see significant
235 declines in metabolic stroke rates (Table 1). This could be due to metabolic risk
236 factors often being overlooked in African stroke patients, with the overall stroke
237 proportion not decreasing.^{31 32 33} Furthermore, Southern Africa has the highest
238 prevalence of metabolic syndrome in Africa.³⁴ Addressing metabolic risk factors is

239 likely the most effective way to mitigate the impact of strokes in Africa.^{29,35}
240 Lesotho, another Southern African country, saw the most significant increase in both
241 ASR of DALYs and deaths from 1990 to 2019 (Supplemental Table 2). Clinical
242 studies indicate that CVDRF is highly prevalent across all age and gender groups in
243 Lesotho,³⁶ highlighting the importance of strengthening preventive measures in
244 Lesotho and similar regions in Southern Africa. Overall, enhancing national economic
245 levels,³⁷ developing plans to protect against and manage metabolic risk factors,³⁸ and
246 individuals paying more attention to metabolic factors related to stroke³⁸ were crucial
247 for reducing DALYs and mortality rates caused by metabolic stroke.

248 As SDI increases, ASR of DALYs and mortality rates shown a similar
249 exponential decline, and the AAPC trend is consistent. This indicates that developing
250 countries with weak socioeconomic foundations face a greater challenge of a
251 sustained burden of metabolic stroke compared to developed nations.³⁹ In
252 high-income regions such as North America, high-income Asia Pacific, Western
253 Europe, and Australia, individuals aged 85 and above accounted for over 70% of
254 metabolic stroke mortality in 2019. These high-income countries face more severe
255 aging problems,⁴⁰ with the proportion of elderly people increasing annually.⁴⁰ As
256 people age, they accumulate metabolic-related diseases (such as diabetes,
257 hypertension, and high cholesterol)⁴¹, increasing their risk of stroke. Additionally, the
258 physiological frailty of individuals aged 85 and above results in poorer stroke
259 outcomes.⁴² These findings suggest the need to strengthen the prevention and
260 management of metabolic-related diseases in developing countries. In high-income
261 countries, should pay more attention to the issue of population aging.

262 Our study also found that, regardless of data collection year and outcome
263 measures, the proportion of behavior risk-related stroke is consistently higher in men
264 than in women, while the proportion of metabolic and environmental risk-related
265 stroke is lower in men. However, strokes related to these three risk factors result in
266 more severe DALYs and mortality rates in men. This may be due to men's insufficient
267 management of metabolic-related diseases (such as hypertension and diabetes),

268 combined with the additive effect of high behavioral risks,⁴³ leading to a higher risk
269 of stroke and more severe outcomes. Increasing men's awareness and management of
270 metabolic diseases, along with regular health screenings and early intervention, can
271 lower stroke incidence and severity.⁴⁴

272

273 **Limitations**

274 Despite the comprehensive nature of our analysis, several limitations should be
275 acknowledged. The reliance on secondary data sources may introduce reporting biases.
276 The variability in healthcare infrastructure and diagnostic capabilities across regions
277 could affect the accuracy of our findings.

278

279 **Conclusion**

280 Our study highlights significant global progress in reducing the stroke burden
281 attributable to metabolic risks, particularly in high SDI regions. However, persistent
282 disparities across different SDI levels, genders, and age groups necessitate continued
283 efforts to address these gaps. Comprehensive public health strategies targeting
284 metabolic risks, with a focus on high-risk populations, are essential to further mitigate
285 the global stroke burden.

286

287 **Funding:** National Natural Science Foundation of China Regional Science
288 Foundation Project (82060857); The Science and Technology Program of Jiangxi
289 Provincial Health and Health Commission (SKJP220219475).

290

291 **Disclosure:** The authors declare no conflict of interest.

292 Reference:

- 293 1. Katan M, Luft A. Global burden of stroke. Paper/Poster presented at: Seminars in
294 neurology; 2018;
- 295 2. Lackland DT, Roccella EJ, Deutsch AF, Fornage M, George MG, Howard G, Kissela
296 BM, Kittner SJ, Lichtman JH, Lisabeth LD. Factors influencing the decline in stroke
297 mortality: a statement from the American Heart Association/American Stroke
298 Association. *Stroke*. 2014;45:315-353.
- 299 3. Mukherjee D, Patil CG. Epidemiology and the global burden of stroke. *World*
300 *neurosurgery*. 2011;76:S85-S90.
- 301 4. Collaborators GS. Global, regional, and national burden of stroke and its risk factors,
302 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *The*
303 *Lancet Neurology*. 2021;20:795.
- 304 5. Guzik A, Bushnell C. Stroke epidemiology and risk factor management. *CONTINUUM:*
305 *Lifelong Learning in Neurology*. 2017;23:15-39.
- 306 6. Mosenzon O, Cheng AY, Rabinstein AA, Sacco S. Diabetes and stroke: what are the
307 connections? *Journal of Stroke*. 2023;25:26-38.
- 308 7. Yuan S, Huang X, Ma W, Yang R, Xu F, Han D, Huang T, Peng M, Xu A, Lyu J.
309 Associations of HDL-C/LDL-C with myocardial infarction, all-cause mortality,
310 haemorrhagic stroke and ischaemic stroke: a longitudinal study based on 384 093
311 participants from the UK Biobank. *Stroke and Vascular Neurology*. 2023;8.
- 312 8. Lawes CM, Bennett DA, Feigin VL, Rodgers A. Blood pressure and stroke: an
313 overview of published reviews. *Stroke*. 2004;35:776-785.

- 314 9. Ye J, Hu Y, Chen X, Yin Z, Yuan X, Huang L, Li K. Association between the
315 weight-adjusted waist index and stroke: a cross-sectional study. *BMC Public Health*.
316 2023;23:1689.
- 317 10. Jørgensen L, Engstad T, Jacobsen BK. Bone mineral density in acute stroke patients:
318 low bone mineral density may predict first stroke in women. *Stroke*. 2001;32:47-51.
- 319 11. Fabjan TH, Hojs R. Stroke and renal dysfunction. *European journal of internal*
320 *medicine*. 2014;25:18-24.
- 321 12. Murray CJ. The global burden of disease study at 30 years. *Nature medicine*.
322 2022;28:2019-2026.
- 323 13. Mensah GA, Fuster V, Murray CJ, Roth GA, Diseases GBoC, Collaborators R. Global
324 burden of cardiovascular diseases and risks, 1990-2022. *Journal of the American*
325 *College of Cardiology*. 2023;82:2350-2473.
- 326 14. Stevens GA, Alkema L, Black RE, Boerma JT, Collins GS, Ezzati M, Grove JT, Hogan
327 DR, Hogan MC, Horton R. Guidelines for accurate and transparent health estimates
328 reporting: the GATHER statement. *The Lancet*. 2016;388:e19-e23.
- 329 15. Feigin VL, Stark BA, Johnson CO, Roth GA, Bisignano C, Abady GG, Abbasifard M,
330 Abbasi-Kangevari M, Abd-Allah F, Abedi V. Global, regional, and national burden of
331 stroke and its risk factors, 1990–2019: a systematic analysis for the Global Burden of
332 Disease Study 2019. *The Lancet Neurology*. 2021;20:795-820.
- 333 16. IfH M. Evaluation. Global Burden of Disease Collaborative Network. Global Burden of
334 Disease Study 2016 (GBD 2016) Results. *Institute for Health Metrics and Evaluation*
335 *Seattle*. 2017.

- 336 17. Jiang X, Ling J, Xiong Q, Chen W, Zou L, Ling Z. Global, regional, and national burden
337 of atrial fibrillation/flutter related to metabolic risks over three decades: estimates from
338 the global burden of disease study 2019. *Eur Heart J Qual Care Clin Outcomes*. 2024.
339 doi: 10.1093/ehjqcco/qcae033
- 340 18. Bersano A, Kraemer M, Touzé E, Weber R, Alamowitch S, Sibon I, Pantoni L. Stroke
341 care during the COVID-19 pandemic: experience from three large European countries.
342 *European journal of neurology*. 2020;27:1794-1800.
- 343 19. Hoyer C, Ebert A, Huttner HB, Puetz V, Kallmünzer B, Barlinn K, Haverkamp C,
344 Harloff A, Brich J, Platten M. Acute stroke in times of the COVID-19 pandemic: a
345 multicenter study. *Stroke*. 2020;51:2224-2227.
- 346 20. Chen S, Liu M, Xie F. Global and national burden and trends of mortality and
347 disability-adjusted life years for silicosis, from 1990 to 2019: results from the Global
348 Burden of Disease study 2019. *BMC pulmonary medicine*. 2022;22:240.
- 349 21. Clegg LX, Hankey BF, Tiwari R, Feuer EJ, Edwards BK. Estimating average annual
350 per cent change in trend analysis. *Statistics in medicine*. 2009;28:3670-3682.
- 351 22. Kim HJ, Fay MP, Feuer EJ, Midthune DN. Permutation tests for joinpoint regression
352 with applications to cancer rates. *Statistics in medicine*. 2000;19:335-351.
- 353 23. Wickham H. ggplot2. *Wiley interdisciplinary reviews: computational statistics*.
354 2011;3:180-185.
- 355 24. Yarberry W, Yarberry W. Dplyr. *CRAN recipes: DPLYR, stringr, lubridate, and regex in*
356 *R*. 2021:1-58.
- 357 25. Arel-Bundock V, Enevoldsen N, Yetman C. countrycode: An R package to convert

- 358 country names and country codes. *Journal of Open Source Software*. 2018;3:848.
- 359 26. Yu G, Yu MG. Package'scatterpie'. *R Package Version*. 2018;1.
- 360 27. Kim AS, Cahill E, Cheng NT. Global stroke belt: geographic variation in stroke burden
361 worldwide. *Stroke*. 2015;46:3564-3570.
- 362 28. Lindmark A, Eriksson M, Darehed D. Socioeconomic status and stroke severity:
363 Understanding indirect effects via risk factors and stroke prevention using innovative
364 statistical methods for mediation analysis. *PLoS One*. 2022;17:e0270533.
- 365 29. Hankey GJ. Population impact of potentially modifiable risk factors for stroke. *Stroke*.
366 2020;51:719-728.
- 367 30. Kõrv L, Vibo R, Kõrv J. Declining first-ever stroke incidence rates in 15-to 54-year-old
368 residents of Tartu, Estonia, from 1991 to 2017. *International Journal of Stroke*.
369 2023;18:462-468.
- 370 31. Kengne AP, Anderson CS. The neglected burden of stroke in Sub-Saharan Africa.
371 *International Journal of Stroke*. 2006;1:180-190.
- 372 32. BeLue R, Okoror TA, Iwelunmor J, Taylor KD, Degboe AN, Agyemang C, Ogedegbe
373 G. An overview of cardiovascular risk factor burden in sub-Saharan African countries:
374 a socio-cultural perspective. *Globalization and health*. 2009;5:1-12.
- 375 33. Peer N, Baatiema L, Kengne A-P. Ischaemic heart disease, stroke, and their
376 cardiometabolic risk factors in Africa: current challenges and outlook for the future.
377 *Expert review of cardiovascular therapy*. 2021;19:129-140.
- 378 34. Faijer-Westerink HJ, Kengne AP, Meeks KA, Agyemang C. Prevalence of metabolic
379 syndrome in sub-Saharan Africa: A systematic review and meta-analysis. *Nutrition*,

- 380 *Metabolism and Cardiovascular Diseases*. 2020;30:547-565.
- 381 35. Akinyemi RO, Ovbiagele B, Adeniji OA, Sarfo FS, Abd-Allah F, Adoukonou T, Ogah
382 OS, Naidoo P, Damasceno A, Walker RW. Stroke in Africa: profile, progress,
383 prospects and priorities. *Nature Reviews Neurology*. 2021;17:634-656.
- 384 36. González Fernández L, Firima E, Gupta R, Sematle MaP, Khomolishoele M, Molulela
385 M, Bane M, Meli R, Tlahali M, Lee T. Prevalence and determinants of cardiovascular
386 risk factors in Lesotho: a population-based survey. *International Health*.
387 2024;16:313-324.
- 388 37. Owolabi MO, Thrift AG, Martins S, Johnson W, Pandian J, Abd-Allah F, Varghese C,
389 Mahal A, Yaria J, Phan HT. The state of stroke services across the globe: Report of
390 World Stroke Organization–World Health Organization surveys. *International Journal*
391 *of Stroke*. 2021;16:889-901.
- 392 38. Zhang F, Liu L, Zhang C, Ji S, Mei Z, Li T. Association of metabolic syndrome and its
393 components with risk of stroke recurrence and mortality: a meta-analysis. *Neurology*.
394 2021;97:e695-e705.
- 395 39. Inam M, Samad Z, Vaughan EM, Almas A, Hanif B, Minhas AM, Jarrar Z, Habib FZ,
396 Sheikh S, Zhu D. Global cardiovascular research: Gaps and opportunities. *Current*
397 *Cardiology Reports*. 2023;25:1831-1838.
- 398 40. Mitchell E, Walker R. Global ageing: successes, challenges and opportunities. *British*
399 *journal of hospital medicine*. 2020;81:1-9.
- 400 41. Mengozzi A, Pugliese NR, Chiriaco M, Masi S, Virdis A, Taddei S. Microvascular
401 ageing links metabolic disease to age-related disorders: the role of oxidative stress

- 402 and inflammation in promoting microvascular dysfunction. *Journal of Cardiovascular*
403 *Pharmacology*. 2021;78:S78-S87.
- 404 42. Kanai M, Noguchi M, Kubo H, Nozoe M, Kitano T, Izawa KP, Mase K, Shimada S.
405 Pre-stroke frailty and stroke severity in elderly patients with acute stroke. *Journal of*
406 *Stroke and Cerebrovascular Diseases*. 2020;29:105346.
- 407 43. Oh JE. Association between smoking status and metabolic syndrome in men. *The*
408 *Korean journal of obesity*. 2014;23:99-105.
- 409 44. Goldstein LB, Bushnell CD, Adams RJ, Appel LJ, Braun LT, Chaturvedi S, Creager
410 MA, Culebras A, Eckel RH, Hart RG. Guidelines for the primary prevention of stroke: a
411 guideline for healthcare professionals from the American Heart Association/American
412 Stroke Association. *Stroke*. 2011;42:517-584.
- 413
414

415 Table 1: The Global Distribution of Metabolic-related Stroke from 1990 to 2019.


Region	ASR of Deaths (per 100,000 persons)			ASR of DALYs (per 100,000 persons)		
	1990	2019	AAPC (%)	1990	2019	AAPC (%)
Global	98.05 (81.38, 111.82)	62.73 (52.26, 71.7)	-1.57 (-1.68, -1.46)	1984.33 (1641.39, 2284.5)	1321.94 (1104.76, 1507.9)	-1.70 (-1.81, -1.58)
Gender						
Female	92.41 (76.01, 105.37)	54.52 (45.02, 63.04)	-2.02 (-2.15, -1.89)	1816.30 (1495.85, 2091.62)	1115.70 (934.35, 1280.44)	-1.88 (-2.01, -1.77)
Male	103.85 (85.09, 121.23)	72.36 (58.32, 84.31)	-1.36 (-1.47, -1.25)	2164.03 (1760.47, 2525.56)	1550.15 (1276.80, 1795.45)	-1.27 (-1.39, -1.17)
SDI Level						
High SDI	59.92 (49.74, 67.59)	23.15 (18.51, 26.85)	-3.47 (-3.61, -3.32)	1151.72 (970.98, 1305.46)	504.31 (417.8, 582.61)	-3.03 (-3.17, -2.89)
High-middle SDI	132.95 (110.1, 151.1)	74.27 (60.54, 85.6)	-2.32 (-2.58, -2.06)	2508.28 (2089.24, 2859.93)	1455.28 (1190.78, 1664.52)	-2.21 (-2.48, -1.94)
Middle SDI	112.49 (90.39, 132.19)	79.62 (64.58, 92.54)	-1.25 (-1.38, -1.12)	2311.85 (1863.53, 2737.07)	1616.17 (1323.16, 1867.44)	-1.29 (-1.4, -1.19)
Low-middle SDI	91.42 (73.5, 106.87)	77.38 (65.18, 88.39)	-0.65 (-0.7, -0.61)	1991.79 (1600.25, 2324.2)	1661.69 (1396.19, 1905.33)	-0.71 (-0.76, -0.66)
Low SDI	94.03 (74.58, 111.7)	77.32 (63.24, 90.72)	-0.77 (-0.82, -0.71)	2073.62 (1628.83, 2476.56)	1660.92 (1356.02, 1953.27)	-0.88 (-0.94, -0.82)
Location						
Africa	108.88 (88.12, 127.26)	92.06 (76.27, 105.76)	-0.58 (-0.68, -0.48)	2292.61 (1849.11, 2673.82)	1886.79 (1567.5, 2178.47)	-0.69 (-0.79, -0.6)
Central Africa	114.35 (86.55, 141.93)	102.13 (79.21, 126.64)	-0.49 (-0.63, -0.35)	2479.81 (1869.75, 3099.59)	2146.35 (1676.75, 2665.89)	-0.6 (-0.75, -0.46)
Central Sub-Saharan Africa	118.85 (89.74, 149.94)	97.4 (74.22, 127.62)	-0.78 (-0.86, -0.7)	2570.82 (1913.87, 3276.18)	2031.16 (1536.57, 2635.3)	-0.91 (-0.99, -0.83)
Eastern Africa	98.5 (77.03, 118.57)	72.25 (56.51, 86.68)	-1.22 (-1.28, -1.16)	2165.11 (1677.52, 2624.4)	1551.89 (1242.26, 1855.31)	-1.32 (-1.38, -1.25)
Eastern Sub-Saharan Africa	97.06 (74.78, 118.74)	79.27 (62.51, 95.19)	-0.81 (-0.86, -0.76)	2168 (1672.15, 2682.88)	1737.23 (1376.82, 2077.23)	-0.89 (-0.94, -0.84)
Northern Africa	130.06 (101.48, 157.76)	101.16 (82.95, 119.17)	-0.68 (-0.75, -0.61)	2489.67 (1950.94, 3014.86)	1909.34 (1579.42, 2235.51)	-0.78 (-0.85, -0.71)
Southern Africa	88.72 (71.02, 104.85)	100.48 (82.97, 115.45)	0.36 (0.01, 0.7)	2005.63 (1631.81, 2354.15)	2129.77 (1774.3, 2468.35)	0.21 (-0.1, 0.53)
Southern Sub-Saharan Africa	71.16 (55.49, 84.77)	87.07 (72.94, 98.07)	0.42 (-0.1, 0.93)	1604.53 (1290, 1891.92)	1781.8 (1496.86, 2009.13)	0.23 (-0.25, 0.73)
Western Africa	110.16 (86.96, 130.51)	93.49 (78.52, 109.33)	-0.58 (-0.66, -0.5)	2334.51 (1846.25, 2749.69)	1941.65 (1631.68, 2277.26)	-0.66 (-0.74, -0.58)
Western Sub-Saharan Africa	109.32 (86.48, 129.64)	95.42 (79.56, 111.35)	-0.49 (-0.59, -0.39)	2320.66 (1829.62, 2736.63)	1988.59 (1649.66, 2323.94)	-0.56 (-0.67, -0.45)
America	47.55 (38.94, 57.16)	27.2 (22.29, 32.21)	-2.1 (-2.21, -2.0)	1012.22 (842.44, 1190.0)	585.66 (491.24, 680.08)	-2.05 (-2.16, -1.94)

	54.25)	31.2)	-1.98)	1154.55)	672.33)	-1.93)
North America	32.79 (26.87, 37.36)	20.16 (16.08, 23.56)	-2.08 (-2.36, -1.8)	670.46 (562.55, 762.88)	442.62 (363.78, 514.99)	-1.76 (-1.99, -1.54)
High-income North America	32.79 (26.87, 37.36)	20.17 (16.09, 23.56)	-2.08 (-2.36, -1.8)	670.56 (562.63, 763)	442.68 (363.82, 515.05)	-1.76 (-1.99, -1.54)
Caribbean	69.62 (57.18, 80.82)	50.39 (41.1, 59.57)	-1.07 (-1.14, -1)	1468.02 (1191.26, 1721.56)	1119.98 (895.05, 1348.33)	-0.9 (-0.99, -0.81)
Central Latin America	46.59 (38.89, 53.09)	26.42 (22.12, 30.1)	-2.09 (-2.22, -1.96)	939.46 (787.02, 1072.63)	554.05 (469.58, 635.03)	-1.98 (-2.13, -1.84)
Southern Latin America	70.09 (56.92, 81.57)	32.08 (26.01, 36.89)	-2.56 (-2.66, -2.45)	1425.89 (1134.55, 1664.44)	672.28 (555.71, 776.58)	-2.52 (-2.59, -2.45)
Tropical Latin America	95.79 (80.28, 108.95)	39.25 (31.97, 44.92)	-2.97 (-3.03, -2.9)	2015.14 (1681.99, 2287.76)	818.1 (679.74, 928.97)	-3.05 (-3.11, -2.99)
Andean Latin America	36.82 (28.53, 45.4)	26.12 (20.76, 32.09)	-1.45 (-1.67, -1.23)	797.63 (608.93, 982.96)	547.58 (435.34, 672.39)	-1.49 (-1.73, -1.25)
Asia	109.73 (88.35, 129)	76.5 (62.47, 88.43)	-1.34 (-1.47, -1.2)	2253.17 (1816.98, 2665.1)	1589.65 (1311.49, 1830.92)	-1.3 (-1.41, -1.18)
East Asia	141.48 (110.69, 172.74)	94.7 (73.8, 113.49)	-1.44 (-1.73, -1.16)	2748.98 (2135.5, 3387.8)	1821.95 (1418.66, 2190.71)	-1.48 (-1.72, -1.24)
Central Asia	114.4 (95.48, 129.75)	97.45 (82.98, 109.79)	-1.04 (-1.35, -0.74)	2498.46 (2093.95, 2834.32)	2004.81 (1713.88, 2251.1)	-1.29 (-1.61, -0.96)
South Asia	69.16 (54.64, 81.69)	58.14 (48.03, 66.95)	-0.74 (-0.83, -0.65)	1533.1 (1213.94, 1815.64)	1255.96 (1036.97, 1453.99)	-0.84 (-0.91, -0.76)
Southeast Asia	134.35 (107.96, 157.97)	116.21 (98.02, 133.41)	-0.5 (-0.66, -0.33)	2933.01 (2358.39, 3425.11)	2500.54 (2104.75, 2894.53)	-0.54 (-0.67, -0.41)
High-income Asia Pacific	76.49 (63.54, 86.97)	20.37 (15.91, 24.35)	-4.72 (-4.9, -4.54)	1495.48 (1244.17, 1699.49)	479.44 (392.86, 559.45)	-4.12 (-4.3, -3.93)
Western Pacific Region	124.25 (98.32, 148.17)	78.71 (62.23, 93.67)	-1.6 (-1.79, -1.42)	2456.18 (1948.33, 2964.63)	1599.54 (1261.08, 1900.22)	-1.5 (-1.66, -1.35)
Europe	103.08 (86.83, 115.32)	44.98 (37.38, 50.9)	-3.26 (-3.5, -3.02)	1923 (1646.08, 2136.3)	900.19 (768.77, 1005.92)	-3.08 (-3.34, -2.81)
Central Europe	157.31 (135.5, 175.52)	71.3 (60.64, 79.45)	-3.03 (-3.22, -2.85)	2890.46 (2506.8, 3205)	1315.56 (1132.13, 1462.32)	-3.06 (-3.25, -2.87)
Eastern Europe	162.13 (137.67, 181.88)	95.11 (80.44, 107.11)	-2.63 (-3.12, -2.13)	3021.29 (2609.33, 3341.03)	1873.22 (1603.11, 2084.29)	-2.4 (-2.9, -1.89)
Western Europe	65.76 (54.93, 73.9)	19.9 (15.97, 23.25)	-4.22 (-4.35, -4.09)	1135.49 (961.15, 1267.8)	382.11 (316.22, 443.03)	-3.87 (-4.01, -3.73)
Europe	103.08 (86.83, 115.32)	44.98 (37.38, 50.9)	-3.26 (-3.5, -3.02)	1923 (1646.08, 2136.3)	900.19 (768.77, 1005.92)	-3.08 (-3.34, -2.81)
European Region	103.22 (86.9, 115.51)	46.37 (38.57, 52.39)	-3.16 (-3.4, -2.92)	1938.93 (1659.85, 2155.18)	939.12 (802.32, 1049.54)	-2.96 (-3.22, -2.7)
Australasia	48.18 (39.93, 54.72)	16.56 (13.05, 19.8)	-3.84 (-3.96, -3.72)	842.69 (711, 949.56)	316.14 (256.74, 371.75)	-3.54 (-3.69, -3.39)

Oceania	106.91 (80.23, 134.98)	93.01 (69.28, 119.19)	-0.57 (-0.62, -0.51)	2370.38 (1771.18, 3007.55)	2107.87 (1590.05, 2703.96)	-0.47 (-0.53, -0.41)
---------	---------------------------	--------------------------	-------------------------	-------------------------------	-------------------------------	-------------------------

416 SDI: Socio-demographic Index; ASR: Age-Standardized Rate; AAPC: Average
417 Annual Percent Change. ASR of Deaths (per 100,000 persons) refers to the
418 Age-Standardized Rate of deaths per 100,000 persons; ASR of DALYs (per 100,000
419 persons) refers to the Age-Standardized Rate of DALYs per 100,000 persons; AAPC:
420 The Average Annual Percent Change between 1990 to 2019.
421

422 Figure 1: The Global Distribution of Metabolic-related Stroke from 1990 to 2019 in
423 Country Level.




424

425 ASR: Age-Standardized Rate; AAPC: Average Annual Percent Change. ASR of
426 Deaths (per 100,000 persons) refers to the Age-Standardized Rate of deaths per
427 100,000 persons; ASR of DALYs (per 100,000 persons) refers to the
428 Age-Standardized Rate of DALYs per 100,000 persons; AAPC: The Average Annual
429 Percent Change between 1990 to 2019 in 204 countries around the world.1990:
430 Age-standardized mortality and DALYs rates for metabolic-related stroke in 204
431 countries around the world in 1990; 2019: Age-standardized mortality and DALYs
432 rates for metabolic-related stroke in 204 countries around the world in 2019.


433

434 Figure 2:
 435 Age-Standardized Mortality Rates and DALYs for Metabolic-Related Stroke by SDI,
 436 1990-2019.



437
 438 ASR: Age-Standardized Rate; AAPC: Average Annual Percent Change. ASR of
 439 Deaths (per 100,000 persons) refers to the Age-Standardized Rate of deaths per
 440 100,000 persons; ASR of DALYs (per 100,000 persons) refers to the
 441 Age-Standardized Rate of DALYs per 100,000 persons; AAPC: The Average Annual
 442 Percent Change between 1990 to 2019.
 443 a, f: Age-standardized DALYs rate(a) and ASMR (f) for metabolic-related stroke by 5
 444 SDI level during 1990-2019 in both genders.
 445 b, g: Age-standardized DALYs rate(b) and ASMR (g) for metabolic-related stroke by
 446 5 SDI level during 1990-2019 in female.
 447 c, h: Age-standardized DALYs rate(c) and ASMR (h) for metabolic-related stroke by
 448 5 SDI level during 1990-2019 in male.
 449 d, i: Age-standardized DALYs rate (d) and ASMR (i) for metabolic-related stroke by
 450 SDI around 204 countries during 1990-2019 in both genders.
 451 e, j: AAPC of Age-standardized DALYs rate(e) and ASMR (j) for metabolic-related
 452 stroke by SDI around 204 countries during 1990-2019 in both genders.


453 Figure 3: Mortality Rates and DALYs for Metabolic-Related Stroke by Age,
 454 1990-2019



455

456 SDI: Socio-demographic Index; AAPC: Average Annual Percent Change. Death rate
 457 of per 100,000 persons refers to the Age group special rate of deaths per 100,000
 458 persons; DALYs rate of per 100,000 persons refers to the Age group special rate of
 459 DALYs per 100,000 persons; AAPC: The Average Annual Percent Change between
 460 1990 to 2019.

462 Figure 4: Mortality Rates and DALYs for Stroke by Risk Factors in 2019.



463

464 SDI: Socio-demographic Index; H SDI: High SDI; HM SDI: High-middle SDI; M
 465 SDI: Middle SDI; LM SDI: Low-middle SDI; L SDI: Low SDI; DALYs rate: the
 466 Age-Standardized Rate of DALYs per 100,000 persons; Deaths rate: the
 467 Age-Standardized Rate of deaths per 100,000 persons.


468 a: The proportion of contributions from different risk factors to stroke varies across
 469 years, SDI level, and outcome indicators.

470 b, c: The proportion of contributions from different risk factors to the ASMR of stroke
 471 in 2019 varies across gender (b: male; c: female) and SDI level.

472 d: The proportion of contributions from different risk factors to stroke varies across
 473 years, gender, and outcome indicators.

474

475 Figure 5: Mortality Rates and DALYs for Subgroup of Stroke in 1990 and 2019.



476

477 SDI: Socio-demographic Index; H SDI: High SDI; HM SDI: High-middle SDI; M
 478 SDI: Middle SDI; LM SDI: Low-middle SDI; L SDI: Low SDI; DALYs rate: the
 479 Age-Standardized Rate of DALYs per 100,000 persons; Deaths rate: the
 480 Age-Standardized Rate of deaths per 100,000 persons.



481 a, b: The ASR of DALYs (a) and deaths (b) of stroke different subtypes varies across
 482 years and SDI level.

483 c, d: The ASR of DALYs (c) and deaths (d) of stroke different subtypes in 2019 varies
 484 across gender and SDI level.



485 e, f: The age specific rate of DALYs (e) and deaths (f) of stroke different subtypes
 486 varies across gender and age group.

487



1990



2019
















AAPC












The ASRs of DALYs (per 100,000 persons)

The ASRs of Deaths (per 100,000 persons)









Intracerebral hemorrhage
 Ischemic Stroke
 Subarachnoid hemorrhage