

1 **Title:** Digital technology as a tool to provide social support to individuals with cancer in low-
2 and middle-income countries: a scoping review

3 **Running Title:** Digital technology for cancer-related social support in LMICs

4
5 **Authors:** Hallie Dau, PhD MPH^{1,2}, Fazila Kassam MSc HBSc³, Beth A. Payne PhD^{2,4}, Hana
6 Miller^{2,5}, Gina Ogilvie, MD MSc FCFP DrPH^{1,2,6}

7
8 **Affiliations:** ¹ School of Population and Public Health, University of British Columbia, Vancouver,
9 Canada

10 ² Women's Health Research Institute, Vancouver, Canada

11 ³ Faculty of Medicine, University of British Columbia, Vancouver, BC, Canada

12 ⁴ BC Children's Hospital Research Institute, Vancouver, BC, Canada

13 ⁵ McGill University, Montreal, Canada

14 ⁶ BC Centre for Disease Control, Vancouver, Canada

15

16

17 **Corresponding author:** Hallie Dau
18 BC Women's Hospital and Health Centre
19 Room H203J - 4500 Oak Street
20 Vancouver, BC, V6H 3N1
21 (778) 951-5291
22 Hallie.Dau@cw.bc.ca

23

24 **Funding:** None

25

26 **Conflict of interest:** None

27

28 **Author Contributions:** HD: conceptualization, methodology, writing – original draft preparation;
29 FK: analysis, writing – original draft preparation, writing – review and editing; BP: writing – review
30 and editing; HM analysis, writing – review and editing; GO: supervision

31 Abstract (300/300):

32
33 BACKGROUND: Cancer is a rising cause of morbidity and mortality in low- and middle-income
34 countries (LMICs). Individuals diagnosed with cancer in LMICs often have limited access to cancer
35 prevention, diagnosis, and treatment services. Social support after a cancer diagnosis is associated with
36 positive health outcomes in the long term. Digital technologies, such as the Internet and mobile
37 phones, could be used to provide support to individuals with cancer in a more accessible way. This
38 scoping review aims to understand how digital technology is currently being utilized by individuals
39 with cancer for social support in LMICs.

40
41 METHODS: Four electronic databases were searched up to June 2024 to identify studies that reported
42 on the use of digital technology for cancer social support in LMICs. Articles were included if they
43 were published in English, included adults diagnosed with any type of cancer, and reported the use of
44 digital technology for social support. Study characteristics, population demographics, and
45 technological interventions reported were extracted.

46
47 RESULTS: In all, 15 articles from 12 studies were included in the scoping review. The results were
48 centralized in four countries, and the most common cancer type reported was breast. Online health
49 communities, Internet-based resources, mobile applications, and telecommunication were the four
50 digital technologies reported. Overall, the articles demonstrated that the use of digital technology for
51 social support can be beneficial for individuals diagnosed with cancer in LMICs.

52
53 CONCLUSION: There is a limited understanding of how digital technology can be used to support
54 individuals with cancer in LMICs. Future research is needed to explore how digital technology can be
55 utilized by underrepresented regions to offer avenues of support for regionally common cancer types

56 such as cervical. Fundamentally, this scoping review highlights the need for additional research on the
57 use of digital technology to support individuals with a cancer diagnosis in LMICs.

58
59 Keywords: cancer, digital, technology, low- and middle-income country; support

60
61
62

63 INTRODUCTION

64 Cancer is rising within low- and middle-income countries (LMICs) with the number of new cases
65 annually in LMICs rapidly increasing. By 2030, it is anticipated that three-quarters of global cancer-
66 related deaths will occur in LMICs.(1) In 2022 the Lancet Oncology Commission published a series
67 highlighting the urgent need to focus on the increasing rates of cancer in Africa, an area that has been
68 historically understudied.(2) The increase in cancer rates in Africa and similar regions can be attributed
69 to urbanization, increased exposure to carcinogenic risk factors, an aging population, sedentary
70 lifestyles, and limited access to cancer prevention, diagnosis, and treatment services.(2, 3) The burden
71 of cancer in LMICs is a major public health challenge that requires urgent attention and action. Efforts
72 to address this challenge should focus on improving cancer prevention, early detection, access to
73 quality care, and risk factor reduction.

74 An opportunity to increase the quality of cancer care lies in sources of global connectivity. As of 2022,
75 82% of people in low-middle income countries have access to and use the internet.(4) More
76 importantly, three-quarters of the world's population owns a mobile phone - 49% of people in low-
77 income countries and 65% of individuals in lower-middle-income countries.(4) With this, the rise of
78 social media has allowed individuals to connect more easily with one another across the globe. As of
79 2022, Facebook (2.96 billion users) is the most popular social network followed by YouTube (2.51
80 billion users) and WhatsApp (2 billion users).(5) The opportunities for connection and healthcare
81 access offered by technology should be leveraged to increase access to essential health services such
82 as cancer support. The utilization of technology for cancer support is aligned with the World Health
83 Organization's prioritization of self-care interventions. As there is currently a global healthcare worker
84 shortage, self-care interventions, particularly those based in digital technology, allow individuals to

85 actively engage in advancing their own health and well-being in a convenient and cost-effective manner
86 that reduces the strain on the healthcare system.(6)

87 Prior research has demonstrated that social support soon after a cancer diagnosis is associated with
88 favorable health outcomes for individuals with cancer in the long term.(7, 8) Additionally, social
89 support positively influences a patient's help-seeking behavior.(9) Unfortunately, in many LMICs,
90 social support is often lacking due to stigma related to diagnosis, poor health system infrastructure,
91 and gender discrimination.(9) Digital technologies such as mobile phones and the internet offer the
92 opportunity for cancer patients in LMICs to receive social support in a more accessible way. Prior
93 research has demonstrated that online cancer communities can decrease psychological distress,(10, 11)
94 improve support,(12, 13) and create feelings of empowerment.(14-16) To date, there is a large scale
95 of evidence on the benefits of digital support for individuals with cancer.(17-20) However, there is
96 limited knowledge about how social support interventions could be applied to individuals with cancer
97 in LMICs. It is important to better understand how digital technologies can improve access to social
98 support and ultimately improve cancer outcomes in LMICs. As the rates of cancer continue to rise in
99 LMICs, the urgent need to address the lack of comprehensive cancer social support programs is
100 amplified. The goal of this scoping review is to understand how digital technology is currently being
101 utilized by individuals with cancer for social support in LMICs.

102 METHODS

103 *Inclusion & exclusion criteria*

104 Articles were included in the review if they (i) were published in English; (ii) were conducted in a
105 LMIC (as defined by to the World Bank criteria);(21) (iv) included adults (≥ 18 years of age) diagnosed
106 with any type of cancer; and (v) reported the use of digital technology for cancer social support.

107 Articles were excluded from the review if they (i) focused solely on children; and (ii) were review
108 articles.

109 *Search strategy & extraction*

110 The electronic databases, MEDLINE, PsychInfo, CINAHL, and CABI Global Health were searched
111 up to June 2024 to identify eligible studies. Search terms are provided in **Supplementary 1**. Covidence
112 systematic review software(22) was used to screen and extract all data from the included studies. Two
113 reviewers independently screened studies and one author extracted data from included studies. Studies
114 were first screened by title and abstract, subsequently, included studies underwent full-text review.
115 The relevant data was extracted from the final included studies. Disagreements were discussed by the
116 study team and resolved by reaching consensus.

117 The study population characteristics extracted by the authors included age, gender, education status,
118 employment status, marital status, income, cancer stage, cancer type, and treatment type, when
119 applicable. In addition, the type of digital technology utilized was also extracted. Following extraction,
120 the data was grouped by the type of digital technology reported. The digital technology was compared
121 and contrasted with regards to its usability and reported outcomes.

122 Social support was based on Cooke et al. and defined as “any emotional, informational, instrumental
123 or appraisal support given to cancer survivors or patients to help them cope with biological,
124 psychological or social stressors.”(23) The definition for digital health technology is based on the
125 World Health Organization’s definition for digital health which defines it as “the use of information
126 and communications technology in support of health and health-related fields.”(24) Digital health
127 technology includes web-based resources, mobile applications, social media platforms, general internet
128 use, and telecommunication.

129 RESULTS

130 In all, 7848 original articles were identified through the search strategy and 15 were included in the
131 final review (**Figure 1**). Three articles were from the same larger study and were combined into one
132 study, resulting in 12 unique studies.(25-27) **Table 1** summarizes the characteristics of the included
133 studies. The majority of included studies were either cross-sectional(28-33) (n=6) in design or a
134 randomized control trial(25-27, 34-36) (n=4). All but one article(26) used quantitative methods.
135 Among the 12 unique studies included in this review, only four countries were studied: China(37) (25-
136 27, 29, 32-34, 36, 38) (n=8), Iran(30, 35) (n=2), Kenya(31) (n=1), and Serbia(28) (n=1). Breast(25-27,
137 29, 31, 37, 38) (n=5) was the most common cancer studied.

138 There were four primary types of digital technologies reported for cancer social support in the 15
139 included studies: online health communities, Internet-based resources, mobile applications, and
140 telecommunication.

141 Cross-sectional studies by Zhou et al. and Lei et al. investigated the role of online health communities
142 (OHCs). Zhou et al. conducted a cross-sectional survey among 162 participants engaged in OHCs in
143 China. They found that e-health literacy was positively associated with the informational and
144 emotional support a cancer survivor received by actively interacting in the OHC.(33) Lei et al. (2021)
145 conducted a social network analysis of user activity and user-generated content in three forums and
146 found that while a few members of the community are very well connected with many other members,
147 most other members have fewer connections. Salient topics across all three forums were related to
148 disease treatment, examination, diagnosis, social life, and healthy behavior change.

149 Three studies reported on Internet-based resources for cancer social support. The first, Shaikh et al.,
150 developed a website to address the unmet needs, including social support, for women with metastatic
151 breast cancer in Kenya.(31) Over a two-year period, the website received 7,864 unique visitors. Among

152 the 114 women with metastatic breast cancer they interviewed during their needs assessment, 90.4%
153 had moderate to high supportive care needs, with psychological being the highest (63%). Given the
154 high need for social support, the authors concluded that interactive Internet-based resources could be
155 useful for women living with metastatic breast cancer in countries with limited health services.(31)
156 Salajegheh et al.(30) and Wang(32) were both cross-sectional studies that surveyed individuals with
157 cancer about their Internet usage. Salajegheh et al. reported that among the 350 individuals with cancer
158 that they surveyed, the Internet was one of the least likely ways individuals sought out information.(30)
159 Wang surveyed pancreatic cancer internet users and found that social media use has a positive impact
160 on mental health, namely anxiety and depression.(32)

161 Five unique studies, all conducted in China, reported on the use of mobile applications for cancer
162 support. (26, 27, 34, 36, 38, 39) Chen et al, Wang et al., and Wu et al. evaluated ways WeChat, a Chinese
163 social media and messaging application, could be used to help support individuals with cancer in
164 China.(36-38) All three interventions provided information and support through the mobile
165 application. Wang et al. included the WeChat platform as well as a micro-website developed by the
166 researchers,(36) Wu et al. recruited nurses to lead and interact with participants through the platform,
167 (38) and Chen et al. developed an interactive platform.(37) At the end of the study period, both Wang
168 et al. and Wu et al. reported that those in the WeChat group had higher psychological well-being than
169 those in the control.(36, 38) Additionally, Wang et al. and Chen et al. reported improved quality of
170 life by the end of the study period.(36, 37) For example, in Wang et al. the quality of life score (range
171 0-100) improved from 51.1 in the pre-intervention to 63.1 in the post-intervention.(36) Wu et al.
172 reported that those who received the intervention reported a higher level of social support outside of
173 family.(38)

174 Both Zhu et al. (25-27) and Lin et al.(34) measured participant’s self-efficacy after using a mobile
175 application created by the study team. Both applications include a discussion forum for participants
176 to engage in. Zhu et al. found that the program improved emotional well-being,(26) self-efficacy,(27)
177 quality of life, (27) and social support among individuals with breast cancer. (27) Lin et al. found that
178 individuals with gynecological cancer in the intervention group had lower levels of illness uncertainty
179 and higher quality of life levels.(34)

180 Two studies reported on the use of telecommunications.(28, 30, 35) Klikovac established a telephone
181 service to provide psychological support to cancer patients of all types and stages, as well as their
182 family members in Serbia. They found that most of the calls were regarding emotional support for
183 sadness, hopelessness, depression and informational support for disease prognosis, treatment, and
184 side effects of treatment.(28) Rahimi et al. conducted a double-blind randomized control trial to
185 explore how peer support via telephone and social networks impacts colorectal cancer patients. They
186 reported that participants who received informational, emotional, and appraisal support from their
187 peers via telephone two times a week and virtual social networks three times a week had overall higher
188 well-being than the control group.(35)

189 DISCUSSION

190 In all, 15 articles and 12 studies were included in this scoping review. The majority of the studies were
191 conducted in China. Online health communities, Internet-based resources, mobile applications, and
192 telecommunication were the four primary types of digital technology utilized for cancer social support.
193 Overall, the review found that the use of digital technology for social support can be beneficial for
194 individuals diagnosed with cancer in LMICs. More specifically, we found that digital technology can
195 improve quality of life, reduce anxiety and depression, and allow individuals to connect with other
196 individuals diagnosed with cancer.

197 Only four countries were represented in this review. The majority of the included studies were from
198 China,(37) (25-27, 29, 32-34, 36, 38) a country that has the most internet users worldwide,(40) as well
199 as the second highest gross-domestic product globally.(41) There was only one study from the African
200 region (Kenya)(31) and no research from South America and Central Asia. This likely is because cancer
201 and other non-communicable diseases have largely been understudied in LMICs with the primary
202 focus being on communicable diseases.(2, 42) As such, there is a gap in understanding how digital
203 technology can be used effectively to provide social support for cancer patients across the broad
204 category of LMICs. It is important to study this topic more widely since many LMICs are similar in
205 that they lack robust healthcare systems that can adequately care for all the physical and social needs
206 of individuals diagnosed with cancer. (1, 3, 43) More widespread and varied research on digital
207 platforms can better lead to the adaptation and adoption of programs across more resource-limited
208 countries. Social support through digital technology could reduce the overall burden of care on the
209 local healthcare system.

210 Similar to a deficiency in the regions studied, there is a limited understanding of how digital technology
211 can support various cancer types. The majority of the included studies reviewed breast cancer.
212 However, the top five types of cancer in LMICs are breast, cervical, liver, colorectal, and lip/oral.(44)
213 Given these cancers differ in their treatment of care, the way in which technology can be used for
214 social support may differ as well. Evidently, cancers associated with significant morbidity and mortality
215 in LMICs such as cervical(45) have not been the focus of digital health interventions and thus, current
216 evidence may not be applicable to all cancer patients from these regions. Therefore, there needs to be
217 a greater understanding of how digital technology can best provide social support for not only a variety
218 of cancer types, but those most common to LMICs.

219 The results of this review demonstrate that despite the surging cases of cancer in LMICs, there has
220 been limited exploration of the opportunities to provide social support to individuals with cancer
221 using digital technologies. More specifically, there is a need to better understand how existing and
222 commonly utilized technological interventions can be implemented for cancer social support in
223 LMICs. To date, the majority of research has focused on mobile applications followed by general
224 Internet use. This is only a small fraction of what could be harnessed to support individuals with
225 cancer in LMICs. For example, with regards to social media, WeChat is the only platform that has
226 been studied for digital social support for individuals with cancer in LMICs.(38) However, WeChat is
227 designed primarily for Chinese users and as such the findings are not applicable to other countries.
228 Consequently, there is an urgent need for research on how the more commonly used and growing
229 platforms like WhatsApp, Facebook, and YouTube can be adapted for digital social support for cancer
230 patients in LMICs. For example, research from the United States and Australia has shown that social
231 media can provide positive social support benefits for individuals diagnosed with cancer(46, 47) and
232 that there is an interest in it, particularly among young adults with cancer.(47-49) Replicating these
233 findings in an LMIC setting should be a research priority.

234 This scoping review was strengthened by its comprehensive search strategy that, in turn, identified
235 multiple gaps in relation to the use of digital technology for social support for individuals diagnosed
236 with cancer in LMICs. Additionally, the review adhered to best practices to reduce bias in results and
237 included two authors to review potential articles. This review was limited in that the authors only
238 searched for studies published in English. As such, publication bias may have occurred.

239 Furthermore, the authors did not consider unpublished or grey literature which may have provided
240 additional results.

241 Despite the vast advancements in technology, there is limited evidence for its use to deliver social
242 support to cancer patients in LMICs. Only 12 studies were identified in this scoping review, and they
243 were limited to four countries, the majority being China. Future research should consider including
244 more nations in underrepresented regions, cancer types that carry the greatest burden in LMICs, and
245 widely used and acceptable digital platforms. Furthermore, longitudinal studies are needed to
246 determine how long the benefits of digital technology last after a diagnosis. As individuals continue to
247 spend increasing amounts of time in the digital space, it is important for the cancer community to
248 focus their attention on harnessing this technology for better support and overall care. This in turn
249 will help reduce the burden of care on local healthcare systems and lead to more positive health
250 outcomes for individuals diagnosed.
251

REFERENCES

- 252
253
- 254 1. Pramesh C, Badwe RA, Bhoo-Pathy N, Booth CM, Chinnaswamy G, Dare AJ, et al.
255 Priorities for cancer research in low-and middle-income countries: a global perspective. *Nature*
256 *Medicine*. 2022;28(4):649-57.
 - 257 2. Ngwa W, Addai BW, Adewole I, Ainsworth V, Alaro J, Alatise OI, et al. Cancer in sub-
258 Saharan Africa: a Lancet Oncology Commission. *The Lancet Oncology*. 2022;23(6):e251-e312.
 - 259 3. Shah SC, Kayamba V, Peek Jr RM, Heimbürger D. Cancer control in low-and middle-
260 income countries: is it time to consider screening? *Journal of global oncology*. 2019;5:1-8.
 - 261 4. International Telecommunication Union. *Measuring digital development: Facts and*
262 *Figures 2022*. 2022.
 - 263 5. Statista. Most popular social networks worldwide as of January 2023, ranked by
264 number of monthly active users 2023 [Available from:
265 <https://www.statista.com/statistics/272014/global-social-networks-ranked-by-number-of-users/>.
 - 266 6. World Health Organization. Self-care interventions for health 2022 [Available from:
267 <https://www.who.int/news-room/fact-sheets/detail/self-care-health-interventions>.
 - 268 7. Ruiz-Rodríguez I, Hombrados-Mendieta I, Melguizo-Garín A, Martos-Méndez MJ. The
269 Importance of Social Support, Optimism, and Resilience on the Quality of Life of Cancer
270 Patients. *Frontiers in Psychology*. 2022:634.
 - 271 8. Schroevers MJ, Helgeson VS, Sanderman R, Ranchor AV. Type of social support matters
272 for prediction of posttraumatic growth among cancer survivors. *Psycho-Oncology*.
273 2010;19(1):46-53.
 - 274 9. McCutchan G, Weiss B, Quinn-Scoggins H, Dao A, Downs T, Deng Y, et al.
275 Psychosocial influences on help-seeking behaviour for cancer in low-income and lower middle-
276 income countries: a mixed-methods systematic review. *BMJ global health*. 2021;6(2):e004213.
 - 277 10. Rodgers S, Chen Q. Internet Community Group Participation: Psychosocial Benefits for
278 Women with Breast Cancer. *Journal of Computer-Mediated Communication*. 2017;10(4).
 - 279 11. Adikari A, de Silva D, Ranasinghe WKB, Bandaragoda T, Alahakoon O, Persad R, et al.
280 Can online support groups address psychological morbidity of cancer patients? An artificial
281 intelligence based investigation of prostate cancer trajectories. *PLOS ONE*.
282 2020;15(3):e0229361.
 - 283 12. Sillence E. Giving and receiving peer advice in an online breast cancer support group.
284 *Cyberpsychology, Behavior, and Social Networking*. 2013;16(6):480-5.
 - 285 13. van Eenbergen MC, van de Poll-Franse LV, Krahmer E, Verberne S, Mols F. Analysis of
286 Content Shared in Online Cancer Communities: Systematic Review. *JMIR Cancer*. 2018;4(1):e6-
287 e.
 - 288 14. Tjørnhøj-Thomsen T. Online interaction. Effects of storytelling in an internet breast
289 cancer support group. 2008.
 - 290 15. van Uden-Kraan CF, Drossaert CHC, Taal E, Shaw BR, Seydel ER, van de Laar MAFJ.
291 Empowering Processes and Outcomes of Participation in Online Support Groups for Patients
292 With Breast Cancer, Arthritis, or Fibromyalgia. *Qual Health Res*. 2008;18(3):405-17.
 - 293 16. Sharf BF. Communicating Breast Cancer On-Line: Support and Empowerment on the
294 Internet. *Women & Health*. 1997;26(1):65-84.
 - 295 17. Pyle D, Perry A, Lamont-Mills A, Tehan G, Chambers SK. A scoping review of the
296 characteristics and benefits of online prostate cancer communities. *Psycho-Oncology*.
297 2021;30(5):659-68.

- 298 18. Setoyama Y, Yamazaki Y, Namayama K. Benefits of peer support in online Japanese
299 breast cancer communities: differences between lurkers and posters. *Journal of medical Internet*
300 *research*. 2011;13(4):e122.
- 301 19. Chung JE. Social Networking in Online Support Groups for Health: How Online Social
302 Networking Benefits Patients. *Journal of Health Communication*. 2014;19(6):639-59.
- 303 20. Vilhauer RP. Perceived Benefits of Online Support Groups for Women with Metastatic
304 Breast Cancer. *Women & Health*. 2009;49(5):381-404.
- 305 21. World Bank. Low & middle income | Data 2020 [Available from:
306 <https://data.worldbank.org/country/XO>].
- 307 22. Covidence systematic review software. Melbourne, Australia: Veritas Health Innovation;
308 2021.
- 309 23. Cooke BD, Rossmann MM, McCubbin HI, Patterson JM. Examining the definition and
310 assessment of social support: A resource for individuals and families. *Family relations*.
311 1988;211-6.
- 312 24. World Health Organization. Global diffusion of eHealth: making universal health
313 coverage achievable: report of the third global survey on eHealth: World Health Organization;
314 2017.
- 315 25. Zhu H, Chen X, Yang J, Wu Q, Zhu J, Chan SW-C. Mobile breast cancer e-support
316 program for Chinese women with breast cancer undergoing chemotherapy (part 3): Secondary
317 data analysis. *JMIR mHealth and uHealth*. 2020;8(9).
- 318 26. Zhu J, Ebert L, Guo D, Yang S, Han Q, Chan SW-C. Mobile Breast Cancer e-Support
319 Program for Chinese Women With Breast Cancer Undergoing Chemotherapy (Part 1):
320 Qualitative Study of Women's Perceptions. *JMIR mHealth and uHealth*. 2018;6(4):e85.
- 321 27. Zhu J, Ebert L, Liu X, Wei D, Chan SW-C. Mobile Breast Cancer e-Support Program for
322 Chinese Women With Breast Cancer Undergoing Chemotherapy (Part 2): Multicenter
323 Randomized Controlled Trial. *JMIR mHealth and uHealth*. 2018;6(4):e104.
- 324 28. Klikovac T. The First Telephone Line for the Psychological Support to Oncological
325 Patients and Their Family Members in Serbia. *Srpski arhiv za celokupno lekarstvo*. 2015;143(11-
326 12):726-30.
- 327 29. Lei Y, Xu S, Zhou L. User Behaviors and User-Generated Content in Chinese Online
328 Health Communities: Comparative Study. *Journal of medical Internet research*.
329 2021;23(12):e19183.
- 330 30. Salajegheh M, Moazed V, Janati Z, Esmaeeli K. Studying health information behaviors
331 of people with cancer. *International Archives of Health Sciences*. 2020;7(1):31-6.
- 332 31. Shaikh AJ, Dhillon N, Shah J, Kathomi C, Kiragu A, Asirwa FC, et al. Supporting
333 Kenyan women with advanced breast cancer through a network and assessing their needs and
334 quality of life. *Supportive care in cancer : official journal of the Multinational Association of*
335 *Supportive Care in Cancer*. 2022;30(2):1557-67.
- 336 32. Wang Y, Bao S, Chen Y-b. How does social media use influence the mental health of
337 pancreatic cancer patients: a chain mediating effect of online social support and psychological
338 resilience. *Frontiers in Public Health*. 2023;11(June).
- 339 33. Zhou J, Wang C. Improving cancer survivors' e-health literacy via online health
340 communities (OHCs): A social support perspective. *Journal of Cancer Survivorship*.
341 2020;14(2):244-52.

- 342 34. Lin H, Ye M, Lin Y, Chen F, Chan S, Cai H, et al. Mobile App for Gynecologic Cancer
343 Support for Patients With Gynecologic Cancer Receiving Chemotherapy in China: Multicenter
344 Randomized Controlled Trial. *J Med Internet Res*. 2023;25:e49939.
- 345 35. Rahimi M, Mahdizadeh M, Chamanzari H, Mahdizadeh S-M. The Effect of Peer Support
346 with Telecommunication on Subjective Well-being in Colorectal Patients: A Randomized
347 Controlled Clinical Trial. *International Journal of Community Based Nursing & Midwifery*.
348 2021;9(2):127-38.
- 349 36. Wang Y, Yang L, Xu W, Wang C, Chen J, Wu Y, et al. Effects of a WeChat-based
350 multimodal psychoeducational intervention on psychological well-being and quality of life in
351 acute leukaemia patients in China: a randomised controlled trial. *Journal of cancer survivorship* :
352 research and practice. 2022;16(6):1461-77.
- 353 37. Chen X, Chaimongkol N, Hengudomsub P. Effects of a Phone-Based Support Program
354 for Women With Breast Cancer Undergoing Chemotherapy: A Pilot Study. *SAGE Open Nurs*.
355 2024;10:23779608241231176.
- 356 38. Wu Q, Kue J, Zhu X, Yin X, Jiang J, Chen J, et al. Effects of Nurse-Led Support Via
357 WeChat, a Smartphone Application, for Breast Cancer Patients After Surgery: A Quasi-
358 Experimental Study. *Telemedicine journal and e-health : the official journal of the American*
359 *Telemedicine Association*. 2020;26(2):226-34.
- 360 39. Zhu J, Ebert L, Zhimin X, Qu S, Chan SW-C, Zhu J, et al. Development of a mobile
361 application of Breast Cancer e-Support program for women with breast cancer undergoing
362 chemotherapy. *Technology & Health Care*. 2017;25(2):377-82.
- 363 40. Statista. Countries with the largest digital populations in the world as of January 2023
364 2023 [Available from: [https://www.statista.com/statistics/262966/number-of-internet-users-in-](https://www.statista.com/statistics/262966/number-of-internet-users-in-selected-countries/)
365 [selected-countries/](https://www.statista.com/statistics/262966/number-of-internet-users-in-selected-countries/)].
- 366 41. The World Bank. GDP (current US\$) 2024 [Available from:
367 https://data.worldbank.org/indicator/NY.GDP.MKTP.CD?most_recent_value_desc=true].
- 368 42. Rubagumya F, Carson L, Mushonga M, Manirakiza A, Murenzi G, Abdihamid O, et al.
369 An analysis of the African cancer research ecosystem: tackling disparities. *BMJ Glob Health*.
370 2023;8(2).
- 371 43. Organization WH. Tracking universal health coverage: 2017 global monitoring report.
372 2017.
- 373 44. International Agency for Research on Cancer. Low middle income. World Health
374 Organization,; 2021.
- 375 45. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global
376 cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36
377 cancers in 185 countries. *CA: a cancer journal for clinicians*. 2021;71(3):209-49.
- 378 46. Kashian N, Jacobson S. Factors of Engagement and Patient-Reported Outcomes in a
379 Stage IV Breast Cancer Facebook Group. *Health Communication*. 2020;35(1):75-82.
- 380 47. Lazard AJ, Collins MKR, Hedrick A, Varma T, Love B, Valle CG, et al. Using Social
381 Media for Peer-to-Peer Cancer Support: Interviews With Young Adults With Cancer. *JMIR*
382 *Cancer*. 2021;7(3):e28234.
- 383 48. Donovan E, Martin SR, Seidman LC, Zeltzer LK, Cousineau TM, Payne LA, et al. The
384 Role of Social Media in Providing Support from Friends for Adolescent and Young Adult
385 (AYA) Patients and Survivors of Sarcoma: Perspectives of AYA, Parents, and Providers. *J*
386 *Adolesc Young Adult Oncol*. 2021;10(6):720-5.

387 49. McNeil R, Egdsdal M, Drew S, McCarthy MC, Sawyer SM. The changing nature of social
388 support for adolescents and young adults with cancer. *European Journal of Oncology Nursing*.
389 2019;43:101667.
390

Table 1. Characteristics of included studies

Author	Year	Country	Study Type	Sample Size	Cancer Type	Digital technology utilized
Chen et al.(37)	2024	China	Pre-test-post-test	20	Breast	Mobile application
Klikovac(28)	2015	Serbia	Cross-sectional	2,748	Multiple cancers	Telephone
Lei et al.(29)	2021	China	Cross-sectional	Not reported	Lung, Breast	Online health community
Lin et al.(34)	2023	China	Randomized control trial	168	Gynecologic	Mobile application
Rahimi et al.(35)	2021	Iran	Randomized control trial	60	Colorectal	Telephone and virtual social networks
Salajegheh et al.(30)	2020	Iran	Cross-sectional	350	Multiple cancers	Internet, social networks, telephone, satellite, radio
Shaikh et al.(31)	2021	Kenya	Cross-sectional	114	Breast	Internet
Wang(36)	2022	China	Randomized control trial	72	Leukemia	Mobile application
Wang(32)	2023	China	Cross-sectional	425	Pancreatic	Internet, social networks
Wu et al.(38)	2020	China	Quasi-experimental	60	Breast	Mobile application
Zhou et al.(33)	2020	China	Cross-sectional	162	Multiple cancers	Online health community
Zhu et al.(25-27)	2018	China	Qualitative, Randomized control trial	13 (qualitative) 114 (quantitative)	Breast	Mobile application

391 Figure 1. PRISMA flow diagram

392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427

