1 2

Documented transboundary transmission of mpox between the Central African Republic and the Democratic Republic of the Congo

3

Emmanuel Hasivirwe Vakaniaki^{1,2}, Eddy Kinganda-Lusamaki^{1,3,4}, Sydney Merritt⁵, Francois 4 Kasongo¹, Emile Malembi⁶, Lygie Lunyanga¹, Sylvie Linsuke¹, Megan Halbrook⁵, Ernest 5 Kalthan⁷, Elisabeth Pukuta¹, Adrienne Amuri Aziza¹, Jean Claude Makangara Cigolo^{1,3,8}, 6 Raphael Lumembe^{1,3}, Gabriel Kabamba^{1,3}, Yvon Anta¹, Pierrot Bolunza⁹, Innocent Kanda⁹, 7 Raoul Nganzobo¹⁰, Thierry Kalonji⁶, Justus Nsio¹¹, Patricia Matoka¹¹, Dieudonné Mwamba¹², 8 Christian Ngandu¹², Souradet Y. Shaw¹³, Robert Shongo⁶, Joule Madinga¹, Yap Boum¹⁴, 9 Laurens Liesenborghs², Eric Delaporte⁴, Ahidjo Ayouba⁴, Nicola Low^{8,15}, Steve Ahuka 10 Mundeke^{1,3}, Lisa E. Henslev¹⁶, Jean-Jacques Muvembe Tamfum^{1,3}, Emmanuel Nakoune¹⁴, 11 Martine Peeters^{4*}, Nicole A. Hoff^{5*}, Jason Kindrachuk^{17,18}, Anne W. Rimoin^{5*}, Placide Mbala-12 Kingebeni^{1,3*} 13

14

15 1. Institut National de Recherche Biomédicale, Kinshasa, Democratic Republic of Congo

16 2. Department of Clinical Sciences, Institute of Tropical Medicine, Antwerp, Belgium

Service de Microbiologie, Département de Biologie Médicale, Cliniques Universitaires de
 Kinshasa, Kinshasa University, Democratic Republic of Congo

TransVIHMI (Recherches Translationnelles sur le VIH et les Maladies Infectieuses
 endémiques et émergentes), Université de Montpellier, French National Research Institute
 for Substaminale Développent (IRD), INSERM, Montpellier, France

5. Department of Epidemiology, Jonathan and Karin Fielding School of Public Health,
University of California, Los Angeles, CA, USA

24 6. Hemorrhagic Fevers and Monkeypox Program, Ministry of Health, Democratic Republic of25 the Congo

26 7. Ministry of Health and Populations, Bangui, Central African Republic

- 8. Graduate School for Cellular and Biomedical Sciences, University of Bern, Bern,
 Switzerland
- 29 9. Provincial Health Division, South Ubangi, Democratic Republic of Congo
- 30 10. Mbaya Health Zone, South Ubangi, Democratic Republic of Congo
- 31 11. National Border Hygiene Program, Ministry of Health, Democratic Republic of Congo
- 32 12. National Institute of Public Health, Ministry of Health, Democratic Republic of Congo

33 13. Department of Community Health Sciences, Max Rady College of Medicine, University of

- 34 Manitoba, Winnipeg, Manitoba, Canada
- 35 14. Pasteur Institute of Bangui
- 36 15. Institute of Social and Preventive Medicine (ISPM), University of Bern, Switzerland
- 37 16. Zoonotic and Emerging Disease Research Unit, National Bio and
- 38 Agro-Defense Facility, USDA Agricultural Research Service (ARS), Manhattan,
- 39 Kansas.
- 40 17. Department of Medical Microbiology & Infectious Diseases, University of Manitoba,
- 41 Winnipeg, Manitoba, Canada
- 42 18. Manitoba Centre for Proteomics & Systems Biology, Department of Internal Medicine,
- 43 University of Manitoba, Winnipeg, Manitoba, Canada
- 44
- 45
- 46
- 47
- 48
- 40
- 49
- 50
- 51
- 52 *Corresponding author:*
- 53 Anne Rimon, PhD, MPH
- 54 Professor, Department of Epidemiology, Gordon-Levin Endowed Chair in Infectious Diseases
- 55 and Public Health
- 56 UCLA Fielding School of Public Health,
- 57 650 Charles E. Young Dr, CHS 41-275, Los Angeles, CA 90095, USA
- 58 Phone: +1(310) 409-6299
- 59 E-mail: <u>arimoin@ucla.edu</u>
- 60 Institutional webpage: <u>www.ucla.edu</u>
- 61
- 62 *Co-corresponding author:*
- 63 Placide Mbala-Kingebeni, MD, PhD
- 64 Associate Professor, School of Medicine, University of Kinshasa (UNIKIN)
- 65 Department of Epidemiology and Global Health; Head of Pathogen Genomics Laboratory,
- 66 Institute National de la Recherche Biomédicale (INRB), Kinshasa, Democratic Republic of the
- 67 Congo (DRC)
- 68 Phone: +243 822 851 584
- 69 E-mail: mbalaplacide@gmail.com
- 70 Institutional webpage: https://inrb.net/

71	
72 73	
74	ABSTRACT
75	
76	Four confirmed mpox cases in South Ubangi province, Democratic Republic of the Congo, were
77	linked to documented transboundary transmission from Central African Republic. Viral genome
78	sequencing shows that the MPXV sequences belong to subclade Ia. This demonstrates the
79	borderless nature of mpox and highlights the need for vigilant regional surveillance.
80	
81	
82 83 84	Keywords: mpox; Democratic Republic of the Congo; Central African Republic; surveillance; cross-border transmission; subclade Ia; MPXV
85	
86	
87	
88	
89	
90	
91	
92	
93	
94	
95	
96	
97	
98	
99	
100	
101	
102	

103

104

105 INTRODUCTION

106 Mpox is a zoonotic viral infectious disease first identified in humans in the Democratic Republic 107 of the Congo (DRC) in 1970 (1). Monkeypox virus (MPXV) is endemic in forested regions of 108 Central and West Africa (2), subclassified into two clades: Clade I and Clade II, where subclade 109 IIb was responsible for a global epidemic in 2022 (3). We have recommended subdivision of 110 Clade I into subclades Ia and Ib, the latter associated with sustained human-to-human 111 transmission in DRC (4). Clade I is endemic in Central Africa, particularly in the DRC and 112 infections are associated with greater disease severity than Clade II (5, 6). Zoonotic spillover has 113 been the primary driver of Clade I MPXV infections; however, sustained human-to-human 114 transmission is increasing and spread of Clade I MPXV through sexual contact has been reported 115 in multiple regions of the DRC (4, 7).

116 The DRC has faced the greatest mpox burden among endemic regions. The number of reported 117 cases has been increasing since the cessation of smallpox vaccination campaigns in 1980 (8). 118 This increase has accelerated from 2022 onwards (9), including into regions with no previously 119 reported cases (10). While cross-border travel has been demonstrated as a factor involved in 120 transmission of Clade II MPXV, less is known for Clade I cases in regional settings. Given the recent geographic expansion for Clade I MPXV and increasing observation of sustained human-121 122 to-human transmission, there are considerable concerns about geographic expansion of mpox 123 through cross-border transmission (10).

The Central African Republic (CAR) shares a >1,000-mile border with DRC. CAR has reported 40 confirmed mpox outbreaks (95 suspected from 2001-2021), increasing from 0-2 annually from 2001-2017 to nine annually since 2018 (11). Most of these outbreaks occurred in two regions along the DRC border. Here, we report suspected cross-border transmission of mpox between CAR and DRC in 2023 and potential links with MPXV circulation in DRC.

129

130 THE STUDY

An alert for a suspected mpox case was issued by the South Ubangi Provincial Health Division,
DRC, in January 2023 following the death of a fisherman who presented with respiratory distress
and skin lesions in Mbaya Health Zone. South Ubangi Province directly borders CAR, separated

by the Ubangi River. The National Programme for Control of Mpox and Viral Haemorrhagic
Fevers (PNLMPX-VHF) and the National Institute for Biomedical Research (INRB) deployed a
multidisciplinary team to investigate the case with the South Ubangi Provincial Health Division.
As part of the investigation, the national and provincial health teams conducted supplemental
training sessions to raise awareness of the clinical signs and transmission modes for mpox.

139 Figure 1 shows a documented chain of MPXV transmission from CAR to Mbaya Health Zone. 140 The index case was an adult old male resident of Bangui, CAR, who moved frequently between CAR and DRC via the Ubangi River (cluster 1, case 1: C1). Following suspected exposure to 141 142 wildlife meat and the potential consumption of rodents in CAR in early January 2023, with 143 symptom onset began shortly after (day 0) and including fever, headache, chills, and cutaneous rash. After traveling to the DRC, he was admitted to Mbaya General Reference Hospital two 144 145 days later, where he died on day 9 post-symptom onset after experiencing respiratory distress. 146 No samples were collected from this probable mpox case nor any information on his immune 147 status.

Fifteen contacts of the patient were identified in Mbaya Health Zone, including five health care workers and 10 close contacts (four close friends and six family contacts). The investigation revealed a cluster of three symptomatic family contacts. Two had samples taken, which were positive for MPXV by PCR; the adult female partner of C1 (C2) and their child (C3). A second child of C1 and C2 (C4) was treated locally for suspected mpox through traditional methods and was not investigated further. No suspected mpox cases were identified among family or close contacts of C1 in Bangui, CAR.

During the investigation, a second mpox cluster was identified, which likely originated from a different source. Three suspected mpox cases were identified at the same hospital and at the same time as the first cluster, with no known epidemiologic link. An adult female patient (cluster 2, case 1: D1) was hospitalized for nonspecific signs of mpox at the same time as C1 and reported contact with a child with mpox symptoms ~28 days prior. The two additional suspected cases were the child of D1 (D2), with subsequent PCR-confirmed mpox, and a contact of D2 (D3) who tested PCR negative. (**Figure 2**).

162 Confirmed mpox clinical symptoms among the confirmed cases included multiple pustular and 163 papular lesions (C2), disseminated pustules (C3), discreet lacrimation in the left eye and post-164 inflammatory hyperpigmentation (D1), and hyperpigmented, disseminated macules and steep-165 edged ulcerations (D2).

166 Subsequent viral genome sequencing was performed on three PCR-positive samples: two family 167 members in the first mpox cluster (C2 and C3) and one from the second cluster (D2) (Figure 3). 168 Phylogenetic analysis included 98 clade I MPXV previously published genomes. The 169 phylogenetic tree was generated by maximum likelihood using the K3Pu+F+I model (12, 13). 170 The three sequences, all clade Ia, clustered with MPXV genome sequences isolated from CAR. 171 The genetic distance between the new cases and the date between the CAR and DRC samples is 172 too long to conclude direct links between the two countries. Nevertheless, the new samples from 173 South-Ubangi cluster in the same subgroup (group II) according to previous classification of 174 MPXV sequences in clade Ia (14).

The PNLMPX-VHF investigated a further 109 suspected mpox cases from 15 health zones in
South Ubangi province from January to November 2023. Cases were predominantly male (60
males; 46 females; 3 unavailable) with an average age of 21 years (range 35 months to 63 years).
Of these, 61 cases (56 %) were confirmed positive by PCR at INRB. Three mpox negative cases
were confirmed positive for varicella zoster virus (3/48; 6%).

180

181 **DISCUSSION**

182 This investigation describes two clusters of mpox cases in Mbaya health zone in South Ubangi 183 province, DRC, presumably resulting from two different introductions. Epidemiological links to 184 cross-border travel from CAR were found in the first cluster and genomic links with previous 185 cases from CAR for both. MPXV transmission in DRC is currently driven by both zoonotic and 186 human to human contact, thus increasing the complexity of containment and mitigation efforts. 187 There are considerable concerns about wider regional and international expansion of mpox 188 including the concentration of MPXV outbreaks in CAR and DRC along the Ubangi River, recent increases in sustained human-to-human transmission, resource limitations for 189 190 identification and treatment of mpox, disease stigma, population displacement due to conflict, 191 and transient cross-border transit.

In August 2022, a first regional meeting involving six neighbouring countries of Central and West Africa was held in Kinshasa, DRC to establish a regional mpox surveillance network - the Mpox Threat Reduction Network. Through this network multiple teams that included experts from the Health Ministries of the DRC and CAR which were able to communicate rapidly on suspected cases crossing borders in South Ubangi while also reporting to the International Health Regulations focal point in the DRC and CAR. This investigation highlights the importance of long-term collaborative partnerships for sustained mpox surveillance and containment in endemic regions.

218

219

220 ACKNOWLEDGEMENTS

221 Figures included in this manuscript were generated using Biorender.com.

222

223 FUNDING

The project or effort depicted is sponsored by the Department of Defense Threat Reduction

Agency. The content of the information does not necessarily reflect the position or the policy of

the federal government, and no official endorsement should be inferred.

227 This work was also supported by the International Mpox Research Consortium (IMReC)

through funding from the Canadian Institutes of Health Research and International Development

229 Research Centre (Grant No. 202209MRR-489062-MPX-CDAA-168421) and the Agence

230 Française de Développement through the PANAFPOX and AFROSCREEN project (grant

agreement CZZ3209), coordinated by ANRS Maladies infectieuses émergentes in partnership

232 with Institut de Recherche pour le Développement (IRD). E.L. received a PhD grant from the

- 233 French Foreign Office.
- 234

235 **REFERENCES:**

236 1. Ladnyj ID, Ziegler P, Kima E. A human infection caused by monkeypox virus in

Basankusu Territory, Democratic Republic of the Congo. Bull World Health Organ.

238 1972;46(5):593-7.

239 2. Titanji BK, Tegomoh B, Nematollahi S, Konomos M, Kulkarni PA. Monkeypox: A

240 Contemporary Review for Healthcare Professionals. Open Forum Infect Dis. 2022

- 241 Jul;9(7):ofac310.
- 3. Happi C, Adetifa I, Mbala P, Njouom R, Nakoune E, Happi A, et al. Urgent need for a
 non-discriminatory and non-stigmatizing nomenclature for monkeypox virus. PLoS Biol. 2022
 Aug;20(8):e3001769.

245 4. Vakaniaki EH, Kacita C, Kinganda-Lusamaki E, O'Toole A, Wawina-Bokalanga T,

Mukadi-Bamuleka D, et al. Sustained Human Outbreak of a New MPXV Clade I Lineage in the
Eastern Democratic Republic of the Congo. Nat Med. 2024 Jun 13.

- Weaver JR, Isaacs SN. Monkeypox virus and insights into its immunomodulatory
 proteins. Immunol Rev. 2008 Oct;225:96-113.
- 250 6. McCollum AM, Damon IK. Human monkeypox. Clin Infect Dis. 2014 Jan;58(2):260-7.

251 7. Kibungu EM, Vakaniaki EH, Kinganda-Lusamaki E, Kalonji-Mukendi T, Pukuta E, Hoff 252 NA, et al. Clade I-Associated Mpox Cases Associated with Sexual Contact, the Democratic 253 Republic of the Congo. Emerg Infect Dis. 2024 Jan;30(1):172-6. 254 8. MSPHP. REPORT ON THE EPIDEMIOLOGICAL SITUATION OF MONKEYPOX 255 VIRUS—S1-S6 2024. Ministry of Public Health, Hygiene and Prevention of the Democratic 256 Republic of the Congo; 2024. 257 9. WHO. Disease Outbreak News: Mpox - Democratic Republic of the Congo; 2024 14 258 June 2024. 259 10. Mbala-Kingebeni P, Rimoin AW, Kacita C, Liesenborghs L, Nachega JB, Kindrachuk J. 260 The time is now (again) for mpox containment and elimination in Democratic Republic of the Congo. PLOS Glob Public Health. 2024;4(6):e0003171. 261 262 Besombes C, Mbrenga F, Schaeffer L, Malaka C, Gonofio E, Landier J, et al. National 11. 263 Monkeypox Surveillance, Central African Republic, 2001-2021. Emerg Infect Dis. 2022 264 Dec;28(12):2435-45. 265 Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder: 12. 266 fast model selection for accurate phylogenetic estimates. Nat Methods. 2017 Jun;14(6):587-9. 267 13. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et al. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic 268 269 Era. Mol Biol Evol. 2020 May 1;37(5):1530-4. 270 14. Berthet N, Descorps-Declere S, Besombes C, Curaudeau M, Nkili Meyong AA, Selekon B, et al. Genomic history of human monkey pox infections in the Central African Republic 271 272 between 2001 and 2018. Sci Rep. 2021 Jun 22;11(1):13085. 273 274 275 276 277 278 279 280 281 282 283 284 285



Figure 1. Suggested transmission route of MPXV between Bangui, Central African Republic
(CAR) and Mbaya Health Zone, South Ubangi Province, Democratic Republic of the Congo,
January 2023



Figure 2. Suggested transmission chain from Central African Republic to the Democratic
Republic of the Congo and second identified chain at Mbaya General Reference Hospital, DRC.
Abbreviations: HCW, healthcare workers; HZ, health zone



340

341

342 Figure 3: Phylogenetic analysis of MPXV sequences from Mbaya rural health zone. 343 Phylogenetic analysis of MPXV genome sequences from samples described in this study and 344 clade I MPXV sequences from Central Africa. Scale bar is in substitutions per site; bootstrap support values are shown at branch points. DNA was extracted at INRB using a Qiagen DNA 345 Mini kit from blood samples and subsequently screened for MPXV with an Orthopoxvirus-346 347 specific real-time PCR assay. Whole genome sequencing was attempted on samples from the index case by Next Generation Sequencing (NGS). The library preparation was performed using 348 349 Illumina DNA Prep with Enrichment and the libraries were enriched for MPXV using 350 biotinylated custom probes synthesized by Twist Biosciences. Note that 23MPX0134C(D2),

351 23MPX0135V(C2), and 23MPX0136V(C3) are samples from crust or vesicles from separate individuals.