Estimating the generation time for influenza transmission using household data in the United States

4 Louis Yat Hin Chan¹, Sinead E. Morris^{1,9}, Melissa S. Stockwell², Natalie M. Bowman³, Edwin Asturias⁴,

5 Suchitra Rao⁴, Karen Lutrick⁵, Katherine D. Ellingson⁵, Huong Q. Nguyen⁶, Yvonne Maldonado⁷, Son H.

McLaren², Ellen Sano², Jessica E. Biddle¹, Sarah E. Smith-Jeffcoat¹, Matthew Biggerstaff¹, Melissa A.
 Rolfes¹, H. Keipp Talbot⁸, Carlos G. Grijalva⁸, Rebecca K. Borchering¹, Alexandra M. Mellis¹, RVTN-

- 8 Sentinel Study Group
- 9 1. Centers for Disease Control and Prevention
- 10 2. Columbia University Irving Medical Center
- 11 3. University of North Carolina at Chapel Hill
- 12 4. University of Colorado School of Medicine and Children's Hospital Colorado
- 13 5. University of Arizona
- 14 6. Marshfield Clinic Research Institute
- 15 7. Stanford University
- 16 8. Vanderbilt University Medical Center
- 17 9. Goldbelt Professional Services

18 * Influenza Division, Centers for Disease Control and Prevention, 1600 Clifton Road, Atlanta, GA 30329,

19 US (<u>LouisChan@cdc.gov</u>)

20 Keywords

- 21 Generation interval
- Serial interval
- 23 Incubation period
- 24 Pre-symptomatic transmission
 - Household transmission
 - Respiratory diseases

27 Abstract

25

- 28 The generation time, representing the interval between infections in primary and secondary cases, is
- 29 essential for understanding and predicting the transmission dynamics of seasonal influenza, including
- 30 the real-time effective reproduction number (Rt). However, comprehensive generation time estimates
- 31 for seasonal influenza, especially post the 2009 influenza pandemic, are lacking.
- 32 We estimated the generation time utilizing data from a 7-site case-ascertained household study in the
- United States over two influenza seasons, 2021/2022 and 2022/2023. More than 200 individuals who
- 34 tested positive for influenza and their household contacts were enrolled within 7 days of the first illness
- 35 in the household. All participants were prospectively followed for 10 days completing daily symptom
- diaries and collecting nasal swabs, which were tested for influenza via RT-PCR. We analyzed these data

- 37 by modifying a previously published Bayesian data augmentation approach that imputes infection times
- of cases to obtain both intrinsic (assuming no susceptible depletion) and realized (observed within
- 39 household) generation times. We assessed the robustness of the generation time estimate by varying
- 40 the incubation period, and generated estimates of the proportion of transmission before symptomatic
- 41 onset, infectious period, and latent period.
- 42 We estimated a mean intrinsic generation time of 3.2 (95% credible interval, Crl: 2.9-3.6) days, with a
- 43 realized household generation time of 2.8 (95% Crl: 2.7-3.0) days. The generation time exhibited limited
- sensitivity to incubation period variation. Estimates of the proportion of transmission that occurred
- 45 before symptom onset, the infectious period, and the latent period were sensitive to variation in
- 46 incubation periods.
- 47 Our study contributes to the ongoing efforts to refine estimates of the generation time for influenza.
- 48 Our estimates, derived from recent data following the COVID-19 pandemic, are consistent with previous
- 49 pre-pandemic estimates, and will be incorporated into real-time Rt estimation efforts.

50 Introduction

- 51 The generation time, a crucial parameter in understanding the dynamics of infectious diseases, is
- 52 defined as the time interval between infections in primary and secondary cases. In the context of
- 53 seasonal influenza, estimation of the generation time becomes increasingly important for predicting the
- 54 trajectory of outbreaks and informed public health decision-making during an influenza season. This
- 55 interval represents the time between typical influenza infections, reflecting when most transmission is
- 56 likely to happen.
- 57 Estimating the generation time is challenging because few investigations can accurately detect the exact
- 58 time of infection. The generation time is often inferred from the serial interval, defined as the time
- 59 between symptom onsets of primary and secondary cases (Svensson 2007), due to the practicality of
- 60 observing symptom onsets rather than infections. However, this alternative measure may not always
- 61 approximate the generation time due to its dependence on the incubation period, defined as the
- 62 duration from infection to symptom onset, and the possibility of asymptomatic infections.
- 63 Accurate estimation of the generation time is important for predicting the real-time effective
- reproduction number (Rt), a metric used to describe transmission intensities through time (Gostic, et al.
- 65 2020). During the 2023/2024 influenza season, the Centers for Disease Control and Prevention (CDC)
- 66 estimated the current epidemic growth status for influenza infections in the U.S. as either growing or
- 67 declining based on the Rt (Centers for Disease Control and Prevention 2024) (Centers for Disease
- 68 Control and Prevention 2024). For this estimate, the generation time was approximated with a serial
- 69 interval from a study by Cowling et al. (Cowling, et al. 2009) that utilized data collected in Hong Kong in
- 70 2007, prior to the 2009 H1N1 influenza pandemic.
- 71 To improve our understanding of seasonal influenza outbreaks, there is a need for more contemporary
- 72 generation time estimates, especially following the COVID-19 pandemic. This analysis provides updated
- 73 generation time estimates derived from an influenza household transmission study (Rolfes, et al. 2023)
- conducted during the 2021/2022 and 2022/2023 influenza seasons in the U.S. We employ a model using
- 75 a published Bayesian data augmentation approach (Hart, Abbott, et al. 2022) (Hart, Maini and
- 76 Thompson 2021) (Hart, Miller, et al. 2022) to impute missing event times, including infections and

- 77 symptom onsets of cases, and estimate generation times. We estimate both the intrinsic generation
- time, which assumes no susceptible depletion, as well as the realized household generation time
- observed within the household setting. We also estimate the serial interval. We derived estimates
- 80 across the two seasons, virus types (influenza A and B), and household sizes to understand potential
- 81 differences and robustness to model assumptions. The insights gained from these sensitivity analyses
- 82 contribute to our understanding of the reliability of our estimates across different data stratifications
- and assumptions, providing evidence that the generation time has remained substantially unchanged
- 84 over the last decade or two.
- 85 We also estimate other transmission parameters, including the proportion of transmission before
- 86 symptomatic onset, the infectious period, and the latent period. This helps in assessing pre- and post-
- 87 symptomatic transmission, thereby providing insights to inform effective disease control strategies.
- 88 These insights are crucial for preventing pre-symptomatic transmission through interventions such as
- 89 isolation.

90 Material and methods

91 Household data

- 92 Participants included in this analysis were enrolled in a 7-site case-ascertained household study, the
- 93 Respiratory Virus Transmission Network Sentinel (RVTN-S), conducted in the U.S. over two consecutive
- 94 influenza seasons: 2021/2022 and 2022/2023 (Rolfes, et al. 2023). After informed consent was obtained,
- 95 the study enrolled individuals identified with influenza infections via polymerase chain reaction (PCR)
- 96 testing and their household contacts within 7 days of the initial illness onset within the household.
- 97 Households were only enrolled if the index case who first presented for clinical testing was the first
- 98 symptomatic or positive person in the household, with no other members of the household
- 99 symptomatic on the first day of index case symptoms. Participants, including both index cases and
- 100 household contacts, were then prospectively followed for 10 days, during which they completed daily
- 101 symptom diaries and collected daily nasal swabs, which were tested for influenza via RT-PCR.
- 102 The dataset encompasses detailed information regarding symptoms and viral testing, including four
- 103 main variables used in the model: whether individuals tested positive for influenza, their symptomatic
- status, dates of positive test results, and dates of symptom onset. Using the test positivity and
- symptomatic status, we stratified individuals into three types: symptomatic infected, asymptomatic
- 106 infected, and uninfected. Both the dates of positive test results and symptom onset were used as upper
- 107 bounds for the date of infection for each individual.
- 108 In the primary analysis, we excluded households with multiple co-primary cases, i.e., more than one
- 109 individual exhibiting the same date of the earliest symptom onset concurrently. To assess robustness,
- we also performed a separate stratified analysis that included households both with and without
- 111 multiple co-primary cases.

112 Estimating the generation time

- 113 We employed a Susceptible-Exposed-Infectious-Recovered (SEIR) model with Bayesian data
- augmentation, originally developed by Hart et al. (Hart, Maini and Thompson 2021) for analyzing COVID-
- 115 19 contact tracing data. The model was also used in two subsequent studies of household data in the
- 116 United Kingdom (Hart, Abbott, et al. 2022) (Hart, Miller, et al. 2022). The SEIR model, referred to as the
- mechanistic model, which includes compartments for asymptomatic, pre-symptomatic and symptomatic

- 118 infectious stages (Hart, Maini and Thompson 2021). Each stage may have different relative
- 119 infectiousness, or transmission rates. Upon infection and entry into the non-infectious exposed stage,
- 120 individuals may progress to become infectious through one of two pathways: either by remaining
- asymptomatic or by developing symptoms following a pre-symptomatic stage. Consequently,
- 122 transmissions may occur before symptom onset, depending on the length of the incubation period.
- 123 We estimated both intrinsic and realized generation times by integrating data augmentation Markov
- 124 Chain Monte Carlo (MCMC) techniques (Hart, Abbott, et al. 2022), to impute infection times and
- symptom onset of cases. The intrinsic generation time assumes no depletion of susceptible individuals,
- 126 providing an estimate of the time it takes for an infected individual to infect others in the community
- 127 with an unlimited supply of susceptible individuals. The realized household generation time reflects the
- actual time interval observed within households, restricted by the depletion of susceptible individuals
- 129 over time. Susceptible depletion refers to the gradual reduction in the number of individuals within a
- population that have not yet been infected with a virus. For example, within the SEIR framework,
- 131 members may become infected, develop immunity, and subsequently be removed from the susceptible
- pool. Considering this distinction allows for a more thorough understanding of influenza transmission
- 133 dynamics, capturing both theoretical and observed aspects of transmission.
- 134 We also estimated several other crucial transmission parameters, including the proportion of
- 135 transmission before symptomatic onset, the ratio of pre-symptomatic to symptomatic transmission
- 136 rates (i.e., relative infectiousness of symptomatic infected individuals before symptom onset compared
- to after), as well as the latent period, the pre-symptomatic infectious period, and the symptomatic
- 138 infectious period, all under the SEIR framework.
- 139 In adapting the model for our influenza study, we used estimates for the incubation period of influenza
- 140 A from a systematic review by Lessler et al. (Lessler, et al. 2009). In sensitivity analyses, we explored
- variations derived from parallel estimates for influenza B (Lessler, et al. 2009) and for influenza
- 142 A(H1N1)pdm09 (Tuite, et al. 2010).
- 143 Regarding the relative infectiousness of asymptomatic infected individuals compared with symptomatic
- 144 infected individuals, we assumed a value of 0.57 (i.e., asymptomatic infected individuals were 43% less
- 145 infectious than those symptomatic infected) based on the mean estimate from a recent study (Tsang, et
- al. 2023), and we also conducted sensitivity analyses using values of 0.11 and 1.54 based on the
- 147 corresponding 95% credible interval (CrI).
- 148 To compare posterior distributions of estimates, we calculated the overlapping index, a measure of
- distribution similarities (Pastore 2018) (Pastore and Calcagnì 2019). A value close to 1 indicates high
- similarity, implying no substantial differences, while a value close to 0 indicates low similarity, implying
- 151 substantial differences. We compared the estimates of generation time across multiple data
- 152 stratifications and sensitivity analyses to the primary results excluding households with multiple co-
- 153 primary cases.
- 154 The model was implemented in R (version 4.3.1) with 1,000,000 Markov chain Monte Carlo iterations,
- discarding the initial 20% as burn-in and obtaining posterior distributions by thinning every 100
- 156 iterations. The code for the model is available at <u>https://github.com/CDCgov/influenza-</u>
- 157 generation time-us.

158 Ethics statement

- 159 This study was reviewed and approved by the IRB at Vanderbilt University Medical Center (see 45 C.F.R.
- 160 part 46.114; 21 C.F.R. part 56.114).

161 **Results**

162 The household data

- 163 During the data cleaning process, we excluded 93 individuals who did not have at least two valid PCR
- tests and 2 individuals who were the only household members. In the primary analysis, we further
- 165 excluded 23 individuals from 6 households that had co-primary cases. The final cleaned dataset,
- 166 covering both seasons, comprised 820 individuals from 246 households (Table 1).
- 167 As shown in Table 1, more households were enrolled in the 2022/2023 season. In both seasons,
- 168 influenza A viruses predominantly circulated. In the 2021/2022 season, influenza A(H3N2) virus was
- 169 identified in 78% of individuals, and influenza A(H1N1) virus was identified in 1% of individuals. In the
- 170 2022/2023 season, the percentages changed to 64% and 6%, respectively. Since households consisting
- of 3 or 4 members were the majority, we stratified the data into two groups: those with 2 or 3
- members, and those with 4 or greater, to ensure comparability in quantity.

Data stratifications	Number of individuals (households)	Symptomatic infected %	Asymptomatic infected %	Uninfected %
All data excluding households with multiple co- primary cases (primary analysis)	820 (246)	59.4% (487/820)	7.2% (59/820)	33.4% (274/820)
Season 2021/2022	308 (90)	59.4% (183/308)	7.5% (23/308)	33.1% (102/308)
Season 2022/2023	512 (156)	59.4% (304/512)	7.0% (36/512)	33.6% (172/512)
Influenza A	683 (209)	61.1% (417/683)	7.5% (51/683)	31.5% (215/683)
Influenza B	137 (37)	51.1% (70/137)	5.8% (8/137)	43.1% (59/137)
Household size of 2 or 3	393 (152)	62.6% (246/393)	5.3% (21/393)	32.1% (126/393)
Household size of 4 or greater	427 (94)	56.4% (241/427)	8.9% (38/427)	34.7% (148/427)
All data including households with multiple co-primary cases	843 (252)	60.3% (508/843)	7.0% (59/843)	32.7% (276/843)

173 Table 1. Characteristics of household data.

174 Consistent estimates of the generation time across data stratifications and

175 parameter assumptions

- 176 In the primary analysis using all data excluding households with multiple co-primary cases from both
- seasons, we estimated a mean intrinsic generation time of 3.2 (95% credible interval, CrI: 2.9-3.6) days
- 178 (Figure 1A, 1B and Table 2). The corresponding mean (intrinsic) serial interval was 3.2 (95% CrI: 2.8-3.5)
- days, with a standard deviation (SD) of 2.2 (95% Crl: 1.8-2.6) days. The mean realized household
- 180 generation time was 2.8 (95% Crl: 2.7-3.0) days, nearly half a day shorter than the mean intrinsic
- 181 generation time.

- 182 We found no substantial differences in the mean intrinsic generation time estimates across multiple
- data stratifications (Figure 1C and Supplemental Table S1). The overlapping indices for both 2021/2022
- and 2022/2023 were high at 71% and 87%, respectively, aligning with the primary analysis above.
- 185 Influenza A data showed a notable high overlapping index of 94%, reflecting its dominance, as influenza
- 186 A was identified in 83% of the individuals. Conversely, using the data exclusively from influenza B yielded
- a similar mean but with a wider credible interval due to the smaller sample size, i.e., influenza B was
- identified in only 17% of the individuals, resulting in a lower overlapping index of 47%. Upon examining
- 189 household sizes, although we found slightly longer mean intrinsic and realized household generation
- times in smaller households compared to larger ones, the overlapping index for household sizes of 2 or 3
- 191 members and 4 or more members were moderately high at 61% and 74%, respectively. Incorporating
- 192 households with co-primary cases remained consistent with a moderately high overlapping index of
- 193 64%, indicating the similarity between exclusion and inclusion of multiple co-primary cases.
- 194 The mean intrinsic generation time exhibited limited sensitivity to variations in the incubation period
- 195 (Figure 1D and Supplemental Table S2). In the primary analysis shown above, we used an incubation
- 196 period with a mean of 1.55 days and a standard deviation (SD) of 0.66 days by fitting previously
- 197 published estimates (Lessler, et al. 2009) to a gamma distribution (Supplemental Figure S1). When
- considering a shorter incubation period, which yielded a mean of 0.61 days and a SD of 0.25 days
- 199 (Lessler, et al. 2009), the mean intrinsic generation time remained unchanged with an overlapping index
- of 86%. Conversely, with a longer incubation period, which yielded a mean of 4.30 days and a SD of 1.25
- days (Tuite, et al. 2010), the mean intrinsic generation time increased slightly with an intermediate
- 202 overlapping index of 56%.
- 203 Our estimates were not sensitive to changes in the relative infectiousness of asymptomatic infected
- 204 individuals, due to the limited number of asymptomatic infected individuals in this study (Supplemental
- Figure S6). We found no substantial differences in the mean intrinsic generation time estimates
- 206 (Supplemental Table S2), as indicated by overlapping indices of 94% and 97% when using the values of
- 207 0.11 and 1.54 compared to the primary value of 0.57 (Tsang, et al. 2023).



208

Figure 1. (A) Distributions of intrinsic generation time and serial interval using the posterior samples. The lines represent the
median, and the shaded areas denote the 95% credible intervals (Crl). The blue color represents the intrinsic generation time
distribution, while the orange color represents the serial interval distribution. (B) Posterior distribution of mean intrinsic and
realized household generation time. (C) Posterior distributions of mean intrinsic generation time across seasons, virus types,
household sizes, and with multiple co-primary cases. The incubation period, derived from influenza A, had a mean of 1.55 days
and a standard deviation (SD) of 0.66 days (Lessler, et al. 2009). Only for influenza B, we assumed the shorter incubation period
to yield a mean of 0.61 days and a standard deviation (SD) of 0.25 days (Lessler, et al. 2009). (D) Posterior distributions of mean

216 intrinsic generation time estimated using the full dataset across different incubation periods.

	Mean	SD
Intrinsic generation time (days)	3.2 (2.9-3.6)	2.1 (1.8-2.5)
Realized household generation time (days)	2.8 (2.7-3.0)	1.6 (1.5-1.8)
Serial interval (days)	3.2 (2.8-3.5)	2.2 (1.8-2.6)

217 Table 2. Posterior mean (95% CrIs) of estimates in primary analysis using the full dataset. The incubation period, derived from

218 influenza A, had a mean of 1.55 days and a standard deviation (SD) of 0.66 days (Lessler, et al. 2009). The relative infectiousness

of asymptomatic infected individuals compared with symptomatic infected individuals was assumed to be 0.57 (Tsang, et al.
 2023).

221 Sensitivity of other transmission parameters to the incubation period

222 In our sensitivity analyses, where we varied the assumed incubation period from the mean of 1.55 days

and SD of 0.66 days, we found significant influences on several crucial pre-symptomatic transmission

- parameters and the duration of various symptomatic infectious stages (Table 3 and Supplemental FigureS5).
- Notably, given the shorter incubation period (mean of 0.61 days and SD of 0.25 days), the proportion of
- transmission before symptomatic onset was lower at 3% (95% Crl: 0-6%), and the ratio of pre-
- 228 symptomatic to symptomatic transmission rates indicated a lower relative infectiousness of
- symptomatic infected individuals before symptom onset compared to after. This indicates that the
- 230 majority of transmission occurred after individuals developed symptoms. Consequently, this was

- 231 reflected in a shorter latent period of 0.4 (95% Crl: 0.2-0.6) days and pre-symptomatic infectious period
- 232 of 0.2 (95% Crl: 0.0-0.4) days, or a longer symptomatic infectious period of 2.6 (95% Crl: 2.3-3.0) days.
- 233 Conversely, given the longer incubation period (mean of 4.30 days and SD of 1.25 days), the proportion
- of transmission before symptomatic onset was higher at 76% (95% Crl: 65-87%), and the ratio of pre-234
- 235 symptomatic to symptomatic transmission rates was higher. This resulted in a longer latent period of 0.9
- (95% Crl: 0.2-1.6) days and pre-symptomatic infectious period of 3.4 (95% Crl: 2.7-4.1) days, while the 236
- 237 symptomatic infectious period was shorter at 1.2 (95% Crl: 0.7-1.9) days.

Incubation period	Shorter	Primary	Longer	
Mean intrinsic generation time	3.2 (2.8-3.6)	3.2 (2.9-3.6)	3.4 (3.1-3.7)	
(days)				
Proportion of transmission before	0.03 (0.00-0.06)	0.16 (0.07-0.25)	0.76 (0.65-0.87)	
symptomatic onset				
Ratio of pre-symptomatic and	0.6 (0.1-1.7)	0.7 (0.2-1.9)	1.3 (0.5-3.2)	
symptomatic transmission rates				
Latent period (days)	0.4 (0.2-0.6)	0.9 (0.2-1.4)	0.9 (0.2-1.6)	
Pre-symptomatic infectious period	0.2 (0.0-0.4)	0.7 (0.2-1.3)	3.4 (2.7-4.1)	
(days)				
Symptomatic infectious period	2.6 (2.3-3.0)	2.0 (1.7-2.4)	1.2 (0.7-1.9)	
(days)				

238 Table 3. Posterior mean (95% Crls) of estimates of generation time and transmission parameters given different assumed

239 incubation periods. The primary incubation period, derived from influenza A, had a mean of 1.55 days and a standard deviation

240 (SD) of 0.66 days (Lessler, et al. 2009). The shorter incubation period, derived from influenza B, yielded a mean of 0.61 days with

241 a SD of 0.25 days (Lessler, et al. 2009), while the longer incubation period, derived from influenza A(H1N1)pdm09, had a mean

242 of 4.30 days with a SD of 1.25 day (Tuite, et al. 2010). The relative infectiousness of asymptomatic infected individuals compared 243

with symptomatic infected individuals was assumed to be 0.57 (Tsang, et al. 2023).

Discussion 244

245 Estimates of generation time

This study employed a Bayesian data augmentation approach (Hart, Maini and Thompson 2021, Hart, 246

247 Abbott, et al. 2022, Hart, Miller, et al. 2022) to estimate both intrinsic and realized generation times

248 using data collected from a U.S. household study during the post COVID-19 pandemic influenza seasons,

249 2021/2022 and 2022/2023. Our findings indicate that the intrinsic generation time, reflecting

250 transmission dynamics within community settings, ranged from 2.9 to 3.6 days, while the realized

251 household generation time, restricted to household settings, ranged from 2.7 to 3.0 days. These

252 estimates of the generation time for influenza fall within the uncertainty bounds of pre-pandemic

253 studies, including directly using viral shedding data (Carrat, et al. 2008) and other contact tracing data

254 (Fraser, et al. 2009) (te Beest, et al. 2013) (Lau, et al. 2015), with estimates varying between 2 and 4

255 days, suggesting that there has not been substantial change since the 2009 H1N1 influenza pandemic.

- 256 Additionally, the overlapping indices higher than 70% suggested no substantial differences between the
- 257 two influenza seasons.
- 258 Both seasons of this study were atypical, being the first seasons since the COVID-19 pandemic, during
- 259 which the immunity to influenza had potentially decreased. The 2022/2023 season, in particular,
- 260 experienced an early influenza activity peak along with RSV and COVID-19 outbreaks. Despite these
- 261 unusual circumstances, both seasons dominated by influenza A(H3N2) were tested in our sensitivity

- analyses, which were based on different parameter assumptions. The generation time estimates
- remained similar to those from earlier studies, suggesting that virus transmission dynamics within
- households have not changed substantially and may not vary widely between types. However, further
- work is needed to fully explore estimates for influenza that were less prevalent in this study (i.e.,
- 266 influenza B and A(H1N1)).
- 267 Our finding that the realized household generation time was shorter than the intrinsic generation time
- 268 could be attributed to the depletion of susceptible individuals over time. As household members
- 269 become infected and develop immunity, although individuals may still be infectious, there are no
- susceptible contacts still exposed to each case. This depletion terminates transmission chains,
- 271 diminishing the potential for further infections. This process, along with factors such as closer proximity
- and longer exposure times inherent to household settings, can increase the chance of transmission
- 273 within households, leading to a shorter observed generation time.
- 274 Our updated estimates, particularly for the intrinsic generation time, may be useful for ongoing
- 275 modeling efforts which require estimated generation times, such as real-time influenza Rt estimation
- 276 (Centers for Disease Control and Prevention 2024) (Centers for Disease Control and Prevention 2024)
- 277 (Gostic, et al. 2020). Our estimates are slightly shorter than the serial interval estimated by Cowling et
- al. (Cowling, et al. 2009) at 3.6 days (95% confidence interval, CI: 2.9-4.3). This shorter interval suggests
- a more rapid spread, potentially leading to higher estimated Rt and emphasizes the need for prompt
- and effective interventions to control transmission.

281 Reliability of other transmission parameters

- 282 Comparing our other parameter estimates with prior research, we found the mean serial interval to be
- 283 3.2 days, within the 95% CI of 2.9 to 4.3 days reported by Cowling et al. (Cowling, et al. 2009). Our
- estimate also falls within the 3-to-4-day range of uncertainty reported in previous household studies
- (Cauchemez, Donnelly, et al. 2009, Petrie, et al. 2013, Levy, et al. 2013, Xu, et al. 2015, Cowling, et al.
- 286 2010, Suess, et al. 2012, Boëlle, et al. 2011, Tsang, et al. 2016).
- 287 We estimated a latent period of less than a day, which is shorter than the 1 to 3 days reported in other
- studies (Tuite, et al. 2010) (Cori, et al. 2012). It is possible that this shorter latent period could be
- influenced by undocumented exposures outside households. For instance, both the index case and
- 290 infected household member may have been exposed to influenza elsewhere, with the index case
- 291 developing symptoms before the household member. Consequently, when the household member
- becomes sick, we attribute it to the index case within the household, but this infection could have
- originated from previous exposure outside the household, which has not been accounted for in this
- analysis. Nonetheless, we excluded households with multiple co-primary cases to reduce these effects,
- ensuring accurate assessment of transmission dynamics within each household.
- 296 Furthermore, there was substantial uncertainty in our estimates for the symptomatic infectious period,
- which ranged from 1 to 3 days, given different assumed incubation periods. Likewise, there has been a
- wide range of estimates in earlier studies, including those less than a day (Cori, et al. 2012) and more
- than 3 days (Tuite, et al. 2010) (Cauchemez, Carrat, et al. 2004).
- 300 The estimates of the pre-symptomatic transmission parameters and the duration of various
- 301 symptomatic stages should be interpreted with caution due to the inherent uncertainty in accurately
- 302 estimating the incubation period. Our estimates incorporated values from various studies based on

different types or subtypes of influenza (Lessler, et al. 2009) (Tuite, et al. 2010). This variability in the

- incubation period contributes to a wide range of pre-symptomatic transmission parameters. Without a
- reliable input for the incubation period, accurately determining these transmission dynamics becomes
- 306 challenging.

307 Implications for preventing transmission

- 308 Understanding the proportion of transmission that occurs prior to symptoms is critical to informing
- 309 effective disease control strategies and assessing the potential impact of post-symptomatic mitigation
- measures, such as isolation of cases. We estimated that between 3% and 76% of transmission may occur
- before a person develops symptoms. This wide range was influenced by our assumptions of the
- incubation period, which were taken from a variety of previously published estimates. Longer incubation
- period assumptions yielded higher estimates of the percentage of transmission that occurred before
- 314 symptoms. Similarly, attempts to estimate individual-level pre-symptomatic transmission using viral
- kinetics data have revealed substantial heterogeneity (Morris, et al. 2024). This highlights that pre-
- 316 symptomatic transmission of influenza does occur, aligning with findings for other respiratory pathogens
- 317 like SARS-CoV-2 (Buitrago-Garcia, et al. 2022).
- Given the wide range of pre-symptomatic transmission, relying solely on isolation of symptomatic
- individuals may reduce but not eliminate influenza transmission. Although isolation measures initiated
- 320 after symptom onset are likely to mitigate at least some influenza spread, given the relatively low levels
- of asymptomatic infection and pre-symptomatic transmission among the majority of individuals, there
- 322 remains large heterogeneity (Morris, et al. 2024). A layered approach, including isolation of ill or
- infected people, maintaining good respiratory hygiene, and promoting influenza vaccination, may be
- 324 most effective to reduce transmission within households (Centers for Disease Control and Prevention
- 2024). This underscores the importance of vaccination as the primary recommendation to prevent
- 326 influenza-associated morbidity and mortality, especially for individuals at increased risk of influenza
- 327 complications.

328 Modeling details and limitations

- 329 While our reliance on household data might introduce limitations, such as the presence of multiple co-
- primary cases, sensitivity analyses confirmed the robustness of our generation time estimates to various
- data stratifications and model assumptions. Nevertheless, there are several limitations to our study.
- First, we did not account for vaccination status. This omission is less likely to impact our estimates for
- influenza given similarities in viral load dynamics between infected vaccinated and unvaccinated
- individuals (Morris, et al. 2024) (Suess, et al. 2012). Nevertheless, it is possible that different vaccination
- 335 statuses could be associated with different generation time estimates.
- 336 Second, we did not account for potential exposures outside households. When we excluded household
- members who did not have at least two valid PCR tests, the household sizes were reduced in our
- analysis. While this might not directly affect our generation time estimates, it could lead to an
- 339 overestimation of overall infectiousness due to the absence of unobserved uninfected members.
- 340 Third, it is essential to acknowledge the influence of the COVID-19 pandemic on our influenza data.
- 341 Generalizing these findings to pre-pandemic or post-pandemic periods should be done with caution.
- 342 During the period when this study was conducted, individuals may have been more likely to adopt
- 343 preventive measures against transmission within the home, such as self-isolating, practicing good

- respiratory and hand hygiene, wearing masks, and reducing contact with household members. These
- non-pharmaceutical interventions, alongside changes in human behavior and heightened awareness of
- infection control, could have impacted the spread of influenza.

347 Conclusions

- Through comprehensive data collected during the 2021/2022 and 2022/2023 influenza seasons in the
- U.S., we provide updated estimates of the generation time, essential for informing influenza modeling
- and public health strategies. Despite the significant changes in public behavior and preventive measures
- 351 due to the COVID-19 pandemic, our study did not detect substantial changes in the generation time of
- influenza since the 2009 influenza pandemic. This finding is particularly significant given that the study
- 353 period followed the extreme measures implemented to prevent COVID-19, which also reduced the
- 354 transmission of influenza and other respiratory infections. Our findings contribute to our understanding
- 355 of influenza transmission dynamics within households and underscore the importance of ongoing
- 356 research for effective outbreak management.

358 Author contributions

	1	2	3	4	5	6	7	8	9	10	11	12	13	14
LYHC	Y	-	Y	-	Y	Y	-	-	Y	-	Y	Y	Y	Y
SEM	-	-	-	-	Y	Y	-	-	-	-	-	-	-	Y
MSS	-	Y	-	-	-	-	-	-	-	-	-	-	-	Y
NMB	-	Y	-	-	Y	-	-	-	-	-	-	-	-	Y
EA	-	Y	-	-	-	-	-	-	-	-	-	-	-	Y
SR	-	Y	-	-	-	-	-	-	-	-	-	-	-	Y
KL	-	Y	-	-	-	-	-	-	-	-	-	-	-	Y
KDE	-	Y	-	-	-	-	-	-	-	-	-	-	-	Y
HQN	-	Y	-	-	-	-	-	-	-	-	-	-	-	Y
YM	-	Y	-	-	-	-	-	-	-	-	-	-	-	Y
SHM	-	Y	-	-	-	-	-	-	-	-	-	-	-	Y
ES	-	Y	-	-	-	-	-	-	-	-	-	-	-	Y
JEB	-	Y	-	-	-	-	-	-	-	-	-	-	-	Y
SESJ	-	Y	-	-	-	-	-	-	-	-	-	-	-	Y
MB	-	-	-	-	-	-	-	-	-	Y	-	-	-	Y
MAR	-	Y	-	-	-	-	-	-	-	-	-	-	-	Y
НКТ	-	Y	-	Y	-	-	-	-	-	-	-	-	-	Y
CGG	-	Y	-	Y	-	-	-	-	-	-	-	-	-	Y
RKB	Y	-	Y	-	Y	Y	-	-	-	Y	-	-	-	Y
AMM	Y	Y	Y		Y	Y	Y	Y	-	Y		-	Y	Y

359

379

... choose from the following items: <u>https://credit.niso.org</u> /

360 <u>https://www.elsevier.com/researcher/author/policies-and-guidelines/credit-author-statement</u>

- 361 1. Conceptualization
- 362 2. Data curation
- 363 3. Formal analysis
- 364 4. Funding acquisition
- 365 5. Investigation
- 366 6. Methodology
- 367 7. Project administration
- 368 8. Resources
- 369 9. Software
- 370 10. Supervision
- 371 11. Validation
- 372 12. Visualization
- 373 13. Roles/Writing original draft
- 374 14. Writing review & editing.

375 Acknowledgments

- The authors thank the following members of the Respiratory Virus Transmission Network –
 Sentinel (RVTN-S) study teams:
 Vanderbilt University Medical Center: Chris Lindsell, Judy King, John Meghreblian,
 - Samuel Massion, Brittany Creasman, Lauren Milner, Andrea Stafford Hintz, Jorge

380		Celedonio, Ryan Dalforno, Maria Catalina Padilla-Azain, Daniel Chandler, Paige Yates,
381		Brianna Schibley-Laird, Alexis Perry, Ruby Swaimn, Mason Speirs, Erica Anderson,
382		Suryakala Sarilla, Amelia Dodds, Dayton Marchlewski, Timothy Williams, Afan Swan,
383		Onika Abrams, Jackson Resser, Ine Sohn, Cara Lwin, Hsi-nien (Jubilee) Tan, Stephen
384		Yeargin, James Grindstaff, Heather Prigmore, Jessica Lai, Zhouwen Liu, James D.
385		Chappell, Marcia Blair, Rendie E. McHenry, Bryan P. M. Peterson, Lauren J. Ezzell.
386	b.	Columbia University: Lisa Saiman, Raul A. Silverio Francisco, Anny L. Diaz Perez, Ana M.
387		Valdez de Romero.
388	С.	Stanford University: Rosita Thiessen, Marcela Lopez, Alondra A. Aguilar, Emma Stainton,
389		Grace K-Y. Tam, Jonathan Altamirano, Leanne X. Chun, Rasika Behl, Samantha A.
390		Ferguson, Yuan J. Carrington, Frank S. Zhou.
391	d.	Marshfield Clinic Research Institute: Edward A. Belongia, Hannah Berger, Vicki Moon,
392		Gina Burbey, Leila Deering, Brianna Freund, Garrett Heuer, Sarah Kopitzke, Carrie
393		Marcis, Jennifer Meece, Jennifer Moran, DeeAnn Hertel, Joshua Petrie, Miriah Rotar,
394		Carla Rottscheit, Elisha Stefanski, Sandy Strey, Melissa Strupp.
395	e.	University of Arizona: Ferris Alaa Ramadan, Flavia Maria Nakayima Miiro, Josue Ortiz,
396		Mokenge Ndiva Mongoh.
397	f.	University of North Carolina: Ayla Bullock, Amy Yang, Quenla Haehnel, Jessica Lin,
398		Julienne Reynolds, Katherine "Katie" Murray, Miriana Moreno Zivanovich, Anna
399		McShea, Brittney Figueroa, Melody Liu.
400	g.	University of Colorado: Kathleen Grice, Cameron Bendalin, Sonia Chavez, Jolie Granger.
401	• We also	o acknowledge the households for their participation.

402 • L. Y. H. C. thanks the CDC Steven M. Teutsch Prevention Effectiveness (PE) Fellowship.

403 Data sharing statement

The household data are available upon reasonable request and upon completion of required approvals.
 The R code for estimating the generation time is available at https://github.com/CDCgov/influenza-

406 generation time-us.

407 Disclaimer

- 408 The conclusions, findings, and opinions expressed by authors contributing to this article do not
- 409 necessarily reflect the official position of the U.S. Department of Health and Human Services, the Public
- 410 Health Service, the Centers for Disease Control and Prevention, or the authors' affiliated institutions.

411 Declaration of Generative AI and AI-assisted technologies in the

412 writing process

- 413 During the preparation of this work the authors used ChatGPT in order to enhance the clarity,
- 414 coherence, and correctness of the writing, and to check for grammatical errors. After using this tool, the
- 415 authors reviewed and edited the content as needed and take full responsibility for the content of the
- 416 publication.

417 Declaration of interest

- M. S. S. reports a leadership role as Associate Director of the American Academy of Pediatrics'
 Pediatric Research in Office Settings (PROS), paid to Trustees of Columbia University. All other
 authors report no potential conflicts.
- N. M.B. reports grant/contracts from NIH to the University of North Carolina School of Medicine, Doris Duke Charitable Foundation, and North Carolina Collaboratory; participation on a DSMB or advisory board for the Snowball Study Technical Interchange; a leadership or fiduciary role on the American Society of Tropical Medicine and Hygiene Scientific Committee; and other financial or nonfinancial interests with the COVID-19 Equity Evidence Academy (RADx-UP CDCC) Steering Committee and North Carolina Occupational Safety and Health Education Research Center.
- E. A. reports serving as a former consultant for Hillevax and Moderna, presenting a Merck supported lecture at the Latin American Vaccine Summit, and receipt of grant/research support
 from Pfizer for pneumococcal pneumonia studies.
- 430 S. R. reports grant support from BioFire.
- H. Q. N. reports grant/research support from CSL Seqirus, GSK, and ModernaTX, and honorarium for participating in a consultancy group for ModernaTX outside the submitted work.
- S. H. M. reports grants/contracts from NIH, the American Academy of Pediatrics, and the Doris
 Duke Charitable Foundation.
- E. S. reports grants or contracts to institution from Vanderbilt University Medical Center
 (originating at CDC #75D30121C11656).
- H. K. T. has received research funding from the CDC.
- C. G. G. reports participation on an advisory board for Merck, and receipt of grant/research
 support from AHRQ, CDC, US Food and Drug Administration, NIH, and Syneos Health.
- All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest.
 Conflicts that the editors consider relevant to the content of the manuscript have been
 disclosed.

443 Financial support

- The parent household transmission study (the Respiratory Virus Transmission Network –
 Sentinel) was funded by the Centers for Disease Control and Prevention (Centers for Disease
 Control and Prevention; contracts 75D30121C11656 and 75D30121C11571) and National Center
 for Advancing Translational Sciences (NCATS) Clinical Translational Science Award (CTSA)
 Program, Award Number 5UL1TR002243-03.
- S. H. M. reports support to institution (Trustees of Columbia University) from Vanderbilt
 University Medical Center (project 75D30121C11656).
- M. S. S. reports a subcontract from Vanderbilt University Medical Center (funding originated from CDC) paid to Trustees of Columbia University.

453

455 References

- Boëlle, Pierre-Yves, Séverine Ansart, Anne Cori, and Alain-Jacques Valleron. 2011. "Transmission
 parameters of the A/H1N1 (2009) influenza virus pandemic: a review." *Influenza and Other Respiratory Viruses* (Wiley Online Library) 5 (5): 306-316.
- Buitrago-Garcia, Diana, Aziz Mert Ipekci, Leonie Heron, Hira Imeri, Lucia Araujo-Chaveron, Ingrid
 Arevalo-Rodriguez, Agustín Ciapponi, et al. 2022. "Occurrence and transmission potential of
 asymptomatic and presymptomatic SARS-CoV-2 infections: Update of a living systematic review
 and meta-analysis." *PLOS Medicine* (Public Library of Science) 19 (5): e1003987.
- Carrat, Fabrice, Elisabeta Vergu, Neil M. Ferguson, Magali Lemaitre, Simon Cauchemez, Steve Leach, and
 Alain-Jacques Valleron. 2008. "Time Lines of Infection and Disease in Human Influenza: A Review
 of Volunteer Challenge Studies." *American Journal of Epidemiology* (Oxford University Press) 167
 (7): 775-785.
- 467 Cauchemez, Simon, Christl A. Donnelly, Carrie Reed, Azra C. Ghani, Christophe Fraser, Charlotte K. Kent,
 468 Lyn Finelli, and Neil M. Ferguson. 2009. "Household Transmission of 2009 Pandemic Influenza A
 469 (H1N1) Virus in the United States." *New England Journal of Medicine* (Mass Medical Soc) 361
 470 (27): 2619-2627.
- 471 Cauchemez, Simon, Fabrice Carrat, Cecile Viboud, Alain Jacques Valleron, and Pierre-Yves Boëlle. 2004.
 472 "A Bayesian MCMC approach to study transmission of influenza: application to household
 473 longitudinal data." *Statistics in Medicine* (Wiley Online Library) 23 (22): 3469-3487.
- 474 Centers for Disease Control and Prevention. 2024. *Current Epidemic Growth Status (Based on Rt) for* 475 *States and Territories.* March 22. Accessed March 25, 2024. https://www.cdc.gov/forecast 476 outbreak-analytics/about/rt-estimates.html.
- 477 —. 2024. Preventing Spread of Respiratory Viruses When You're Sick. March 1. Accessed June 11, 2024.
 478 https://www.cdc.gov/respiratory-viruses/prevention/precautions-when-sick.html.
- 479 —. 2024. Technical Blog: Improving CDC's Tools for Assessing Epidemic Growth. March 5. Accessed
 480 March 25, 2024. https://www.cdc.gov/forecast-outbreak-analytics/about/technical-blog 481 rt.html.
- 482 Cori, A., A.J. Valleron, F. Carrat, G. Scalia Tomba, G. Thomas, and P.Y. Boëlle. 2012. "Estimating influenza
 483 latency and infectious period durations using viral excretion data." *Epidemics* 4 (3): 132-138.
- Cowling, Benjamin J, Vicky J Fang, Steven Riley, J S Malik Peiris, and Gabriel M Leung. 2009. "Estimation
 of the serial interval of influenza." *Epidemiology* (LWW) 20 (3): 344--347.
- Cowling, Benjamin J., Kwok Hung Chan, Vicky J. Fang, Lincoln L.H. Lau, Hau Chi So, Rita O.P. Fung,
 Edward S.K. Ma, et al. 2010. "Comparative Epidemiology of Pandemic and Seasonal Influenza A
 in Households." *New England Journal of Medicine* (Mass Medical Soc) 362 (23): 2175-2184.
- Fraser, Christophe, Christl A. Donnelly, Simon Cauchemez, William P. Hanage, Maria D. Van Kerkhove, T.
 Déirdre Hollingsworth, Jamie Griffin, et al. 2009. "Pandemic Potential of a Strain of Influenza A
 (H1N1): Early Findings." *Science* (American Association for the Advancement of Science) 324
 (5934): 1557-1561.

493 Gostic, Katelyn M., Lauren McGough, Edward B. Baskerville, Sam Abbott, Keya Joshi, Christine Tedijanto, 494 Rebecca Kahn, et al. 2020. "Practical considerations for measuring the effective reproductive 495 number, Rt." PLOS Computational Biology (Public Library of Science) 16 (12): e1008409. Hart, William S, Elizabeth Miller, Nick J Andrews, Pauline Waight, Philip K Maini, Sebastian Funk, and 496 497 Robin N Thompson. 2022. "Generation time of the alpha and delta SARS-CoV-2 variants: an 498 epidemiological analysis." The Lancet Infectious Diseases (Elsevier) 22 (5): 603--610. 499 Hart, William S, Philip K Maini, and Robin N Thompson. 2021. "High infectiousness immediately before 500 COVID-19 symptom onset highlights the importance of continued contact tracing." eLife (eLife 501 Sciences Publications, Ltd) 10: e65534. 502 Hart, William S, Sam Abbott, Akira Endo, Joel Hellewell, Elizabeth Miller, Nick rews, Philip K Maini, 503 Sebastian Funk, and Robin N Thompson. 2022. "Inference of the SARS-CoV-2 generation time 504 using UK household data." eLife (eLife Sciences Publications, Ltd) 11: e70767. 505 Lau, Max S.Y., Benjamin J. Cowling, Alex R. Cook, and Steven Riley. 2015. "Inferring influenza dynamics and control in households." Proceedings of the National Academy of Sciences (National Acad 506 507 Sciences) 112 (29): 9094-9099. 508 Lessler, Justin, Nicholas G Reich, Ron Brookmeyer, Trish M Perl, Kenrad E Nelson, and Derek AT Cummings. 2009. "Incubation periods of acute respiratory viral infections: a systematic review." 509 510 The Lancet infectious diseases (Elsevier) 9 (5): 291--300. 511 Levy, Jens W., Benjamin J. Cowling, James M. Simmerman, Sonja J. Olsen, Vicky J. Fang, Piyarat

- Levy, Jens W., Benjamin J. Cowling, James M. Simmerman, Sonja J. Olsen, Vicky J. Fang, Piyarat
 Suntarattiwong, Richard G. Jarman, Brendan Klick, and Tawee Chotipitayasunondh. 2013. "The
 Serial Intervals of Seasonal and Pandemic Influenza Viruses in Households in Bangkok, Thailand."
 American Journal of Epidemiology (Oxford University Press) 177 (12): 1443-1451.
- Morris, Sinead E., Huong Q. Nguyen, Carlos G. Grijalva, Kayla E. Hanson, Yuwei Zhu, Jessica E. Biddle,
 Jennifer K. Meece, et al. 2024. "Influenza virus shedding and symptoms: Dynamics and
 implications from a multi-season household transmission study." *medRxiv* (Cold Spring Harbor
 Laboratory Press) 2024--03.
- Pastore, Massimiliano. 2018. "Overlapping: a R package for Estimating Overlapping in Empirical
 Distributions." *Journal of Open Source Software* (The Open Journal) 3 (32): 1023.
- Pastore, Massimiliano, and Antonio Calcagnì. 2019. "Measuring Distribution Similarities Between
 Samples: A Distribution-Free Overlapping Index." *Frontiers in Psychology* (Frontiers Media SA)
 10: 1089.
- Petrie, Joshua G., Suzanne E. Ohmit, Benjamin J. Cowling, Emileigh Johnson, Rachel T. Cross, Ryan E.
 Malosh, Mark G. Thompson, and Arnold S. Monto. 2013. "Influenza Transmission in a Cohort of Households with Children: 2010-2011." *PLOS ONE* (Public Library of Science) 8 (9): e75339.

Rolfes, Melissa A., H. Keipp Talbot, Huong Q. McLean, Melissa S. Stockwell, Katherine D. Ellingson, Karen Lutrick, Natalie M. Bowman, et al. 2023. "Household Transmission of Influenza A Viruses in 2021-2022." JAMA (American Medical Association) 329 (6): 482-489.

Suess, Thorsten, Cornelius Remschmidt, Susanne B. Schink, Brunhilde Schweiger, Alla Heider, Jeanette
 Milde, Andreas Nitsche, et al. 2012. "Comparison of Shedding Characteristics of Seasonal
 Influenza Virus (Sub)Types and Influenza A(H1N1)pdm09; Germany, 2007–2011." *PLOS ONE* (Public Library of Science) 7 (12): e51653.

- 534 Svensson, Åke. 2007. "A note on generation times in epidemic models." *Mathematical Biosciences* 535 (Elsevier) 208 (1): 300-311.
- te Beest, Dennis E, Jacco Wallinga, Tjibbe Donker, and Michiel van Boven. 2013. "Estimating the
 generation interval of influenza A (H1N1) in a range of social settings." *Epidemiology* (LWW) 24
 (2): 244--250.
- 539Tsang, Tim K, Lincoln LH Lau, Simon Cauchemez, and Benjamin J Cowling. 2016. "Household transmission540of influenza virus." *Trends in microbiology* (Elsevier) 24 (2): 123--133.
- Tsang, Tim K., Can Wang, Vicky J. Fang, Ranawaka A. P. M. Perera, Hau Chi So, Dennis K. M. Ip, Gabriel
 M. Leung, J. S. Malik Peiris, Simon Cauchemez, and Benjamin J. Cowling. 2023. "Reconstructing
 household transmission dynamics to estimate the infectiousness of asymptomatic influenza
 virus infections." *Proceedings of the National Academy of Sciences* 120 (33): e2304750120.
- Tuite, Ashleigh R., Amy L. Greer, Michael Whelan, Anne-Luise Winter, Brenda Lee, Ping Yan, Jianhong
 Wu, et al. 2010. "Estimated epidemiologic parameters and morbidity associated with pandemic
 H1N1 influenza." *CMAJ* 182 (2): 131--136.
- Xu, Cuiling, Kwok-Hung Chan, Tim K. Tsang, Vicky J. Fang, Rita O. P. Fung, Dennis K. M. Ip, Simon
 Cauchemez, Gabriel M. Leung, J. S. Malik Peiris, and Benjamin J. Cowling. 2015. "Comparative
 Epidemiology of Influenza B Yamagata- and Victoria-Lineage Viruses in Households." *American Journal of Epidemiology* (Oxford University Press) 182 (8): 705-713.
- 552

554 Supplementary material

555 The incubation period distribution

- 556 The incubation period distribution was modeled using estimates for influenza A from a systematic
- review by Lessler et al. (Lessler, et al. 2009), with a mean of 1.55 days and a standard deviation (SD) of
- 558 0.66 days. These estimates were fitted to a gamma distribution to characterize the distribution of the
- incubation period (Supplemental Figure S1).



560

Figure S1. Incubation period distribution. The black circles and blue lines represent the data (Lessler, et al. 2009), and the (A)
 cumulative distribution function and (B) probability density function of a gamma distribution fitted to the data.

563 The observed household serial interval of single infection pairs

564 We found that the observed household serial interval, calculated without modeling, solely using data

565 from households with single infection pairs (i.e., single primary case to single secondary case) and

566 without potential transmission chains, had a mean of 3.7 days (and a SD of 2.3 days). This was longer

than the mean intrinsic serial interval of 3.2 (95% CrI: 2.8-3.5) days when considering households of all

568 sizes with all potential transmission chains (Table 2). This does not necessarily indicate that the intrinsic

- 569 value was shorter than the realized household one. Rather, it is mainly due to the restriction of single
- 570 infection pairs or mostly smaller household sizes of 2 members.
- 571 In the main text, we found slightly longer mean intrinsic and realized household generation times in
- 572 smaller households compared to larger ones (Figure 1C and Supplemental Table S1). Larger households
- 573 with more exposure and potential transmission chains could have a shorter interval, while smaller
- 574 households could have a longer interval.

575 Specification of parameters for the mechanistic model

- 576 In the mechanistic model (Hart, Abbott, et al. 2022), two parameters, namely the ratio of the mean
- 577 latent and incubation period and the mean symptomatic infectious period, were estimated directly
- 578 (Supplemental Figure S2), while the proportion of transmission before symptomatic onset was
- 579 calculated by weighting the pre-symptomatic period by the ratio of pre-symptomatic and symptomatic
- 580 transmission rates and dividing it by the sum of the (pre-symptomatic and symptomatic) infectious
- 581 periods. The mean latent and mean pre-symptomatic periods were calculated by dividing the incubation
- 582 period by the ratio of mean latent and incubation period.



583

Figure S2. Posterior and prior distributions of estimated parameters. Solid and dashed lines represent posterior and prior
 distributions, respectively.

586 Variability in estimates across data stratifications

- 587 Although the generation time or serial interval of influenza B may be longer than that of influenza A
- 588 (Levy, et al. 2013), this was not the case in our findings from the two seasons (Supplemental Table S1,
- 589 Figure S3 and S4). However, we note that the mean intrinsic generation time exhibited a wider credible
- 590 interval when using data exclusively from influenza B compared to influenza A, which likely reflects the
- 591 dominance of influenza A during the study timeframe and the smaller sample size of influenza B.

Data stratifications	Mean intrinsic generation time (95% Crls)	Overlapping index (%, compared to the primary analysis)
All data excluding households with multiple co-primary cases (primary analysis in Table 1)	3.2 (2.9-3.6)	100
Season 2021/2022	3.3 (2.8-4.0)	71
Season 2022/2023	3.2 (2.8-3.6)	87
Influenza A	3.2 (2.9-3.6)	94
Influenza B	3.2 (2.3-4.5)	47

Household size of 2 or 3	3.4 (2.9-4.0)	61
Household size of 4 or greater	3.1 (2.7-3.6)	74
All data including households with	3.1 (2.7-3.4)	64
multiple co-primary cases		

592 Table S1. The posterior mean (95% CrIs) of mean intrinsic generation time across seasons, virus types, household sizes, and with

593 multiple co-primary cases. The incubation period, derived from influenza A, had a mean of 1.55 days and a standard deviation

594 (SD) of 0.66 days (Lessler, et al. 2009). Only for influenza B, we assumed the shorter incubation period to yield a mean of 0.61 595 days and a standard deviation (SD) of 0.25 days.





Figure S3. Posterior distributions of parameters across data stratifications: (A) seasons, (B) virus types, (C) household sizes, and
 (D) with multiple co-primary cases.





Figure S4. Posterior distributions of parameters across data stratifications: (A) seasons, (B) virus types, (C) household sizes, and
 (D) with multiple co-primary cases.

602 Sensitivity analyses

- 603 Similar to the sensitivity analyses using the full dataset, we found that the incubation period had a
- 604 limited effect on the intrinsic generation time when exclusively using data from households circulating
- 605 influenza A (Supplemental Figure S6, Panel B) or households circulating influenza B (Supplemental Figure
- 606 S6, Panel C).

607 Consistent with the previous study (Hart, Abbott, et al. 2022), assuming a higher relative infectiousness

of asymptomatic infected individuals resulted in slightly lower estimates of the overall infectiousness of

609 infectors (Supplemental Figure S6, Panel D).

Sensitivity analyses	Mean intrinsic generation time (95% Crls)	Overlapping index (%, compared to the primary analysis)
Primary analysis (in Table 1)	3.2 (2.9-3.6)	100
Longer incubation period	3.2 (2.8-3.6)	86
Shorter incubation period	3.4 (3.1-3.7)	56
Lower relative infectiousness	3.2 (2.9-3.6)	94
Higher relative infectiousness	3.2 (2.9-3.6)	97

610 Table S2. The posterior mean (95% CrIs) of mean intrinsic generation time given different incubation periods or relative

611 infectiousness of asymptomatic infected individuals. The primary incubation period, derived from influenza A, had a mean of

612 1.55 days and a standard deviation (SD) of 0.66 days (Lessler, et al. 2009). For the shorter incubation period derived from

613 influenza B, we assumed a mean of 0.61 days and a SD of 0.25 days (Lessler, et al. 2009). For the longer incubation period

614 derived from influenza A(H1N1)pdm09, we assumed a mean of 0.61 days and a SD of 0.25 days (Tuite, et al. 2010).



615

616 Figure S5. Posterior distributions of parameters given different assumptions: (A-C) incubation periods, and (D) relative

617 infectiousness of asymptomatic infected individuals. Panel (A) presents results obtained using data from households with both
618 influenza A and B, whereas Panels (B) and (C) present results obtained using data solely from households with influenza A and B,
619 respectively.



620

621 Figure S6. Posterior distributions of parameters given different assumptions: (A-C) incubation periods, and (D) relative

622 infectiousness of asymptomatic infected individuals. Panel (A) presents results obtained using data from households with both
 623 influenza A and B, whereas Panels (B) and (C) present results obtained using data solely from households with influenza A and B,

624 respectively.