

Single-cell RNA Sequencing of Peripheral Blood Mononuclear Cells in Patients with Single Ventricle/Hypoplastic Left Heart Syndrome

Hui-Qi Qu¹, Kayleigh Ostberg¹, Diana J Slater¹, Fengxiang Wang¹, James Snyder¹, Cuiping Hou¹, John J Connolly¹, Michael March¹, Joseph T Glessner^{1,2,3}, Charlly Kao¹, Hakon Hakonarson^{1,2,3,4,5*}.

¹The Center for Applied Genomics, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, 19104, USA.

²Department of Pediatrics, The Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, 19104, USA.

³Division of Human Genetics, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, 19104, USA.

⁴Division of Pulmonary Medicine, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, 19104, USA.

⁵Faculty of Medicine, University of Iceland, 101 Reykjavik, Iceland

* Corresponding author:

Hakon Hakonarson M.D., Ph.D.

Center for Applied Genomics (CAG)

3615 Civic Center Blvd

Abramson Building

Philadelphia, PA 19104, United States of America

Telephone: 267-426-0088

Fax: 267-426-0363

Email: hakonarson@email.chop.edu

Abstract

Background: Single ventricle and hypoplastic left heart syndrome (SV/HLHS) patients require lifelong medical monitoring and management to address potential complications and optimize their health. The consequence of SV/HLHS had detrimental effects on multiple organ systems, including on peripheral blood mononuclear cells (PBMCs) and can weaken the immune system, exacerbating the risk of infection and various cardiovascular complications.

Methods: Using single-cell RNA sequencing (scRNA-seq), we studied PBMCs from 33 pediatric patients (10 females and 23 males) with SV/HLHS. By a pair-wise study design, the SV/HLHS patients were compared to 33 controls without heart diseases.

Results: Four cell types account for the top 62% cumulative importance of disease effects on gene expression in different cell types, i.e., [T cells, CD4+, Th1/17], [T cells, CD4+, TFH], [NK cells], and [T cells, CD4+, Th2]. Significant sex differences were observed in [T cells, CD4+, TFH], with less prominent effects in female patients. A total of 6659 genes in different cell types were significantly differentially expressed (DE). Hierarchical clustering by WGCNA analysis of the DE genes revealed that DE genes in NK cells are most closely related to those in SV/HLHS. A total of 822 genes showed cell specific DE with opposite directions in different cell types, highlighting overrepresented MYC and IFN- γ activity in T cell and NK cell populations, as well as underrepresentation in monocytes and Treg cells.

Conclusion: This study elucidates the complex transcriptome landscape in PBMCs in patients with SV/HLHS, emphasizing the differential impacts on various cell types. New insights are gained into the precise modulation of MYC and IFN- γ activity in SV/HLHS, which may help balance immune responses and reduce harmful inflammation, and promote effective tissue repair and infection control.

Keywords: monocytes; NK cells; peripheral blood mononuclear cells; single-cell RNA sequencing; Treg cells

1. Introduction

Single ventricle and hypoplastic left heart syndrome (SV/HLHS) are complex congenital heart defects, necessitating restructuring of circulation to optimize blood flow and oxygenation¹. Affected patients require lifelong medical monitoring and management to address potential complications and optimize their health. Peripheral blood mononuclear cells (PBMCs) have been implicated in the pathophysiology of various cardiovascular complications, including heart failure, arrhythmias, and vascular dysfunction. In particular, infection is a significant complication for individuals with SV/HLHS. The frequent use of central lines, catheters, and other medical devices in these patients serves as potential sources of infection². In addition to bloodstream infections, patients are also at an increased risk for respiratory infections due to impaired immune function and potential respiratory complications from heart failure³. The chronic impact of SV/HLHS on PBMCs can weaken the immune system, exacerbating the risk of infection and various cardiovascular complications. Immunosuppression heightens susceptibility to both bloodstream and respiratory infections, further complicating clinical management. The altered immune profile contributes to cardiovascular issues such as heart failure, arrhythmias, and vascular dysfunction. Understanding the molecular and cellular

mechanisms underlying these complications is crucial. Key research questions include identifying the specific gene expression changes in PBMCs of patients with SV/HLHS compared to healthy controls and elucidating the underlying molecular pathways in PBMCs that contribute to the altered immune profile in SV/HLHS patients. Transcriptome profiling of PBMCs can offer valuable insights into the pathogenesis of SV/HLHS-related complications.

Using single-cell RNA sequencing (scRNA-seq), we examined the impact of SV/HLHS on PBMCs, uncovering sex differences across various cell types. By focusing on differentially expressed (DE) genes in SV/HLHS, we identified cell type-specific expression of a number of genes related to SV/HLHS. Additionally, we highlighted co-expressed gene sets related to SV/HLHS, providing new insights into the molecular underpinnings of the disease.

2. Methods

2.1 Subjects

This study included 33 pediatric patients (10 females and 23 males) with SV/HLHS. Among these patients, 20 had European ancestry, 9 had African ancestry, 1 had Asian ancestry, and 3 had other ancestries. The ages ranged from 0 to 21 years, with a median age of 6 years. Fifteen patients were diagnosed with SV, 24 were diagnosed with HLHS, and 18 underwent the Fontan procedure. Using a pair-wise study design, these patients were compared to 10 female and 23 male de-identified children without heart disease. All experimental protocols were approved by the Institutional Review Board (IRB) of the Children's Hospital of Philadelphia (CHOP) with the IRB number: IRB 16-013278. Informed consent was obtained from all subjects. If subjects are under 18, consent was also obtained from a parent and/or legal guardian with assent from the child if 7 years or older.

2.2 PBMC scRNA-seq

Blood samples were collected from each participant in EDTA-coated tubes and immediately processed at the Center for Applied Genomics (CAG) at CHOP. PBMCs were isolated using Ficoll density gradient centrifugation. scRNAseq was conducted with the 10X Chromium Single Cell Gene Expression assay (10x Genomics, Single Cell 3' v3). Sequencing was carried out on the Illumina HiSeq2500 platform using SBS v4 chemistry. The resulting data from the Chromium single-cell RNA sequencing were processed and analyzed using Cell Ranger 7.1.0 software (10x Genomics), with reads aligned to the GRCh38 reference genome.

Each pair of subjects was considered an independent experiment. The scRNA-seq data from the 33 independent pairs of children were analyzed using the Seurat R package^{4,5} for pair-wise comparison, employing SCTransform for normalization and scaling. To enhance comparability among samples, Harmony⁶ was used to align datasets post-principal component analysis (PCA). Uniform manifold approximation and projection (UMAP) (Becht et al. 2019) was applied to group cells into clusters. Analysis included 15 cell types, identified using singleR and the `celldex::DatabaseImmuneCellExpression Data()` function⁷ (Supplementary Figure 1).

2.3 Assessing cell types affected by SV/HLHS using machine learning

To assess the impact of SV/HLHS on different cell types, data were structured into a matrix where rows represented individual genes and subject pairs, and columns denoted different cell

types. Each cell in the matrix contained the log₂ fold change (log₂FC) in gene expression for each cell type between each case/control pair. Missing values, due to insufficient cell numbers for a specific gene in a cell type, were imputed using the mean log₂FC from other experiments. If a gene was not expressed in a particular cell type, it was assigned a value of 0.

A RandomForestRegressor from scikit-learn⁸ was selected for its ability to manage high-dimensional data and generate feature importance metrics. The dataset was split into training and testing sets with the 0.8-0.2 train-test split. The number of trees (n_estimators) was set to 100, max_depth to none, min_samples_split to 2, and random_state to 42. The model's robustness was assessed using 5-fold cross-validation (CV) to prevent overfitting, with the mean squared error (MSE) calculated for each fold to evaluate performance. Post-training, feature importance values were used for indicating the relative contribution of each cell type to the model's predictions. Cell types were then ranked according to their importance scores. To estimate the statistical significance of feature importance values, the permutation_importance function from sklearn.inspection was employed. This function shuffled feature values to perform 1000 permutations for each feature. The p-value for each feature was computed as the proportion of permuted importance values that were equal to or greater than the original importance values.

2.4 Differential expression (DE) analysis

Based on the results of the Seurat R package^{4,5}, a gene with DE was defined as FDR<0.05 in at least 2 pairs of samples, and with log₂FCs in the same direction within the same cell type. Distribution of DE genes by cell type is visualized with the Venn diagram and the UpSetR package⁹.

2.5 Weighted Gene Co-expression Network Analysis (WGCNA) of DE genes

WGCNA analysis was done in 6319 DE genes with expression detected in each of the 66 PBMC samples. The average expression level per cell type for each gene in each sample was calculated using log-normalized values with the Seurat R package^{4,5}. The WGCNA analysis was performed using the WGCNA R package^{10,11}. Hallmark gene set overrepresentation analysis (ORA) were done using the clusterProfiler R package¹² and the msigdb R package¹³.

3. Results

3.1 Cell types affected by SV/HLHS

The CV MSE scores, the mean MSEs, and the standard deviations are shown in Table 1. The test MSEs are close to the mean CV MSEs in all three groups, indicating consistent model performance. The 15 cell types along with their corresponding feature importance values are shown in Table 2. Among the 15 cell types, two cell types, i.e. [T cells, CD8+, naive, stimulated] and [T cells, CD4+, naive, stimulated], were informative in less than 50% of the subject pairs. All the other 13 cell types were informative in at least 27 (82%) out of the 33 pairs of subjects.

Four cell types account for the top 62% cumulative importance in disease effects on gene expression in different cell types, i.e., [T cells, CD4+, Th1/17], [T cells, CD4+, TFH], [NK cells], and [T cells, CD4+, Th2]. All four cell types have permutation P-values<0.001 in all subject pairs. Significant sex differences were observed in [T cells, CD4+, TFH], which was not affected in female patients (Table 2, Figure 1).

3.2 DE genes

A total of 6659 genes in different cell types showed significant DE in the same direction in at least two independent pairs of samples (Supplementary Table 1). Gene sets that play critical roles in energy production, immune regulation, cell survival and proliferation, and maintenance of cellular integrity are significantly overrepresented (Table 3, Supplementary Figure 2). The distribution of the DE genes by cell type is shown in Figure 2. Among these genes, 822 genes showed opposite directions in different cell types (Supplementary Table 2).

3.3 WGCNA of DE genes

Among the 6659 DE genes, 6319 genes have expression detected in each of the 66 PBMC samples. Among the 15 cell types, 9 cell types are informative in all the 66 PBMC samples, including [Monocytes, CD14+], [NK cells], [T cells, CD4+, memory Treg], [T cells, CD4+, naïve], [T cells, CD4+, TFH], [T cells, CD4+, Th1], [T cells, CD4+, Th1/17], [T cells, CD4+, Th17], and [T cells, CD4+, Th2]. The genes were grouped into different modules based on the similarity of their expression patterns across samples (Supplementary Table 3). The Hallmark gene sets highlighted in each module are shown in Supplementary Table 4. The correlations of module eigengenes with cell types, sex, and disease status are shown in Fig.3. Among these modules, the turquoise and pink modules are significantly overexpressed in SV/HLHS. The turquoise module is significantly overexpressed in [NK cells] (Fig.3). Two Hallmark gene sets, MITOTIC_SPINDLE and G2M_CHECKPOINT, are highlighted in the turquoise module (Table 4). Genes in the pink module exhibited variable expression patterns in relation to cell types and disease correlation. Specifically, these genes showed higher expression in [Monocytes, CD14+], but their expression in [Monocytes, CD14+] was downregulated in SV/HLHS. Additionally, their expression in T cells were overexpressed in SV/HLHS. No Hallmark genes were significantly overrepresented within this gene set.

4. Discussion

4.1 Immune cells affected by SV/HLHS

This study showed four cell types of T cells to be most affected in SV/HLHS, i.e., [T cells, CD4+, Th1/17], [T cells, CD4+, TFH], [NK cells], and [T cells, CD4+, Th2]. Th1/17 cells producing both IFN- γ (Th1) and IL-17 (Th17). These cell types and cytokines may help combat infections, but may also contribute to immune dysregulation involved in myocardial damage, fibrosis, and vascular dysfunction¹⁴. In these cells, the Interferon-Induced Proteins with Tetratricopeptide Repeats (IFIT) family of proteins, including *IFIT1*, *IFIT1B*, *IFIT2*, and *IFIT3*, are significantly overexpressed in SV/HLHS, indicating an upregulated response to interferon signaling¹⁵. TFH cells are critical for the formation of germinal centers and the production of high-affinity antibodies¹⁶. The activation of TFH cells in SV/HLHS may be due to recurrent infections and chronic inflammation. Their prolonged activation can exacerbate cardiovascular complications, increasing the risk of autoimmunity. Interestingly, our study showed that TFH cells in females were less affected, which may explain their better clinical outcomes. Th2 cells are involved in the immune responses, including regulation of allergic responses¹⁷. Chronic activation of these cell types can contribute to an inflammatory environment and exacerbate immune dysregulation.

4.2 NK cells and related DE genes

This study identified 6659 DE genes, which are overrepresented with critical roles in energy production, immune regulation, cell survival and proliferation, and maintenance of cellular integrity. By WGCNA analysis of the DE genes, hierarchical clustering revealed that genes expressed in NK cells are most closely related to SV/HLHS (Fig.3). NK cells are a vital component of the innate immune system, targeting and killing virally infected cells¹⁸. Genes in the lightgreen (HALLMARK_TNFA_SIGNALING_VIA_NFKB overrepresented in this module, with higher expression in NK cells, upregulated in SV/HLHS), midnightblue (HALLMARK_ALLOGRAFT_REJECTION overrepresented in this module, with higher expression in NK cells, downregulated in SV/HLHS), and turquoise (HALLMARK_MITOTIC_SPINDLE and HALLMARK_G2M_CHECKPOINT overrepresented in this module, with higher expression in NK cells, upregulated in SV/HLHS) modules were significantly correlated with NK cells (Supplementary Table 4). Upregulation of genes involved in the TNF-alpha signaling pathway via the NF-kB pathway regulates cell proliferation, differentiation, and survival, and leads to the activation of inflammatory responses¹⁹. The aggregate effects of downregulated expression of the gene set HALLMARK_ALLOGRAFT_REJECTION in NK cells may result in reduced cytotoxic activity and impaired ability to target and eliminate infected cells, weakening overall immune surveillance and response¹³. The overexpression of the genes of HALLMARK_MITOTIC_SPINDLE and HALLMARK_G2M_CHECKPOINT in NK cells can increase proliferation and cell cycle activity²⁰.

4.3 Cell type specific DE genes

Cell-type specific DEs are commonly identified in this study, underscoring cellular heterogeneity consequent to SV/HLHS (Fig.2). In particular, 822 (12%) genes showed cell-type specific DE with opposite directions in different cell types (Supplementary Table 2). Among these, a number of genes are related to MYC targets, variant 1 and interferon gamma response.

The HALLMARK_MYC_TARGETS_V1 genes are direct targets of the MYC transcription factor, which is a critical regulator of cell growth, proliferation, and metabolism²¹. In SV/HLHS, these genes tend to be upregulated in [T cells, CD4+, naive], [T cells, CD4+, TFH], [T cells, CD4+, Th1], and [T cells, CD8+, naive]; and downregulated in [Monocytes, CD14+], [Monocytes, CD16+], [T cells, CD4+, memory TREG], [T cells, CD4+, naive, stimulated], [T cells, CD4+, Th17], and [T cells, CD4+, Th2]. The differential regulation of MYC targets underscores the functional specialization between T cells and monocytes. The upregulation in naive CD4+ T cells, TFH cells, Th1 cells, and naive CD8+ T cells, particularly upon activation, require robust proliferative capacity, which is supported by MYC target upregulation²². In contrast, monocytes, which are more involved in immediate and direct immune responses, do not rely heavily on such proliferation mechanisms²³. The downregulation in memory Treg cells, which are crucial for maintaining immune tolerance and preventing autoimmunity²⁶, may lead to a reduced number for these cells to control excessive immune responses, possibly leading to autoimmunity or prolonged inflammation.

The HALLMARK_INTERFERON_GAMMA_RESPONSE genes are upregulated in response to interferon-gamma (IFN- γ), a cytokine crucial for innate and adaptive immunity²⁴. In SV/HLHS, these genes tend to be upregulated in [T cells, CD4+, naive], [T cells, CD4+, naive, stimulated], [T cells, CD4+, TFH], [T cells, CD4+, Th17], [T cells, CD4+, Th1], and [T cells, CD8+, naive]; and downregulated in [Monocytes, CD14+] and [Monocytes, CD16+]. The upregulation of IFN-

γ response genes in T cells suggests a robust activation state and readiness for immune response in these cells, indicating a potential for strong pro-inflammatory signaling. Conversely, monocytes play a crucial role in inflammation and tissue repair²³. Monocytes differentiate into macrophages and dendritic cells, which help clear debris, secrete growth factors, and regulate inflammation, thus promoting healing and regeneration²⁵. The reduced IFN- γ response in monocytes might imply a diminished capacity to manage inflammation and repair tissue²³.

Precisely modulating MYC and IFN- γ activity in SV/HLHS, based on the insights gained from our study, may balance immune responses, reduce harmful inflammation, and promote effective tissue repair and infection control. More specifically, at the cellular level, this precise modulation could involve targeting overactive T cell populations while bolstering the proliferation and function of monocytes, balancing effective infection control with minimizing harmful inflammation.

Conclusion

This study elucidates the complex transcriptome landscape in PBMCs in SV/HLHS, emphasizing the differential impacts on various cell types. The significant correlation between gene expression in NK cells and SV/HLHS highlights a cellular emphasis. A number of gene sets in NK cells offer insights into the underlying mechanisms and provide targets for intervention in SV/HLHS. Additionally, specific genes that exhibit differential expression in opposite directions across various cell types present unique targets and opportunities to fine-tune immunity in SV/HLHS. Future research should explore therapeutic interventions aimed at modulating immune responses by targeting the critical gene sets identified in this study.

Ethics Approval and Consent to Participate

All experimental protocols were approved by the Institutional Review Board (IRB) of the Children's Hospital of Philadelphia (CHOP) with the IRB number: IRB 16-013278. Informed consent was obtained from all subjects. If subjects are under 18, consent was also obtained from a parent and/or legal guardian with assent from the child if 7 years or older.

Declaration of Conflicting Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The study was supported by the Institutional Development Funds from the Children's Hospital of Philadelphia to the Center for Applied Genomics, and The Children's Hospital of Philadelphia Endowed Chair in Genomic Research to HH.

Table 1 Different parameters and train-test split ratios tested for the RandomForestRegressor

Subjects	Cross-Validation MSE Scores	Mean CV MSE	Standard Deviation of CV MSE	Test MSE
All	[4.62e-03, 5.20e-03, 4.88e-03, 5.13e-03, 5.23e-03]	5.01e-03	2.31e-04	4.81e-03
Females	[6.00e-03, 5.75e-03, 6.37e-03, 6.24e-03, 5.32e-03]	5.94e-03	3.72e-04	5.76e-03
Males	[5.54e-03, 5.45e-03, 5.73e-03, 5.56e-03, 5.68e-03]	5.59e-03	1.00e-04	5.38e-03

Table 2 Cell types with corresponding feature importance values

Cell type	Importance (all)	Permutation P-value* (all)	Importance (female)	Permutation P-value* (female)	Importance (male)	Permutation P-value* (male)
T cells, CD4+, Th1/17	0.2857	0.000	0.2874	0.000	0.2866	0.000
T cells, CD4+, TFH	0.1330	0.000	0.0491	0.939	0.1325	0.000
NK cells	0.1126	0.000	0.1269	0.000	0.0909	0.000
T cells, CD4+, Th2	0.0860	0.000	0.0488	0.000	0.0820	0.000
Monocytes, CD14+	0.0639	1.000	0.0633	1.000	0.0634	1.000
T cells, CD4+, naïve Treg	0.0528	0.000	0.0332	0.000	0.0832	0.000
T cells, CD4+, Th17	0.0500	0.000	0.0876	0.000	0.0410	0.000
Monocytes, CD16+	0.0494	1.000	0.0501	1.000	0.0484	1.000
T cells, CD8+, naïve	0.0320	1.000	0.0274	0.000	0.0316	0.984
T cells, CD4+, naïve	0.0294	0.000	0.1192	0.000	0.0262	0.000
B cells, naïve	0.0285	0.000	0.0328	0.000	0.0253	1.000
T cells, CD4+, memory Treg	0.0259	0.000	0.0229	0.000	0.0349	0.000
T cells, CD8+, naïve, stimulated [#]	0.0190	1.000	0.0194	1.000	0.0182	1.000
T cells, CD4+, Th1	0.0189	1.000	0.0223	1.000	0.0206	1.000
T cells, CD4+, naïve, stimulated [#]	0.0131	0.000	0.0094	0.000	0.0151	0.000

*Based on 1000 permutations. [#][T cells, CD8+, naïve, stimulated] and [T cells, CD4+, naïve, stimulated] were informative in less than 50% of the subject pairs.

Table 3 Hallmark gene sets overrepresented in differentially expressed genes in SV/HLHS

HALLMARK	GeneRatio	BgRatio	pvalue	p.adjust	qvalue
MYC_TARGETS_V1	164/2073	200/4383	3.64E-25	9.10E-24	5.75E-24
OXIDATIVE_PHOSPHORYLATION	164/2073	200/4383	3.64E-25	9.10E-24	5.75E-24
INTERFERON_GAMMA_RESPONSE	144/2073	200/4383	3.74E-13	6.23E-12	3.93E-12
TNFA_SIGNALING_VIA_NFKB	137/2073	200/4383	4.74E-10	5.92E-09	3.74E-09
PROTEIN_SECRETION	74/2073	96/4383	1.62E-09	1.62E-08	1.02E-08
MTORC1_SIGNALING	132/2073	200/4383	3.83E-08	3.19E-07	2.02E-07
HEME_METABOLISM	130/2073	200/4383	1.89E-07	1.35E-06	8.54E-07
INTERFERON_ALPHA_RESPONSE	67/2073	97/4383	9.69E-06	6.06E-05	3.82E-05
ALLOGRAFT_REJECTION	123/2073	200/4383	2.55E-05	1.41E-04	8.93E-05
TGF_BETA_SIGNALING	40/2073	54/4383	5.25E-05	2.63E-04	1.66E-04
IL2_STAT5_SIGNALING	120/2073	199/4383	1.12E-04	5.07E-04	3.21E-04
UNFOLDED_PROTEIN_RESPONSE	70/2073	113/4383	1.07E-03	4.47E-03	2.82E-03
REACTIVE_OXYGEN_SPECIES_PATHWAY	34/2073	49/4383	1.39E-03	5.35E-03	3.38E-03
APOPTOSIS	94/2073	161/4383	2.63E-03	9.39E-03	5.93E-03
ANDROGEN_RESPONSE	61/2073	100/4383	3.71E-03	1.24E-02	7.80E-03
DNA_REPAIR	87/2073	150/4383	4.82E-03	1.51E-02	9.52E-03
COMPLEMENT	112/2073	200/4383	7.15E-03	2.10E-02	1.33E-02
ADIPOGENESIS	110/2073	200/4383	1.54E-02	4.21E-02	2.66E-02
PI3K_AKT_MTOR_SIGNALING	61/2073	105/4383	1.60E-02	4.21E-02	2.66E-02
MITOTIC_SPINDLE	109/2073	199/4383	1.84E-02	4.59E-02	2.90E-02

Table 4 Hallmark gene sets overrepresented in the turquoise module correlated with SV/HLHS and NK cells

HALLMARK	qvalue	Genes
MITOTIC_SPINDLE	3.13E-04	<i>LRPPRC/SPTBN1/CLASP1/RAB3GAP1/FLNB/PPP4R2/RASA2/SMC4/RFC1/CENPE/ARHGAP10/RICTOR/RANBP9/CD2AP/LATS1/EZR/RAPGEF5/PCM1/ARFGEF1/ABL1/ABI1/KIF5B/PCGF5/PXN/ARHGEF7/KLC1/AKAP13/PAFAH1B1/CDC27/CSNK1D/MAPRE1/STAU1/TIAM1/PCNT/MYH9</i>
G2M_CHECKPOINT	3.13E-04	<i>MTF2/CUL3/SMARCC1/STAG1/SMC4/YTHDC1/HNRNPD/CENPE/SLC12A2/HUS1/DBF4/CUL1/EZH2/WRN/RAD21/RAD23B/ABL1/KIF5B/NUP98/RBM14/CUL5/SLC38A1/CCNT1/SLC7A1/CUL4A/ARID4A/KNL1/GSPT1/SLC7A5/PAFAH1B1/CDC27/SS18/CHAF1A/DDX39A/NUP50</i>

Supplementary Table 1 Average log₂FC of the DE genes by the 15 cell types

*Positive value means higher expression in cases.

Supplementary Table 2 DE genes with opposite directions in different cell types

*Positive value means higher expression in cases.

Supplementary Table 3 The DE genes grouped into different modules

Supplementary Table 4 Significant Hallmark gene sets overrepresented in each module

Figures




Fig.1 Cell types affected in SV/HLHS. X-axis represents feature importance in RandomForestRegressor.




Fig.2 Distribution of DE genes by cell type






Fig.3 WGCNA module relationships

Supplementary Figure 1 Cell types of PBMCs in a case and control pair.



Supplementary Figure 2 Hallmark Overrepresentation Analysis for DE genes



References

1. Ohye RG, Schranz D, D'Udekem Y. Current therapy for hypoplastic left heart syndrome and related single ventricle lesions. *Circulation* 2016;134(17):1265-79.
2. García H, Cervantes-Luna B, González-Cabello H, et al. Risk factors for nosocomial infections after cardiac surgery in newborns with congenital heart disease. *Pediatrics & Neonatology* 2018;59(4):404-09.
3. Healy F, Hanna B, Zinman R. Pulmonary complications of congenital heart disease. *Paediatric respiratory reviews* 2012;13(1):10-15.
4. Butler A, Hoffman P, Smibert P, et al. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nature biotechnology* 2018;36(5):411-20.
5. Satija R, Farrell JA, Gennert D, et al. Spatial reconstruction of single-cell gene expression data. *Nature Biotechnology* 2015;33(5):495-502. doi: 10.1038/nbt.3192
6. Korsunsky I, Millard N, Fan J, et al. Fast, sensitive and accurate integration of single-cell data with Harmony. *Nature methods* 2019;16(12):1289-96.
7. Aran D, Looney AP, Liu L, et al. Reference-based analysis of lung single-cell sequencing reveals a transitional profibrotic macrophage. *Nature immunology* 2019;20(2):163-72.
8. Kramer O, Kramer O. Scikit-learn. *Machine learning for evolution strategies* 2016:45-53.
9. Conway JR, Lex A, Gehlenborg N. UpSetR: an R package for the visualization of intersecting sets and their properties. *Bioinformatics (Oxford, England)* 2017;33(18):2938-40.
10. Zhang B, Horvath S. A general framework for weighted gene co-expression network analysis. *Statistical applications in genetics and molecular biology* 2005;4(1)
11. Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis. *BMC bioinformatics* 2008;9:1-13.
12. Yu G, Wang L-G, Han Y, et al. clusterProfiler: an R package for comparing biological themes among gene clusters. *Omics: a journal of integrative biology* 2012;16(5):284-87.
13. Dolgalev I. msigdb: MSigDB gene sets for multiple organisms in a tidy data format. *R package version* 2020;7(1)
14. Leung S, Liu X, Fang L, et al. The cytokine milieu in the interplay of pathogenic Th1/Th17 cells and regulatory T cells in autoimmune disease. *Cellular & molecular immunology* 2010;7(3):182-89.
15. Zhou X, Michal JJ, Zhang L, et al. Interferon induced IFIT family genes in host antiviral defense. *International journal of biological sciences* 2013;9(2):200.
16. Ramiscal RR, Vinuesa CG. T-cell subsets in the germinal center. *Immunological reviews* 2013;252(1):146-55.
17. Bosnjak B, Stelzmueller B, Erb KJ, et al. Treatment of allergic asthma: modulation of Th2 cells and their responses. *Respiratory research* 2011;12:1-17.
18. Moretta A, Marcenaro E, Parolini S, et al. NK cells at the interface between innate and adaptive immunity. *Cell Death & Differentiation* 2008;15(2):226-33.
19. Pugazhenth S, Zhang Y, Bouchard R, et al. Induction of an inflammatory loop by interleukin-1 β and tumor necrosis factor- α involves NF- κ B and STAT-1 in differentiated human neuroprogenitor cells. *PloS one* 2013;8(7):e69585.
20. Oshi M, Le L, Tokumaru Y, et al. Cell proliferation-related pathway scores (G2M Checkpoint, E2F Targets, MYC Targets V1 and V2, Mitotic Spindle, p53 pathway) as predictive and prognostic biomarkers in pancreatic cancer. *Cancer Research* 2021;81(13_Supplement):1966-66.
21. Carroll PA, Freie BW, Mathsyaraja H, et al. The MYC transcription factor network: balancing metabolism, proliferation and oncogenesis. *Frontiers of medicine* 2018;12:412-25.
22. Chapman NM, Boothby MR, Chi H. Metabolic coordination of T cell quiescence and activation. *Nature reviews immunology* 2020;20(1):55-70.

23. Trzebanski S, Jung S. Plasticity of monocyte development and monocyte fates. *Immunology Letters* 2020;227:66-78.
24. Lin F-C, Young HA. The talented interferon-gamma. *Advances in Bioscience and Biotechnology* 2013;4(07):6-13.
25. Ogle ME, Segar CE, Sridhar S, et al. Monocytes and macrophages in tissue repair: Implications for immunoregenerative biomaterial design. *Experimental Biology and Medicine* 2016;241(10):1084-97.
26. Eggenhuizen PJ, Ng BH, Ooi JD. Treg enhancing therapies to treat autoimmune diseases. *International journal of molecular sciences* 2020;21(19):7015.