

Lineage B Genotype III of Dengue Virus Serotype 3 (DENV-3III_B) is responsible for Dengue Outbreak in Dire

Dawa City, Ethiopia, 2023

Abebe Asefa Negeri^{1*}, Dawit Hailu Alemayehu², Saro Abdella Abraham¹, Tsigereda Kifle Wolde¹, Gutema Bulti¹, Alemnesh Hailemariam Bedasso¹, Danile Tsega¹, Ebise Abosex¹, Eyilachew Zenebe Awule³, **Diana Rojas-Gallardo⁴**, Asefa Konde Korkiso¹, Kalkidan Melaku², Raffael Joseph², Abaysew Ayele², Mesfin Mengesha Tsegaye², Anne Piantadosi⁵, Getachew Tollera¹, Alemseged Abdissa², Mesay Hailu Dangisso¹, Adane Mihret², Andargachew Mulu², Tesfaye Gelanew^{2*}

¹Ethiopian Public Health Institute, Addis Ababa Ethiopia

²Armauer Hansen Research Institute, Addis Ababa, Ethiopia

³Dire Dawa City Administration Health Bureau

⁴**Population Biology, Ecology and Evolution Graduate Program, Emory University, Atlanta, GA, USA**

⁵Emory University School of Medicine, Atlanta, Georgia, USA

* Correspondence: tesfayegtaye@gmail.com; tesfaye.gelanew@ahri.gov.et, Tel.: +251980153960;

abebea84@gmail.com (Abebe Assefa Negeri)

Abstract

Background: The eastern parts of Ethiopia, including Dire Dawa City, have faced annual dengue fever (DF) outbreaks since 2013, resulting in substantial healthcare and economic consequences. However, there remains a lack of comprehensive evidence regarding the specific dengue virus (DENV) serotypes and genotypes associated with those outbreaks.

Methodology: On December 14, 2023, the National Arbovirus Laboratory at the Ethiopian Public Health Institute received seventy serum samples from 70 patients suspected of DF during the outbreak in Dire Dawa City. Samples positive for DENV with adequate volume and CT values underwent sequencing of the CprM region of the DENV genome. The obtained sequences were typed using the Dengue Virus Typing Tool and underwent phylogenetic analysis to evaluate sequence diversity and relationships to DENV strains from other global DENV-endemic regions.

Results: Overall, 32 (45.7%) of the patients displayed one or more early warning signs indicative of severe dengue. Among the 13 patients who were hospitalized for 2 to 10 days, all except two had at least one symptom indicative of severe dengue. Out of 67 samples with adequate volume, 44 (65.6%) were positive for DENV RNA using RT-PCR, and 21 successfully underwent CprM sequencing. All of them belonged to DENV-3, genotype III, major lineage B (DENV-3III_B), but represented two different minor lineages (DENV-3III_B.2 and DENV-3III_B.3). Phylogenetic analysis demonstrated that both lineages were closely related to sequences from the Afar region of Ethiopia collected in 2023, indicating that the outbreaks were related and were due to multiple co-circulating lineages.

Conclusions: DENV-3III_B was identified as the cause of the 2023 DF outbreak in Dire Dawa City. The identification of two co-circulating lineages in this outbreak underscores the complex nature of DENV transmission in the African continent and highlights the need for increased viral genomic surveillance.

Keywords: DENV, DF, Serotype, Genotype, CprM, DENV-3/GIII, Dire Dawa City, Ethiopia

1. Introduction

Dengue fever (DF), caused by four genetically distinct serotypes of dengue virus (DENV-1 to DENV-4), has become a significant global health challenge. Over the past two decades, the incidence of dengue has increased tenfold, rising from 500,000 reported cases worldwide in 2000 to a staggering 5.2 million cases in 2019. In 2019, dengue outbreaks spread across 129 countries, marking an unprecedented peak. Although there was a slight decline in cases between 2020 and 2022 due to the COVID-19 pandemic and reduced reporting rates, 2023 witnessed a resurgence in dengue cases globally[1],[2],[3].

In areas where DF cases have been identified as endemic and/or epidemic, especially in tropical and subtropical countries, nonspecific febrile illnesses are common. Febrile illnesses can be caused by a variety of infectious agents such as malaria parasites, alphaviruses, and flaviviruses, complicating surveillance and response programs for outbreaks and endemic diseases[4]. In addition, the *Aedes* mosquito species, particularly *Ae. Aegypti* and *Ae. Albopictus* that transmit DENV, are also responsible for the transmission of other arboviruses (e.g., Zika virus, Yellow Fever virus, and Chikungunya virus), and are common in tropical and subtropical countries[5].

In Ethiopia, the first laboratory-confirmed outbreak of DF occurred in Deri Dawa City in 2013. Since then, Ethiopia has experienced nearly annual outbreaks of DF in multiple regions of the country (Figure 1). In the Afar Region, DF outbreaks occurred in 2019 in Gewane District[6]. In the Ethiopian Somalia region, the first outbreak was reported in Godey town between 2014 and 2015 followed by outbreaks that occurred in **Kabridahar in 2017[7]** and **Warder Woreda in 2021[8]**. These recurrent outbreaks have impacted an already fragmented health system[9] and the economy of these parts of the country.

In April 2023, an outbreak of DF was initially reported in the Afar region, the northeastern part of Ethiopia. The first affected districts were Logia and Mille. Since April 4, 2023, however, it spread to all seven districts and towns in the Afar region **(Figure 1)**. As of June 26, 2023, a total of 6,133 suspected and confirmed cases were reported, resulting in nine associated deaths (with a case: fatality ratio of 0.5%)[10]. It was

documented that 8 of the 10 epidemic samples examined by RT-PCT triplex [for Dengue, Zika, and Chikungunya] belonged to serotype DENV-3 [10].

As a suspected expansion of the outbreak in the Afar region, a DF outbreak was recorded in Dire Dawa City in December 2023 (Figure 1). However, there is no data regarding the DENV serotypes and genotypes associated with the outbreak. Whether the outbreak was caused by DENV-3 as in the Afar region remains unknown. In the present study, we present data on the serotypes, genotypes, and lineages of DENV strains responsible for the outbreak in Dire Dawa City during 2023 and discuss the epidemiological and clinical consequences.

2. Materials and Methods

2.1. Ethical considerations

2.1. Ethical considerations

As our study aimed to better understand DENV strains linked to DF outbreaks, we obtained an ethical approval waiver (protocol number: PO-036-24) from the All-African Leprosy Rehabilitation and Training Center/Armauer Hansen Research (ALERT/AHRI) Ethics Committee. Additionally, we obtained permission from the Dire Dawa City Health Bureau and health facilities' authorities to investigate these outbreak samples. Since the samples used in this study were obtained from the dengue outbreak during the 2023 outbreak investigation and confirmation undertaken to respond to and control the public health emergencies, no informed consent was obtained during the sample collection.

2.2. Study Settings and Patient Data

On December 14, 2023, the Ethiopia Public Health Institute (EPHI) received 70 serum samples from an outbreak of DF in Dire Dawa City Administration (Figure 1). These samples were collected from DF suspected cases and shipped to the National Arbovirus Laboratory of EPHI for laboratory confirmation of the causative agent as part of an outbreak response investigation (Figure 2). The reported clinical signs and symptoms for

each suspected case were extracted from medical charts available at the health facilities they were admitted 104
using a case-based surveillance request form. Mild symptoms include sudden onset of high fever, vomiting, 105
nausea, rash, arthralgia (joint pain), headache, chills, myalgia (muscle pain), sore throat, dizziness, runny nose, 106
pruritus (itchiness), adenosis, diarrhea and shivering while warning signs or indicative of severe dengue 107
include: belly pain or tenderness, persistent vomiting (at least 3 times/24 hrs), bleeding from the nose or gum, 108
restless or irritable, feeling weak, conjunctiva congestion and convulsions (Supplementary Table 1 and Table 109
2). The category of mild symptoms and warning signs indicative of severe dengue is based on the WHO severe 110
dengue classification[11]. 111




Figure 1. Map showing the Dire Dawa City Administration where the Dengue outbreak occurred, 114
and study DENV strains originated. The shaded areas indicate the epicenters of dengue outbreaks 115
in Ethiopia since 2013: Dire Dawa (red), Afar (red-violet), and Somalia (light blue) 116

2.3. Blood sample collection and processing 117

Each of the 70 dengue-suspected patients submitted blood samples for laboratory diagnosis of the 118
causative agent for febrile illness (Supplementary Table 1). The blood samples were allowed to clot, and the 119
sera were separated. The sera were then shipped in a triple package cold chain to the National Arbovirus 120
Laboratory of the EPHI. At the laboratory, the sera were kept frozen at -20°C until they were used for DENV 121
viremia detection using reverse transcriptase (RT)- PCR (RT-PCR). 122

2.4 Viral RNA Extraction and PCR Detection of DENV 123

Total RNA was isolated from each serum sample using the QIAmp viral RNA kit (QIAGEN, INC, 124
Valencia, CA). The RNA was eluted in 50 μl of AVE buffer, which was provided with the kit. DENV Viremia 125
Detection was done using RealStar® Dengue RT-PCR kit (Altona Diagnostics, Hamburg-Altona, Germany). 126
Briefly, 20 μl of the master mix (containing primers and probes) was mixed with 10 μl of the RNA sample or 127
control. The mixture was centrifuged for 30 seconds at approximately 1000 x g (~3000 rpm). The reactions 128
were incubated for 20 minutes at 55°C , followed by 40 cycles of 95°C for 1 min, 55°C for 45 seconds, 72°C 129
for 15 seconds, and final incubation at 68°C for 10 min. The PCR results were interpreted based on the 130
manufacturer's protocol. A threshold cycle (Ct) value of ≤ 36 for DENV-specific RNA and a Ct value of ≤ 30 131
for the Internal Control were considered positive samples for DENV. 132

2.5 Semi-Nested RT-PCR Amplification (Amplicon size 511 bp) 133

RNA samples with a Ct-value of 31 or lower and sufficient volume were selected for CprM RT-PCR 134
amplification following the procedure described elsewhere [12]. Briefly, the DENV CprM region (652) bp 135
was first amplified using forward primer (FP): 5'-TCAATATGCTGAAACGCGCGAGAAACCG-3' (132– 136
153) and the reverse primer (RP): 5'- GCGCCTTCNGNNGACATCCA -3' (764–783) and the high-capacity 137
cDNA reverse transcription kit (Invitrogen). The reactions were incubated first at 98°C for 10 seconds 138

followed by 35 cycles of 98°C for 10 seconds, 60°C for 20 seconds, 72°C for 30 seconds, and final incubation 139
at 72 °C for 5 minutes. Semi-nested PCR was also used to amplify 511 bp of the CprM region of the viral 140
genome using amplified product as a template, and the same FP: 5'- 141
TCAATATGCTGAAACGCGCGAGAAACCG-3' (132–153) but a different RP: 5'- 142
TTGCACCAACAGTCAATGTCTTCAGGTTC-3' (614-642)[12]. The semi-nested amplification was 143
performed in an Applied Biosystems Veriti 96-Well Thermal Cycler (ThermoFisher Scientific) using the 144
following cycling parameters: 94°C for 2 min; 40 cycles of 94°C for 30 s, 70°C for 1 s (ramp rate 20%), 55°C 145
for 45 s (ramp rate 20%), and 65°C for 3 min, 20 s; and final extension for 10 min at 65°C [13]. Following 146
the amplification, 5 µl of the first amplicon and semi-nested amplicon size of 652 bp and 511 bp, respectively 147
were visualized on ethidium bromide-stained agarose gel under UV light. Only the 511 bp DENV amplified 148
PCR products were purified using a QIAquick PCR purification Kit (QIAGEN, Germany) and were sequenced 149
in both strands using a Big Dye Terminator Cycle Sequencing kit (Applied Biosystems), and the same primers 150
used for nested-PCR. The CprM sequences were confirmed by BLAST. The forward and reverse sequences 151
were aligned and manually edited using Bio Edit software to obtain the consensus sequence. The obtained 152
sequences were analyzed using **Dengue Virus Typing Tool at Genome Detective server** [13] to identify the 153
serotypes, genotypes, and lineages of DENV associated with the outbreak. 154

2.6 Phylogenetic analysis 155

The CprM consensus sequences for 21 DENV strains from Dire Dawa City were deposited in GenBank 156
under accession numbers PP751832-PP751852. Nucleotide BLAST searches were performed for each of 157
the 21 new sequences, and all DENV-3III CprM gene and whole genome sequences with >98% similarity 158
were downloaded. Duplicates and identical sequences from the same country and collection date were 159
removed. Representative sequences for DENV-3 genotypes I, II, and V were also included as references. 160
The final data set include a total of 123 DENV-3 CprM gene sequences collected between 2006-2023 across 161
the globe. 162

For phylogenetic reconstruction, sequences were aligned using MAFFT (version 7.520)[14], and a maximum likelihood tree was constructed with the IQ-TREE software (version 2.2.2.7)[15] using the TIM2e+G4 substitution model (according to AICc) with a bootstrap value of 10,000. Finally, the resulting tree was visualized and annotated using iTOL (Interactive Tree Of Life)[16].

3. Results

3.1. Patient characteristics

Among the 70 suspected dengue fever (DF) patients, 38 (54%) were female, as indicated in Table 1 and Table S1. The median age was 30 with an interquartile (IQ) range of 17 years. Most (81%) of the suspected cases visited health facilities as outpatients (Table 1 and Supplementary Table 1), while the remaining 13 cases required inpatient hospitalization for a duration of 2 to 10 days (Supplementary Table 1).

Table 1. Status of admission and demographic characteristics of dengue fever suspected cases (n = 70) reported at health facilities in the Dire Dawa City Administration, 2023.

Variable	Specific variable	N (%)
Age	Median age 30; IQR 17	-
Sex	Male	32(46)
	Female	38(54)
Patient admission status	Outpatient	57(80)
	Inpatient	13 (20)

The clinical signs/symptoms observed in these (n = 70) DF suspected cases, whether admitted as outpatients or inpatients, covered a wide range of symptoms consistent with febrile illness. Nearly all of them exhibited three or more clinical symptoms consistent with mild DF (Supplementary Table 1). However, regardless of hospitalization status, 32 (45.7%) of these patients displayed one or more early warning signs

indicative of severe dengue[11]. Among the 13 patients who were hospitalized for 2 to 10 days, all except two had at least one symptom indicative of severe dengue (Supplementary Table 1). The most commonly observed clinical presentations were acute fever (n = 66; 98.15%), headache (n = 65; 97.01%) and chills (n = 47; 70.17%) followed by aches and pains (n = 31; 46.27%) and vomiting/nausea (n = 31; 46.27%) (Table 2). In addition, all of the cases with early warning signs of severe dengue were later confirmed to harbor DENV viremia by RT-PCR. Supplementary Table 1 summarizes the details of the demographic and clinical characteristics of study participants.

Table 2. Clinical characteristics of dengue fever suspected cases (n = 70) reported at health facilities in the Dire Dawa City Administration, 2023.

Clinical symptoms ¹	Yes ²	No
A. Mild symptoms		
Fever: sudden onset of high fever	66(98.15%)	1(1.49%)
Chills	47(70.17%)	20(29.85%)
Vomiting or Nausea	31(46.27%)	36(53.73%)
Aches and Pains: Typically, behind the eyes, muscle, joint, or bone pain	31(46.27%)	36(53.73%)
Myalgia	28(45.9%)	33(54.1%)
Running nose	21(31.34%)	46(68.66%)
Diarrhoea	13(19.4%)	54(89.60%)
Rash	8(13.11%)	53(86.89%)
Sore throat	6(8.89%)	61(91.04%)
Dizzy	5(7.46%)	62(92.54%)
Shivering	4(5.97%)	63(94.03%)
Pruritus	2(2.99%)	65(97.01%)

Adenoids	1(1.52%)	65(98.48%)
----------	----------	------------

B. Warning signs indicative of severe dengue

Weak (Feeling tired)	26(38.81%)	41(61.19%)
Restless, or Irritable	15(22.39%)	52(77.61%)
Arthralgia	9(13.43%)	58(86.57%)
Persistent Vomiting (at least 3 times/24 Hrs)	9(13.43%)	58(86.57%)
Bleeding from the nose or gums	6(8.96%)	61(91.04%)
Belly Pain or Tenderness.	5(7.46%)	62(92.54%)
Conjunctival congestion	3(4.48%)	64(95.52%)

¹ Categorization of mild symptoms and warning signs indicative of severe dengue are based on WHO classification. ² Clinical data for 4 cases out of the 70 suspected dengue fever cases were missed or incomplete.

3.2. DENV Viremia Detection

Among 70 specimens, 3 samples failed to pass the DENV nucleic acid amplification quality control.

Among the 67 samples that passed quality control, 65.67% (44/67) had DENV RNA detected(Figure 2).




Figure 2. Flow chart showing enrollment of suspected Dengue and/ or other febrile illnesses, including malaria patients and molecular detection and characterization of strains isolated during the 2023 dengue outbreak in Dire Dawa, Ethiopia.

Supplementary Table 2 summarizes laboratory investigation of dengue-confirmed patients, including DENV serotypes, genotypes, lineage, and sublineages with their corresponding GenBank Accession Numbers.

3.3. DENV sequence classifications

Based on rRT-PCR $C_T < 31$, 27 samples out of 44 DENV viremia-positive samples were selected for CprM sequencing, and 21 were successful (Figure 2) allowing classification of their serotypes, genotypes, and lineages (Supplementary Table 2). The sequence quality for 6 samples was not sufficient, and the DENV serotypes could not be determined (Figure 2). According to the new proposed DENV nomenclature [13], the Dengue Virus Typing Tool assigned all 21 sequences to DENV-3, genotype III, major lineage B (DENV-3III_B), and the minor lineages were assigned as follows: 17 sequences belonged to or were most closely related to minor lineage 2 (DENV-3III_B.2), and 4 sequences belonged to or were most closely related to minor lineage 3 (DENV-3III_B.3) (Figure 3).

3.4 Phylogenetic analysis

We investigated phylogenetic relationships between the 21 new DENV-3III_B CprM sequences obtained in this study and other publicly available DENV-3III_B CprM sequences from various regions including Africa (Kenya, Gabon, Burkina Faso, Mozambique, Tanzania, and Ethiopia), Asia (Middle East, India, Sri Lanka, Thailand, and China), the Americas (Jamaica, Martinique, Dominican Republic and USA), and Europe (Italy).

As expected, the 17 DENV-3III_B minor lineage 2 (DENV-3III_B.2) sequences clustered separately from the 4 DENV-3III_B minor lineage 3 (DENV-3III_B.3) or related sequences (Figure 3). The DENV-3III_B.2 sequences from our study clustered with other sequences from 2023 collected across different continents, including one from an outbreak in Lazio, Italy, and one from China attributed to a case imported from Burkina Faso. On the other hand, the DENV-3III_B.3 sequences clustered with two sequences from

4. Discussion

Knowledge of the genetic diversity of DENV strains responsible for outbreaks is essential for efficient management, vaccine development, and outbreak prevention. Though Ethiopia's eastern region, including Dire Dawa City, has seen many DF outbreaks since 2013, there is scanty information regarding the diversity of circulating DENV strains. In this study, we used genetic analysis of the DENV CprM region to demonstrate that the 2023 outbreak in Dire Dawa City was due to two co-circulating lineages of DENV-3III_B, each of which was closely related to sequences from a related outbreak in the Afar region of Ethiopia in 2023. Our results are consistent with a recent study that analyzed 7 full DENV-3III genome sequences from the Afar outbreak and found evidence for multiple introductions. However, the paucity of regional DENV sequence data makes it difficult to determine the timing and origin of these introductions. Recently, other DENV-3III outbreaks have been reported in East Africa, notably in South Sudan in 2022[18] and in Kenya in 2019[19]. We were not able to assess the genetic relatedness between the current strains and outbreak strains from these neighboring countries due to the fact they employed other targets for sequencing such as the E gene. These results underscore the need for further genomic surveillance of DENV in East Africa, ideally using full-length genome sequences.

In contrast to the prior dengue outbreaks in Ethiopia which were more mild[7],[20],[21],[22], in this study 46% (n = 32) of the 70 DF-suspected cases displayed early warning signs indicative of severe dengue, while the remaining cases showed mild symptoms. All of these cases with early warning signs of severe dengue were confirmed to harbor DENV viremia by RT-PCR.

We propose three scenarios that might have led to the occurrence of severe cases of DF during the 2023 outbreak in Dire Dawa City. One possibility is the observed severe dengue cases could be secondary DENV-3III infections that happened after primary infection with a different serotype. In support of this, prior DF outbreaks reported from Dire Dawa City were associated with DENV-1 and DEV-2 strains [7],[20],[21],[22],(Mesfin et al, in Press Viruses, MDPI), although previous attempts to determine serotypes

were restricted to a small number of DENV strains. Thus, the recurrent occurrence of dengue outbreaks associated with different DENV serotypes calls for the need for close monitoring and management of future DF outbreaks to effectively control and manage secondary infections.

The second possibility is that DENV-3III is more pathogenic, as suggested by prior research that connected outbreaks of DENV-3III to Dengue Hemorrhagic Fever (DHF) and Dengue Shock Syndrome (DSS) [23].

The third possibility is that the severe cases may be the result of co-infection with malaria parasites, as both infections are co-endemic in Dire Dawa City, and there is a chance that co-infection can worsen disease severity and outcomes. This is supported by the recent surge in malaria[24],[25] and dengue cases[26],[27] in Dire Dawa City which posed significant challenges for healthcare providers. Given the co-endemicity and overlapping of symptoms (e.g., fever), there is a risk of co-infection with both dengue and malaria and misdiagnosing dengue as malaria (and vice versa). Thus, an integrated approach involving surveillance, education, and differential diagnosis is crucial for better patient care in Dire Dawa City Administration. Healthcare professionals need to be aware of the clinical features, diagnostic tests, and management protocols for malaria and dengue. In addition, the Dire Dawa City Administration should avail appropriate laboratory tests (such as rapid diagnostic tests and PCR) at all health facilities that facilitate accurate diagnosis of dengue and malaria and prompt treatment and management of severe cases.

Among the five distinct genotypes (I–V) of DENV-3, all 21 sequenced DENV strains from the present study belonged to DENV-3III, which is the most widespread and was associated with large outbreaks in Asia, Africa, and the Americas[28],[29],[30]. The first autochthonous case of DENV-3III in Africa was reported in Mozambique in 1985[31]. Several man-made and natural factors, including (i) migration and conflict-related population displacement, (ii) frequent migration via trade routes that can spread the virus to new areas, and (iii) recurrent flooding that is made worse by climate change that provides ideal breeding grounds for Aedes mosquitoes may have contributed to the rapid geographic expansion of DENV-3III in Africa, resulting in outbreaks in areas like Dire Dawa City[3],[28],[32],[23],[33].

Even though this work offers important insights into DENV strains responsible for the 2023 DF outbreak in Dire Dawa City where severe cases and deaths were recorded, it suffers limitations. First, the study's capacity to detect genetic variation was limited due to the use of a short genomic target (CprM, 511 base pairs) for the characterization of the outbreak strains. Expanding the genomic coverage could have enhanced our understanding of DENV diversity and relationships to other DENV-3III strains across the globe. Second, DENV CprM sequencing was performed on a small proportion of DF-suspected cases reported during the outbreak, and a larger sample size may have identified greater genetic diversity. Third, the study lacks information on other febrile illness-causing pathogens like malaria parasite examination with microscopy or rapid diagnostic test results, for the studied DF-suspected cases, so cases of co-infection may have been missed. Future research addressing these limitations will therefore improve our comprehension of disease dynamics, provide a thorough understanding of DENV strains and their evolutionary history, expand our understanding of DENV outbreaks, and provide guidance for more effective preventative measures.

5. Conclusions

In summary, our results demonstrate that the 2023 DF outbreak in Dire Dawa City, Ethiopia, was due to multiple co-circulating lineages of DENV-3III. The severe cases recorded during this outbreak may be due to either the genetic makeup of DENV-3III, secondary DENV infection, or co-infection of DENV and malaria. Additionally, the intercontinental transmission of DENV-3III underscores the importance of vigilant surveillance and preparedness to mitigate the impact of such outbreaks. Thus, collaborative efforts across borders are crucial for effective prevention and control strategies.

Supplementary Materials: Supplementary Table 1: Demographic and clinical parameters of dengue suspected patients and Supplementary Table 2: Laboratory investigation of dengue confirmed patients, including DENV serotype and genotype with their corresponding GenBank Accession Numbers.

Author Contributions: Conceptualization, A.A.N. and S.A.A.; study design, S.A.A., A.Mu. And T.G.; methodology, A. Mu., S.A.A., T.G., H.A.; software, T.G.; validation, T.G., DM.RG., A.A.N., and Z.Z.; formal analysis, T.G., DM.RG., and A.Ay.; Laboratory investigations, D.H.A., T.K.W., G.B., A.H.B, D.T., E.A,

EZA, A.As, A.K.K., K.M., and R.J.; resources, A.Mu, A.Mi., S.A.A., A.Ab., G.T., M.H; Field data and 309
samples acquisition, E.Z.A., D.T., and A.K.K. data curation, T.G., and A.A.N.; writing—original draft 310
preparation, T.G., and A.A.N.; writing—review and editing, T.G., A.P., A.Mu., and A.A.N.; visualization, 311
T.G., **DM.RG**, A.Ay., and A.P.; supervision, S.A.A. All authors have read and agreed to the published version 312
of the manuscript. 313

Funding: This research received no external funding. 314

Acknowledgments 315

We would like to thank all study participants in this study. We are also grateful for the support and cooperation 316
we received from physicians, nurses of Dire Dawa City health facilities, and administrative authorities of Dire 317
Dawa City Administration Health Hureau and the health facilities administration. The authors acknowledge 318
EPHI's support of the field study and AHRI and EPHI's reagent support of the molecular analysis. 319

Conflicts of Interest 321

The authors declare no conflicts of interest. 322

Data Availability Statement 323

The datasets generated and/or analyzed during the current study are available in the manuscript and the 324
Supplementary Materials. **The sequence data presented in this study are openly available in the GenBank** 325
database, accession numbers PP751832-PP751852. 326

References 328

1. Dengue- Global situation. <https://www.who.int/emergencies/disease-outbreak-news/item/2023-DON498>. 329
Accessed 7 Apr 2024. 330
2. Yang X, Quam MBM, Zhang T, Sang S. Global burden for dengue and the evolving pattern in the past 331
30 years. *Journal of Travel Medicine*. 2021;28:taab146. 332

3. Geographical expansion of cases of dengue and chikungunya beyond the historical areas of transmission in the Region of the Americas. <https://www.who.int/emergencies/disease-outbreak-news/item/2023-DON448>. Accessed 7 Apr 2024. 333
334
335
4. Roy SK, Bhattacharjee S. Dengue virus: epidemiology, biology, and disease aetiology. *Can J Microbiol*. 2021;67:687–702. 336
337
5. Dengue and severe dengue. <https://www.who.int/news-room/fact-sheets/detail/dengue-and-severe-dengue>. Accessed 25 Feb 2024. 338
339
6. Mekuriaw W, Kinde S, Kindu B, Muluaem Y, Hailu G, Gebresilassie A, et al. Epidemiological, Entomological, and Climatological Investigation of the 2019 Dengue Fever Outbreak in Gewane District, Afar Region, North-East Ethiopia. *Insects*. 2022;13:1066. 340
341
342
7. Gutu MA, Bekele A, Seid Y, Mohammed Y, Gemechu F, Woyessa AB, et al. Another dengue fever outbreak in Eastern Ethiopia—An emerging public health threat. *PLOS Neglected Tropical Diseases*. 2021;15:e0008992. 343
344
345
8. Mesfin Z, Ali A, Abagero A, Asefa Z. Dengue Fever Outbreak Investigation in Werder Town, Dollo Zone, Somali Region, Ethiopia. *Infection and Drug Resistance*. 2022;15:7207–17. 346
347
9. Mohan A, Fakhor H, Nimavat N, Wara UU, Lal PM, Costa AC dos S, et al. Dengue and COVID-19: A risk of coepidemic in Ethiopia. *J Med Virol*. 2021;93:5680–1. 348
349
10. Sisay C, Tadesse A, Workie F, Yizengaw A, Ali A, Yusuf J, et al. Description of Serotype 3 Dengue Fever Virus: Clinical, Surveillance and Geographical Expansion in the Northeastern Ethiopia. *Science Journal of Public Health*. 2023. <https://doi.org/10.11648/j.sjph.20231104.12>. 350
351
352
11. Htun TP, Xiong Z, Pang J. Clinical signs and symptoms associated with WHO severe dengue classification: a systematic review and meta-analysis. *Emerg Microbes Infect*. 10:1116–28. 353
354
12. Lanciotti RS, Calisher CH, Gubler DJ, Chang GJ, Vorndam AV. Rapid detection and typing of dengue viruses from clinical samples by using reverse transcriptase-polymerase chain reaction. *Journal of Clinical Microbiology*. 1992;30:545–51. 355
356
357
13. Hill V, Cleemput S, Fonseca V, Tegally H, Brito AF, Gifford R, et al. A new lineage nomenclature to aid genomic surveillance of dengue virus. 2024;:2024.05.16.24307504. 358
359
14. Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Mol Biol Evol*. 2013;30:772–80. 360
361
15. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular Biology and Evolution*. 2015;32:268–74. 362
363
364

16. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. *Nucleic Acids Research*. 2021;49:W293–6. 365
366
17. Mekonnen F, Khan BA, Nibret E, Munshea A, Tsega D, Endalamaw D, et al. Introduction of Dengue Virus Serotype 3 in the Afar Region, Ethiopia. 2024;:2024.07.19.24310689. 367
368
18. Eldigail MH, Abubaker HA, Khalid FA, Abdallah TM, Musa HH, Ahmed ME, et al. Association of genotype III of dengue virus serotype 3 with disease outbreak in Eastern Sudan, 2019. *Virology Journal*. 2020;17:118. 369
370
371
19. Muthanje EM, Kimita G, Nyataya J, Njue W, Mulili C, Mugweru J, et al. March 2019 dengue fever outbreak at the Kenyan south coast involving dengue virus serotype 3, genotypes III and V. *PLOS Glob Public Health*. 2022;2:e0000122. 372
373
374
20. Biru M, Geleta D. Dengue Fever Outbreak Investigation and Response in Dire Dawa City Administration, Ethiopia, 2017. *Journal of Medicine*. 2020. 375
376
21. Zerfu B, Kassa T, Legesse M. Epidemiology, biology, pathogenesis, clinical manifestations, and diagnosis of dengue virus infection, and its trend in Ethiopia: a comprehensive literature review. *Tropical Medicine and Health*. 2023;51:11. 377
378
379
22. Sisay C, Waldetensai A, Seyoum M, Tayachew A, Wossen M, Keneni D, et al. Detection of serotype 1- Dengue fever outbreak in Dire Dawa city, Eastern Ethiopia. *Ethiopian Journal of Public Health and Nutrition (EJPHN)*. 2022;5:49–54. 380
381
382
23. Messer WB, Gubler DJ, Harris E, Sivananthan K, Silva AM de. Emergence and Global Spread of a Dengue Serotype 3, Subtype III Virus - Volume 9, Number 7—July 2003 - *Emerging Infectious Diseases journal* - CDC. <https://doi.org/10.3201/eid0907.030038>. 383
384
385
24. A malaria outbreak in Ethiopia came from an invasive Asian mosquito. 2022. 386
<https://www.sciencenews.org/article/malaria-outbreak-ethiopia-invasive-mosquito-asia>. Accessed 11 Mar 2024. 387
388
25. Emiru T, Getachew D, Murphy M, Sedda L, Ejigu LA, Bulto MG, et al. Evidence for a role of *Anopheles stephensi* in the spread of drug- and diagnosis-resistant malaria in Africa. *Nat Med*. 2023;29:3203–11. 389
390
26. Degife LH, Worku Y, Belay D, Bekele A, Hailemariam Z. Factors associated with dengue fever outbreak in Dire Dawa administration city, October, 2015, Ethiopia - case control study. *BMC Public Health*. 2019;19:650. 391
392
393
27. Nigussie E, Atlaw D, Negash G, Gezahegn H, Baressa G, Tasew A, et al. A dengue virus infection in Ethiopia: a systematic review and meta-analysis. *BMC Infectious Diseases*. 2024;24:297. 394
395
28. Maquart M. Circulation of Dengue Virus Type 3 Genotype III in Africa Since 2008. *JHVRV*. 2016;4. 396

29. Waman VP, Kale MM, Kulkarni-Kale U. Genetic diversity and evolution of dengue virus serotype 3: A comparative genomics study. *Infection, Genetics and Evolution*. 2017;49:234–40. 397
398
30. Titir SR, Paul SK, Ahmed S, Haque N, Nasreen SA, Hossain KS, et al. Nationwide Distribution of Dengue Virus Type 3 (DENV-3) Genotype I and Emergence of DENV-3 Genotype III during the 2019 Outbreak in Bangladesh. *Trop Med Infect Dis*. 2021;6:58. 399
400
401
31. Gubler DJ, Sather GE, Kuno G, Cabral JR. Dengue 3 Virus Transmission in Africa. *The American Journal of Tropical Medicine and Hygiene*. 1986;35:1280–4. 402
403
32. Amarasinghe A, Kuritsky JN, Letson GW, Margolis HS. Dengue Virus Infection in Africa. *Emerg Infect Dis*. 2011;17:1349–54. 404
405
33. Eltom K, Enan K, El Hussein ARM, Elkhidir IM. Dengue Virus Infection in Sub-Saharan Africa Between 2010 and 2020: A Systematic Review and Meta-Analysis. *Front Cell Infect Microbiol*. 2021;11:678945. 406
407
408
- Disclaimer/Publisher’s Note:** The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the journal and/or the editor(s) and should not be construed as official, or as reflecting true views of the Armauer Hansen Research Institute and Ethiopian Public Health Institute. 409
410
411
412




Figure 1




Figure 2

Tree scale: 0.1

Lineage

- B.2
- B Related to but not part of 2
- B.3
- B Related to but not part of 3
- B Could not assign

Continent

- Africa
- Europe
- Asia
- America




Figure 3