1

Lung functions among children and adolescents with sickle cell disease receiving care at Jaramogi Oginga Odinga Teaching and Referral Hospital Kisumu, Kenya

5	Lung functions among children and adolescents with sickle cell disease
6	
/	
8	Fredrick O. Olewe ^{1,2*}
9	Constance N. Tenge ¹ ¶
10	Irene K. Marete ¹ ¶
11	
12	
13	1 Department of Child Health and Pediatrics School of Medicine, Moi University
14	2 Center for Research in Therapeutic Science (CREATES)- Strathmore University
15	
16	
17	* <u>fred84doc@gmail.com</u> (FO)
18	
19	These authors contributed equally
20	

21 Abstract

Sickle cell disease is a genetic disorder associated with lifelong symptoms of anemia, vaso-oclusive crisis and organ damage. Pulmonary complications and declining lung functions are contributors to morbidity and mortality. Determination of lung functions can enable early diagnosis and institution of appropriate intervention.

26 The objectives were; To determine the prevalence, patterns, and the factors associated with abnormal lung

27 functions in children and adolescents aged 6-17 years with confirmed diagnosis Sickle Cell Disease.

28 This was a cross sectional study done at Jaramogi Oginga Odinga Teaching and Referral Hospital in

29 Kisumu, Kenya. A total of 138 participants were recruited. A structured data collection tool was used to

30 collect Socio-demographics and clinical characteristics. The spirometry was done using NDD Easy-On PC

31 spirometer and results entered into database, cleaned and analyzed for proportions, frequency, mean, range

32 and odds ratio at 95% confidence level.

33 Females were 65(47%), males 73(53%), and the mean age was 10.99 years with SD of 3.15. A total of 34 79(57%) were from rural and fuel used was fired wood 41%. Hospital admission were 84(61%), vasooclusive crisis 104 (75%), acute chest syndrome 61(44%) and blood transfusions 56(40%) in the last 12 35 months. Prevalence of abnormal lung functions was 28 % (39), restrictive pattern 26(67%) and obstructive 36 37 pattern 13(33%). Urban setting (OR 24.101 p value 0.163), being female (OR 18.911 p value 0.069), and 38 blood transfusion (OR 11.683 p value 0.195) had high odds ratio while, stable hydroxyurea use (OR 0.525 39 p value 0.678) and hospital admission (OR 0.048 p value 0.121) had low odds ratio, but not statistically significant. 40

One third had abnormal lung function mostly restrictive followed by obstructive pattern. Residence in urban
setting, being female and blood transfusion had high odds ratio, while stable hydroxyurea use and hospital
admission had low odds ratio but no statistically significant.

44

46 Introduction

47

Sickle cell Disease (SCD) is a genetic disorder of the hemoglobin molecule that forms part of the red blood cell. The red blood cell is made up of a hem group and four globin chains which comprise of two beta and two alfa chains (B2a2). The gene clusters on chromosome 16 and 11 control globin chain production. The allele which is responsible for production of SCD is autosomal recessive and is found on the short arm of chromosome 11 (1).

In Sub Saharan Africa, around 240,000 children are born with the disease each year (2). Most of these children die without ever being diagnosed as there is limited newborn screening and early treatment in Sub-Saharan Africa. SCD is responsible for 50-90% of under-five mortalities in low income countries. This is because there is limited data on the natural history of the disease in our setting which makes early identification and treatment of the complications difficult leading to poor prognosis (3)

It is predicted that the numbers of children with this disorder will steadily increase over the next 40 years if mechanisms are not put in place to control the disease. Given this predicted rise in numbers of children with SCD, major investment in clinical research is necessary to fully understand the natural history of sickle cell disease in our setting and the associated respiratory complications (4).

62 Children with homozygous state (Hb SS which usually have a more severe course of the disease or Hb SC 63 with a milder course of the disease) are usually symptomatic with lifelong symptoms. which can be acute 64 or chronic with chronic organ damage which can present as kidney failure, stroke, stunting cardiomyopathy 65 and chronic lung disease which present with respiratory symptoms (5) (3)

66 Usually, the severity of the disease is attributed to a number of factors such as genetic modifiers, nutritional 67 status, access healthcare among others (6). However, the real burden and the prevalence of associated 68 morbidities such as obstructive and restrictive lung disease which can eventually progress to chronic lung 69 disease remains elusive (4).

There have also been contradicting findings from previous studies on whether the pattern of lung function abnormality contributes to future morbidity and mortality among SCD patients. In contrast, it has been argued that, the pattern of abnormal lung function test is not associated with previous or future painful crisis or acute chest syndrome and that, the abnormal lung functions are sometimes over estimated in certain circumstances (7).

75 It has been difficult comparing results between older studies which is hindered by the fact that there were
76 many different reference values and criteria used to define lung function abnormalities (8)

Acute and chronic lung complications in sickle cell disease are major causes of morbidity and mortality, accounting for up to 20-30% of all sickle cell deaths. This indicates that screening should be done early and required interventions instituted in order to reduce the burden of respiratory related complications among sickle cell patients (9)

The most prevalent abnormal spirometry findings among children and adolescents with SCD varies globally. In the US in a study done by Cohen and colleagues, noted that 30% of children had an abnormal lung function pattern. The most common abnormal pattern was obstruction which was found in 16%, with fewer patients having restriction at 7% (7). In Italy, Tassel found out that the most prevalent pattern was restrictive at 13% with obstruction at 5%, which is similar to what has been noted in some of the African countries (10)

In Sub Saharan Africa, which is a low income setting, restrictive spirometry pattern is the most prevalent pathological pattern. However, among pediatric patients with SCD living in high- income countries, obstructive lung function is more frequent than restrictive. This is a study which was done to compare the abnormal lung functions among children with SCD in Africa and United Kingdom (Arigliani, et al., 2019)

In the general populations, it has been established that a number of factors affect lung function, both environmental and intrinsic factors. In addition, it has been noted that lung function decline over time among children living with SCD. Regarding risk factors for abnormal lung function in children with SCD,

94 it has been pointed out that there is an association with history of acute chest syndrome (ACS), whereas the 95 rate of painful crises seemed to be unrelated to respiratory abnormalities, it has also been observed that 96 patients with more than 3 episodes of ACS had lower FEV1 than the predicted value (12), (13), (14).

In addition, Malnutrition was also associated with increased risk of restrictive spirometry pattern in children with SCD from Central Africa, while high body mass index (BMI) percentile was found to be an independent predictor of increase in forced expiratory volume at one second (FEV1) in patients from High Income Countries. Furthermore, Sex as an independent variable has been noted to have no significant influence on the respiratory function abnormalities in children with sickle cell disease in the same study. Arigliani and colleagues also noted that lung function decline with age in both children from low income and high income countries with differences in the patterns of abnormities (12) (8)(15)(16), (17)

In a study done in West Africa, ACS, hydroxyurea use and painful crisis were found to be significantly
associated with changes in lung functions. Furthermore, the use of hydroxyurea as a disease modifying
drug in sickle cell disease reduces the progression of loss of lung function in children with SCD (18) (19).
Additionally, a History of asthma and wheezing has also been associated with obstructive lung function
abnormality as was observed in a study by Manuel and colleagues (20), (21).

109 Methods and materials

110 Study site

This was a descriptive cross-sectional hospital-based study carried out at Jaramogi Oginga Odinga Teaching and Referral Hospital (JOOTRH) in Kisumu a level five hospital in western Kenya. Recruitment was done at the two outpatient clinics. The clinic at the Obama Children's hospital catered for children aged 6 to 13 years while those who are between 14 to 17 years were recruited from the hemato-oncology clinic within the same hospital which has a well-established sickle cell clinic.

117	Study	popu	lation
-----	-------	------	--------

- 118 Children with SCD receiving care at JOOTRH
- 119 Study period
- 120 The study was done over a period of six months from 1st October 2023 to 30th March 2024
- 121 Eligibility criteria
- 122 Inclusion criteria
- 123 1. Children between 6-17 years with clinical features of SCD and with a confirmatory test and receiving
- 124 care at JOOTRH whose parent or guardian consented and those who above 7 years gave a written
- 125 assent
- 126 Exclusion criteria
- 127 1. Confirmed cases of or on treatment for pneumonia, pulmonary tuberculosis or any other respiratory
- 128 disease at the time of recruitment
- Children who were not able to follow instruction for the study procedures due to illness or any other
 reason
- 131 3. Children who were on bronchodilator or steroids for active treatment of asthma

132

133 Sample size

134 The sample size was calculated using the Fisher's formula based on a p of 0.3 from the study by Cohen

which showed a prevalence of abnormal lung function at 30% (7).

$$\mathbf{n} = \frac{Z^2 \ \hat{p} \left(1 - \hat{p}\right)}{\mathbf{e}^2}$$

- 137 $1.96^{2*}0.3^{*}0.7 = 323$
- 138 0.05²
- 139

140 Given that the total number of unique children above 6-17 years seen per month was approximately 30,

141 therefore, the approximate total population of interest during data collection period was estimated to be 240

142 children.

143	Corrected sample size for finite population
144 145 146	$\frac{323}{1+(n-1)/N}$
147 148 149	$\frac{323}{1+(323-1)/240}$ =137.9=138

150 Recruitment

151 The Principal Investigator (PI) and the research assistants were trained on spirometry by a Respiratory 152 Society of Kenva (RESOK) trainer and certified to perform and interpreted the spirometry results. After being attended to by the clinician on duty for routine clinic visit, potential participants were directed to the 153 154 PI or the research assistant for screening who identified children who met the study requirements. Parents 155 or guardians who were willing to let their children join the study proceeded to give written consent and 156 written assent was obtained from children above 7 years. The PI or the research assistant took a short medical history of the participants including demographics such as age and sex, residence rural or urban, 157 158 cooking space whether indoor or outdoor, fuel used for cooking whether firewood, gas or charcoal. Medical information was also obtained which included hydroxyurea use, number of vaso-occlusive crisis (VOC), 159 160 ACS, blood transfusion, and hospital admission in the past 12 months. Anthropometric measurements such as weight and height was obtained and used to calculate BMI and recorded on a structured data collection 161 tool which was validated before the study. Where information was not clear, this was clarified from the 162 patients' folders. 163

164 Two NDD Easy-on PC spirometer device were available for the study where one acted as a backup. The 165 spirometry test was presented as a blowing game with computer incentives of blowing off the candles on a 166 birthday cake first for purposes of practice and during the actual procedure where blowing off all the candles

167 corresponded with achieving the required forced expiration. The participants were encouraged to blow and168 where necessary demonstration was done alongside the participant.

The patients were put in a sitting position with their arms resting on the arm rest of the chair and their back 169 170 straight and given time to relax and be composed before the spirometry procedure. A clean mouth piece 171 was attached to the spirometer device, patient's nose clipped and he or she was instructed to take in normal 172 three breathes before taking in a deep breath and then blowing out forcefully for 6 seconds, as the instructor 173 verbally encouraged the patient and the patient looking at the candle on the screen. The procedure was 174 repeated 3 times to obtain the highest values with minimal curve variability, which was in conformity with 175 standard international guidelines of European Respiratory Society and American Thoracic Society in terms of quality and reproducibility (23). Quality control was done to ensure consistency and the best values were 176 used for interpretation of the results in comparison to the predicted reference values. Where there was 177 178 difficulty interpreting the results, a pulmonologist was consulted for assistance

179 Independent variables

Some of the independent variables in the study were age, sex, BMI which was categorized as; normal, underweight, overweight or obesity, vaso-occlusive crisis, acute chest syndrome, blood transfusions, number of hospital admissions, stable hydroxyurea use, history of asthma or wheeze, type of cooking fuel used, residence either rural or urban and cooking area either outdoor or indoor.

184 Dependent variable

The dependent variable was abnormal lung function of which three types of outcomes were considered which were; restrictive i.e. lower FEV1<80% of lower limit of normal and obstructive i.e. lower FVC <80% or FEV1/FVC ratio <70% of the lower limit of normal or mixed with binary out comes for each and was coded as 0 absent or 1 present for each case.

189 Data Management and Analysis

190 Data entry and cleaning

191 Data was entered into database and cleaned to confirm accuracy and conformity with the source document

- 192 before analysis.
- 193 Data analysis

Data analysis was done according to the statistical analysis plan (SAP) in place to compute the frequency and proportion for categorical data, mean, SD and range for continuous data. A bivariate analysis was done for association, odds ratio at 95% confidence level. Fishers exact tests where applicable to test for statistically significance.

198 Ethical Approval

The study procedures were started after obtaining ethical approval from the Institutional Research Ethics
Committee (IREC) of Moi University/ Moi Teaching and Referral Hospital and from Jaramogi Oginga
Odinga Teaching and Referral Hospital Institutional Scientific Ethics and Research Committee (ISERC).
Administrative approval was obtained from the Hospital allowing the conduct of the study within the
facility.

Written consent was obtained from parents/guardian of all the children below 18 years and assent from children above 7 years.

206 Confidentiality

The participant's information was de-identified to remove any personally identifiable information in order
 to maintain confidentiality. The folders were kept in lockable drawers which were accessible to the PI and
 other authorized study staff only

210

211 Results

- A total of 151 participants were screened, 13 of them were screen excluded while 138 met the inclusion
- criteria and considered for the final analysis
- 214 Socio demographic characteristics
- 215 Characteristics of the participants' care giver
- Out of 138 participants, 103(75%) of the participants were brought to the hospital by their mothers, 13
- 217 (9%) were brought by their fathers while only 8(6%) were brought by others who were neither their mother,
- father or guardian. Out of the total number of parents, guardians and others who brought the participants to
- the hospital, 59(43%) of the parents or guardians of the children surveyed had attained secondary education,
- 42(30%) had Tertiary level while 3(2%) had no education.

221 Socio-demographic characteristic of the participants

- Table 1 shows the demographic characteristics; Sex distribution shows males were of 73(53%) while the
- 223 female were 65(47%).
- 224 225 226 227 228 229 230 231 232 232 233 234 235

Table 1: sociodemographic characteristics

Demographics		Freq (n)	Percent (%)
C	M-1-	72	52
Sex	Male	/3	53
	Female	65	47
Residence	Bungoma	1	1
	Homabay	4	3
	Kisumu	132	95
	Vihiga	1	1
Type of residence	Rural	79	57
	Urban	59	43
Fuel used	Firewood	56	41
	Charcoal	58	42
	Gas	24	17
Cooking space	Indoor	125	91
	Outdoor	13	9
BMI	overweight	0	0
	Underweight	126	91
	Normal	12	9

- 255 Anthropometric measurements

Anthropometric measurements showed that the mean age was 10.99 years with an SD of 3.15. The mean
weight of the children was 29.06 kg (with the minimum weight of 14.1 kg and maximum weight of 62 kg)
and SD of 9.8. The children had a mean height of 138.19 cm, (minimum height was108 cm and maximum
height as 178 cm) and SD of 15.5.

262 Clinical characteristics

263	Looking at the clinical features in table 2, 102(74%) of the children reported being on stable dose of
264	hydroxyurea within the las 3 months, 34(25%) reported no VOC while 71(51%) had 1-2 VOC,
265	additionally,77(56) had no ACS while 52 (38%) had one to two ACS in the past 12 months. In addition,
266	82(59) had received no blood transfusion and only 10(4%) had received more than three blood transfusions
267	in the past 12 months, while 84(61%) had been admitted to the hospital within the last 12 months

268

Table 2 pati	ents clinical	factors
--------------	---------------	---------

Clinical characteristics	Freq (n)	Percentage
Stable hydroxyurea dose		
Yes	102	74
No	36	26
No of VOCs within 12 months		
None	34	25
one-Two	71	51
>three	33	24
No of ACS within 12 months		
None	77	56
one-Two	52	38
>three	9	6
No of Blood transfusion within 12 months		
None	82	59
One-two	46	33
>three	10	4
Admission history within 12 months		
Yes	84	61
No	54	39

270 Prevalence of abnormal lung function

- 271 Out of the total number of the children, 39(28%) of them had abnormal lung functions, while 99(72%) had
- 272 normal lung function test

273 Patterns of abnormal lung functions

- Table 3 shows patterns of abnormal lung function. Out of the 39 children with abnormal lung functions
- 26(67%) had restrictive lungs function pattern while the rest 13(33%) had obstructive lung function pattern
- with none of the children having obstructive pattern.

277 Table 3: patterns of abnormal lung functions

Pattern of abnormal lung function	n	Percentage (%)
Restrictive	26	67
Obstructive	13	33
Mixed	0	0

²⁷⁸

279 Factors associated with abnormal lung functions

Table 4 shows factors associated with abnormal lung functions. The findings showed that being female appears to have a substantially high odds ratio of 18.91 compared to males, though the pvalue 0.069, 95% CI (0.794, 450.23). There is no significant association between weight status (underweight or normal) with abnormal lung functions, normal weight low odds ratio of 0.3036 and a p-value of 0.301 95% CI (0.032, 2.902). Living in an urban setup had remarkably high odds ratio 24.10 compared to rural settings, with a p-value 0.163 95% CI (0.276, 2104.596).

		Odds Ratio	P Value	[95% Conf. Interval]
				Lower, upper
Sex	Male	REF		
	Female	18.911	0.069	0.794 , 450.25
Weight	Underweight	REF		
	Normal	0.304	0.301	0.032, 2.902
	overweight	0		
Type of residence	Rural	REF		
	Urban	24.102	0.163	0.276, 2104.596
Type of fuel used at home	firewood	REF		
	charcoal	0.196	0.389	
	Gas	0.134	0.486	
No. of ACS in past 12 months	0	REF		
	1-2	2.444	0.533	0.147, 40.578
	≥3	1 (empty)		
No of Blood Transfusions received in the last 12	0	REF		
	1-2	11.683	0.195	0.285, 479.720
	≥3	1 (empty)		
Stable hydroxyurea dose for the last 3 months	No	REF		
	Yes	0.525	0.678	0.025, 10.927
admission in the past in the last 12 months	No	REF		
	Yes	0.048	0.121	0.001, 2.225

286 Table 4: Factors associated with abnormal lung functions

288 Clinical factors and patterns of abnormal lung functions

289 Table 5 shows clinical factors and patterns of abnormal lung functions where more than 3 blood transfusion

in 12 mothers showed statistically significant association with both restrictive and obstructive abnormal

291 patterns of lung function.

292 Table 5: Clinical factors and patterns of abnormal lung functions

Factors		Restrictive (Low FVC)	Obstructive (Low FEV1)	P-value
No of Vaso-Occlusive Crisis	0	n(%)	n(%)	
	1-2	7(70)	3(30)	
	3	13(85)	2(15)	0.614
Number of Acute chest syndrome (ACS)	0	11(75)	3(25)	
- <u>-</u>	1-2	16(78)	4(22)	
	3	12(73)	5(27)	0.816
Number of blood transfusions received	0	2(100)	0(0)	
	1-2	14(80)	3(20)	
	3	16(82)	4(20)	0.031
Patient on stable hydroxyurea dose for the last 3 months	Yes	0(0)	2(100)	
	No	22(79)	6(21)	0.566
History of asthma	Yes	8(73)	3(27)	
	No	1(100)	0(0)	0.573
History of Wheezing	Yes	29(76)	9(24)	
	No	3(100)	0(0)	0.314
Admitted in the past 12 month	Yes	27(75)	9(25)	
	No	19(79)	5(21)	0.434

297 Discussion

298

299 Prevalence of abnormal lung functions

Children living in low and middle income countries still have a substantial burden of abnormal lung functions. This was demonstrated in this study where 39(28%) of the children had an abnormal lung functions mostly of restrictive followed by obstructive pattern. Factors such us being female, blood transfusion and living in urban setting had high odds ratios while hospital admissions and stable hydroxyurea use had very low odds ratios.

305 There is poor access to health services and noncompliance to the sickle cell treatment leading to 306 complications such as VOC, ACS and hypoxic injuries. These frequent insults from hypoxia VOC and ACS 307 contribute to the interstitial lung damage leading to abnormal lung functions (8). There was also a female 308 predominance where out of 66, 20(30.3%) had abnormal lung function. This female predominance can be explained by the cultural practices of cooking with firewood in an enclosed cooking space as women are 309 310 expected to cook hence exposure smoke which leads to lung injuries hence abnormal lung functions. The 311 findings are similar with the study done in the USA by Cohen et al that found a prevalence of 30% among 312 children and adolescents (7). However, in a study done by Dei-Adomakoh and colleagues in adults with sickle cell, there was 70.4% prevalence which indicated that the decline in lung function among children is 313 314 a continuous process and worsens as they grow older (14). The difference can be explained by the fact that 315 the study populations were different in characteristic majorly adults as oppose to children and adolescents 316 in our case.

In Nigeria, in a study that was done by Arigliani to compare lung functions among Nigerian children and those from the United Kingdom, there was a prevalence of 38% in Nigeria, and 27% in United Kingdom. This is slightly higher in Nigeria than our study which was at 28%, but finding in the UK were similar to this study despite the fact that this was a cohort study where the participants were followed up and UK has a better health care system than Africa which alludes to the fact that developing abnormal lung function

could be intrinsic to having sickle cell disease. In this Study also, confounders were excluded as appropriate
 (11). These findings cannot over emphasize the need for including lung function test as part of routine care
 for children and adolescents with sickle cell disease

In Kuwait, A Adekite found a similar prevalence of 28.5%. Besides, they used a different phenotype of HbSC with very high fetal hemoglobin levels with mild symptoms and still had comparable prevalence to this study which involved HbSS phenotype. This equally showed that there is more work to be done to establish the exact mechanism of lung damage in sickle cell disease (24)

329 Patterns of abnormal lung function in children with SCD

330 There has been a contentious argument as to what is the most prevalent pattern of abnormal lung function 331 among children with sickle cell disease and as to whether this varies with regions. In this study, we found out that restrictive pattern was the most prevalent and double at 26(67%) while obstructive pattern was 332 333 13(33%). There was no mixed pattern observed in this study. The restrictive pattern seen in this study can be explained by the fact that most of the participants were underweight which is associated with restrictive 334 335 lung functions. This is in contrast to a study by Cohen, where they found out that the most prevalent pattern 336 was obstructive at 16% (7). This was possibly due to the urban setting of their studies as previously it has 337 been shown that children in such set up are more predisposed to hyperactive airway disease. In Ghana, Kuti 338 and Adegoke found that most of the children had restrictive lung disease followed by obstructive and a few 339 had mixed pattern (25). This is in congruent to this study where we found out a similar trend except for 340 mixed pattern which was not found among these children, which could be because we used only spirometry 341 which was not adequate in characterizing the patterns further.

There seem to be regional variation of the patterns as in the study done by Manuel et al which found exact reverse of our findings in this current study. In their study, the most prevalent pattern was obstructive at 19% followed by restrictive at 9% (20), they equally did not report on the mixed pattern. Their study setting was more urban which is similar to the study by Cohen and colleagues but with different results. In this

study, reversibility testing was not done to further characterize the obstructive pattern into either reversibleor irreversible which would give a glimpse of the possible cause as this was outside the study aims.

The previously differing opinions of the pattern of abnormal lung functions among children with sickle cell 348 349 disease is slowly taking shape as more studies indicate that in Africa and some European countries, the 350 most prevalent pattern is restrictive lung disease. This pattern can be explained by the repeated insults to 351 the lungs as a target organ as had been alluded to earlier. Furthermore, it was noted by Kuti and Adegoke 352 in Nigeria in their study that 22.1% of the children with SCD had restrictive pattern while 5.8% had obstructive and only 1.9% had mixed pattern (25). Though the findings are slightly different, it gives a 353 354 similar picture to the findings in this study where 19% had restrictive, 9% obstructive with none having 355 mixed pattern.

356 Factors associated with abnormal lung functions in children with SCD

It is now known that lung functions decline over time among children living with Sickle Cell Disease, however, there is paucity of data on some of the contributing factors to this rapid decline. However, the factors that contribute to this decline still remains elusive to the medical fraternity. Different studies have found different factors to be contributory, furthermore, most factors are usually not statistically significant despite the high odds ratio. This was no different from this study.

362 In this study being female had an increased risk of having an abnormal lung function compared to male OR 363 18.91 CI 0.7943,450.25, however, this was not statistically significant with a p value of 0.069. This is 364 similar to the study done in Central Africa by Arigliani which associated being female with increased risk 365 of having abnormal lung functions (12). The increased likelihood in females compared to males in this 366 study can be explained by the fact that in greater Africa and mostly in rural areas cooking is done indoors 367 with firewood which produces a lot of smoke by females increasing their exposure to pollutants leading to 368 lung damage hence developing the abnormal lung functions. This was in contrast to the previous studies 369 which showed that sex has no influence on the risk of having an abnormal lung function among children

with sickle cell disease (9), (26). This difference could not be explained since there were more males in
this study compared to females at 73(52.55%) and 65(47.45%) respectively. However, it can be
hypothesized that the difference could be attributed to the cooking environment where mostly occupied by
females.

When it comes to the BMI, most studies have found an association between obesity and abnormal lung functions mostly obstructive lung disease as in the case of a study by Koumbourlis where BMI had a statistically significant influence with a p value of 0.02 (27). In this study having a normal BMI was shown to be protective, however, this was not statistically significant with a p value of 0.301, with OR of 0.304 at 95% CI (0.0318, 2.9019). This similarity is due to the fact that we used similar populations of study. In another study however, BMI was found to have no association with abnormal lung function (14). This difference could be explained by the fact that they used slightly different study population.

In this study, it was postulated that urban residence may have some significant contributions in children with sickle cell anemia, this could be similar to populations without SCD. This was attributed to the industrial and motor vehicle emissions causing air pollution in town centers which contributes to lung injury. It was found out that living in urban set up had an increased risk of having an abnormal lung functions at more than 24 times compared to rural set up. However, this was not statistically significant OR 24.1015 CI 95% (0.2760, 2104) with a p value of 0.163.

In Nigeria, use of solid fuels was found to have no significant association with abnormal lung functions in children with sickle cell disease (28). This was similar to this study where it was found out that type of fuel had no significant association with abnormal lung functions p value 0.389 for charcoal and 0.486 for gas. However, compared to use of fired wood, charcoal and gas were protective with OR of 0.1959 and 0.1344 respectively. This can be attributed to the amount of smoke produced by firewood within an enclosed indoor cooking place which predisposes the house occupants to air pollutions hence increasing chances of having lung damage and abnormal lung functions.

394 Given the hypothesized mechanism of lung injury leading to fibrosis and interstitial lung injury with 395 ultimate impaired lung function in children with SCD, the number of Acute Chest Syndrome (ACS) has 396 been associated with the decline in lung function in this kind of population (29). In this study, children who 397 had more than 2 episodes of ACS in the past 12 months had a slightly increased likelihood of having an 398 abnormal lung functions twice as high compared to those who had none, however, this was not statistically 399 significant OR 2 .44 CI (0.1473, 40.5778) at 95% confidence level. In a study by Arigliani, having more 400 than 3 episodes of ACS was associated with impaired lung function (12). However, Cohen and colleagues 401 found no association of ACS with abnormal lung functions. This difference can be explained by the fact 402 that Cohen study was done in a high income country with good health care services to children with SCD which limited the effect of ACS due to timely and adequate management of the acute complications of 403 404 patients with sickle cell disease. Kuti also found out that ACS was associated with abnormal lung functions 405 in Nigerian children with SCD with an OR of 8.5 (25), (7).

406 Sometimes children with SCD, can have complications which warrant urgent transfusions. This can be 407 occasional or at times they can be dependent on chronic transfusion. As an independent factor, transfusion 408 was associated with abnormal lung functions with a p value of 0.195. However, in the presence of other 409 factors, children who received more than 2 transfusions within the last 12 months had greatly increased risk 410 of having an abnormal lung function even though it was not statistically significant OR 11.683 95% CI (0.285, 479) P value 0.195 this increased likelihood may not be due to transfusion itself but could be 411 412 attributed to other issues such as noncompliance to medications and clinic appointments predisposing them 413 to multiple complications which contribute to their abnormal lung functions. This was different from the 414 study by Adekile in Nigeria which found out that there was no association between FEV1 and 415 transfusions (24). This difference was because they narrowed only to the effect on FEV as opposed to 416 generally looking at the abnormal lung function as was the case in this study.

417 The mainstay of therapeutic management of SCD patient is by use of hydroxyurea which is based on the 418 weight and patients' response. Lung functions have been shown to improve after starting hydroxyurea

419 treatment (18). This raised the questions as to whether hydroxyurea use can slow or prevent the 420 development of abnormal lung function. In this study, being on a stable dose of hydroxyurea for three 421 months shown to be protective, however, it was not statistically significant with an OR 0.566 at 95% CI 422 (0.02, 10.92) with a p value of 0.678. This was similar to a study by J. Stewart who found no significant 423 association with a p value 0.24 (30). Similarly Arigliani found no significant association in Central Africa 424 (12). This shows that there is still work to be done in order to ascertain the contribution of hydroxyurea to 425 the decline in lung functions among children with SCD. In a meta-analysis by Taksande, hydroxyurea use 426 was found to be protective, but not statistically significant which is similar to this study (21).

427 Following the devastating complications of SCD, most children needs frequent and repeated hospital admission for appropriate and proper care. This is not possible for some families from low income countries 428 429 thus leads to inadequate medical care. In this study, hospital admission within the past year was protective 430 but not significant OR 0.047 P value 0.121 CI (0.001, 2.224), this can be explained by the fact that children 431 with more hospital admissions received adequate and appropriate medical care which reduced their chances of developing the complications associated with development of abnormal lung functions. This was 432 433 different from a study by Kuti which found a significant association of abnormal lung functions with hospitalization p value <0.05 in Nigeria (25), but similar to the findings of the study by Arigliani and 434 435 colleagues (8). It is therefore necessary for improved health care services for children and adolescents with 436 SCD in order to improve their general well-being and reduce risk of developing an abnormal lung functions. 437 This study had a limitation of wider confidence interval which can be improved in future by doing cohort 438 studies in Africa with larger sample sizes.

439

440

442 Conclusion

- 443 The prevalence of abnormal lung function among children with Sickle Cell Disease was 28%
- 444 The most prevalent pattern of abnormal lung function was restrictive followed by obstructive and no mixed
- 445 pattern
- 446 Urban setting, being female and the number of blood transfusion received within the past 12 months were
- 447 associated with high odds ratio while hospital admissions and stable hydroxyurea used were associated with
- 448 low odds ratio.

449 Acknowledgment

450 I would like to thank Prof. Bernhards Ogutu, Dr. Vincent Were, Dr. John Orimbo, Dr. Allan Otieno,

451 Caroline Atieno, Pamela Atieno Nashon Okello, Sophy Omondi, Grace Omukoko, Ishmael Sports and Jane

452 Olum for their valuable contribution towards the realization of the study goals.

453 References

- Tenge CN. Preventing birth of sicklers as an effective strategy in a sickle cell disease control programme. East Afr Med J. 2019;96(1):2241–9.
- 456 2. Williams TN. Sickle Cell Disease in Sub-Saharan Africa. Hematol Oncol Clin North Am. 2016
 457 Apr 1;30(2):343–58.
- Uyoga S, Macharia AW, Mochamah G, Ndila CM, Nyutu G, Makale J, et al. The epidemiology of sickle cell disease in children recruited in infancy in Kilifi, Kenya: a prospective cohort study.
 Lancet Glob Heal. 2019 Oct 1;7(10):e1458–66.
- 4. Kato GJ, Piel FB, Reid CD, Gaston MH, Ohene-Frempong K, Krishnamurti L, et al. Sickle cell disease. Nat Publ Gr [Internet]. 2018;4. Available from: www.nature.com/nrdp
- 463 5. Watenga EK, Songok JJ, Tenge CN. The prevalence of sickle cell trait among adolescents in
 464 Bungoma County, Kenya. East Afr Med J. 2019;96(5):2683–91.
- 465 6. Rees DC, Brousse VAM, Brewin JN. Determinants of severity in sickle cell disease. Blood Rev
 466 [Internet]. 2022;56(June):100983. Available from: https://doi.org/10.1016/j.blre.2022.100983
- Cohen RT, Strunk RC, Rodeghier M, Rosen CL, Kirkham FJ, Kirkby J, et al. Pattern of lung
 function is not associated with prior or future morbidity in children with sickle cell anemia. Ann
 Am Thorac Soc. 2016;13(8):1314–23.
- 470 8. Arigliani M, Gupta A. Management of chronic respiratory complications in children and
 471 adolescents with sickle cell disease. Eur Respir Rev [Internet]. 2020;29(157):1–9. Available from: 472 http://dx.doi.org/10.1183/16000617.0054-2020
- 9. Dei-Adomakoh YA, Afriyie-Mensah JS, Forson A, Adadey M, Ndanu TA, Acquaye JK. Lung
 Function Abnormalities in Sickle Cell Anaemia. Adv Hematol. 2019;2019.
- Tassel C, Arnaud C, Kulpa M, Fleurence E, Kandem A, Madhi F, et al. Leukocytosis is a risk factor for lung function deterioration in children with sickle cell disease. Respir Med. 2011
 May;105(5):788–95.
- 478 11. Arigliani M, Castriotta L, Zubair R, Dogara LG, Zuiani C, Raywood E, et al. Differences in lung function between children with sickle cell anaemia from West Africa and Europe. Thorax. 2019 Dec 1;74(12):1154–60.
- 481 12. Arigliani M, Kitenge R, Castriotta L, Ndjule P, Barbato V, Cogo P, et al. Lung function in children with sickle cell disease from Central Africa. Thorax. 2019 Jun 1;74(6):604–6.
- 13. Ndukwu CI, Ozoh OB, Ale BM, Ayuk AC, Elo-Ilo JC, Awokola BI. Spirometry Abnormalities
 and Its Associated Factors Among Primary School Children in a Nigerian City. Clin Med Insights
 Pediatr. 2021;15:117955652110018.
- 486 14. Dei-Adomakoh YA, Afriyie-Mensah JS, Forson A, Adadey M, Ndanu TA, Acquaye JK. Lung
 487 Function Abnormalities in Sickle Cell Anaemia. Adv Hematol. 2019;2019.
- 15. Cook J, Jefferis O, Matchere P, Mbale E, Rylance J. Sickle-cell disease in Malawian children is associated with restrictive spirometry: a cross-sectional survey. Int J Tuberc Lung Dis [Internet].
 2013 Sep 1 [cited 2022 Apr 24];17(9):1235–8. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23827770
- 16. Rucheet Patel, M.D.* Olufolake Adisa MD. AMF. Decline of Lung Function in Children with

- 493 Sickle Cell Disease Is Not Associated with Restrictive Defects. 2017 [cited 2022 Aug 28];
 494 Available from: www.atsjournals.org.
- Willen SM, Cohen R, Rodeghier M, Kirkham F, Redline SS, Rosen C, et al. Age is a predictor of a small decrease in lung function in children with sickle cell anemia. Am J Hematol. 2018 Mar 1;93(3):408–15.
- 498 18. Kotwal N, Pillai DK, Darbari DS, Sun K, Koumbourlis AC. Spirometric Changes After Initiation 499 of Hydroxyurea in Children With Sickle Cell Anemia. J Pediatr Hematol Oncol [Internet]. 2021 500 Dec 30 [cited 2022 Apr 24]; Available from: http://www.ncbi.nlm.nih.gov/pubmed/34966092
- 501 19. Meier ER, Miller JL. Sickle cell disease in children. Drugs. 2012;72(7):895–906.
- Manuel Arteta1, Andrew Campbell1, Mehdi Nouraie2, Sohail Rana2, Onyinye Onyekwere2,
 Gregory Ensing1, et, al. Abnormal pulmonary function and associated risk factors in children and
 adolescents with sickle cell anemia. Physiol Behav. 2017;176(1):139–48.
- Taksande A, Jameel PZ, Pujari D, Taksande B, Meshram R. Variation in pulmonary function tests
 among children with sickle cell anemia: A systematic review and meta-analysis. Pan Afr Med J.
 2021;39(140).
- 508 22. Moore VC. Spirometry: Step by step. Breathe. 2012;8(3):233–40.
- Beydon N, Davis SD, Lombardi E, Allen JL, Arets HGM, Aurora P, et al. An Official American
 Thoracic Society/European Respiratory Society Statement: Pulmonary function testing in
 preschool children. Am J Respir Crit Care Med. 2007;175(12):1304–45.
- 512 24. Adekile AD, Azab AF, Owayed A, Khadadah M. Correlates of Pulmonary Function in Children
 513 with Sickle Cell Disease and Elevated Fetal Hemoglobin. Med Princ Pract. 2018;27(1):49–54.
- 514 25. Kuti B, Adegoke S. Authors' Response Pulmonary function abnormalities in Nigerian children
 515 with sickle cell anemia: Prevalence, pattern, and predictive factors. Pediatr Respirol Crit Care
 516 Med. 2019;3(1):18.
- 517 26. Achigbu KI, Odetunde OI, Chinawa JM, Achigbu EO, Ikefuna AN, Emodi IJ, et al. Pulmonary
 518 function indices in children with sickle cell anemia in Enugu, south-east Nigeria. Saudi Med J.
 519 2015 Aug 3;36(8):928–34.
- 520 27. Koumbourlis AC. Lung function in sickle cell disease: An elusive relationship. Vol. 51, Pediatric
 521 Pulmonology. John Wiley and Sons Inc.; 2016. p. 665–7.
- Arigliani M, Dogara LG, Zubair R, Castriotta L, Pusiol A, Zuiani C, et al. Lung function
 impairment in pediatric patients with sickle cell anemia from Nigeria is associated with low steady
 state hemoglobin. Am J Hematol. 2019 Mar 1;94(3):E70–1.
- Al Biltagi M, Bediwy AS, Toema O, Al-Asy HM, Saeed NK. Pulmonary functions in children and adolescents with sickle cell disease. Pediatr Pulmonol. 2020 Aug 1;55(8):2055–63.
- 527 30. Stewart JAT, Willen SM, Cohen R, Rodeghier M, Kirkham F, Redline SS, et al. BMI percentile is
 528 an independent predictor of increase in lung function in children with sickle cell anemia. Am J
 529 Hematol. 2019 May 1;94(5):E136–8.
- 530