Overlap between ultra-processed food and food that is high in fat, salt or sugar: analysis of 11 annual waves of the UK National Diet and Nutrition Survey 2008/09-2018/19

Authors: Viktorija Kesaite¹, Yanaina Chavez-Ugalde¹, Martin White¹, Jean Adams¹

Affiliations:

1. MRC Epidemiology Unit, University of Cambridge School of Clinical Medicine, Cambridge, United Kingdom

Corresponding author: Viktorija Kesaite

E-mail: Viktorija.Kesaite@mrc-epid.cam.ac.uk

Word count: 3486

Number of tables: 4

Number of figures: 1

Orcid IDs:

Viktorija Kesaite: https://orcid.org/0000-0002-0053-7583

Yanaina Chavez-Ugalde: <u>https://orcid.org/0000-0001-6191-2722</u>

Jean Adams: https://orcid.org/0000-0002-5733-7830

Martin White: <u>https://orcid.org/0000-0002-1861-6757</u>

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

1 Abstract

2 While many countries use guidance and policies based on nutrients and food groups to support 3 citizens to consume healthy diets, fewer have explicitly adopted the concept of ultra-processed foods 4 (UPF). UPF consumption is associated with many adverse health outcomes in cohort studies. In the 5 UK, a nutrient profiling model (NPM) is used to identify foods high in fat, salt or sugar (HFSS) and 6 several policies target these. It is not known how well the NPM also captures UPF. We aimed to 7 quantify the proportion of food and drink items consumed in the UK that are HFSS, UPF, both or 8 neither and describe the food groups making the largest contributions to each category. We analysed 9 data from the National Diet and Nutrition Survey (NDNS), between 2008/09 and 2018/19, using 10 descriptive statistics. We used three metrics of food consumption: all foods, % of energy in all foods 11 (reflecting that different foods are consumed in different portion sizes and are of different energy 12 densities), and % of food weight in all foods (reflecting that some UPFs have few calories but are 13 consumed in large volumes). We found that, 33.4% of foods, 47.4% of energy, and 16.0% of food 14 weight were HFSS; 36.2%, 59.8% and 32.9% respectively were UPFs; 20.1%, 35.1% and 12.6% were 15 both; and 50.5%, 27.9% and 63.7% were neither. In total, 55.6% of UPF foods, 58.7% of energy from 16 UPFs and 38.3% of food weight from UPF consumed were also HFSS. The most common food groups 17 contributing to foods that were UPF but not HFSS were low calorie soft drinks and white bread. The 18 UK NPM captures at best just over half of UPFs consumed in the UK. Expanding the NPM to include 19 ingredients common in UPFs would capture a larger percentage of UPFs and could incentivise "de-20 formulation" of UPF products.

What is already known on	In many high-income countries, including the UK, consumption of		
this topic	ultra-processed food (UPF) and foods high in fat, salt or sugar (HFSS)		
	constitute more than half of all calories consumed.		
	Despite emerging evidence of the negative health impacts of UPF,		
	current UK diet and obesity policy focuses on reducing consumption of		
	HFSS foods. HFSS foods are identified using a Nutrient Profiling Model		
	(NPM).		
What this study adds	To our knowledge, this is the first study to examine the extent to which		
	the UK NPM also identifies UPFs. We found that the NPM captures over		
	half of UPFs based on all foods consumed and percent of energy, but		
	only about a third based on food weight. Amongst all foods consumed,		
	the most common food groups that were UPF but not HFSS were low		
	calorie soft drinks and white bread. Other types of bread (e.g. brown,		
	wholemeal) and high fibre breakfast cereals were also common.		
How this study might	Further consideration and potential inclusion of those UPFs that are		
affect research, practice,	not under the scope of the current definition of HFSS (e.g. foods		
or policy	containing artificial sweeteners) could enable existing policies to have		
	a wider reach.		

23 INTRODUCTION

Numerous studies have shown that diets high in some components (e.g. sodium, sugar, fat and calories) and low in others (e.g. whole grains and fruits) increase the risk of non-communicable diseases such as cardiovascular disease and cancer (1). Recent estimates suggest that food and drinks (hereafter: 'foods') high in fat, salt, or sugar (HFSS) constitute more than half of total energy intake among adults in the UK (2, 3). Similar estimates are observed in the consumption of ultra-processed foods (UPF) (4).

In the UK, a Nutrient Profiling Model (NPM) is used to identify HFSS foods and restrict advertising on television (5), out-of-doors in some local authorities and where foods can be placed in grocery stores (6). The NPM was originally developed in 2004/05, and defines foods as being HFSS based on energy, saturated fat, total sugar, sodium, fibre, protein, and fruit, vegetable, and nut content per 100g (7). An updated version was developed in 2018 (8), based on revised guidance on sugar and fibre intake (9) however, despite consultations, has not yet been adopted into practice (10).

been negatively associated with adverse health outcomes (12). The Nova system is the most common method of classifying foods based on their degree of food processing used in the academic literature and has been identified as the most applicable to the UK (10). Recent Euromonitor statistics suggest that the highest volume of sales of UPFs globally are in Western Europe, North America, and Australasia, with increasing trends observed elsewhere (13, 14). Findings based on a systematic review indicate that the US and the UK had the highest proportion of energy intake from UPFs (>50%)(4).

The UK's approach to public health nutrition, where nutrients and food groups such as fruit and
vegetables, are used to drive guidance and policy reflects similar approaches in many countries.
However, a growing number of countries have now also included specific reference to avoidance of
UPFs in their dietary guidance (15).

47 As policy makers contend with whether to focus policy and dietary guidance on UPFs, an important 48 question is: how well do current approaches, such as the UK's NPM, also identify UPFs? If most UPFs 49 are also HFSS then the benefits of introducing additional UPF-focused policy and guidance may be 50 minimal. We are aware of only one other study examining the overlap between UPFs and other 51 methods of identifying less-healthy foods (3). This recent US study found that around three-quarters 52 of UPF foods purchased were also HFSS and that extending the HFSS definition to include non-53 nutritive sweeteners, flavourings, colours and additives would capture nearly 100% of UPFs (3). To 54 the best of our knowledge no previous work has explored the overlap between HFSS, as identified by

55 the UK NMP, and UPF in the UK.

56 In this study, our aim was to quantify the proportion of foods consumed in the UK that are HFSS, UPF,

57 both or neither.

58 METHODS

We used data from eleven waves (2008/9 – 2018/19) of the UK National Diet and Nutrition Survey
(NDNS).

61 Data description

- NDNS is a continuous cross-sectional survey conducted every year in the UK. It collects data on food consumption, nutrient intake, and nutritional status of the general population aged 1.5 years and over residing in private UK households. Each year, a multistage probability design is used to generate a new random sample. Selected household addresses are clustered into small geographical units called Primary Sampling Units. Households are randomly selected from these units, and participants are randomly selected from each household. Full data is collected from about 500 children and 500 adults each year.
- 69 In 2008-19, all NDNS participants self-reported their food and beverage consumption over three or
- 70 four days using a written food diary, with portion sizes recorded using standard household measures
- 71 and product labels. Parents and guardians reported on behalf of children under 11 years. Food diary
- 72 data are entered into the Diets In Nutrients Out (DINO) database and food codes are used to link
- 73 foods to the NDNS nutrient composition database, which includes nutritional information on more
- than 6000 foods (16). The response rate to food diaries is 50% or more. The NDNS provides sample
- 75 weights to reduce the risk of non-random selection (16).

76 Categorisation of foods as HFSS and UPF

77 We categorised foods as HFSS or not based on the UK NPM 2004/5 and, in a sensitivity analysis, using 78 the 2018 NPM. We used the Nova framework to classify foods as UPF or not, using assignments from 79 previous work (17) (18). In this, two researchers independently classified foods into the four Nova 80 groups. To assign Nova group, NDNS foods were coded where possible based on their NDNS main food 81 group (n=56). If there was uncertainty about whether all foods in a main food group would have the 82 same Nova category, foods were categorised where possible by food subgroups (n=136). In cases of 83 further uncertainty, individual food items (n=4,555) were classified (e.g., composite dishes which 84 include individual foods from more than one main or sub-group). Any disagreement between the 85 researchers on the classifications were resolved through discussion. This process achieved a high 86 degree of inter-rater reliability - 97% in the first and 99.8% in the second round after discussion (19).

Food supplements (i.e., vitamins and minerals) are not classified by the Nova system and we did not
include them or alcoholic beverages.

89 Data analyses

90 Reflecting our aims, our analyses were descriptive. Firstly, we conducted a food-level analysis. This 91 used all reported foods consumed by NDNS participants in 2008/09-2018/19 with foods reported 92 multiple times being included multiple times, which effectively weights the findings for consumption. 93 We described the percentage of foods that were UPF, HFSS, neither or both. We present this first as 94 a food level analyses including all foods and, separately, the relative contribution of UPFs and HFSS to 95 total food energy (kcal) and total food weight (g) intake. We then present examples of food groups 96 making the greatest contributions to each category based on all foods, total food energy (kcal), and 97 total weight (g), and finally present a person-day-level analysis which replicates the food level analysis 98 adjusting for non-response bias. We include an analysis based on total food energy to reflect that 99 different foods are consumed in different portion sizes and have different energy densities and so 100 make different relative contributions to the diet. This enables comparisons with other studies (3). We 101 also include a metric based on total food weight to reflect that some UPFs make little contribution to 102 energy (e.g., artificially sweetened beverages), and list the NDNS main food groups making the largest 103 contributions to each of these four categories (UPF, HFSS, both and neither).

Secondly, we conducted person-level analyses. These replicate the food level analyses but by including population survey weights we additionally account for non-response bias. As a robustness check, we replicated the food-level analyses of foods consumed for men and women separately. The study protocol is available at <u>https://osf.io/hr6f3/</u>, the code and the libraries for calculating HFSS scores and the analyses is available at <u>https://github.com/VKesaite/HFSS-and-UPF</u>. All calculations were carried out in Python version 3.11.4. There were no substantive changes to the protocol.

110 **RESULTS**

111 A total of 15,655 individuals (7,207 males and 8,448 females) were included in the analysis. They

- 112 reported consuming 1,730,158 foods, representing 4,555 unique food names. We report findings
- 113 based on the 2004/05 NPM here and on the 2018 NPM in supplemental material.

114 Food-level analysis

115 Figure 1 shows Venn diagrams describing the proportions of (a) foods, (b) food energy and (c) food

- 116 weight that were HFSS (right inner circle), UPF (left inner circle), both (inner circle overlap) and neither
- 117 (outer circle). The proportion of UPFs that are HFSS is also shown. In total, 33.4% of foods, 47.4% of

energy and 16.0% of food weight was HFSS, whilst 36.2% of food, 59.8% of energy and 32.9% of food
weight was UPF. We found that 55.6% of UPFs, 58.7% of energy from UPFs, and 38.3% of UPF weight
was also HFSS. Robustness checks in males and females separately showed similar findings (Table S1).
Analysis using the 2018 version of the NPM led to broadly similar results, but in general a slightly
smaller proportion of UPF was categorised as HFSS compared to the 2004/05 NPM (Table S1).

123 Table 1 lists the ten main food groups making the greatest contributions to the categories HFSS only, 124 UPF only, both and neither for all foods. Tables 2 and 3 show similar data for food energy and food 125 weight respectively. The most common main food groups contributing to foods and food energy that 126 were HFSS but not UPF reflected products high in sugars (i.e. 'sugars, preserves and sweet spread') 127 and dairy products (or replacements) such as cheese, butter, oil, butter and oil replacements and 128 whole milk. When analysed by food weight, the most common food groups contributing to the HFSS 129 only category were dairy products such as whole milk, cheese, butter, and cream products, and also 130 sugars (i.e. 'sugars, preserves and sweet spread'), bacon and ham products, and starchy products (i.e. 131 'pasta, rice & other cereals').

The most common main food groups contributing to foods in the category UPF but not HFSS often included non-nutritive sweeteners (i.e. 'soft drinks low calorie', 'artificial sweeteners') or were breads including white, brown, granary and wholemeal bread. Fried and roast potatoes and potato products were also a common food group in this category. When considering food weight and food energy, the most common food groups in the UPF only category were similar to total food category.



139 Figure 1. The percentage of (a) all foods, (b) food energy in kcal, and (c) food weight in grams consumed that was derived from foods that are high in fat,

salt or sugar, ultra-processed foods, both or neither; UK National Diet and Nutrition Survey 2008/9-2018

Note. HFSS = high in fat, salt or sugar (using the 2004/05 Nutrient Profiling Model); UPF = ultra-processed food

143 Table 1. The ten main food groups (% contribution) making the greatest contributions to the categories: high in fat, salt or sugar only, ultra-processed

144 food only, both and neither at the food level for all foods; UK National Diet and Nutrition Survey 2008/9-2018/19

All foods consumed				
HFSS only	UPF only	HFSS and UPF	Neither HFSS nor UPF	
Sugars, preserves & sweet spread (26.6%)	Soft drinks, low calorie (21.3%)	Miscellaneous [*] (14.0%)	Tea, coffee & water (31.0%)	
Miscellaneous [*] (13.1%)	White bread (17.5%)	Biscuits (10.1%)	Vegetables not raw (13.0%)	
Cheese (10.9%)	Miscellaneous [*] (5.9%)	Reduced fat spread (9.6%)	Semi-skimmed milk (12.3%)	
Butter (10.4%)	Chips, fried & roast potatoes, potato products (5.8%)	Soft drinks, not low calorie (8.2%)	Fruit** (9.8%)	
Other margarine, fats & oils [‡] (9.2%)	High fibre breakfast cereals (5.5%)	Crisps & savoury snacks (6.1%)	Salad & other raw vegetables (7.9%)	
Whole milk (7.9%)	Brown, granary & wheatgerm bread (5.5%)	Chocolate confectionery (6.1%)	Pasta, rice & other cereals (4.3%)	
Bacon & ham (4.1%)	Yoghurt, fromage, frais, & dairy desserts (5.4%)	Buns, cakes, pastries & fruit pies (5.4%)	Miscellaneous [*] (3.4%)	
Pufa margarine & oils (3.8%)	Wholemeal bread (5.3%)	Bacon & ham (3.6%)	Other potatoes, potato salads & dishes ^{‡‡} (2.6%)	
Nuts & seeds (2.9%)	Vegetables not raw (4.0%)	Other breakfast cereals ⁺⁺ (3.5%)	Fruit juice (2.6%)	
Other milk & cream ⁺ (2.7%)	Artificial sweeteners (3.5%)	Sugars, preserves & sweet spreads (3.5%)	Chicken & turkey dishes (2.3%)	

145

146 Note: HFSS = high in fat, salt or sugar (using the 2004/05 Nutrient Profiling Model); UPF = ultra-processed food ; main food groups are those used in the National Diet and

147 Nutrition Survey; *Miscellaneous includes e.g. mayonnaise, sour cream dip, soups; **Fruit includes e.g. raw fruit, fruit stewed with and without sugar, fruit canned in fruit

and sugar syrup; ⁺Other milk & cream includes e.g. plant-based milk alternatives, coffee creamer, crème fraiche; ⁺⁺Other breakfast cereals includes e.g. special flakes, rice

149 crispies; *Other margarine, fats & oils includes e.g. vegetable suet, palm oil, chicken fat; **Other potatoes, potato salads & dishes includes potato curry, mash potato with

150 butter and cream.

151 Table 2. The ten main food groups (% contribution) making the greatest contributions to the categories: high in fat, salt or sugar only, ultra-

152 processed food only, both and neither at the food level for food energy (kcal); UK National Diet and Nutrition Survey 2008/9-2018/19

Food energy (kcal) in all foods consumed					
HFSS only	UPF only	HFSS and UPF	Neither HFSS nor UPF		
Cheese (22.2%)	White bread (27.8%)	Biscuits (11.2%)	Pasta, rice & other cereals (16.2%)		
Butter (14.6%)	Chips fried & roast potatoes & potato products (12.7%)	Buns, cakes, pastries & fruit pies (10.0%)	Fruit** (11.9%)		
Sugars, preserves & sweet spreads (12.9%)	Brown granary & wheatgerm bread (8.4%)	Chocolate confectionery (7.8%)	Semi skimmed milk (10.8%)		
Whole milk (8.6%)	Wholemeal bread (7.3%)	Crisps & savoury snacks (7.2%)	Chicken & turkey dishes (9.3%)		
Bacon and ham (8.2%)	High fibre breakfast cereals (6.9%)	Pasta, rice & other cereals (7.1%)	Other potatoes, potato salads & dishes ⁺⁺ (7.7%)		
Other margarine, fats & oils [‡] (6.0%)	Pasta, rice & other cereals (5.5%)	Soft drinks, not low calorie (7.0%)	Vegetables not raw (7.0%)		
Nuts & seeds (4.3%)	Vegetables not raw (4.3%)	Miscellaneous [*] (5.8%)	Beef, veal & dishes (5.5%)		
Other milk & cream ⁺ (3.0%)	Yoghurt, fromage, frais & dairy desserts (3.9%)	Sausages (5.4%)	Eggs, and egg dishes (4.7%)		
Buns, cakes, pastries (2.2%)	Miscellaneous [*] (3.5%)	Reduced fat spread (5.1%)	Fruit juice (4.3%)		
Pufa margarine & oils (2.1%)	Coated chicken (3.3%)	Meat pies & pastries (4.1%)	Chips fried & roast potatoes, and potato products (4.1%)		
Pasta, rice & other cereals (2.0%)	Other milk & cream ⁺ (1.9%%)	Other breakfast cereals (3.8%)	Whole milk (4.0%)		

153

154 Note: HFSS = high in fat, salt or sugar (using the 2004/05 Nutrient Profiling Model); UPF = ultra-processed food ; main food groups are those used in the National Diet and

155 Nutrition Survey; *Miscellaneous includes e.g. mayonnaise, sour cream dip, soups; **Fruit includes e.g. raw fruit, fruit stewed with and without sugar, fruit canned in fruit

and sugar syrup; [†]Other milk & cream includes e.g. plant-based milk alternatives, coffee creamer, crème fraiche; ^{††}Other breakfast cereals includes e.g. special flakes, rice

157 crispies; *Other margarine, fats & oils includes e.g. vegetable suet, palm oil, chicken fat; **Other potatoes, potato salads & dishes includes potato curry, mash potato with

158 butter and cream.

159 Table 3. The ten main food groups (% contribution) making the greatest contributions to the categories: high in fat, salt or sugar only, ultra-processed

160 food only, both and neither at the food level for food weight (g); UK National Diet and Nutrition Survey 2008/9-2018/19

161

Food weight (g) in all foods consumed				
HFSS only	UPF only	HFSS and UPF	Neither HFSS nor UPF	
Whole milk (34.4%)	Soft drinks, low calorie (35.4%)	Soft drinks, not low calorie (34.3%)	Tea, coffee & water (57.3%)	
Cheese (15.2%)	White bread (9.9%)	Pasta, rice & other cereals (6.2%)	Semi-skimmed milk (7.9%)	
Sugars, preserves, & sweet spreads (9.3%)	Soft drinks, not low calorie (6.5%)	Miscellaneous [*] (5.7%)	Fruit** (6.7%)	
Bacon and ham (9.1%)	Miscellaneous [*] (6.1%)	Buns, cakes, pastries & fruit pies (5.6%)	Vegetables, not raw (5.6%)	
Butter (5.4%)	Chips fried & roast potatoes, & potato products (5.5%)	Biscuits (5.1%)	Fruit juice (3.7%)	
Other milk & cream ⁺ (2.4%)	Yoghurt, fromage, frais & dairy desserts (4.7%)	Sausages (4.3%)	Pasta, rice, & other cereals (3.3%)	
Pasta, rice & other cereals (2.0%)	Vegetables not raw (4.3%)	Yoghurt, fromage, frais & dairy desserts (3.4%)	Other potatoes, potato salads & dishes ^{‡‡} (3.1%)	
Nuts & seeds (2.0%)	Tea, coffee & water (3.4%)	Chocolate confectionery (3.3%)	Whole milk (2.1%)	
Other margarine, fats & oils [‡] (1.8%)	Pasta, rice & other cereals (3.4%)	Crisps & savoury snacks (3.0%)	Salad & other raw vegetables (2.0%)	
White fish coated or fried (1.7%)	Brown granary & wheatgerm bread (3.1%)	Ice cream (2.9%)	Chicken & turkey dishes (1.9%)	

162

163 Note: HFSS = high in fat, salt or sugar (using the 2004/05 Nutrient Profiling Model); UPF = ultra-processed food ; main food groups are those used in the National Diet and

164 Nutrition Survey; *Miscellaneous includes e.g. mayonnaise, sour cream dip, soups; **Fruit includes e.g. raw fruit, fruit stewed with and without sugar, fruit canned in fruit and 165 sugar syrup; [†]Other milk & cream includes e.g. plant-based milk alternatives, coffee creamer, crème fraiche; [‡]Other margarine, fats & oils includes e.g. vegetable suet, palm

166 oil, chicken fat; ^{‡‡}Other potatoes, potato salads & dishes includes potato curry, mash potato with butter and cream.

167 In the category of foods that were UPF and HFSS, the most common main groups reflected 168 miscellaneous foods (e.g., condiments and soups), bakery products (e.g. biscuits and 'buns, cakes, 169 pastries & fruit pies'), reduced fat spread, and 'soft drinks, not low calorie'. When considering by food 170 energy and food weight, products high in sugar (biscuits, 'buns, cakes, pastries & fruit pies', chocolate 171 confectionery), 'crisps and savoury snacks', and 'pasta, rice & other cereals' made substantial

- 172 contributions to the category that was both UPF and HFSS.
- 173 The most common main food group contributing to the category of foods that were neither HFSS nor
- 174 UPF was tea, coffee and water. Cooked and raw vegetables, semi-skimmed milk and fruit were also
- were also prominent. Similar findings were seen for food weight. By food energy, the food groups
- making the largest contribution to this category included pasta, rice & other cereals, fruit, semi-
- 177 skimmed milk, and chicken and turkey dishes.

178 Person-level analysis

- 179 The results of the person-level analysis are summarised in Table 4. Additionally adjusting for non-
- 180 response bias led to small changes compared to the estimates in Figure 1. At person-level, 55.1% of
- 181 UPF foods are also HFSS (c.f. 55.6% at food level), 53.8% of UPF energy is also HFSS (c.f. 58.7%), and
- 182 40.0% of UPF food weight is also HFSS (c.f. 38.3%).

183 Table 4. The percentage of per capita daily foods, food energy in kcal, and food weight in grams

- consumed that was derived from foods that are high in fat, salt or sugar, ultra-processed foods,
 both and neither; UK National Diet and Nutrition Survey 2008/9-2018/19
- 186

	All foods consumed; % (SD)	Food energy (kcal); % (SD)	Food weight (g); % (SD)
Neither UPFs nor HFSS	52.9 (14.4)	31.7 (13.6)	66.8 (18.6)
All UPF	33.6 (15.6)	54.9 (15.7)	29.7 (18.3)
UPF only	15.0 (8.9)	24.6 (10.5)	18.2 (13.4)
All HFSS	32.0 (10.6)	43.7 (13.2)	15.0 (10.9)
HFSS only	13.5 (8.1)	13.4 (8.7)	3.5 (4.3)
HFSS and UPFs	18.5 (10.0)	30.3 (13.8)	11.5 (10.0)
UPF that is also HFSS	55.1 (15.5)	53.9 (17.3)	40.0 (19.0)

187

188 *Note:* HFSS = high in fat, salt or sugar (using the 2004/05 Nutrient Profiling Model); UPF = ultra-processed food;

- $189 \qquad {\rm data\ are\ weighted\ to\ correct\ for\ non-random\ non-response\ of\ households}.$
- 190
- 191 Analysis using the 2018 version of the NPM led to broadly similar results, but overall a slightly smaller
- 192 proportion of UPF was categorised as HFSS compared to the 2004/05 NPM (Table S2).

193 **DISCUSSION**

194 Summary of main findings

- This is the first study to assess the extent to which foods consumed in the UK are HFSS, UPFs, both or neither. Using 11 years of data from NDNS we found that around one third of all foods consumed (33.4%) were HFSS, one third (36.2%) were UPF, one fifth were both (20.1%) and half (50.5%) neither. We found that 55.6% of UPFs, 58.7% of food energy and 38.3% of food weight consumed were also HFSS. This means that policies focused on HFSS reduction also target between one third and just over a half of UPFs.
- 201 When considering all foods consumed, common foods that were UPF but not HFSS (and so would not
- 202 be included in HFSS-focused policies) included low calorie soft drinks, manufactured breads, and high
- 203 fibre breakfast cereals. Food groups making substantial contributions to all foods consumed in the
- 204 category HFSS and UPF were miscellaneous (e.g., condiments and soups), biscuits, reduced fat spread,
- 205 and 'soft drinks, not low calorie'.

206 Strengths and limitations

To the best of our knowledge, this is the first study to classify all foods consumed in the UK as both HFSS and UPF and compare these classifications. We used food-level dietary data, with 3 or 4 food diary days per person from the NDNS (20, 21). Food diaries reduce the recall bias common in food frequency questionnaires and 24-hour recalls (21, 22). We used three metrics of food consumption – total number of foods, food energy, and food weight. Analysing food energy recognises that different foods are consumed in different portion sizes while assessing food weight identifies, in particular, foods that have a high weight (e.g. drinks) but low energy ingredients (e.g. non-nutritive sweeteners).

Whilst the accuracy of food classification using the Nova system has been criticised as poorly replicable (23), we and others have achieved a high degree of inter-rater reliability (19, 24). Furthermore, by including all foods reported by NDNS participants (excluding alcohol and supplements), we effectively weighted by frequency of consumption. Additional adjustment for non-response bias by including survey weights in individual-level analyses did not change the overall pattern of findings indicating that our findings are likely to be generalisable to the UK population.

220 Despite their strengths, food diaries are self-reported leading to the potential for measurement error.

221 Further, whilst we included 11 years of data, the most recent was from 2019. Given the increasing

number of new foods commercially available (25), we may not have captured recent changes in food

223 consumption. Moreover, combining 11 years of data might have masked important changes over time,

but ensured a large sample size. NDNS did not collect data from new participants in 2020 and subsequently changed dietary data collection method meaning that newer comparable data is not available.

227 There are also some potential limitations with the nutritional measures used in this study. The Nova 228 classification has been criticised for its lack of consideration of nutritional content and quality. With 229 some foods, ultra-processing might yield nutritional and health benefits such as micronutrient 230 fortification, increased affordability, extended shelf life and reduced waste (26). A recent study 231 indicated that while consumption of some UPFs, such as artificially and sugar-sweetened beverages, 232 was associated with an increased risk of cancer and cardiometabolic diseases, no evidence of an 233 association was found for others such as UPF bread and cereals (1). Our second measure of nutritional 234 quality, HFSS has also been criticised as an oversimplified measure of nutritional quality (27).

235 Interpretation and implications

We found that, at best, 58.7% of UPFs consumed over 11 years of the UK NDNS were identified as HFSS. This is a lower degree of overlap than a recent US study which found that 75.4% of UPF were also HFSS (3). These differences may reflect the different food supply, purchase and consumption patterns seen in the UK and USA.

Our findings suggests that current UK policy which focuses on HFSS captures at best just over half of UPFs, but some key UPFs are not included. Amongst all foods consumed, common food groups that were UPF but not HFSS included 'low calorie soft drinks', bread, and miscellaneous products (e.g., condiments and soups).

244 Extending the definition of HFSS to include non-nutritive sweeteners would increase the proportion 245 of UPF captured by HFSS to just over two-thirds. Similar findings were reported by Popkin et al (2024) 246 (3). Whilst some recent UK public health nutrition policies have increased the number of products 247 available that contain non-nutritive sweeteners (28), the World Health Organization has recently 248 noted that these may not be useful for weight loss (29). There may also be other ways that the HFSS 249 algorithm could be adapted to capture more UPFs. However, there remains a lack of consensus among 250 policymakers and scientists on whether the evidence on the health harms of UPF is strong enough to 251 justify regulation (10).

When analysing by food energy, rather than total number of foods the 'other milk & cream' food group was among the top ten food groups contributing to the category of UPF but not HFSS. This includes plant-based milk alternatives. Some plant-based alternatives can be healthier than equivalent meat

255 and dairy products (30) and some plant-based alternatives are UPF. Whilst there is uncertainty about 256 the environmental impacts of UPF as a category (31), reducing consumption of animal-based products 257 at a population level will be essential to achieve net-zero (32). UPF plant-based alternatives could be 258 a stepping stone to less processed plant-based alternatives (33, 34). This highlights the trade-offs 259 common in individual dietary choices as well as policymaking and why simply dichotomising foods into 260 those that should be consumed or avoided is challenging, however that dichotomisation is achieved. 261 Nevertheless, food profiling methods that consider both human and planetary health would allow for 262 a more holistic approach, and regulating foods on sustainability grounds is likely to have health-co-263 benefits (35).

264 We found that many food groups are common contributors to both HFSS only, UPF only, both and neither. For example, regardless of the metric of consumption (foods, food energy or food weight), 265 266 the main food group 'miscellaneous' is in the top ten contributors to many categories. This reflects 267 that not all foods within main food groups necessarily share the same UPF or HFSS status. For example, 268

homemade soups could be HFSS whilst manufactured soups are likely to be UPF and may be HFSS.

269 Much policy in the UK has focused on food reformulation, whereby manufacturers are incentivised to 270 change the composition of foods to ensure they are not HFSS (36). Other policies have focused 271 specifically on reducing sugar and calorie content (37). Whilst there is evidence that some of these 272 policies have prompted reformulation (e.g. removal of sugar from soft drinks) (38) and have been 273 associated with overall reduction in sugar consumption(39) and health benefits (40), reformulated 274 foods which are no longer HFSS often remain UPF (e.g. artificially sweetened soft drinks). An 275 alternative approach using policies that incentivise de-formulation (41) (i.e. removing components 276 and additives commonly used in UPFs, such as non-nutritive sweeteners) could help reduce 277 production and consumption of UPF. Accompanying such policy approaches with those that 278 incentivise whole food consumption could offer important synergies.

279 The difference in results we find by analysing the data by all foods, food energy and food weight reflect 280 differences in the food energy and weight of different foods. For example, low calorie soft drinks make 281 up a substantial proportion of UPF only foods and food weight, but not food energy. This reflects that 282 these products are low in energy but may be consumed in large portions.

283 The 2018 NPM was developed to take account of updated guidance on sugar and fibre consumption 284 (9). Whilst this generated broadly similar results to using the 2004/05 NPM, in most cases a slightly 285 smaller proportion of UPF were categorised as HFSS (e.g. 55.6% of UPF foods consumed were HFSS

using the 2004/05 NPM and 54.4% using the 2018 NPM). These small differences are unlikely to be ofpolicy relevance.

288 Our results reinforce the well-established finding that current UK diets are sub-optimal for health. 289 After adjustment for non-response bias, we found that 54.9% of per capita daily energy was derived 290 from UPFs and 43.7% from HFSS foods. This is comparable to previous findings. For instance, a 291 previous study in the UK based on NDNS data from 2008-14 found that 56.8% of energy was derived 292 from UPFs (42). Similarly, in Canada 47.7% of energy was derived from UPFs (43), and in the US 57.5% 293 was derived from UPFs (44). Studies from other countries find a smaller proportion of energy 294 attributable to UPFs. For example, a study from Italy found that 17.3% of energy was derived from 295 UPFs (45). Current UK dietary guidance advises citizens to eat HFSS foods "less often and in small 296 amounts" (46).

297 CONCLUSION

In this analysis of 11 years of data from the UK NDNS, we found that the NPM, used to identify foods that are HFSS for regulatory purposes identifies, at best, 58.7% of UPFs. If UK policymakers decide that regulation of UPFs is necessary, additional action will be required to extend current policy. This could involve extending the current NPM to include ingredients, such as non-nutritive sweeteners, that are common in foods that are UPF but not currently identified as HFSS. Further work is required to confirm that UPF consumption is causally associated with health harms and to determine the environmental impacts of UPF.

305 ACKNOWLEDGEMENTS

- 306 We are grateful to Birdem Amoutzopoulos and David Collins for reviewing the code for categorising
- 307 foods high in fat, salt or sugar.

308 FUNDING

309 This research has been funded by the Economic and Social Research Council (ESRC).

COMPETING INTERESTS

311 There are no competing interests.

312 **RESEARCH ETHICS APPROVAL**

313 No ethics approval was required.

314 **REFERENCES**

- Cordova R, Viallon V, Fontvieille E, Peruchet-Noray L, Jansana A, Wagner K-H, et al.
 Consumption of ultra-processed foods and risk of multimorbidity of cancer and cardiometabolic
 diseases: a multinational cohort study. The Lancet regional health Europe. 2023:100771.
- Mytton OT, Forouhi NG, Scarborough P, Lentjes M, Luben R, Rayner M, et al. Association
 between intake of less-healthy foods defined by the United Kingdom's nutrient profile model and
 cardiovascular disease: A population-based cohort study. PLOS Medicine. 2018;15(1):e1002484.
- 321 3. Popkin BM, Miles DR, Taillie LS, Dunford EK. A policy approach to identifying food and
 beverage products that are ultra-processed and high in added salt, sugar and saturated fat in the
 United States: a cross-sectional analysis of packaged foods. The Lancet Regional Health Americas.
 2024;32:100713.
- Marino M, Puppo F, Del Bo' C, Vinelli V, Riso P, Porrini M, Martini D. A systematic review of
 worldwide consumption of ultra-processed foods: Findings and criticisms. Nutrients.
 2021;13(8):2778.
- 328 5. Rayner M, Scarborough P, Lobstein T. The UK Ofcom Nutrient Profiling Model: Defining 329 'healthy'and 'unhealthy'foods and drinks for TV advertising to children. London: OfCom. 2009.
- 330 6. Davies S. Time to solve childhood obesity. Department of Health Social Care. 2019.
- 331 7. Department of Health. Nutrient profiling technical guidance. 2011.
- 8. Public Health England. UK Nutrient Profiling Model 2018 review. London, UK: Public HealthBingland; 2018.
- 334 9. Scientific Advisory Committee on Nutrition. Carbohydrates and health. London; 2015.
- 335 10. Office for Health Improvement and Disparities. The Scientific Advisory Committee on
 336 Nutrition (SACN) position statement on processed foods and health. UK, 2023.
- Monteiro CA, Cannon, G., Lawrence, M., Costa Louzada, M.L. and Pereira Machado, P. Ultra processed foods, diet quality, and health using the NOVA classification system. Rome, FAO. 2019.
- Lane MM, Gamage E, Du S, Ashtree DN, McGuinness AJ, Gauci S, et al. Ultra-processed food
 exposure and adverse health outcomes: umbrella review of epidemiological meta-analyses. Bmj.
 2024;384:e077310.
- 342 13. Vandevijvere S, Jaacks LM, Monteiro CA, Moubarac JC, Girling-Butcher M, Lee AC, et al.
 343 Global trends in ultraprocessed food and drink product sales and their association with adult body
 344 mass index trajectories. Obesity Reviews. 2019;20:10-9.
- 345 14. Vandevijvere S, Jaacks LM, Monteiro CA, Moubarac JC, Girling-Butcher M, Lee AC, et al.
 346 Global trends in ultraprocessed food and drink product sales and their association with adult body
 347 mass index trajectories. Obes Rev. 2019;20 Suppl 2:10-9.
- Koios D, Machado P, Lacy-Nichols J. Representations of Ultra-Processed Foods: A Global
 Analysis of How Dietary Guidelines Refer to Levels of Food Processing. International journal of health
 policy and management. 2022;11(11):2588-99.

Bates B, Lennox A, Prentice A, Bates C, Page P, Nicholson S, Swan G. National Diet and
Nutrition Survey: results from years 1, 2, 3 and 4 (combined) of the Rolling Programme (2008/20092011/2012): a survey carried out on behalf of Public Health England and the Food Standards Agency.
2014.

17. Chavez-Ugalde Y, De Vocht F, Jago R, Adams J, Ong KK, Forouhi N, et al. Manuscript title:
Ultra-processed food consumption in UK adolescents: distribution, trends, and sociodemographic
correlates using the National Diet and Nutrition Survey 2008/09 to 2018/19. medRxiv.
2023:2023.06.05.23290977.

18. Chavez-Ugalde IY, de Vocht F, Jago R, Adams J, Ong KK, Forouhi NG, et al. Ultra-processed
food consumption in UK adolescents: distribution, trends, and sociodemographic correlates using
the National Diet and Nutrition Survey 2008/09 to 2018/19. European Journal of Nutrition. 2024:115.

363 19. Colombet Z., O'Flaherty M., Chavez-Ugalde Y. NOVA classification of the National Diet and
 364 Nutrition Survey, waves 1 to 11 (2008/09 to 2018/19): GitHub; 2023 [Available from:
 365 <u>https://github.com/zoecolombet/NOVA_NDNS_code</u>.

Black AE, Prentice AM, Goldberg GR, Jebb SA, Bingham SA, Livingstone MB, Coward WA.
 Measurements of total energy expenditure provide insights into the validity of dietary
 measurements of energy intake. Journal of the American Dietetic Association. 1993;93(5):572-9.

Bingham SA. Limitations of the Various Methods for Collecting Dietary Intake Data. Annals of
 Nutrition and Metabolism. 1991;35(3):117-27.

Park Y, Dodd KW, Kipnis V, Thompson FE, Potischman N, Schoeller DA, et al. Comparison of
self-reported dietary intakes from the Automated Self-Administered 24-h recall, 4-d food records,
and food-frequency questionnaires against recovery biomarkers. The American journal of clinical
nutrition. 2018;107(1):80-93.

375 23. Gibney MJ. Ultra-processed foods: Definitions and policy issues. Current Developments in376 Nutrition. 2019;3(2).

377 24. Steele EM, O'Connor LE, Juul F, Khandpur N, Galastri Baraldi L, Monteiro CA, et al. Identifying
378 and Estimating Ultraprocessed Food Intake in the US NHANES According to the Nova Classification
379 System of Food Processing. The Journal of Nutrition. 2023;153(1):225-41.

Boukid F, Sogari G, Rosell C. Edible insects as foods: mapping scientific publications and
product launches in the global market (1996-2021). Journal of Insects as Food and Feed.
2023;9(3):353-68.

383 26. McClements DJ. Designing healthier and more sustainable ultraprocessed foods.
384 Comprehensive Reviews in Food Science and Food Safety. 2024;23(2):e13331.

Volatier J-L, Biltoft-Jensen A, De Henauw S, Gibney MJ, Huybrechts I, McCarthy SN, et al. A
new reference method for the validation of the nutrient profiling schemes using dietary surveys.
European journal of nutrition. 2007;46:29-36.

Luick M, Bandy LK, Harrington R, Vijayan J, Adams J, Cummins S, et al. The impact of the UK
soft drink industry levy on the soft drink marketplace, 2017–2020: An interrupted time series
analysis with comparator series. PloS one. 2024;19(6):e0301890.

391 29. World Health Organization. Use of non-sugar sweeteners: WHO guideline. Geneva: World392 Health Organization; 2023.

30. Crimarco A, Springfield S, Petlura C, Streaty T, Cunanan K, Lee J, et al. A randomized
crossover trial on the effect of plant-based compared with animal-based meat on trimethylamine-Noxide and cardiovascular disease risk factors in generally healthy adults: Study With Appetizing
Plantfood-Meat Eating Alternative Trial (SWAP-MEAT). Am J Clin Nutr. 2020;112(5):1188-99.

397 31. Kesse-Guyot E, Allès B, Brunin J, Fouillet H, Dussiot A, Berthy F, et al. Environmental impacts
398 along the value chain from the consumption of ultra-processed foods. Nature Sustainability.
399 2023;6(2):192-202.

Willett W, Rockström J, Loken B, Springmann M, Lang T, Vermeulen S, et al. Food in the
Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable
food systems. The Lancet. 2019;393(10170):447-92.

33. Northcott T, Lawrence M, Parker C, Baker P. Ecological regulation for healthy and
sustainable food systems: responding to the global rise of ultra-processed foods. Agriculture and
Human Values. 2023.

406 34. Anastasiou K, Baker P, Hadjikakou M, Hendrie G, Lawrence M. A conceptual framework for
407 understanding the environmental impacts of ultra-processed foods and implications for sustainable
408 food systems. Journal of Cleaner Production. 2022:133155.

409 35. Laine JE, Huybrechts I, Gunter MJ, Ferrari P, Weiderpass E, Tsilidis K, et al. Co-benefits from
410 sustainable dietary shifts for population and environmental health: an assessment from a large
411 European cohort study. The Lancet Planetary Health. 2021;5(11):e786-e96.

Gressier M, Swinburn B, Frost G, Segal AB, Sassi F. What is the impact of food reformulation
on individuals' behaviour, nutrient intakes and health status? A systematic review of empirical
evidence. Obesity Reviews. 2021;22(2):e13139.

415 37. Stanner S, Spiro A. Public health rationale for reducing sugar: Strategies and challenges.416 Nutrition Bulletin. 2020;45(3):253-70.

Scarborough P, Adhikari V, Harrington RA, Elhussein A, Briggs A, Rayner M, et al. Impact of
the announcement and implementation of the UK Soft Drinks Industry Levy on sugar content, price,
product size and number of available soft drinks in the UK, 2015-19: A controlled interrupted time
series analysis. PLoS medicine. 2020;17(2):e1003025.

39. Rogers NT, Cummins S, Jones CP, Mytton O, Rayner M, Rutter H, et al. Estimated changes in
free sugar consumption one year after the UK soft drinks industry levy came into force: controlled
interrupted time series analysis of the National Diet and Nutrition Survey (2011–2019). Journal of
epidemiology and community health. 2024:jech-2023-221051.

425 40. Cobiac LJ, Rogers NT, Adams J, Cummins S, Smith R, Mytton O, et al. Impact of the UK soft
426 drinks industry levy on health and health inequalities in children and adolescents in England: An
427 interrupted time series analysis and population health modelling study. Plos Medicine.
428 2024;21(3):e1004371.

429 41. Srour B. Food processing and risk of non-communicable diseases: findings from the
430 NutriNet-Santé cohort: Université Paris-Nord-Paris XIII; 2019.

431 42. Rauber F, Louzada MLdC, Steele EM, Millett C, Monteiro CA, Levy RB. Ultra-processed food
432 consumption and chronic non-communicable diseases-related dietary nutrient profile in the UK
433 (2008–2014). Nutrients. 2018;10(5):587.

434 43. Moubarac J-C, Batal M, Louzada ML, Martinez Steele E, Monteiro CA. Consumption of ultra-435 processed foods predicts diet quality in Canada. Appetite. 2017;108:512-20.

436 44. Martínez Steele E, Popkin BM, Swinburn B, Monteiro CA. The share of ultra-processed foods
437 and the overall nutritional quality of diets in the US: evidence from a nationally representative cross438 sectional study. Population health metrics. 2017;15:1-11.

439 45. Ruggiero E, Esposito S, Costanzo S, Di Castelnuovo A, Cerletti C, Donati MB, et al. Ultra440 processed food consumption and its correlates among Italian children, adolescents and adults from
441 the Italian Nutrition & Health Survey (INHES) cohort study. Public Health Nutrition.
442 2024 24(18):C250 74

442 2021;24(18):6258-71.

443 46. England PH. The Eatwell Guide: helping you eat a healthy, balanced diet. Public Health

444 England London, UK; 2016.