

1 Effect of Midwife-Led Pelvic Floor Muscle Training on Prolapse Symptoms and Quality of Life
2 in Women with Pelvic Organ Prolapse in Ethiopia: A Cluster-Randomized Controlled Trial

3
4 Melese Siyoum (MS)^{1,2*}, Rahel Nardos (RN)³, Biniyam Sirak (BS)⁴, Theresa Spitznagle (TS)^{5,6},
5 Wondwosen Teklesilasie (WT)², Ayalew Astatkie (AA)²

6
7 ¹Department of Midwifery, College of Medicine and Health Sciences, Hawassa University,
8 Hawassa, Ethiopia

9 ²School of Public Health, College of Medicine and Health Sciences, Hawassa University,
10 Hawassa, Ethiopia

11 ³Department of Obstetrics and Gynecology, and Women's Health, University of Minnesota, USA

12 ⁴Addis Ababa Fistula Hospital, Ethiopia

13 ⁵Program in Physical Therapy, Washington University in St Louis, USA

14 ⁶Worldwide Fistula Fund, USA

15

16

17

18 ***Corresponding Author**

19 Melese Siyoum

20 Email: melesesiyoum755@gmail.com

21

22 **Source of funding:** University of Minnesota, worldwide Fistula Fund and Maternal Health

23 **Abstract**

24 **Background:** Pelvic organ prolapse (POP) is a common condition that can significantly impact a
25 woman's quality of life. Pelvic floor muscle training (PFMT) is recommended as a first-line
26 conservative treatment for prolapse, but evidence on its effectiveness from low-resource settings
27 is limited. This study aimed to assess the effect of midwife-led pelvic floor muscle training on
28 prolapse symptoms and health-related quality of life (HRQoL) among women with mild to
29 moderate pelvic organ prolapse in Ethiopia.

30 **Methods and Findings:** A community-based, parallel-groups, two-arm cluster-randomized
31 controlled trial was conducted in Dale and Wonsho districts of Sidama Region, Ethiopia. Women
32 with symptomatic POP stages I-III were randomized by cluster to receive either midwife-led
33 PFMT plus lifestyle counseling (intervention group) or lifestyle counseling alone (control
34 group). The participants and counsellors knew what the women were receiving but they were not
35 aware of the other group. The outcome assessors, who collected data at the end of intervention,
36 were blinded to the participants' treatment allocation. The primary outcomes were changes in
37 prolapse symptom score (POP-SS) and prolapse quality of life (P-QoL). Mixed-effects
38 generalized linear model was used to determine the effect of PFMT on prolapse symptoms and
39 P-QoL at 99% confidence level. Adjusted β coefficients were used as effect measures. The level
40 of significance was adjusted for multiple comparisons.

41 A total of 187 women were randomized to intervention (n=86) from 4 clusters and control
42 (n=101) arms from another four clusters. At sixth months, the intervention group showed
43 significantly greater improvements both in prolapse symptoms and P-QoL. The mean change
44 difference in POP-SS was -4.11 (99% CI [-5.38, -2.83]; $p < 0.001$). Similarly, the mean change
45 difference was: -8.86 (99% CI [-13.84, -3.89]; $p < 0.001$) in physical domain of P-QoL; -11.18
46 (99% CI [-15.03, -7.32]; $p < 0.001$) in psychological domain of P-QoL, and -9.01 (99% CI [-
47 10.49, -5.54]; $p < 0.001$) in personal relationship domain of P-QoL. A significantly higher
48 proportion (83.72%) of women in the intervention group perceived their condition as "better"
49 after the intervention as compared to 41.58% in the control group. Women with earlier stages of
50 prolapse (stage I and II) experienced higher benefits compared to those in stage III.

51 **Conclusions:** A midwife-led PFMT combined with lifestyle counseling significantly improves
52 prolapse symptoms and quality of life in mild to moderate POP. This strategy can be integrated

53 into the existing maternal and reproductive health programs to address POP in low-income
54 settings where access to trained specialist is limited.

55 **Trial registration:** Pan African Clinical Trial Registry (PACTR202302505126575).

56 **Keywords:** Pelvic organ prolapse, pelvic floor muscle training, cluster-randomized trial, quality
57 of life, Ethiopia.

58

59 **Author Summary**

60 **Why Was This Study Done?**

- 61 • Pelvic organ prolapse (POP) is a prevalent condition that can severely impact women's
62 quality of life, particularly in low-resource settings where access to specialized care is
63 limited.
- 64 • Prior to this research, there was limited evidence regarding the effectiveness of pelvic
65 floor muscle training (PFMT) as a conservative treatment option for POP in such
66 contexts, highlighting the need for effective interventions.

67 **What Did the Researchers Do and Find?**

- 68 • The study utilized a cluster-randomized controlled trial design involving eight clusters
69 (four clusters in each arm) and a total of 187 women, all drawn from these clusters, with
70 mild to moderate pelvic organ prolapse (POP) in Sidama region, Ethiopia. The
71 intervention group received midwife-led pelvic floor muscle training (PFMT) combined
72 with lifestyle counseling, while the other group received only lifestyle counseling.
- 73 • The key findings from the study showed that women who participated in the midwife-led
74 pelvic floor muscle training (PFMT) combined with lifestyle counseling experienced a
75 significant reduction in their pelvic organ prolapse (POP) symptoms. Specifically, there
76 was an average decrease of 4.11 points in their symptom scores, which indicates a
77 notable improvement in how they felt. Additionally, the women reported substantial
78 enhancements in various aspects of their quality of life, meaning they felt better overall
79 and were able to engage more fully in daily activities

80 **What Do These Findings Mean?**

- 81 • The research demonstrates that midwife-led PFMT can serve as an effective first-line
82 treatment for POP, improving symptoms and quality of life in low-resource settings.
- 83 • However, limitations include reliance on self-reported data, and a relatively short follow-
84 up period may limit generalizability of the findings.

85

86 **Introduction**

87 Symptomatic pelvic organ prolapse (POP) is the descent of one or more of the uterus, cervix,
88 anterior or posterior vaginal wall, or the vaginal apex from its normal position. This descent can
89 be caused by deficient pelvic fascia, or weakness in the ligaments or muscles that support the
90 pelvic organs [1]. These structures can be affected by several risk factors including aging,
91 vaginal childbirth, chronic constipation, hereditary connective tissue weakness, persistent
92 elevated intra-abdominal pressure (IAP), menopause, and heavy lifting [2].

93 Pelvic organ prolapse (POP) is a prevalent condition affecting many women globally, with
94 estimated prevalence rates varying widely from 1% to 65% [3, 4]. This broad range can be
95 attributed to the challenges in accurately estimating prevalence, largely due to the differing
96 methodologies employed in various studies. For instance, researchers may utilize a variety of
97 symptom questionnaires and clinical examination techniques, leading to inconsistencies in the
98 reported prevalence [3, 5].

99 Women with symptomatic POP present with a range of prolapse-related symptoms that are
100 associated with the urinary, sexual, and anorectal symptoms [6]. It is evidenced that depressive
101 symptoms and sexual function disorders are frequently associated with POP [7, 8]. In addition to
102 disrupting daily functioning, these multifaceted symptoms of POP have significant negative
103 impact on patients overall quality of life, body image and sexual function [8]. The severity of
104 symptoms is directly related to the degree of prolapse. As the prolapse descends closer to or
105 beyond the hymen, the symptoms become more frequent and bothersome [9, 10].

106 The current POP treatment options are either surgery or conservative managements approaches
107 [11]. The conservative management options include pelvic floor muscle training (PFMT),
108 pessary, life style modification, or watchful waiting, which are recommended as first-line options
109 if the prolapse is not the indication for corrective surgery [2]. Conservative management has the
110 advantage of causing fewer adverse effects and complications [12]. The existing evidence shows
111 that PFMT has positive effect on prolapse symptom and severity [13]. Due to its effectiveness
112 and safety, PFMT is indicated as a first-line treatment for women with mild to moderate POP
113 [13]. PFMT improves prolapse symptoms by improving pelvic muscle strength, endurance,
114 power, relaxation and combination of these parameters [1].

115 Awareness of POP and conservative treatment option is very poor among women [14] and
116 healthcare professionals [15] even in high-income countries. To enhance the accessibility and
117 variety of service for women, it is essential to understand various models of service delivery
118 [16]. In low- and middle-income countries, evidence about the feasibility and effect of PFMT on
119 prolapse symptoms and health-related quality of life (HRQoL) is limited.

120 In Ethiopia, there is scarcity of evidence on the effectiveness of PFMT, and women often seek
121 healthcare only at advanced stages of prolapse [17] when impact of PFMT may be limited.
122 Additionally, the number of trained specialists such as urogynecologists or physical therapist are
123 limited in Ethiopia. There are only 450 gynecologists registered in the Ethiopian Society of
124 Obstetricians and Gynecologists out of which only 15 are urogynecologists and 13 are
125 urogynecology fellows in 2021 [18]. In areas of the world with developed physical therapy
126 training (PT) programs, (PT) management of pelvic organ prolapse is the gold standard for
127 administering the service [19]. However, it is considered a sub-specialty within the profession
128 and currently the Ethiopian Ministry of Health does not support specialty practice for the
129 profession of Physical Therapy [20]. Fortunately, midwives, who are generally more accessible
130 than specialists [21], are capable of learning how to deliver PFMT services and are well-
131 positioned to provide this support to women in their communities. To address these challenges
132 and improve access to the conservative management options, research into different models of
133 PFMT service delivery is critical particularly in low resource settings.

134 The aim of this study was to assess the effect of PFMT provided by trained midwives on
135 prolapse symptoms and HRQoL among women with mild to moderate POP (stage I-III) in
136 Sidama region, Ethiopia. Our hypothesis was that providing midwife-led PFMT approximately
137 monthly alongside three times per-day home-based exercise by participants for six months will
138 improve prolapse symptoms and prolapse related quality of life.

139

140

141

142

143 **Methodology**

144 **The Consolidated Standards of Reporting Trials (CONSORT) Checklist**

145
146 The CONSORT (Consolidated Standards of Reporting Trials) 2010 statement [22] was used to
147 guide the reporting of this cRCT. The CONSORT checklist consists of a 25-item checklist that
148 focuses on the key elements for reporting the design, conduct, analysis, and interpretation of
149 RCTs. An extension to the CONSORT statement for reporting cluster randomized trials (cRCTs)
150 [23] was also consulted. We have addressed all applicable items from the CONSORT checklist in
151 the reporting of this cRCT (Supplementary File 1).

152

153 **Study design and participants**

154 A community-based, parallel-groups, two-arm, cluster-randomized controlled trial (cRCT) was
155 conducted among women with symptomatic POP stages I-III in D-W HDSS. First, a baseline
156 survey was conducted to determine the burden and risk factors of POP and to identify eligible
157 cases [24]. Women aged over 18 years with at least one symptom of POP, confirmed as stages I-
158 III using the standard POP-Q measure [25], were included in the study. The identified cases were
159 then randomized by community clusters into intervention and control arms. Cluster
160 randomization was preferred to minimize spillover of the intervention to the control arm and due
161 to the feasibility of providing the intervention at group level (nearby health facility). The clusters
162 were *kebeles* which are the smallest administrative units in Ethiopia, similar to neighborhoods or
163 wards. They have a wide range of population sizes, from 500 to several thousand residents [26].
164 The surface area of *kebeles* varies depending on their location and population density, with those
165 in densely populated urban areas being smaller compared to rural *kebeles*. Each *kebele* is further
166 divided into smaller units called *gots* or *villages*. *Kebele* administrations are responsible for
167 providing basic services, such as health, education, and infrastructure, to their residents. They
168 also play a crucial role in population-based surveys and research studies in Ethiopia, often
169 serving as the primary sampling units [27, 28]. *Kebeles* are organized into larger
170 administrative divisions known as *woredas* (Districts).

171 Sample size calculations were performed using Open-Epi version 3.1, based on the following
172 assumptions: a 99% confidence level, 80% power, an exposed-to-unexposed ratio of 1, a POP-SS
173 mean difference of 3.16 with a standard deviation of 4.78 in the intervention group, and a mean
174 difference of 0.12 with a standard deviation of 3.86 in the control group over six months [15].
175 This calculation yielded a required sample size of 96, adjusted for an anticipated 16% refusal and
176 loss to follow-up [15], resulting in an individual-based total sample size of 111.34. This sample
177 size was utilized for all outcomes since we were unable to determine the effect of pelvic floor
178 muscle training (PFMT) on quality of life as measured by the Prolapse Quality of Life
179 Questionnaire. The decision to adjust the confidence level to 99% was based on the adjusted
180 level of significance to 0.0125 than the usual P-value of 0.05 to account for multiple
181 comparisons across various outcomes (POP-SS and three distinct domains of P-QoL).

182 To determine the minimum number of clusters required, the individual-based sample size was
183 multiplied by the intra-cluster correlation coefficient (ICC). Since there was no reported ICC
184 from earlier studies, we used an ICC of 0.03, which was suggested as estimated fairly typical
185 value of the Intra-Cluster Correlation (ICC) [29, 30]. This calculation indicated a need for 1.67
186 clusters per arm. To ensure adequate power [31, 32], a total of eight clusters were selected
187 randomly from the 12 available clusters in the D-W HDSS.

188 After adjusting for cluster effects with a variance inflation factor (VIF) of 1.39, the total effective
189 sample size increased to 155, resulting in an expected average of 19.4 participants per cluster.
190 However, during enrollment, a slightly varying number of eligible cases was identified in each
191 cluster. Excluding additional participants was deemed unethical, and alternative treatment
192 options could not be provided due to the cRCT design. Furthermore, it was believed that
193 including all eligible cases would be manageable and enhance the study's power [29, 30].
194 Consequently, all eligible cases were included, raising the total sample size from 155 to 187
195 participants. The exclusion criteria were: women who were diagnosed to have symptomatic POP
196 stage I-III but planned to leave the study area within six months (study period), severely ill
197 women, women with psychiatry disorders, women with delivered myoma, or women who want
198 to have prolapse surgery within six months.

199

200 **Randomization and Masking**

201 The study site, D-W HDSS, comprises eight kebeles from Dale district and four kebeles from
202 Wonsho district. These were stratified into two strata based on their geographical location (Dale
203 and Wonsho districts). Stratifying on location reduces stratum variability and helps to ensure that
204 they are balanced across study arms [31, 32]. To improve similarity across treatment and control
205 groups, a balanced number of clusters from each stratum were assigned to each arm. First all
206 eligible clusters in each district were listed and assigned a unique identifier number. Then, a
207 lottery method was used to select two of four clusters from Wonsho district and randomly
208 assigned to each arm. Similarly, six of eight clusters from Dale districts were selected and three
209 clusters were assigned to each arm randomly by lottery method. Then, 102 households were
210 selected from each cluster using computer-generated random numbers (based on the list of their
211 house numbers) to identify eligible women. One woman (usually head of the household) was
212 recruited for the study.

213 To reduce the risk of spillover-effect of the intervention, the interventional and control clusters
214 were separated by a buffer zone [31], and when it is impossible due to the limited number of
215 clusters in the D-W HDSS, adjacent clusters were assigned to the same arm. Finally, all eligible
216 study participants from each cluster identified during the baseline survey were included in the
217 study (Figure 1).

218 *Figure 1: Randomization and Sampling procedure used to assess the effect of pelvic floor muscle*
219 *training on prolapse symptoms and quality of life in Sidama Region, Ethiopia, August 2024.*

220
221 The enrollment period was open for eight months from 13 March to 28 October 2023. Each
222 participant was followed for six months after enrollment. The last participants were enrolled on
223 October 28, 2023 so the final follow-up visit was on April 27, 2024. Allocation concealment was
224 conducted by staff of D-W HDSS at Hawassa University. To ensure unbiased allocation, the
225 lottery method was performed by an independent researcher who was not involved in the
226 recruitment, intervention, or outcome assessment processes. Two staff members of Yirgalem
227 Hamline Fistula center, who performed pelvic examination to determine POP stage, recruited
228 eligible participants for intervention.

229 The participants in the intervention group were aware that they were receiving the PFMT
230 program but they were not aware of the control group. Similarly, the control group knew that
231 they are receiving life style counselling but they are not aware of the intervention group. The
232 same is true for the healthcare providers delivering the counselling. Moreover, the outcome
233 assessors, who collected data at the end of intervention, were blinded to the participants’
234 treatment allocation.

235

236 **Procedure**

237 In Ethiopia, each kebele/cluster is served by its own health post, staffed by health extension
238 workers. Additionally, groups of five or more kebeles are supported by a single health center,
239 staffed by formally trained healthcare professionals, including midwives. For this study, the
240 intervention was implemented at eight distinct health centers. The intervention was implemented
241 at health centers due to their capacity to deliver the training and their geographical accessibility
242 to participants, especially when compared to hospitals.. In this study area, all health centers have
243 at least three degree-holding midwives responsible for providing antenatal, delivery, and
244 postnatal care. Each health center is equipped with various rooms, including delivery rooms,
245 cervical cancer screening rooms, and outpatient rooms, which are suitable for conducting the
246 Pelvic Organ Prolapse Quantification (POP-Q) [25] assessments and ensuring confidentiality
247 during individual counseling sessions.

248 From each selected health facility, two female midwives were trained either on pelvic floor
249 muscle training and how to counsel patients, or only on how to counsel for life style modification
250 based on their pre-allocated health facility. The training that included education on theory,
251 simulation, and video aids was given by a fellowship trained urogynaecologist. Then, the trained
252 midwives administered the intervention to the women assigned in the intervention arms.

253 Women who were assigned to intervention received pelvic floor muscle training and prolapse
254 lifestyle advice focusing on risk factors for POP such as constipation, chronic coughing, heavy
255 lifting, and other medical conditions. The control group received only lifestyle modification. In
256 the interventional group, the pelvic examiners link eligible women to the assigned midwife with
257 the POP-Q findings. Then, at the initial visit, women were informed by midwives about the stage

258 of POP they had and type of prolapse (anterior, posterior, or central) which was explained in a
259 private room. Then the advantage of pelvic floor muscle exercise, and how it works, the detail of
260 the exercise and follow-up schedule were explained in detail at group level in a private room for
261 40-60 minutes.

262
263 They were trained to sit or lie comfortably, train pelvic muscles with 10 times 10-second
264 maximum holds, and 10 fast contractions [15, 33]. They were instructed to repeat this at least
265 three times per day (morning, afternoon, and night) [15, 34]. Women were taught how to
266 precontract the pelvic floor muscles against increases in intra-abdominal pressure (Knack
267 exercise). They were trained to exercise 10 long squeezes and lift
268 contractions, each held for 10 seconds, allowing time between the squeeze to relax and lower the
269 muscle for up to 5 seconds. The long squeezes were followed by 10 short squeezes. Women were
270 advised to try to feel the releasing and relaxing of the squeeze after 10 short squeezes, and
271 finally, to do a squeeze with a light cough for 10 times. The long squeezes were for 10 seconds
272 each and the 10 quick squeezes for 1-2 seconds each.

273 To help participants correctly identify and contract their pelvic floor muscles, the participants in
274 the intervention group received pelvic floor muscle training with internal digital biofeedback
275 administered by the trained midwives. During the individual training sessions, midwives used
276 various verbal cues and analogies. For example, participants were instructed to “imagine you are
277 trying to stop the flow of urine or hold back gas. Squeeze as though you are trying to slow down
278 or stop your urine mid-stream” [35]. This relevant analogy helped participants visualize the
279 specific action they needed to perform. Throughout the exercises, midwives provided descriptive
280 feedback to guide participants. They would say things like, “As you squeeze, focus on lifting the
281 muscles around your vagina and anus upwards. You should feel a tightening and lifting
282 sensation” [33, 35, 36]. This encouraged participants to be aware of the specific muscles being
283 engaged during the contractions. We translated the Lifestyle Advice Leaflet from the POPPY
284 trial [15] into Sidaamu Afoo, the local language, to enhance understanding and effectiveness
285 during counseling sessions.

286 They were appointed to attend five visits (at weeks 0, 2, 6, 10, and 15) at a nearby health facility.
287 The appointment frequency was adapted from the United Kingdom National Health Service

288 (NHS) [37]. Initial appointments were scheduled closely together to reinforce correct exercise
289 techniques and ensure participants fully understood all the advice provided. Subsequently,
290 appointments were spaced further apart to promote independent home exercise. A 15-weeks
291 intensive training was recommended to get adequate muscle strength [38]. The change in
292 prolapse symptoms and health related quality of life were analyzed at six months.

293 A life style counselling was given for both arms focusing on the following points [15, 36].

294 ***Bowel Health:*** To avoid constipation, which increases strain on pelvic floor muscles and can
295 exacerbate prolapse symptoms, it's important to consume a balanced diet rich in fruits,
296 vegetables, and adequate fiber. Aim for 1.5 to 2 liters of water daily to maintain bowel regularity.
297 When using the toilet, sit fully rather than hovering, as this promotes muscle relaxation. Position
298 yourself comfortably with your feet apart and hands resting on your thighs or supported. Keep
299 your abdominal muscles relaxed; if you need to bear down, take a deep breath to ease tension.
300 Light pressure on the vaginal bulge towards the back passage may assist with complete bowel
301 emptying.

302 ***Bladder Health:*** It's crucial to urinate only when the bladder signals the need, rather than going
303 "just in case." Sitting fully on the toilet aids in muscle relaxation, and leaning forward slightly
304 after urination can help ensure complete bladder emptying. Avoid straining, as this can worsen
305 prolapse symptoms. Maintaining the recommended fluid intake of 1.5 to 2 liters per day is
306 essential; reducing fluids does not necessarily decrease urinary frequency and may lead to
307 bladder issues.

308 ***Weight Management:*** Maintaining a healthy weight is vital, as excess weight can place
309 additional pressure on the pelvic floor. Weight loss may improve prolapse symptoms.

310 ***Reducing Intra-abdominal Pressure:*** For those with a chronic cough, follow prescribed
311 treatments and stay hydrated to prevent strain from coughing. Proper lifting techniques are also
312 important; adopt good habits even for lighter items, tightening pelvic floor muscles before
313 lifting. When lifting heavy loads, divide the weight and lift from low positions to minimize
314 strain.

315 ***Exercise and Physical Activity:*** It is advisable to avoid activities that worsen symptoms, such as
316 prolonged standing. Try to alternate standing tasks with sitting breaks. Additionally, high-impact

317 exercises, like jumping, jogging, or aerobics that involve both feet leaving the ground
318 simultaneously, should be avoided. Instead, opt for low-impact activities like modified aerobics,
319 cycling, using an elliptical trainer, or brisk walking.

320

321 **Outcomes**

322 Our primary outcomes were pelvic organ prolapse symptoms and health-related quality of life
323 (HRQoL). The POP-SS was measured by the *Sidaamu Afoo* version of the pelvic organ prolapse
324 symptom score (POP-SS) questionnaire [39]. The psychometric properties of the questionnaire
325 have been evaluated in the same setting and found to have acceptable reliability and validity. It is
326 a patient reported-outcome measure that contain seven items with five Likert-scale ranging from
327 0 to 4, with a possible total score of 0 to 28 (the higher score representing worse symptom) [40].
328 Health-related quality of life was measured by using the prolapse quality of life (P-QoL)
329 questionnaire translated from the Amharic version [41]. While the original version has 20 items
330 with nine domains [42], in a previous work designed to adapt the P-QoL locally (*to Amharic*
331 *version*), the dimensions have been condensed into three: physical function (including General
332 health perception (GHP), prolapse impact (PI), role (RL), physical (PL) and social limitation
333 (SL), and severity measures (SM)); psychological function (including emotional disturbances
334 (EMO) and sleep/energy disturbances (SE)); and personal relationships ((PR). Each domain is
335 related to a particular aspect of QoL, and scores in each domain range from 0 to 100, the higher
336 score indicating poor quality of life. This tool has previously been validated in a similar context
337 (northern Ethiopia).

338 The secondary outcome, women's perceived change In prolapse, was evaluated using the patient
339 global impression of change index (PGI-I), a seven-point Likert scale (1 = very much better to 7
340 = very much worse) [43]. In this study, patients were classified as improved if they scored 1, 2,
341 or 3 on the PGI-I scale, no change if they scored 4 and worse if they scored 5, 6 or 7.

342 Intervention adherence was measured in terms of attendance at appointments [15] and the
343 amount of exercise women reported at follow-up time. Women's adherence to training protocol
344 is considered adequate when they complete at least 80% of the training sessions given by
345 midwives [44].

346 The details of the baseline data collection procedures and tools have been published elsewhere
347 [24]. All tools were translated from English and Amharic into the local language (*Sidaamu Afoo*).

348 The baseline data about sociodemographic characteristics and prolapse symptoms were collected
349 through face-to-face interview with women at their home by eight first degree nurses using
350 electronic data collection forms prepared using Kobo tool box. The women were subsequently
351 invited for pelvic examination in a nearby health center to confirm their prolapse status and
352 degree of prolapse by using the POP-Q [25]. The POP-Q was performed by staff members of
353 Yirgalem Hamlin Fistula Center who have special training on Fistula and POP identification.
354 Then women with symptomatic POP stages I-III were interviewed how much the prolapse had
355 affected their quality of life using P-QoL tool. The detail POP-Q procedures and data quality
356 assurance method was published elsewhere [24].

357 At three months and six months of intervention, four first degree nurses went to the house of the
358 study participants and collected the data through face-to-face interviews. The data collection tool
359 included the POP-SS and P-QoL questionnaires used at baseline, PGI-I change and questions
360 used to assess their adherence to training protocol [44].

361

362 **Statistical analysis**

363 All data analyses accounted for the complex sampling design, which included stratification by
364 district (Dale and Wonsho), clustering by kebele, and weighting for unequal probability of
365 selection. The weighting variable was the product of cluster and household selection weights,
366 normalized by dividing each unit's survey weight by the unweighted average of the survey
367 weights. Categorical variables were summarized as frequencies and percentages, while
368 continuous variables were summarized using means and standard deviations or medians and
369 interquartile ranges.

370 The analysis was based on the intention-to-treat principle. In practice, all participants adhered to
371 their allocated treatment and completed the follow-up as scheduled. Although the intervention
372 was randomly assigned at the cluster level, outcomes were evaluated at the individual level. An
373 Individual-level analysis was performed using a mixed-effects generalized linear model

374 (MEGLM), incorporating random effects to account for clustering by health centers and survey
375 weights to adjust for unequal selection probabilities.

376 To complement the primary analysis, a cluster-level analysis was conducted following Hayes and
377 Moulton's framework. Cluster-level means were calculated for key outcomes (mean change in
378 POP-SS, physical domain, psychological domain and personal relationship) and analyzed using
379 linear regression to compare intervention and control clusters (Supplementary file 5).

380 To compare the primary outcomes at six months, we fitted a multilevel mixed-effects generalized
381 linear model (MEGLM) to the change from baseline in POP-SS and each domain of HRQoL at
382 three and six months. The model included fixed effects for treatment group, time, and other
383 relevant covariates (stage of prolapse, duration of prolapse symptoms, residence, parity, age,
384 level of education, history of prolonged labor, wealth index, constipation, and chronic cough), as
385 well as a random intercept for patients within centers and a random slope for time within
386 patients. The relevance of the covariates was determined based on prior research, clinical aspect,
387 and their potential impact on the intervention and outcomes (prolapse symptoms and quality of
388 life). MEGLM was chosen for its flexibility and robustness needed to accurately model the
389 complex and varied data structures often found in cRCT leading to more reliable and
390 interpretable results [45].

391 The difference between intervention and control group was estimated by mean change from
392 baseline at six months. Bonferroni correction for multiple measurement [46] was used to adjust
393 level of significance to 0.0125 as we measured effect of PFMT on four outcomes (POP-SS and
394 three domains of HRQoL). Then a mean change difference between the intervention and control
395 arms was presented with 99% confidence interval and P-values. This adjustment is crucial for
396 maintaining statistical rigor and minimizing the risk of Type I errors in the analysis [46].

397 To evaluate model fit, we initially fitted a null model for each outcome variable (POP-SS and the
398 three domains of Prolapse Quality of Life (P-QoL)), incorporating random effects for both
399 cluster and participant ID. We calculated the intraclass correlation coefficient (ICC) to determine
400 the proportion of variance in the outcome attributable to the clustering structure, which was
401 found to be 4% and less in this study. Subsequently, we fitted models that included the
402 intervention variable to the outcome variables individually. This step allowed us to assess the
403 impact of the intervention on each outcome. To control for effect of confounders and effect

404 modifiers, for each model with added predictors, we evaluated model fitness using the Pseudo
405 Log-Likelihood values for model fitness and Adjusted Wald Test for significance of predictors.
406 Higher (less negative) Pseudo Log-Likelihood values indicate better model fit and significant
407 Wald test indicates importance of the predictors [47]. This approach allowed us to identify the
408 most appropriate models for each outcome. Residual analysis was done to assess
409 homoscedasticity and normality of residuals by using histogram. To check for effect modifiers,
410 we employed statistical tests to assess interactions between PFMT and potential effect modifiers.
411 This approach allowed us to determine whether the effect of PFMT varies across different
412 subgroups. The data set on which the analysis based was provided as a supplementary file 2.

413 **Trial Registration**

414 The trial was registered at the Pan African Clinical Trial Registry (<https://pactr.samrc.ac.za>)
415 database, with the registration number PACTR202302505126575.

416

417 **Results**

418 **Trial profile**

419 A total of 187 eligible women were randomized to the intervention group (n = 86) or the control
420 group (101). All of them completed the follow-up and completed the questionnaire at three and
421 six months. The age of participant ranged from 19 to 80 years with the median of 36 years (IQR:
422 31–50). All were parous, with a median of 4 childbirths (IQR: 4–6). The median hours of daily
423 heavy lifting work were 4 hours (IQR: 3–5). The most common stage of prolapse was Stage II
424 120/187(64.17%). The median duration of feeling prolapse symptom was 48 (30 -60) months
425 with the minimum of five months and maximum of 20 years. At baseline, a slightly higher
426 number of women in menopause, family history of prolapse, and history of home birth were
427 observed in the intervention group. The mean baseline POP-SS and quality of life domain scores
428 were almost similar across the groups. Almost half 92 (49.2%) of the participants had prolapse of
429 multiple vaginal compartments; whereby 161 (86.1%) had anterior prolapse, 113 (60.43%) had
430 posterior prolapse and 28/187 (14.97%) had apical prolapse. Overall, while the intervention and
431 control arms were balanced in terms of most characteristics at baseline, there were some

432 important characteristics that showed significant imbalances. Detailed sociodemographic
 433 characteristics are presented in Table 1.

434 **Table 1: Sociodemographic and Clinical Characteristics of Trial Participants at Baseline in**
 435 **Sidama Region, Ethiopia (August 2024) (n = 187)**

| Variables | Category | Control(n=101) | Intervention (n=86) |
|------------------------|---------------------|----------------|---------------------|
| Districts | | | |
| Number of clusters | Dale | 3 | 3 |
| | Wonsho | 1 | 1 |
| Number of participants | Dale | 64 | 70 |
| | Wonsho | 37 | 16 |
| Clusters | Aposto | 21 | - |
| | Megara | - | 26 |
| | Dagia | - | 16 |
| | Shoye | - | 27 |
| | Wayicho | 26 | - |
| | Dansha sire | 17 | - |
| | Mamana | 37 | - |
| | Bokaso | - | 16 |
| Age at base line | Median (IQR) | 34 (30-37) | 50 (35-60) |
| Age at first Birth | Median (IQR) | 18 (17-19) | 17 (16-20) |
| Number of childbirths | (Median (IQR) | 4 (4-5) | 5 (4-7) |
| Age at last delivery | Median (IQR) | 30 (26-32) | 36 (29-42) |
| Level of Education | No formal education | 67 (66.82%) | 60 (69.47%) |
| | Primary school | 18 (17.94%) | 20 (23.16%) |
| | Secondary and above | 15 (15.25%) | 6 (7.38%) |
| Occupation | Employed | 5 (5.38) | 5 (5.79%) |
| | Un-employed | 96 (94.62%) | 81 (94.21%) |
| Menopause | Yes | 24 (24.22%) | 53 (61%) |
| | No | 77 (75.78%) | 34 (39%) |

| | | | |
|--|--------------------|------------------|-----------------|
| Experience of prolonged labor | Yes | 62 (61%) | 46 (53.16%) |
| | No | 39 (39%) | 40 (46.84%) |
| History of home birth | Yes | 66 (65.47%) | 63 (73.16%) |
| | No | 35 (34.53%) | 23 (26.84%) |
| Family history of prolapse | Yes | 53 (52.47%) | 8 (9%) |
| | No | 48 (47.53%) | 78 (91%) |
| Stage of prolapse | Stage 1 | 11 (11.21%) | 11 (13.16%) |
| | Stage 2 | 65 (64.57%) | 55 (64.21%) |
| | Stage 3 | 24 (24.22%) | 19 (22.63%) |
| Body mass index | <18.5 | 43 (42.15%) | 38 (44.74%) |
| | 18.5 – 24.9 | 55 (54.26%) | 43 (49.47%) |
| | >=25 | 4 (3.59%) | 5 (5.79%) |
| Type (site) of prolapse | Anterior only | 34 (33.66%) | 37 (43.02%) |
| | Posterior only | 15 (14.85%) | 10 (11.63%) |
| | More than one site | 52 (51.49%) | 39 (45.35%) |
| Mean Base-line POP-SS | Mean, SD | 12.17 ± (4) | 13.86 ± (4.4) |
| Baseline Symptoms | | | |
| Protruding mass per vagina | | 97 (96.04%) | 82 (95.35) |
| Vaginal Discomfort/pain | | 97 (96.04%) | 83 (96.51%) |
| Heaviness in the lower abdomen | | 96 (95.05%) | 77 (89.53%) |
| Lower back dragging | | 97 (96.04%) | 79 (91.86%) |
| Strain to urinate | | 85 (84.16%) | 81 (94.19%) |
| Difficulty to empty bladder | | 77 (76.24%) | 81 (94.19%) |
| Difficulty to empty bowel | | 73 (72.28%) | 67 (77.91%) |
| Prolapse quality of life (baseline) | | | |
| Physical function | Median (IQR) | 40.5 (29.7 - 51) | 43 (35 – 56.7) |
| Personal relation | Median (IQR) | 44.44 (33 - 55) | 44.44 (33 - 55) |
| Psychological function | Median (IQR) | 46.69 (33 - 53) | 40 (33 - 53) |

436 **IQR*= *Inter-quartile range*, *SD*=*Standard Deviation*; *POP-SS* = *Pelvic organ prolapse*
437 *symptoms score*. ** *P-values presented with* [¥] *indicate Mann-Whitney U test*, [©] *indicate chi-*
438 *square test and* [®] *indicate independent t-test*.

439 **Adherence and Follow-up**

440 All study participants adhered strictly to their allocated treatment and completed the follow-up as
441 scheduled. No participants changed their allocation, missed follow-up appointments, or failed to
442 complete the treatment regimen. Consequently, while our analysis framework was based on the
443 intention-to-treat principle, the actual adherence and follow-up were perfect, ensuring that all
444 participants were analyzed according to their initial allocation.

445

446 **Prolapse Symptoms and Quality of Life at Six Months**

447 A summary at the cluster level reveals that all clusters experienced at least a minimum mean
448 change of 1 point in the POP-SS score. Notably, a larger change was observed in the intervention
449 clusters compared to the control clusters, suggesting a more significant impact of the PFMT
450 intervention on participants in those areas (Table 2).

451

452

453

454

455

456

457

458

459

460

461 **Table 2: Table 1: Change in Pelvic Organ Prolapse Symptom Scores (POP-SS) and Related**
 462 **Domains by Cluster and Intervention Group**

| Clusters | Group | Post intervention | Outcome variables | | | |
|--------------------|--------------|-------------------|---------------------|-----------------------|------------------------|-----------------------|
| | | | POP-SS | Physical function | Psychological function | Personal relation |
| Aposto (n=21) | Control | Mean \pm SD | 11.65 (\pm 3.79) | 41.20 (\pm 19) | 36.25(\pm 12) | 38.64(\pm 15) |
| | | Mean difference | -1.13 (\pm 2.39) | -3.64 (\pm 8.78) | -8.12 (\pm 12.08) | -3.86 (\pm 8.18) |
| Bokaso (n=16) | Intervention | Mean \pm SD | 8.83 (\pm 4.99) | 35.33(\pm 14.3) | 38.35(\pm 12.86) | 29.63 (\pm 11.92) |
| | | Mean difference | -3.17 (\pm 2.32) | -11.93 (\pm 14.24) | -10.00 (\pm 18.22) | -13.89 (\pm 16.61) |
| Dagia (n=16) | Intervention | Mean \pm SD | 8.67 (\pm 5.07) | 23.55 (\pm 15.92) | 26.68(\pm 14.63) | 29.63(\pm 16.08) |
| | | Mean difference | -6.06 (\pm 3.54) | -19.50 (\pm 11.60) | -11.49 (\pm 18.71) | -16.05 (\pm 15.22) |
| Dansha Sire (n=17) | Control | Mean \pm SD | 12.42 (\pm 5.40) | 37.37 (\pm 16.57) | 35.10(\pm 16.42) | 38(\pm 19.48) |
| | | Mean difference | -1.16 (\pm 1.82) | -1.42 (\pm 4.56) | -3.16 (\pm 8.02) | -2.92 (\pm 7.44) |
| Mamana (n=37) | Control | Mean \pm SD | 9.93 (\pm 4.12) | 39 (\pm 14.97) | 35.33(\pm 13.78) | 39.09(\pm 13.61) |
| | | Mean difference | -1.26 (\pm 2.13) | -14.99 (\pm 2.56) | -5.43 (\pm 56) | -3.70 (\pm 6.88) |
| Megara (n=26) | Intervention | Mean \pm SD | 7.62 (\pm 5.14) | 23.46 (\pm 14.28) | 29.21(\pm 13.23) | 32.18(\pm 18.28) |
| | | Mean difference | -6.0 (\pm 4) | -14.99 (\pm 12.11) | -12.88 (\pm 15.98) | -11.88 (\pm 16.53) |

| | | | | | | |
|-------------------|--------------|-----------------|---------------------|-----------------------|-----------------------|-----------------------|
| Shoye (n=27) | Intervention | Mean \pm SD | 8.8 (\pm 5.01) | 33.39 (\pm 13.77) | 30.24(\pm 14.49) | 28.89(\pm 12.68) |
| | | Mean difference | -5.90 (\pm 4.77) | -11.61 (\pm 17.69) | -12.45 (\pm 13.67) | -11.11 (\pm 13.52) |
| Wayicho (n=26) | Control | Mean \pm SD | 10.41 (\pm 4.26) | 33.24 (\pm 21) | 35.42(\pm 14.43) | 37.93(\pm 16.98) |
| | | Mean difference | -1.72 (\pm 3.20) | -3.72 (\pm 6.86) | -8.97 (\pm 12.00) | -4.60 (\pm 11.30) |

463 *SD=Standard Deviation; POP-SS = Pelvic organ prolapse symptoms score*

464

465 Women in the intervention group reported significant improvements in prolapse symptoms and
 466 across all domains of prolapse-related quality of life (P-QoL) at six months. In the unadjusted
 467 model, the most substantial improvement was observed in the psychological function component
 468 of the P-QoL, with a mean change difference of -11.18 (99% CI [-15.03, -7.32]; $p < 0.001$),
 469 indicating a large effect size. Moreover, a significantly higher number of women in the
 470 intervention group reported their condition as “better” after the intervention compared to the
 471 control group. In the control group, more than 27/101 (26%) of women became worse in 6
 472 months (Table 3).

473 Table 3: Effect of pelvic floor muscle training on prolapse symptom and prolapse quality of life,
474 Sidama Region, Ethiopia, August 2024.

| Outcome | After 6months | | Change difference from baseline (mean, 99% CI, P-value) |
|---|------------------------|-------------------------|---|
| | Intervention (n=86) | Control (n=101) | |
| POP-SS (mean ±SD) | 8.4 (± 5.15) | 10.8 (± 4.48) | |
| Change from baseline (mean ±SD) | -5.44 (± 4.03) | -1.34 (± 2.47) | - 4.11 (-5.08 -3.14) |
| | P < 0.001 ^b | P < 0.001 ^b | P < 0.001 ^ψ |
| Prolapse-Quality of life | | | |
| Physical function (mean ±SD) | 28.89 ± (15.6) | 38.1 ± (18.1) | |
| Change from baseline physical function (mean ±SD) | -14.2 ± (14.85) | -2.44 ± (5.68) | - 11.48 (-15.9, -7.1) |
| | P < 0.001 ^b | P = 0.0076 ^b | P < 0.001 ^ψ |
| Psychological function (mean, ±SD) | 24.96 ± (10.95) | 35.5 ± (14.33) | |
| Change from baseline psychological function (mean, ±SD) | -17.76 ± (17.46) | -6.52 ± (10.39) | -11.24 (-16.8, -5.7) |
| | P < 0.001 ^b | P < 0.001 ^b | P = 0.01 ^ψ |
| Personal relation (mean, ±SD) | 30.17 ± (15.13) | 38.5 ± (15.89) | |
| Change from baseline personal relation (mean, ±SD) | -12.81 ± (15.80) | -3.84 ± (8.51) | -8.97 (-13.9, -4.02) |
| | P < 0.001 ^b | P < 0.001 ^b | P < 0.001 ^ψ |
| Perceived improvement after intervention | | | |
| Better | 72 (83.72%) | 42 (41.57%) | 0.0049 [ⓐ] |
| No change | 10 (11.63%) | 32 (31.68%) | |
| Worse | 4 (4.65%) | 27 (26.73%) | |
| POP-SS at 6th months | | | |
| Protruding mass per vagina | 71 (82.56%) | 100 (99.01%) | 0.009 [ⓐ] |
| Vaginal discomfort/pain | 69(80.23%) | 100 (99.01%) | 0.005 [ⓐ] |
| Heaviness in the lower abdomen | 58 (67.44%) | 93 (92.08%) | 0.06 [ⓐ] |
| Lower back dragging | 64 (74.42%) | 95 (94.06%) | 0.048 [ⓐ] |
| Strain to urinate | 58 (67.44%) | 81 (80.20%) | 0.2 [ⓐ] |

| | | | |
|-----------------------------|-------------|-------------|-------------------|
| Difficulty to empty bladder | 58 (67.44%) | 70 (69.31%) | 0.46 [ⓐ] |
| Difficulty to empty bowel | 46 (53.49%) | 67 (66.34%) | 0.19 [ⓐ] |

475 *CI= confidence interval; SD=Standard Deviation; POP-SS = Pelvic organ prolapse symptoms
476 score; n = sample size. ** P-values presented with ^v indicate paired t-test, ^w was based on mixed-
477 effects generalized linear model and [ⓐ] indicate chi-square test.

478

479

480 **Effect of covariates on pelvic organ prolapses and quality of life**

481 We observed that women in the stage I and II prolapses experienced more marked improvements
482 compared to stage III prolapse both in prolapse symptom and quality of life. Women with higher
483 POP-SS scores, indicating more severe symptoms, experienced less improvement in the physical
484 domain of quality of life compared to those with lower scores. The analysis showed that as the
485 severity of prolapse symptoms increased, the associated improvements in the physical domain of
486 the Prolapse Quality of Life (P-QoL) questionnaire were diminished. Moreover, after adjusting
487 for the severity of prolapse symptoms and the stage of prolapse, the effect of PFMT on the
488 physical domain of quality of life was slightly diminished but showed increased effects in the
489 personal relationship and psychological domains (Table 4).

490

491 Table 4: Adjusted Effect of Pelvic Floor Muscle Training on Prolapse Symptoms and Quality of
492 Life in Sidama Region, Ethiopia, August 2024.

| Predictors | Category | β (crude) | β (adjusted) | 99%CI | P-value |
|-------------------------------|---------------------|-----------------|--------------------|------------------|---------|
| POP-SS | | | | | |
| Intervention | Mean change | - 4.11 | -4.11 | -5.38, -2.83 | < 0.001 |
| Stage | Stage I | -5.22 | -4.94 | -7.38, -2.49 | < 0.001 |
| | Stage II | -3.03 | -2.99 | -4.83, -1.13 | < 0.001 |
| | Stage III | Ref | Ref | | |
| Symptom duration | | 0.0 | 0.03 | 0.005, 0.06 | =0.002 |
| prolong labor | | 1.22 | 0.80 | -0.704, 2.30 | =0.17 |
| Education | Not educated | Ref | Ref | | |
| | Primary school | -0.5 | -0.15 | -1.77, 1.46 | =0.81 |
| | Secondary and above | 0.04 | -0.04 | -2.11, 2.03 | =0.96 |
| Physical Function | | | | | |
| Intervention effect | Mean change | - 11.47 | -8.86 | -13.84, -3.89 | < 0.001 |
| Stage | Stage I | -39 | -34.9 | -42.16, -27.62 | < 0.001 |
| | Stage II | -18.37 | -16 | -22.52, -10.79 | < 0.001 |
| | Stage III | Ref | Ref | | |
| Baseline POP-SS | | 1.45 | 0.64 | 0.21, 1.06 | <0.001 |
| Personal relationship | | | | | |
| Intervention effect | Mean change | -8.97 | -9.01 | (-10.49, -5.54) | < 0.001 |
| Stage of prolapse | Stage I | -24.3 | -23.9 | (-29.65, -18.23) | < 0.001 |
| | Stage II | -15.12 | -15.20 | (-20.69, -9.71) | < 0.001 |
| | Stage III | Ref | Ref | | |
| Baseline POP-SS | | 0.83 | 0.23 | (-0.08, 0.55) | 0.06 |
| Psychological function | | | | | |
| Intervention effect | Mean change | -11.24 | -11.18 | (-15.03, -7.32) | < 0.001 |
| Stage of prolapse | Stage I | -18.57 | -16.80 | (-24,34, -9.26) | < 0.001 |

| | | | | | |
|-----------------|-----------|-------|-------|---------------|-------|
| | Stage II | -5.53 | -5.13 | (-9.93, 0.32) | 0.006 |
| | Stage III | Ref | Ref | | |
| Baseline POP-SS | | 0.33 | 0.02 | (-0.43, 0.46) | 0.9 |

493 *POP-SS = pelvic organ prolapses symptom score, CI= confidence interval, Ref= reference
 494 category. ** The regression table shows a combination of four different regression tables (POP-
 495 SS and three domains of Prolapse quality of life).

496 Adverse Events

497 No adverse events were observed.

498

499 Discussion

500 This study showed that midwife-led pelvic floor muscle training combined with lifestyle
 501 counselling significantly improves prolapse symptoms with a mean change difference of -
 502 4.11(99% CI [-5.38, -2.83]; p < 0.001) in prolapse symptom score and each domain of prolapse-
 503 quality of life with a mean change difference of -8.86 (99% CI[-13.84, -3.89]; p<0.001) in the
 504 physical domain, -9.01 (99% CI [-10.49, -5.54]; p <0.001) in the personal relationship domain,
 505 and -11.18 (99% CI [-15.03, -7.32]; p<0.001) in the psychological domain. A significantly higher
 506 number of women in the intervention group perceived their condition as “better” since the
 507 intervention compared to the control group, highlighting the subjective benefit of PFMT. The
 508 change in POP-SS was more noticeable in women with prolapse stage-I with a mean change
 509 difference of -4.94 (99% CI [-7.38, -2.49]; p<0.001) and stage II with a mean change difference
 510 of -2.99 (99% CI [-4.83, -1.13]; p<0.001) as compared to stage III in reducing POP-SS. Similar
 511 effect was observed on each domain of P-QoL. In addition, severity of baseline prolapse
 512 symptoms was associated with physical function of the P-QoL domain with a mean change
 513 difference of 0.64 (99% CI 0.21, 1.01]; p<0.001) compared to minor prolapse symptoms
 514 indicating that more severe symptoms were linked to less improvement.

515 The results of the present study align with existing literature from developed countries indicating
 516 the benefit of PFMT for the treatment of POP symptoms and quality of life. A multicenter
 517 Randomized Controlled Trial (RCT) study from 25 centers (United Kingdom, Dunedin, New
 518 Zealand and Australia) conducted by Hagen et al. [15] showed that PFMT led to significant

519 reduction in POP symptoms and improved quality of life. A single center RCT and systematic
520 review of RCT findings also show that PFMT is effective in reducing POP-symptoms, prolapse
521 stage and quality of life in women with POP-Q stage I-III [4]. Based on the findings of those
522 studies, the Cochrane reviews [48, 49] and national institute for health and care excellence
523 (NICE) guidelines [50] recommended PFMT as a first line of treatment for incontinence and
524 pelvic organ prolapse, to be provided at primary care facilities [51]. Our finding extends these
525 findings and recommendations to a low-income setting which is of a different cultural and health
526 care context, highlighting the universal applicability of PFMT across diverse settings. Moreover,
527 the observed difference in mean change is statistically significant and clinically important as it
528 exceeded the minimally important change of 1.5 for POP-SS [52]. Again, a remarkable
529 improvement was reported on “a feeling of vaginal discomfort/pain and protruding mass through
530 vagina” which are the major indicators for prolapse surgery and regarded as the most crucial
531 treatment target [53]. The significant reduction in these key symptoms underscores the
532 importance of PFMT. These two symptoms are closely linked to the physical support provided
533 by the pelvic floor muscles. Strengthening these muscles through PFMT may lead to a more
534 noticeable reduction in these specific symptoms compared to others that may be less directly
535 influenced by muscle strength or may involve more complex physiological changes.

536 The existing evidence shows that the improvement in POP-SS and P-QoL after PFMT can be
537 attributable to several mechanisms. Overall, PFMT strengthens the pelvic floor muscle,
538 providing better support for pelvic organs which can reduce/stop degree of prolapse and alleviate
539 associated symptoms like vaginal bulging, discomfort, and lower back pain [44, 54]. Enhanced
540 muscle strength also improves urinary and bowel control, reducing symptoms like straining to
541 urinate and difficulty emptying the bowel [54, 55]. The contraction and relaxation of pelvic floor
542 muscles during PFMT promote blood flow to the pelvic region and stimulate the nerves in the
543 area, improving nerve function [56].

544 The study indicates that women with Stage I and II prolapse experience a greater mean change
545 difference in symptom scores compared to those with Stage III, after adjusting for other factors.
546 Borello-France and colleagues reported that women with stage II prolapse were better able to
547 elevate their pelvic floor than those with stage III or IV [57]. Severity of prolapse symptoms was
548 associated with physical function suggesting that more severe POP-SS causes less improvement.

549 After adjusting for prolapse stage and symptom severity, the effect of PFMT remained significant
550 with slightly reduced effect on physical function but slightly increased effects on personal
551 relationship and psychological domains.

552 The findings of this study have important implications for clinical practice and public health
553 policy in low-resource settings. The first implication is that the result suggest that midwife-led
554 PFMT can be considered as a first-line conservative treatment for mild to moderate stage of
555 prolapse in all settings especially in resource limited settings where access to specialized care is
556 limited. This can be integrated in to the existing maternal and reproductive health programs. The
557 second implication is that in settings where the number of urogynecologists/pelvic physiotherapy
558 expert is limited, like the current Ethiopian context, training midwives to deliver PFMT and
559 lifestyle counseling is feasible and effective to address prolapse symptoms and improve quality
560 of life among women with POP. We recognize that the most recommended form of PFMT is
561 supervised training given by a physiotherapist or urogynecologists who have greater in-depth
562 knowledge [1]. However, this is not feasible in low-resource settings due to shortage of trained
563 specialists (only less than 28 urogynecologists in Ethiopia). It was evidenced that nurse-delivered
564 PFMT have comparable effect with those delivered by specialists when implemented properly
565 [58, 59]. In the present study, the observed mean change difference was relatively larger and the
566 compliance/adherence to training was better compared to previous studies. This adherence might
567 be because midwives working in the nearby health facility and the women in the intervention
568 know each other, and that may create trust and acceptability. This is a nice model that can be
569 scaled up in other settings including high income countries.

570 This study has its own strength and limitations. Among others, the key strength of the study
571 includes the rigorous cluster randomized controlled design which enhances the internal validity
572 of the finding, high adherence rate and complete follow-up which ensure the reliability of the
573 results. The use of validated and culturally adapted measurement tools ensures the relevance and
574 accuracy of the collected data, and also the use of a complex survey design, which accounts for
575 the clustered nature of the data, enhances the potential generalizability of the results to the target
576 population. Above all, to our knowledge, this study is the first to evaluate effect of PFMT on
577 prolapse symptoms in Ethiopia.

578 However, some limitations should be considered. The reliance on self-reported adherence to
579 PFMT and symptom improvement may introduce performance bias since the participants were
580 not blinded to the intervention. This participants' knowledge of their treatment allocation may
581 influence their reported outcomes that could lead to higher treatment effect and reported
582 adherence rate. In addition, the follow-up period was relatively short, which prevents us from
583 determining whether the observed effects are likely to be sustained over time. During
584 randomization, we aimed to balance the two groups by allocating similar clusters using stratified
585 cluster sampling. However, this method cannot account for individual-level variables, which is
586 an inherent limitation of cluster sampling. As a result, the age of the participant, parity and
587 family history of the participants were not balanced in this study. The age discrepancy may stem
588 from inaccuracies in measuring age, as many individuals in rural Ethiopia has no birth
589 certificates.

590

591

592 **Conclusion**

593 Midwife-led pelvic floor muscle training combined with lifestyle counseling significantly
594 improves prolapse symptoms and quality of life across all domains in women with pelvic organ
595 prolapse. The observed treatment effects were significant and clinically important. The effects of
596 PFMT were higher in women with earlier stage of prolapse. Move over, in settings with shortage
597 of urogynecologists, training midwives to deliver PFMT and lifestyle counselling is a feasible
598 and effective approach to improve prolapse symptoms and quality of life. Future studies should
599 incorporate objective measures of adherence and explore long-term effects. Healthcare
600 organizations should advocate and endorse training of midwives and other healthcare providers
601 to deliver PFMT.

602

603 **Declarations**

604 **Ethical Considerations and consent to participate**

605 This study was conducted in accordance with the principles of the Declaration of Helsinki.
606 Ethical approval was obtained from Institutional Review Board of Hawassa University, College
607 of Medicine and Health Sciences (Ref. No. IRB/157/14). Written informed consent was obtained
608 with provisions for participants unable to read and write to sign with their index fingers,
609 confirming that the consent had been read to them. Given the sensitive nature of the condition,
610 data regarding self-reported prolapse symptoms were collected in a private location within the
611 participants' homes. Pelvic examinations were conducted in delivery rooms to ensure
612 participants' physical privacy, while lifestyle counseling and pelvic floor muscle training took
613 place in private rooms. All study information was kept confidential, with participants assigned
614 codes to replace their names and personally identifying information. Access to the key linking
615 these codes to individual identities was restricted to the principal investigator. The risks to
616 participants were minimal. While there was potential for psychological distress related to the
617 pelvic examination, as well as temporary discomfort during physical exams and inconvenience
618 from follow-up visits, these risks were mitigated by allowing participants to withdraw from the
619 study at any time during the consent process. The research team provided travel costs and time
620 compensation, equivalent to approximately four dollars (200 Ethiopian Birr), to the participants.
621 During baseline data collection, around 26 participants with stage IV prolapse were referred to
622 the Yirgalem Hamlin Fistula Center for further management. After the study's completion,
623 women with symptomatic stage III prolapse in the control group linked with the Yirgalem
624 Hamlin Fistula Center for additional care.

625 **Consent for publication**

626 Not Applicable

627 **Authors contribution**

628 Melese Siyoum: Conceptualization, methodology, fund acquisition, mentorship, formal analysis,
629 validation, writing original draft, review and editing on final version.

630 Rahel Nardos: Methodology, mentorship, fund acquisition, validation, reviewing and editing the
631 manuscript

632 Biniyam Sirak: Methodology, fund acquisition, validation, reviewing and editing the manuscript

633 Theresa Spitznagle: Methodology, validation, reviewing and editing the manuscript

634 Wondwosen Teklesilasie: Methodology, validation, reviewing and editing the manuscript

635 Ayalew Astatkie: Conceptualization, methodology, formal analysis, validation, reviewing and
636 editing the manuscript

637 All authors read and approved the final manuscript

638 **Acknowledgment**

639 We would like to thank our sponsors (Hawassa University, University of Minnesota, Worldwide
640 Fistula Fund, and Maternal Health Fund) for their financial support. The study participants and
641 data collectors also deserve gratitude. We also thank Dr. Karen Gold and Joseph Kinahan
642 (Maternal Health Fund), Soja Orłowski (Worldwide Fistula Fund), Katie Brown (University of
643 Minnesota) and Alemu Tamiso (Hawassa University) for their facilitation in transferring funds.
644 Yirgalem Hamlin Fistula Center, Health centers in the study area and Mr. Rekiku Fikre (Hawassa
645 University) deserve acknowledgement for facilitating transportation service support in material.

646 **Funding**

647 The payment for data collectors, participant transportation, and field supervisor per-diem was
648 sponsored by Hawassa University, University of Minnesota, Worldwide Fistula Fund and
649 Maternal Health Fund. The funders have no role in designing the study, intervention, data
650 management and publication.

651 **Competing of interest**

652 All authors declare that they have no competing of interest

653 **Availability of data and materials**

654 The dataset used for this study is available within the manuscript and as supplementary file

655

656

657

658

659 **References**

- 660 1. Bo K, Frawley HC, Haylen BT, Abramov Y, Almeida FG, Berghmans B, et al. An
661 International Urogynecological Association (IUGA)/International Continence Society (ICS) joint
662 report on the terminology for the conservative and nonpharmacological management of female
663 pelvic floor dysfunction. *International urogynecology journal*. 2017;28:191-213.
- 664 2. Bø K, Anglès-Acedo S, Batra A, Brækken IH, Chan YL, Jorge CH, et al. International
665 urogynecology consultation chapter 3 committee 2; conservative treatment of patient with pelvic
666 organ prolapse: pelvic floor muscle training. *International urogynecology journal*.
667 2022;33(10):2633-67.
- 668 3. Hadizadeh-Talasaz Z, Khadivzadeh T, Mohajeri T, Sadeghi M. Worldwide Prevalence of
669 Pelvic Organ Prolapse: A Systematic Review and Meta-Analysis. *Iranian Journal of Public
670 Health*. 2024.
- 671 4. Brown HW, Hegde A, Huebner M, Neels H, Barnes HC, Marquini GV, et al. International
672 urogynecology consultation chapter 1 committee 2: Epidemiology of pelvic organ prolapse:
673 prevalence, incidence, natural history, and service needs. *International Urogynecology Journal*.
674 2022:1-15.
- 675 5. Barber MD, Maher C. Epidemiology and outcome assessment of pelvic organ prolapse.
676 *International urogynecology journal*. 2013;24:1783-90.
- 677 6. Ward RM, Edwards DRV, Edwards T, Giri A, Jerome RN, Wu JM. Genetic epidemiology
678 of pelvic organ prolapse: a systematic review. *American journal of obstetrics and gynecology*.
679 2014;211(4):326-35.
- 680 7. Ai F, Deng M, Mao M, Xu T, Zhu L. Depressive symptoms screening in postmenopausal
681 women with symptomatic pelvic organ prolapse. *Menopause*. 2018;25(3):314-9.

- 682 8. Lowenstein L, Gamble T, Deniseiko Sanses TV, Van Raalte H, Carberry C, Jakus S, et al.
683 Sexual function is related to body image perception in women with pelvic organ prolapse. The
684 journal of sexual medicine. 2009;6(8):2286-91.
- 685 9. Ellerkmann RM, Cundiff GW, Melick CF, Nihira MA, Leffler K, Bent AE. Correlation of
686 symptoms with location and severity of pelvic organ prolapse. American journal of obstetrics and
687 gynecology. 2001;185(6):1332-8.
- 688 10. Swift SE, Tate SB, Nicholas J. Correlation of symptoms with degree of pelvic organ
689 support in a general population of women: what is pelvic organ prolapse? American journal of
690 obstetrics and gynecology. 2003;189(2):372-7.
- 691 11. Aponte MM, Rosenblum N. Repair of pelvic organ prolapse: what is the goal? Current
692 urology reports. 2014;15:1-5.
- 693 12. Wiegersma M, Panman CM, Hesselink LC, Malmberg AG, Berger MY, Kollen BJ, et al.
694 Predictors of success for pelvic floor muscle training in pelvic organ prolapse. Physical Therapy.
695 2019;99(1):109-17.
- 696 13. Giagio S, Innocenti T, Salvioli S, Lami A, Meriggiola MC, Pillastrini P, et al.
697 Completeness of exercise reporting among randomized controlled trials on pelvic floor muscle
698 training for women with pelvic organ prolapse: A systematic review. Neurourology and
699 Urodynamics. 2021;40(6):1424-32.
- 700 14. Abhyankar P, Uny I, Semple K, Wane S, Hagen S, Wilkinson J, et al. Women's
701 experiences of receiving care for pelvic organ prolapse: a qualitative study. BMC women's
702 health. 2019;19:1-12.

- 703 15. Hagen S, Stark D, Glazener C, Dickson S, Barry S, Elders A, et al. Individualised pelvic
704 floor muscle training in women with pelvic organ prolapse (POPPY): a multicentre randomised
705 controlled trial. *The Lancet*. 2014;383(9919):796-806.
- 706 16. Abhyankar P, Wilkinson J, Berry K, Wane S, Uny I, Aitchison P, et al. Implementing
707 pelvic floor muscle training for women with pelvic organ prolapse: a realist evaluation of
708 different delivery models. *BMC Health Services Research*. 2020;20:1-16.
- 709 17. Belayneh T, Gebeyehu A, Adefris M, Rortveit G, Gjerde JL, Ayele TA. Pelvic organ
710 prolapse surgery and health-related quality of life: a follow-up study. *BMC Women's Health*.
711 2021;21:1-11.
- 712 18. Workineh ZA, Ayana BA, Gashaye KT, Wubneh SB, Kassie BA. Surgical management
713 practice of pelvic organ prolapse among Ethiopian gynecologists, 2021: a descriptive study.
714 *BMC Women's Health*. 2023;23(1):654.
- 715 19. Sung VW, Jeppson P, Madsen A. Nonoperative management of pelvic organ prolapse.
716 *Obstetrics & Gynecology*. 2023;141(4):724-36.
- 717 20. Footer CB, Tsegaye HS, Yitnagashaw TA, Mekonnen W, Shiferaw TD, Abera E, et al.
718 Empowering the physiotherapy profession in Ethiopia through leadership development within
719 the doctoring process. *Frontiers in Public Health*. 2017;5:51.
- 720 21. (FMoHE) TFDRoE. National Health Policy. In: Health, editor. Addis Ababa2024.
- 721 22. Moher D, Hopewell S, Schulz KF, Montori V, Gøtzsche PC, Devereaux PJ, et al.
722 CONSORT 2010 explanation and elaboration: updated guidelines for reporting parallel group
723 randomised trials. *Bmj*. 2010;340.
- 724 23. Campbell MK, Piaggio G, Elbourne DR, Altman DG. Consort 2010 statement: extension
725 to cluster randomised trials. *Bmj*. 2012;345.

- 726 24. Siyoum M, Nardos R, Teklesilasie W, Astatkie A. Prevalence and risk factors of pelvic
727 organ prolapse among women in Sidama region, Ethiopia: A community-based survey. *Women's*
728 *Health*. 2024;20:17455057241265078.
- 729 25. Reid F. Assessment of pelvic organ prolapse: a practical guide to the pelvic organ
730 prolapse quantification. *Obstetrics, Gynaecology & Reproductive Medicine*. 2014;24(6):170-6.
- 731 26. Vértesy L, Lemango TB. Comparison of local governments in Hungary and Ethiopia. *DE*
732 *IURISPRUDENTIA ET IURE PUBLICO: JOG-ÉS POLITIKATUDOMÁNYI FOLYÓIRAT=*
733 *JOURNAL OF LEGAL AND POLITICAL SCIENCES*. 2022;13(1-2.):62-75.
- 734 27. Gedefaw AA. Analysis of the contribution of land registration to sustainable land
735 management in East Gojjam Zone, Ethiopia. *Land*. 2023;12(6):1157.
- 736 28. Nefa SA, Ibrahim AM, Mulushewa SS, Wonga MM, Lee H-Y, Siyum GA, et al. Study
737 design for the 2019 baseline survey of newly established longitudinal surveillance woredas as the
738 field lab to serve child health improvement project of Benishangul-Gumuz region, Ethiopia.
739 *Journal of Global Health Science*. 2019;1(2).
- 740 29. Donner A, Birkett N, Buck C. Randomization by cluster: sample size requirements and
741 analysis. *American journal of epidemiology*. 1981;114(6):906-14.
- 742 30. Hemming K, Eldridge S, Forbes G, Weijer C, Taljaard M. How to design efficient cluster
743 randomised trials. *bmj*. 2017;358.
- 744 31. Hayes RJ, Moulton LH. *Cluster randomised trials*: Chapman and Hall/CRC; 2017.
- 745 32. Rutterford C, Copas A, Eldridge S. Methods for sample size determination in cluster
746 randomized trials. *International journal of epidemiology*. 2015;44(3):1051-67.

- 747 33. Roth SM, Ivey FM, Martel GF, Lemmer JT, Hurlbut DE, Siegel EL, et al. Muscle size
748 responses to strength training in young and older men and women. *Journal of the American*
749 *Geriatrics Society*. 2001;49(11):1428-33.
- 750 34. Maxwell M, Berry K, Wane S, Hagen S, McClurg D, Duncan E, et al. Pelvic floor muscle
751 training for women with pelvic organ prolapse: the PROPEL realist evaluation. *Health Services*
752 *and Delivery Research*. 2020;8(47).
- 753 35. Hay-Smith J, Bo K, Berghmans B, Hendriks E, de Bie R, van Doorn EvW. Pelvic floor
754 muscle training for urinary incontinence in women. *The Cochrane database of systematic*
755 *reviews*. 2008;2008(3).
- 756 36. Maxwell M, Semple K, Wane S, Elders A, Duncan E, Abhyankar P, et al. PROPEL:
757 implementation of an evidence based pelvic floor muscle training intervention for women with
758 pelvic organ prolapse: a realist evaluation and outcomes study protocol. *BMC health services*
759 *research*. 2017;17:1-10.
- 760 37. Women's NCCf, Health Cs. *Urinary incontinence: The management of urinary*
761 *incontinence in women*. 2006.
- 762 38. Bø K. Pelvic floor muscle exercise for the treatment of stress urinary incontinence: an
763 exercise physiology perspective. *International Urogynecology Journal*. 1995;6(5):282-91.
- 764 39. Siyoum M, Teklesilasie W, Nardos R, Sirak B, Astatkie A. Reliability and validity of the
765 Sidaamu Afoo version of the pelvic organ prolapse symptom score questionnaire. *BMC Women's*
766 *Health*. 2023;23(1):324.
- 767 40. Hagen S, Glazener C, Sinclair L, Stark D, Bugge C. Psychometric properties of the pelvic
768 organ prolapse symptom score. *BJOG: an International Journal of Obstetrics & Gynaecology*.
769 2009;116(1):25-31.

- 770 41. Belayneh T, Gebeyehu A, Adefris M, Rortveit G, Genet T. Translation, transcultural
771 adaptation, reliability and validation of the pelvic organ prolapse quality of life (P-QoL) in
772 Amharic. *Health and quality of life outcomes*. 2019;17:1-11.
- 773 42. Digesu GA, Khullar V, Cardozo L, Robinson D, Salvatore S. P-QOL: a validated
774 questionnaire to assess the symptoms and quality of life of women with urogenital prolapse.
775 *International Urogynecology Journal*. 2005;16:176-81.
- 776 43. Srikrishna S, Robinson D, Cardozo L. Validation of the Patient Global Impression of
777 Improvement (PGI-I) for urogenital prolapse. *International urogynecology journal*. 2010;21:523-
778 8.
- 779 44. Brækken IH, Majida M, Engh ME, Bø K. Morphological changes after pelvic floor
780 muscle training measured by 3-dimensional ultrasonography: a randomized controlled trial.
781 *Obstetrics & Gynecology*. 2010;115(2 Part 1):317-24.
- 782 45. McCulloch CE, Searle SR, Neuhaus JM. *Generalized, linear, and mixed models*: Wiley
783 Online Library; 2001.
- 784 46. Bland JM, Altman DG. Multiple significance tests: the Bonferroni method. *Bmj*.
785 1995;310(6973):170.
- 786 47. Heeringa SG, West BT, Berglund PA. *Applied survey data analysis*: chapman and
787 hall/CRC; 2017.
- 788 48. Dumoulin C, Hunter KF, Moore K, Bradley CS, Burgio KL, Hagen S, et al. Conservative
789 management for female urinary incontinence and pelvic organ prolapse review 2013: Summary
790 of the 5th International Consultation on Incontinence. *Neurourology and urodynamics*.
791 2016;35(1):15-20.

- 792 49. Hagen S, Stark D. Conservative prevention and management of pelvic organ prolapse in
793 women. *Cochrane Database of Systematic Reviews*. 2011(12).
- 794 50. National Institute for Health and Care Excellence guideline NG123 N. Urinary
795 incontinence and pelvic organ prolapse in women: management London 2019 [
- 796 51. Smith A, Bevan D, Douglas HR, James D. Management of urinary incontinence in
797 women: summary of updated NICE guidance. *Bmj*. 2013;347.
- 798 52. Hagen S, Glazener C, Cook J, Herbison P, Toozs-Hobson P. Further properties of the
799 pelvic organ prolapse symptom score: minimally important change and test-retest reliability.
800 *Neurourol Urodyn*. 2010;29(6):1055-6.
- 801 53. Ge J, Wei X, Zhang H, Fang G. Pelvic floor muscle training in the treatment of pelvic
802 organ prolapse: A meta-analysis of randomized controlled trials. *Actas Urológicas Españolas*
803 (English Edition). 2021;45(1):73-82.
- 804 54. Blanchard V, Nyangoh-Timoh K, Fritel X, Fauconnier A, Pizzoferrato A-C. Importance
805 of a pelvic floor lifestyle program in women with pelvic floor dysfunctions: A pilot study.
806 *Journal of Gynecology Obstetrics and Human Reproduction*. 2021;50(4):102032.
- 807 55. Wiegersma M, Panman CM, Kollen BJ, Berger MY, Lisman-Van Leeuwen Y, Dekker JH.
808 Effect of pelvic floor muscle training compared with watchful waiting in older women with
809 symptomatic mild pelvic organ prolapse: randomised controlled trial in primary care. *Bmj*.
810 2014;349.
- 811 56. Hagen S, Glazener C, McClurg D, Macarthur C, Elders A, Herbison P, et al. Pelvic floor
812 muscle training for secondary prevention of pelvic organ prolapse (PREVPROL): a multicentre
813 randomised controlled trial. *The lancet*. 2017;389(10067):393-402.

- 814 57. Borello-France DF, Handa VL, Brown MB, Goode P, Kreder K, Scheufele LL, et al.
815 Pelvic-floor muscle function in women with pelvic organ prolapse. *Physical therapy*.
816 2007;87(4):399-407.
- 817 58. Waterfield AE. *A community study of pelvic floor muscle function in women: Exeter and*
818 *Plymouth Peninsula Medical School*; 2011.
- 819 59. Child S, Bateman A, Shuttleworth J, Gericke C, Freeman R. Can primary care nurse
820 administered pelvic floor muscle training (PFMT) be implemented for the prevention and
821 treatment of urinary incontinence? *A study protocol*. *F1000Research*. 2013;2.
- 822