- 1 Developing an evaluation framework for public health
- 2 environmental surveillance: Protocol for an international,
- 3 multidisciplinary e-Delphi study
- 4 Public Health Environmental Surveillance Evaluation Framework (PHES-
- 5 **EF**)
- 6 Douglas G. Manuel^{1,2,3,4*} (0000-0003-0912-0845), Carol Bennett^{1,2} (0000-0002-3033-9860), Emma Brown¹
- 7 (0000-0002-0869-9907), David L. Buckeridge⁵ (0000-0003-1817-5047), Yoni Freedhoff² (0000-0001-
- 8 9391-3455), Sarah Funnell⁶, Farah Ishtiaq⁷ (0000-0002-6762-7014), Matthew J. Wade^{8,9} (0000-0001-
- 9 9824-7121), David Moher^{3,10} (0000-0003-2434-4206), on behalf of the PHES-EF Executive Group^
- ¹ Methodological and Implementation Research Program, Ottawa Hospital Research Institute, Ottawa,
- 11 Ontario, Canada
- 12 ² Department of Family Medicine, University of Ottawa, Ottawa, Ontario, Canada
- ³ School of Epidemiology and Public Health, Faculty of Medicine, University of Ottawa, Ottawa, Ontario,
- 14 Canada
- 15 ⁴ C.T. Lamont Primary Health Care Research Centre Program, Bruyère Research Institute, Ottawa,
- 16 Ontario, Canada
- ⁵ School of Population and Global Health, McGill University, Montreal, Quebec, Canada

- ⁶ Department of Family Medicine, Faculty of Health Sciences, Queen's University, Kingston, Ontario,
- 19 Canada
- ⁷ Tata Institute for Genetics and Society, National Centre for Biological Sciences, Bengaluru,
- 21 Karnataka, India
- ⁸ Data Analytics and Surveillance Group, UK Health Security Agency, London, United Kingdom
- ⁹ School of Engineering, Newcastle University, Newcastle-upon-Tyne, United Kingdom
- ¹⁰Centre for Journalology, Methodological and Implementation Research Program, Ottawa Hospital
- 25 Research Institute, Ottawa, Ontario, Canada
- 26 *Corresponding Author
- 27 Email: dmanuel@ohri.ca (DGM)
- 28 ^Membership of the PHES-EF Executive Group is provided in the Acknowledgments.

29 Metadata

30 Funding

- 31 This project is funded by the Canadian Institutes of Health Research-funded network, CoVaRR-Net
- 32 (Coronavirus Variants Rapid Response Network. FRN: 175622), and Health Canada (through the Safe
- 33 Restart Agreement Contribution Program. Arrangement #: 2223-HQ-000098). The Canadian Institutes of
- 34 Health Research and Health Canada have not been involved in the design or conduct of the study and
- 35 the views expressed herein do not necessarily represent the views of either funding organization.

36 Competing interests

37 The authors have declared that no competing interests exist.

38 Data availability

39 Anonymous aggregate data will be made publicly available when the study is completed and published.

40 Abstract

41 Introduction

- 42 Public health environmental surveillance has evolved, especially during the coronavirus disease
- 43 pandemic, with wastewater-based surveillance being a prominent example. As surveillance methods
- 44 diversify and expand, it is essential to have a robust evaluation of surveillance systems. This electronic
- 45 Delphi study will propose an evaluation framework for public health environmental surveillance,
- 46 informed by the expanding practice of wastewater-based surveillance during the pandemic.

47 Methods

48 The Public Health Environmental Surveillance Evaluation Framework (PHES-EF) will be developed in five 49 steps. In Step 1, a multinational and multidisciplinary executive group will be formed to guide the 50 framework development process. In Step 2, candidate items for Round 1 of the electronic Delphi will be 51 generated by conducting relevant scoping reviews and consultation with the study executive group. In 52 Step 3, an international electronic Delphi will be conducted over two rounds to develop consensus on 53 items for the framework. In Step 4, the executive group will reconvene to finalize the evaluation 54 framework, discuss standout items, and determine the dissemination strategies. Lastly, Step 5 will focus 55 on disseminating the evaluation framework to all parties involved with or affected by wastewater-based 56 surveillance using traditional and public-oriented methods.

57 Discussion

58 The electronic Delphi study will provide multidisciplinary and multinational consensus for the evaluation 59 framework for public health environmental surveillance by providing a set of minimum criteria required

- 60 for their evaluation. The evaluation framework is intended to support the sustainability of
- 61 environmental surveillance and improve its implementation, reliability, credibility, and value.

62 Introduction

63	Public health relies on robust surveillance systems to support informed decision-making and timely
64	interventions [1]. Environmental-based surveillance, particularly wastewater-based testing, has
65	historically informed public health responses to threats like typhoid, polio, and illicit drug use [2–6]. The
66	coronavirus disease (Covid-19) pandemic marked a turning point, with an expansion of wastewater-
67	based monitoring and surveillance to an unprecedented scale, highlighting its potential in broader public
68	health contexts [7].
69	Building on the advances and experiences from Covid-19 wastewater-based surveillance (wastewater
70	surveillance) during the pandemic, this study aims to develop a comprehensive evaluation framework.
71	Wastewater surveillance will be an example; however, it will apply to broader public health
72	environmental surveillance systems, including air and surface monitoring. Surveillance targets
73	considered include infectious agents and health risks beyond Covid-19.
74	Wastewater surveillance, once a peripheral tool, gained prominence during the pandemic. Its global
75	implementation in diverse settings [8], from metropolitan to rural areas [9,10], highlighted its strengths.
76	These strengths include the provision of insights into disease spread, unaffected by clinical testing
77	access or population health-seeking behaviours, and offering a non-invasive, resource-efficient
78	surveillance method [11,12].
79	Despite these strengths, the widespread application of wastewater surveillance faces challenges
80	[5,13,14]. Environmental variability affects measurement consistency, and the need for multidisciplinary
81	collaboration brings logistic complexities [14]. Rapidly evolving methodologies have led to various
82	analytic techniques and interpretation approaches, resulting in inconsistencies and uncertainties [7].

- 83 These challenges underscore the need for a structured evaluation framework. Such a framework can
- 84 enhance accountability, transparency, and trust in wastewater surveillance systems, potentially
- 85 improving surveillance performance and identifying best practices [1].

86 Study objectives

- 87 This project will use a multidisciplinary and multinational consensus approach to develop an evaluation
- 88 framework for public health environmental surveillance (Public Health Environmental Surveillance
- 89 Evaluation Framework, PHES-EF). An evaluation framework provides structured guidance to facilitate a
- 90 systematic approach to the evaluation of the program [15–17]. See below for the study's definition of an
- 91 evaluation framework.
- 92 Specific objectives for the framework development process are as follows: (1) engage a multidisciplinary
- 93 executive group of international experts to inform the development of the evaluation framework; (2)
- 94 review the literature to identify current guidance on surveillance evaluation items; (3) prioritize items
- 95 for an evaluation framework for environmental surveillance using an international electronic-Delphi (e-
- 96 Delphi) process; (4) conduct a consensus meeting, which engages experts across disciplines, to create an
- 97 evaluation framework for public health environmental surveillance based on feedback from the
- 98 international e-Delphi process; and, (5) develop a comprehensive dissemination plan for the evaluation
- 99 framework.

100 **Definitions**

- 101 We define key terms as follows:
- Environmental surveillance: Surveillance of substances or organisms in air, water, soil, and
 living organisms to detect changes or threats that may affect human health, ecosystems, or both
 [18].

105	•	Evaluation Framework: A structured system or schema to systematically assess a program
106		[1,15,17]. An evaluation framework presents a clear list of items or elements that should appear
107		in an evaluation, such as constructs or concepts. It provides an organisation or structure for the
108		list of items; defines how the list and organisation were developed; and, potentially, includes
109		details about measuring the evaluation items. Importantly, an evaluation framework facilitates
110		informed decision-making by providing a comprehensive understanding of program
111		effectiveness and areas for improvement.
112	•	Public health surveillance: The ongoing systematic collection, analysis, interpretation and
113		dissemination of health data for the planning, implementation and evaluation of public health
114		interventions and strategies to protect and improve the health of populations [19].
115	٠	Wastewater: Water that has been in contact with people (e.g., for washing) or used for
116		cleansing and sanitation (e.g., for flushing away faecal matter), and is discharged via sewers or
117		other sanitation systems [6].
118	•	Wastewater-based surveillance (wastewater surveillance): A public health approach that
119		examines specific substances or organisms in wastewater (sewage). Wastewater surveillance
120		can be referred to as wastewater-based epidemiology; however, the latter can be specific to the
121		use of wastewater to study and analyse the distribution, patterns, and determinants of health
122		and disease conditions in defined populations [19].

123 Materials and methods

124 Study design

- 125 This protocol was developed *a priori*. The development of PHES-EF is guided by recommendations for
- 126 Delphi techniques in the health sciences [20]. An overview of the study flow is detailed in Fig 1.

A multinational and multidisciplinary executive group will oversee the development of the evaluation

127 Fig 1. Overview of the development process for PHES-EF.

128 Step 1: Establish the executive group

129

130	framework. The group will bring expertise from various fields, including public health, infectious disease,
131	epidemiology; environmental and physical sciences; mathematical sciences; social sciences;
132	communication, knowledge translation and exchange; knowledge users; and engaged public.
133	Content expert and knowledge user executive group members will be identified by examining relevant
134	author lists, internet-based searching, and the study working group's professional networks. In contrast,
135	the engaged member of the public will be identified through an open call for expressions of interest. The
136	composition of the study executive group is detailed in S1 Table.
137	The executive group will be responsible for providing feedback on the framework development process
138	(e.g., framework organization, completeness, additions, redundancy), supporting and advising working
139	groups in their area of expertise, connecting the study to other related projects, recruiting e-Delphi
140	panellists, participating in the consensus meeting, disseminating the evaluation framework, advising
141	and/or participating in post-publication activities, and promoting the adoption of the evaluation
142	framework among their professional networks. Co-chairs will also be responsible for decision-making
143	when the executive group cannot reach consensus, leading executive group activities, and the overall
144	framework development process. Any conflicts of interest among the research team, including executive
145	group members, will be assessed before beginning the study.

146 Step 2: Generate candidate evaluation items

- 147 We will systematically compile a list of e-Delphi candidate items through three processes: scoping
- 148 reviews, content analysis and extraction, and executive group consultation.

149 Scoping reviews

- 150 Two scoping reviews will be performed to provide a comprehensive view of public health environmental
- 151 surveillance system evaluations. The first review will identify frameworks (e.g., sources which guide the
- 152 planning, management, or conduct of processes) and criteria (e.g., principles on which a judgment may
- 153 be based) for evaluating public health or environmental surveillance systems. The second review will
- explore insights from the Covid-19 pandemic, particularly focusing on the principles of equity, trust,
- 155 open science, community engagement, and One Health in the context of wastewater surveillance [21–
- 156 23]. Detailed protocols for the scoping reviews are available on OSF
- 157 (<u>https://doi.org/10.17605/OSF.IO/GSKF6</u>; <u>https://doi.org/10.17605/OSF.IO/C9A3Q</u>). The completed
- 158 scoping reviews will be reported following the Preferred Reporting Items for Systematic Reviews and
- 159 Meta-Analyses extension for scoping reviews (PRISMA-ScR) [24].

160 Content analysis and extraction

161 We will conduct a content analysis on items identified during the data extraction phase for both scoping

- 162 reviews. The content analyses will identify and group criteria for evaluating public health and
- 163 environmental surveillance systems. In the first step, one team member will group all the extracted
- 164 items into initial domains and themes. In the second step, two researchers will independently compare
- the category system of the first researcher to agree on the final category system consensually. In the
- third step, all criteria will be reviewed again to merge or modify duplicate concepts and create a
- 167 parsimonious framework for the initial list of e-Delphi candidate items.

168 Executive group consultation

- 169 The executive group will review the initial list of e-Delphi candidate items and discuss the initial
- 170 structure and presentation of Round 1 items. Their review will focus on creating an evidence-informed
- 171 public health environmental surveillance framework centred around SARS-CoV-2 wastewater
- surveillance, and incorporating lessons learned from the pandemic (e.g., equity, open science,
- 173 community engagement, and One Health). Additional items will be added if the executive group
- 174 determines that key evaluation areas have not been addressed.

175 Step 3: International e-Delphi survey

176 A Delphi technique will be used to develop an internationally accepted evaluation framework for public

177 health environmental surveillance. The Delphi method is an iterative multi-round approach that uses a

- series of sequential surveys, interspersed by controlled feedback, to elicit consensus among a group of
- individuals while maintaining anonymity [25,26]. An e-Delphi method will be used to overcome
- 180 geographic barriers and allow us to engage panellists internationally across various time zones. Delphi
- 181 surveys relating to health sciences have typically been conducted for 2-3 rounds [20,27,28]. Therefore, a
- 182 two-round e-Delphi survey and a subsequent consensus meeting will be conducted to balance feasibility
- 183 with rigour. Each round will be active for approximately 4-6 weeks to allow panellists sufficient time to
- 184 participate in the survey while balancing study timeline constraints [29].

185 Recruitment of panellists

We will recruit a multinational, multidisciplinary panel of wastewater surveillance experts, knowledge users, and engaged members of the public to complete the e-Delphi survey. Content experts are adult individuals with knowledge in their respective subdiscipline groups (see S2 Table). Panellists will be selected to capture the multiple perspectives of those that influence the design, implementation,

190	evaluation, use, and reporting of wastewater surveillance activities, including the following
191	subdisciplines groups: public health, infectious disease, epidemiology; environmental and physical
192	sciences; mathematical sciences; social sciences; and communication, knowledge translation and
193	exchange. Targeted recruitment will also be conducted to identify knowledge users and engaged
194	members of the public. We will use purposive sampling to identify potentially eligible panellists by
195	examining relevant author lists from the scoping reviews, professional networks, and internet-based
196	searching, among other potential methods. Potential panellists may also be recruited using social media,
197	a study website or calls for expressions of interest during relevant meetings, presentations, or
198	correspondence. Snowball sampling may also be employed by executive group members and potential
199	panellists to maximize recruitment.
200	Potentially eligible panellists will receive a recruitment email from the study working group, executive
201	group members, or other potential panellists. A call for panellists to share the e-Delphi with other
202	potentially eligible panellists may be included in recruitment materials and the e-Delphi survey.

203 Inclusion criteria for e-Delphi panellists are outlined in Table 1.

204 Table 1. Eligibility criteria for e-Delphi survey panellists

Discipline group	Discipline subgroup	Inclusion criteria
Content experts	Public health, infectious	An adult ^a who is proficient in English and has a
	disease, epidemiology	graduate degree in one of the listed
	Environmental and physical	specializations ^b , or \geq 3 years of professional
	sciences	experience, or \geq 2 peer-reviewed publications
		relating to wastewater surveillance.
	Mathematical sciences	

	Social sciences Communication, knowledge translation and exchange	
Knowledge users	1	An adult ^a who is proficient in English and who is a professional who does not have specialized training or qualifications in wastewater surveillance, but who uses surveillance to inform policy and action in their workplace.
Engaged public		An adult ^a who is proficient in English and has relevant lived experience.

Definitions: (1) Professional experience – Paid employment or professional practice in a listed specialization (current
 or former); (2) Graduate or professional degree – Master's or Doctor of Philosophy (PhD) relating to a listed discipline

subgroup.

208 ^aAdult: ≥ 18 years of age

^bSee S2 Table for specializations associated with each discipline.

210

211 We will conduct an all-rounds e-Delphi invitation approach, meaning panellists will be invited to every

round regardless of whether they participated in the previous round [30]. Previous Delphi studies have

commonly ranged from 20 to 30 panellists; however, there is no agreed upon standard [28,31]. Studies

have also shown that the number of panellists and their diversity can impact the quality of outcomes

215 [31]. Therefore, assuming only half of registered panellists will participate in at least one round, we will

aim to recruit at least 50 panellists. Preferably, with at least 8-10 people per discipline subgroup. The

study working group will monitor the distribution of registered panellists based on their demographic

information and will try to distribute appropriately across discipline subgroups and other demographicmarkers.

220 Registration to complete the e-Delphi will open approximately four weeks before the first round, and e-221 Delphi round surveys will be active for roughly 4-6 weeks. The tentative start and end dates for 222 recruitment are 04/11/2024 and 21/02/2025, respectively. Recruitment or retention efforts will occur throughout each e-Delphi round and the four weeks of targeted recruitment before Round 1. Email 223 224 reminders to complete the survey will be sent twice per e-Delphi round. Panellists may also receive up 225 to two reminders to complete the screening questionnaire during the targeted recruitment period. The 226 time required to complete each e-Delphi round survey will be approximately one hour. The estimated 227 time commitment will be provided to all potential panellists during recruitment and before starting the 228 consensus process.

229 Panellist withdrawal or termination

230 Panellists will not be able to withdraw from the study due to the safeguards in place to maintain

anonymity; however, they may choose to stop actively participating in the study at any time.

232 Panellist compensation

Content experts and knowledge users who complete all rounds of the e-Delphi process will receive a
\$100 CAD pre-paid Visa card in compensation for the time required to complete the e-Delphi process.
Engaged members of the public who complete all rounds will be entered to win one of five \$100 CAD
pre-paid Visa cards. Remuneration has been shown to encourage participation and retention from
diverse groups, thereby minimizing bias, and motivating thoughtful, high-quality responses, thereby
improving data quality [32].

239 Procedure

- 240 We will conduct a two-round e-Delphi survey to generate consensus on evaluation criteria (see Fig 2).
- 241 The full survey will be pre-tested and validated before administration. Summaries of Round 1 will be
- compiled for the subsequent round.
- Fig 2. E-Delphi and consensus meeting methodology, methods to achieve consensus on core items for
- 244 **PHES-EF.** Consensus: if \geq 70% of panellist votes fall within the same category (irrelevant, equivocal,
- relevant); no consensus if <70% agreement. Item relevancy: *core item*: highly or moderately
- relevant; *further discussion*: slightly relevant, neutral, slightly irrelevant; *excluded item*: moderately or
- 247 highly irrelevant.
- 248 We will use the Surveylet Delphi platform to administer the survey [33]. The platform facilitates
- interactive dialogue between panellists by allowing them to justify their responses and comment on the
- 250 responses of others. Panellists can also save their progress and answer survey questions across multiple
- 251 sessions. Custom survey pathways will be generated for each panellist discipline subgroup (i.e.,
- 252 panellists from different discipline subgroups will be shown a different collection of candidate items).
- 253 Within each custom discipline subgroup survey pathway, panellists can skip items or self-declare that
- they are not qualified to assess certain candidate items. To ensure the independent evaluation of items,
- the order of candidate items may be randomized for each panellist.
- 256 Individuals will complete the screening questionnaire separate from the e-Delphi platform using
- LimeSurvey [34]. The study working group will review eligibility, and if eligible to participate in the study,
- individuals will receive an individualized link for anonymous participation in the e-Delphi survey. Implied
- consent will be obtained electronically for all panellists using the Surveylet Delphi platform before
- 260 participation in the survey. Panellists who are eligible for compensation or wish to receive
- acknowledgement for their contributions will be able to do so by clicking a link at the end of each survey

- round. The link will bring them to a separate LimeSurvey form that will not be linked to their survey
- 263 responses, allowing for e-Delphi responses to remain anonymous.

264 Defining consensus

- 265 We will use a 7-point Likert scale (1 = highly irrelevant to 7 = highly relevant). Panellist responses will be
- 266 categorised as excluded item (irrelevant: 1–2), further discussion (equivocal: 3–5), or core item
- 267 (relevant: 6–7). Consensus for each item is defined as \geq 70% of the panellist votes falling within the
- 268 same category (1–2, 3–5, or 6–7).
- 269 The approach to use a 7-point fully-labeled Likert scale a higher number of categories compared to
- 270 many studies is informed by the consideration that panellists are professionals and engaged members
- of the public with good cognitive skills; therefore, more categories and labels will be more discriminating
- and reproducible [35–38]. The \geq 70% consensus cut-off a lower cut-off compared to many studies is
- informed by the rapidly evolving nature of environmental surveillance for public health and the wide
- range of disciplines invited to the panel; therefore, a high level of agreement may not occur [35–38].

275 Stopping rules

- 276 When consensus is reached for a given item, no subsequent rounds of ranking for that item will be
- 277 performed. Consensus items that are considered equivocal, in additional to remaining non-consensual
- items, will be further deliberated during the consensus meeting.

279 Round 1

280 Panellists will be invited to rate their level of agreement with candidate items generated from scoping

- 281 review results and consultation with the study executive group. Free-text boxes will be included for
- 282 panellists to provide feedback or identify additional candidate items to be included in the next e-Delphi
- survey round.

284 Round 2

Regardless of whether they participated in the previous round, panellists will be invited to participate in Round 2 of the e-Delphi survey. All panelists will be invited to rate new items and re-rate previous items that did not reach consensus. When re-rating their level of agreement, panellists will be presented with their previous round scores alongside the aggregate group results. Anonymous feedback from Round 1 will also be compiled and presented during Round 2. Any newly suggested items during Round 2 will be deliberated on during the consensus meeting.

291 Summary panelist characteristics

- 292 At the start of the e-Delphi survey, panellists will provide basic demographic details to ascertain their
- discipline subgroup, ensuring they receive the correct list of candidate items. To maintain anonymity,
- this demographic data will not contain identifiable information, and only aggregate responses of 5 or
- 295 more panelists will be analysed. We will report aggregate panel characteristics based on gender,
- discipline group or subgroup (depending on the number of panelists), level and type of experience,
- 297 country income level, and World Bank work regions.

298 Quantitative analysis

We will report response rates, agreement levels, consensus, and stability across rounds using median, range, and IQR for each criterion item. The proportion of panelists who skipped items or selected "not qualified to respond" will be reported. Fisher's exact test will be used to evaluate differences in item ratings by panellist characteristics: primary panellist disciplines (see S2 Table), country income level (high-income versus low-and-middle income), and World Bank world regions (see S3 Table).

304 Qualitative analysis

- 305 Content analyses will be conducted on panelists' feedback to extract common themes and insights. A
- 306 summary will describe the evolution of each criterion across rounds, including reasons for item
- 307 modification or elimination.

308 Grouping of subgroups

- 309 Subgroups with fewer than 5 panellists will be combined with similar groups based on panellist numbers
- and group likeness. Decisions on combined subgroups will be made prior to analyses of the items.

311 Exclusion of data

- 312 All panellists that participate in at least one round will have their responses included in analyses unless
- 313 the study working group determines that a panellist's response warrants being excluded from analyses
- for reasons that are newly developed or not previously recognised.

315 Step 4: Consensus meeting

- 316 In preparation for the consensus meeting, the study working group will categorize the remaining
- 317 candidate items for consideration into a preliminary evaluation framework. This will include merging
- and/or modifying the remaining items based on gathered feedback from the e-Delphi rounds.
- An in-person or hybrid (in-person and online) consensus meeting will be held using the Surveylet Delphi
- 320 survey platform after the results of the e-Delphi have been compiled and analysed. If in a hybrid format,
- 321 Microsoft Teams will be used for messaging and videoconferencing. Video recording and transcription
- 322 services may also be used to aid notetaking. The primary objective will be to achieve expert consensus
- 323 on the final list of items for PHES-EF, through review and discussion of salient items. This process will be
- 324 guided by the empirical evidence that was identified during the scoping reviews, and the opinions

- 325 gathered during the e-Delphi process. The secondary objective will be to discuss publication and
- 326 dissemination strategies for the final evaluation framework.
- 327 Steps to produce the final list of items are as follows: (i) present the results of the e-Delphi exercise
- 328 (name, rationale, and score of each item); (ii) discuss the rationale and relevance for including the items
- in the framework; and (iii) vote on equivocal and non-consensual items. We will invite all members of
- the executive group to participate in the consensus meeting. PHES-EF will be developed based on the
- final list of items that receive consensus during the executive group consensus meeting.

332 Step 5: Dissemination of evaluation framework

333 The dissemination of this evaluation framework will start with the publication of the PHES-EF protocol.

The development of the evaluation framework will be reported in a statement document that will

include the rationale and a brief description of the meeting and the panellists involved. While the

336 publication strategy will be finalized based on the consensus meeting discussion, the preliminary

approach will be to post to OSF, as a preprint, before publication in an open-access peer-reviewed

journal. During this time, we will actively seek feedback from experts, knowledge users, and the public

339 within our networks and via social media. PHES-EF will be made available using an open license (CC-BY-

340 SA-4.0 license).

We will develop a website and may create a Discourse server or explore other social media platforms for ongoing user interactions. Active dissemination approaches will include presenting at relevant scientific conferences, holding webinars, and conducting workshops. Additional dissemination practices will be determined during the consensus meeting.

345 Equity, diversity, and inclusion (EDI)

346 "Equity is defined as the removal of systemic barriers and biases enabling all individuals to have equal opportunity to access and benefit from the program" [39]. To achieve this, members of the research 347 348 team are committed to developing a strong understanding of the systemic barriers faced by individuals 349 from underrepresented groups. "Diversity is defined as differences in race, colour, place of origin, 350 religion, immigrant and newcomer status, ethnic origin, ability, sex, sexual orientation, gender identity, 351 gender expression and age" [39]. Panellists will be asked to provide characteristic information (e.g., 352 country of residence, gender, primary discipline) to assess and help ensure (through more targeted 353 recruitment) a large diversity in perspectives. "Inclusion is defined as the practice of ensuring that all 354 individuals are valued and respected for their contributions and are equally supported" [39]. To help 355 ensure that all research team members are integrated and supported, we will have a code of conduct 356 statement in the executive group Terms of Reference. 357 All study working group members will complete EDI-related Canadian Institutes of Health Research

(CIHR) training [40], and the First Nations Principles of OCAP[®] (Ownership, Control, Access, and
Possession) training for research [41]. The executive group will receive an overview of EDI and OCAP[®]
and will be encouraged and supported to complete such training. Furthermore, staff trained in EDI and
OCAP[®] principles will review the framework from an EDI perspective to ensure appropriate language.

362 Knowledge user and public involvement

The international e-Delphi survey will collect responses from content experts, knowledge users, and engaged members of the public to ensure that all parties involved with or affected by wastewater surveillance are represented in the consensus-building approach. Apart from what has already been described, more detailed recruitment or public engagement methods may be informed by the executive

367 group engaged member of the public. Additionally, to ensure accurate and transparent reporting of

- 368 knowledge user and public involvement throughout the study, we will refer to the Guidance for
- 369 Reporting Involvement of Patients and the Public (GRIPP2) checklist [42]. We will document the
- 370 methods used to engage knowledge users and members of the public, report the impacts and outcomes
- of their engagement, and report on lessons learned from the experience.

372 Acknowledgements

373 E-Delphi participation will not satisfy authorship criteria; however, panellists will have the option to

- 374 receive individual acknowledgement for their contributions to the framework, given they provide
- 375 consent to be named. Individuals who participate in the consensus meeting will have the opportunity to
- be named as authors.

377 Research data management

378 The project will follow an open-science approach and make available the data, code, and materials for

all project stages, including the search strategy, search findings, initial curation of evaluation measures,

380 summary of executive group discussions, e-Delphi processes, and agreement procedures. The

381 management of research data will follow a data management and sharing plan.

All personal data will be managed using LimeSurvey. Personal data will only be collected from individuals during the screening questionnaire, and from panellists who wish to receive compensation and/or individual acknowledgement for their contributions to the evaluation framework. LimeSurvey data management practices are overseen by The Ottawa Hospital (TOH) Research Institute Methods Centre that protects privacy and confidentiality while assisting authorized users in collecting and managing survey data. LimeSurvey is a free, open-source, online survey tool. It has been installed on a TOH server in the TOH Demilitarized Zone and all data collected is stored in a TOH database within the

389 TOH firewall. LimeSurvey provides a streamlined process to build and manage online surveys. Users can

- 390 create and publish surveys, collect responses, view analytics, and export survey data for further
- analysis. Because LimeSurvey is installed outside of the TOH firewall in the TOH Demilitarized Zone,
- 392 surveys can be responded to from anywhere using an internet connection.
- 393 Calibrum's Surveylet Delphi platform will be used to manage e-Delphi responses. It may also be used for
- the consensus meeting. The e-Delphi will only collect anonymous data using a modified Delphi
- approach; meanwhile, the consensus meeting may collect identifiable data using a real-time Delphi
- approach. Surveylet complies with the EU-U.S. Privacy Shield Framework and Swiss-U.S. Privacy Shield
- 397 Framework regarding the collection, use, and retention of personal information. The Surveylet platform

also allows its users to be General Data Protection Regulation (GDPR) compliant. Calibrum's servers are

399 protected by high-end firewall systems and vulnerability scans are conducted regularly. Calibrum uses

- 400 Transport Layer Security (TLS) encryption (also known as HTTPS) for all transmitted data. Surveys are
- 401 also protected with passwords and HTTP referrer checking. All data are encrypted at rest and hosted by
- 402 SSAE-16 SOC II certified third-party data centres. Data on deprecated hard drives are destroyed by U.S.
- 403 DOD methods and delivered to a third-party data destruction service.

404 Study record retention

Any data that is exported from either survey company will be stored securely in a password protected
location within TOH's SharePoint. Upon study completion, all study records will be retained securely for
10 years, then destroyed as per the Ottawa Hospital Research Institute (OHRI) requirements.

408 Ethical considerations

- 409 This study has received ethical approval (20230428-01H) from the Ottawa Health Science Network
- 410 Research Ethics Board (OHSN-REB). Implied consent will be obtained from all panellists prior to

411	participating in the e-Delphi survey. Information about the survey and consent practices will be provided
412	on the first page of the survey. Panellists will be informed that by providing their demographic
413	information and proceeding to the next page, they are providing their implied consent to participate in
414	the study.

- Only anonymous panellist responses will be used for analyses. All panellist data and responses will be
- 416 exported from relevant survey platforms to a shared folder on the Ottawa Hospital SharePoint. The
- 417 shared folder contents will be password protected and only REB-approved study working group
- 418 members from the Ottawa Hospital will have access to the data.

419 Discussion

- 420 This paper outlines the protocol for a two-round e-Delphi study that will be used to develop a
- 421 multinational, multidisciplinary evaluation framework for public health environmental surveillance.
- 422 Developing an evaluation framework will contribute to a better understanding of the ideal scope and
- 423 use cases for wastewater surveillance and improve the reliability and credibility of surveillance data for
- 424 public health officials. An evaluation framework will also help promote the sustainability of wastewater
- 425 surveillance systems by providing an effective performance measurement tool.
- 426 We strove to design the PHES-EF protocol using a rigorous consensus approach. The development of the
- 427 PHES-EF study protocol was informed by ACCORD (ACcurate COnsensus Reporting Document): A
- 428 reporting guideline for consensus methods in biomedicine developed via a modified Delphi [43].
- 429 Surveillance evaluation frameworks are uncommonly developed using a consensus approach. PHES-EF
- 430 will be developed for environmental surveillance and informed by wastewater surveillance during the
- 431 pandemic. However, associated scoping reviews and prior surveillance evaluation frameworks are

- related largely to any public health surveillance system [44]. Therefore, PHES-EF may be applicable for
- 433 evaluating public health surveillance beyond environmental systems.

434 Limitations

- 435 Although the Delphi technique uses a systematic process to generate consensus among a panel of
- 436 experts, the method is not without its limitations. One of the primary concerns when using this
- 437 technique is the ability to implement a truly representative expert panel. Despite our plan to use various
- 438 recruitment methods to enroll panellists from all discipline subgroups with varying demographic
- 439 characteristics, consensus results may ultimately be influenced by the final panel composition [45].
- 440 Another potential limitation in using a modified Delphi approach is the potential influence of controlled
- 441 feedback on the convergence of panellist opinions. Although providing controlled feedback between
- 442 survey rounds aids in generating consensus, this approach may bias consensus results by influencing
- 443 individual panellists' opinions [46].
- Furthermore, due to feasibility and resource limitations, only sources that are available in English will be included in associated scoping reviews. Additionally, only individuals who are proficient in English will be eligible to participate in the e-Delphi survey and consensus meeting. Therefore, the findings of this study may not reflect relevant knowledge that is not available in English.

448 Authors' contributions

- 449 **Conceptualization:** Douglas G. Manuel, Carol Bennett.
- 450 **Funding acquisition:** Douglas G. Manuel.
- 451 Methodology: Douglas G. Manuel, Carol Bennett, Emma Brown, David Moher.
- 452 **Supervision:** Douglas G. Manuel.
- 453 Writing original draft: Douglas G. Manuel, Carol Bennett, Emma Brown.
- 454 Writing review and editing: David Moher, Matthew J. Wade, Yoni Freedhoff, David L. Buckeridge,
- 455 Farah Ishtiaq, Sarah Funnell.

456 Acknowledgements

- 457 We thank Mary Jessome (PhD candidate, University of British Columbia) for her contributions to the
- 458 study as a past executive group member with expertise in social sciences. We would also like to thank
- 459 Peter Farrell (Research Librarian, University of Ottawa) for developing the associated scoping review
- 460 search strategies for this study.
- 461 PHES-EF Executive Group (ordered alphabetically according to surname except for the first two462 authors):
- 463 Douglas G. Manuel^{*} (0000-0003-0912-0845), David L. Buckeridge^{*} (0000-0003-1817-5047), Yoni
- 464 Freedhoff (0000-0001-9391-3455), Sarah Funnell, Bernd M. Gawlik (0000-0002-4406-5362), Farah
- 465 Ishtiaq (0000-0002-6762-7014), Amy E. Kirby, Kerrigan McCarthy (0000-0001-8958-9795), David Moher

- 466 (0000-0003-2434-4206), Beate Sander (0000-0003-2128-9133), Jeremy Veillard, Matthew J. Wade
- 467 (0000-0001-9824-7121)
- 468 *Co-chairs of PHES-EF Executive Group

469 Supporting information

- 470 **S1 Table. Executive group composition.** Note: The regional classifications are used by the World Bank
- 471 for organizing economic data, development projects, and policy analysis. These classifications are not
- 472 strictly geographical but also consider economic and developmental similarities and relationships.
- 473 S2 Table. Executive group and e-Delphi panellist disciplines. The list of specializations or related
- 474 concepts is not extensive.
- 475 S3 Table. World region classifications. Note: The regional classifications are used by the World Bank for
- 476 organizing economic data, development projects, and policy analysis. These classifications are not
- 477 strictly geographical but also consider economic and developmental similarities and relationships.

478 **References**

- Woodward G, Manuel DG, Goel V. Developing a balanced scorecard for public health. Toronto:
 Institute for Clinical Evaluative Sciences; 2004.
- McClary-Gutierrez JS, Mattioli MC, Marcenac P, Silverman AI, Boehm AB, Bibby K, et al. SARS-CoV-2
 Wastewater Surveillance for Public Health Action. Emerg Infect Dis. 2021;27: e210753.
 doi:10.3201/eid2709.210753
- Diamond MB, Keshaviah A, Bento AI, Conroy-Ben O, Driver EM, Ensor KB, et al. Wastewater
 surveillance of pathogens can inform public health responses. Nat Med. 2022;28: 1992–1995.
 doi:10.1038/s41591-022-01940-x
- 487 4. Ettore Zuccato, Chiara Chiabrando, Sara Castiglioni, Renzo Bagnati, Roberto Fanelli. Estimating
 488 Community Drug Abuse by Wastewater Analysis. Environ Health Perspect. 2008;116: 1027–1032.
 489 doi:10.1289/ehp.11022

- 490 5. Hrudey SE, Bischel HN, Charrois J, Chik AHS, Conant B, Delatolla R, et al. Wastewater Surveillance for
 491 SARS-CoV-2 RNA in Canada. FACETS. 2022;7: 1493–1597. doi:10.1139/facets-2022-0148
- 492 6. World Health Organization. Environmental surveillance for SARS-COV-2 to complement public health
 493 surveillance. Geneva, Switzerland: World Health Organization; 2023 Sep p. 55. Available:
 494 https://www.who.int/publications/i/item/9789240080638
- 495 7. Manuel DG, Delatolla R, Fisman DN, Fuzzen M, Graber T, Katz GM, et al. The Role of Wastewater
 496 Testing for SARS-CoV-2 Surveillance. Ontario COVID-19 Science Advisory Table; 2021 Aug.
 497 doi:10.47326/ocsat.2021.02.40.1.0
- 498 8. Woytus A. What's in the wastewater? In: St. Louis [Internet]. 22 Sep 2021 [cited 1 Jun 2023].
 499 Available: https://www.stlmag.com/api/content/3f60d498-1b26-11ec-a8ea-1244d5f7c7c6/
- D'Aoust PM, Towhid ST, Mercier É, Hegazy N, Tian X, Bhatnagar K, et al. COVID-19 wastewater
 surveillance in rural communities: Comparison of lagoon and pumping station samples. Sci Total
 Environ. 2021;801: 149618. doi:10.1016/j.scitotenv.2021.149618
- 503 10. Shah S, Gwee SXW, Ng JQX, Lau N, Koh J, Pang J. Wastewater surveillance to infer COVID-19
 504 transmission: A systematic review. Sci Total Environ. 2022;804: 150060.
 505 doi:10.1016/j.scitotenv.2021.150060
- Liu P, Ibaraki M, VanTassell J, Geith K, Cavallo M, Kann R, et al. A sensitive, simple, and low-cost
 method for COVID-19 wastewater surveillance at an institutional level. Sci Total Environ. 2022;807:
 151047. doi:10.1016/j.scitotenv.2021.151047
- Larsen DA, Wigginton KR. Tracking COVID-19 with wastewater. Nat Biotechnol. 2020;38: 1151–1153.
 doi:10.1038/s41587-020-0690-1
- 511 13. Gretchen Vogel. Signals from the sewer—Measuring virus levels in wastewater can help track the
 512 pandemic. But how useful is that? Science. 9 Mar 2022375. Available:
 513 https://www.science.org/content/article/pandemic-signals-sewer-what-virus-levels-wastewater
- 513https://www.science.org/content/article/pandemic-signals-sewer-what-virus-levels-wastewater-514tell-us. Accessed 1 Jun 2023.
- 14. Manuel D, Amadei CA, Campbell JR, Brault J, Zierler A, Veillard J. Strengthening Public Health
 Surveillance Through Wastewater Testing: An Essential Investment for the COVID-19 Pandemic
 and Future Health Threats. World Bank Group. The World Bank; 2022 pp. 1–67.
- 518 15. Fynn JF, Hardeman W, Milton K, Jones AP. A scoping review of evaluation frameworks and their
 applicability to real-world physical activity and dietary change programme evaluation. BMC Public
 Health. 2020: 1000. doi:https://doi.org/10.1186/s12889-020-09062-0
- 16. Nilsen P. Making sense of implementation theories, models, and frameworks. Implement Sci 30.
 2020; 53–79.
- 523 17. Calba C, Goutard FL, Hoinville L, Hendrikx P, Lindberg A, Saegerman C, et al. Surveillance systems
 524 evaluation: a systematic review of the existing approaches. BMC Public Health. 2015;15.
 525 doi:10.1186/S12889-015-1791-5

526 18. Andrews JR, Yu AT, Saha S, Shakya J, Aiemjoy K, Horng L, et al. Environmental Surveillance as a Tool 527 for Identifying High-risk Settings for Typhoid Transmission. Clin Infect Dis. 2020;71: S71–S78. 528 doi:10.1093/cid/ciaa513 529 19. Porta M. A Dictionary of Epidemiology. A Dictionary of Epidemiology. Oxford University Press; 2016. 530 Available: 531 https://www.oxfordreference.com/display/10.1093/acref/9780199976720.001.0001/acref-532 9780199976720 533 20. Spranger J, Homberg A, Sonnberger M, Niederberger M. Reporting guidelines for Delphi techniques 534 in health sciences: A methodological review. Z Evid Fortbild Qual Gesundh Wesen ZEFQ. 172: 1–11. 535 21. How epidemiology has shaped the COVID pandemic. Nature. 2021;589: 491–492. 536 doi:10.1038/d41586-021-00183-z 537 22. Devine D, Gaskell J, Jennings W, Stoker G. Trust and the Coronavirus Pandemic: What are the 538 Consequences of and for Trust? An Early Review of the Literature. Polit Stud Rev. 2021;19: 274-539 285. doi:10.1177/1478929920948684 540 23. World Health Organization. Defining collaborative surveillance: a core concept for strengthening the global architecture for health emergency preparedness, response, and resilience (HEPR). Geneva: 541 542 World Health Organization; 2023. Available: https://iris.who.int/handle/10665/367927 543 24. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for Scoping 544 Reviews (PRISMA-ScR): Checklist and Explanation. Ann Intern Med. 2018;169: 467–473. 545 doi:10.7326/M18-0850 546 25. Nasa P, Jain R, Juneja D. Delphi methodology in healthcare research: How to decide its appropriateness. World J Methodol. 2021;11: 116–129. doi:10.5662/wjm.v11.i4.116 547 548 26. Shang Z. Use of Delphi in health sciences research: A narrative review. Medicine (Baltimore). 549 2023;102: e32829. doi:10.1097/MD.000000000032829 550 27. Guidance on Conducting and REporting DElphi Studies (CREDES) in palliative care: 551 Recommendations based on a methodological systematic review. 2017;31: 684–706. 552 28. Niederberger M, Spranger J. Delphi Technique in Health Sciences: A Map. Front Public Health. 553 2020;8: 457. doi:10.3389/fpubh.2020.00457 554 29. Schmalz U, Spinler S, Ringbeck J. Lessons Learned from a Two-Round Delphi-based Scenario Study. 555 MethodsX. 2021;8: 101179. doi:10.1016/j.mex.2020.101179 556 30. Boel A, Navarro-Compán V, Landewé R, van der Heijde D. Two different invitation approaches for 557 consecutive rounds of a Delphi survey led to comparable final outcome. J Clin Epidemiol. 2021;129: 31-39. doi:10.1016/j.jclinepi.2020.09.034 558 559 31. Gattrell WT, Hungin AP, Price A, Winchester CC, Tovey D, Hughes EL, et al. ACCORD guideline 560 for reporting consensus-based methods in biomedical research and clinical practice: a study 561 protocol. 2022;7: 3.

- 562 32. Permuth-Wey J, Borenstein AR. Financial Remuneration for Clinical and Behavioral Research 563 Participation: Ethical and Practical Considerations. Ann Epidemiol. 2009;19: 280–285. 564 doi:10.1016/j.annepidem.2009.01.004 565 33. Surveylet. United States: Calibrum; Available: https://calibrum.com 566 34. LimeSurvey: An open source survey tool. Hamburg, Germany: LimeSurvey GmbH; Available: 567 https://www.limesurvey.org 568 35. Diamond IR, Grant RC, Feldman BM, Pencharz PB, Ling SC, Moore AM, et al. Defining consensus: A 569 systematic review recommends methodologic criteria for reporting of Delphi studies. J Clin 570 Epidemiol. 2014;67: 401–409. doi:10.1016/j.jclinepi.2013.12.002 571 36. Belton I, MacDonald A, Wright G, Hamlin I. Improving the practical application of the Delphi method in group-based judgment: A six-step prescription for a well-founded and defensible process. 572 573 Technol Forecast Soc Change. 2019;147: 72–82. doi:10.1016/j.techfore.2019.07.002 37. Weijters B, Cabooter E, Schillewaert N. The effect of rating scale format on response styles: The 574 575 number of response categories and response category labels. Int J Res Mark. 2010;27: 236–247. 576 doi:10.1016/j.ijresmar.2010.02.004 38. Lange T, Kopkow C, Lützner J, Günther K-P, Gravius S, Scharf H-P, et al. Comparison of different 577 578 rating scales for the use in Delphi studies: different scales lead to different consensus and show 579 different test-retest reliability. BMC Med Res Methodol. 2020;20: 28. doi:10.1186/s12874-020-0912-8 580 581 39. Government of Canada. Best Practices in Equity, Diversity and Inclusion in Research. In: New 582 Frontiers in Research Fund [Internet]. 22 Jun 2021. Available: https://www.sshrccrsh.gc.ca/funding-financement/nfrf-fnfr/edi-eng.aspx 583 584 40. Government of Canada. Introduction to GBA Plus. 31 Mar 2021 [cited 1 Jun 2023]. Available: 585 https://women-gender-equality.canada.ca/gbaplus-course-cours-586 acsplus/eng/mod00/mod00 01 01.html 587 41. The First Nations Information Governance Centre. The First Nations Principles of OCAP®. In: The 588 First Nations Information Governance Centre [Internet]. 1 Jun 2023 [cited 1 Jun 2023]. Available: 589 https://fnigc.ca/ocap-training/ 590 42. Staniszewska S, Brett J, Simera I, Seers K, Mockford C, Goodlad S, et al. GRIPP2 reporting checklists:
- 42. Staniszewska S, Brett J, Simera I, Seers K, Mockford C, Goodlad S, et al. GRIPP2 reporting checklists:
 tools to improve reporting of patient and public involvement in research. BMJ. 2017;358: j3453.
 doi:10.1136/bmj.j3453
- 43. Gattrell WT, Logullo P, Zuuren EJ van, Price A, Hughes EL, Blazey P, et al. ACCORD (ACcurate
 COnsensus Reporting Document): A reporting guideline for consensus methods in biomedicine
- 595 developed via a modified Delphi. PLOS Med. 2024;21: e1004326.
- 596 doi:10.1371/journal.pmed.1004326

- 44. Yang L, Weston C, Cude C, Kincl L. Evaluating Oregon's occupational public health surveillance
 system based on the CDC updated guidelines. Am J Ind Med. 2020;63: 713–725.
 doi:10.1002/ajim.23139
- 45. Winkler J, Moser R. Biases in future-oriented Delphi studies: A cognitive perspective. Technol
 Forecast Soc Change. 2016;105: 63–76. doi:10.1016/j.techfore.2016.01.021
- 46. Goodman CM. The Delphi technique: a critique. J Adv Nurs. 1987;12: 729–734. doi:10.1111/j.1365 2648.1987.tb01376.x

604




Figure 1




Figure 2