

Impacts of Global School Feeding Programmes on Children's Health and Wellbeing Outcomes: A Scoping Review

Amy Locke^{1*}, Michaela James¹, Hope Jones¹, Rachel Davies¹, Francesca Williams¹, Sinead Brophy¹.

Affiliations

1 National Centre for Population Health and Wellbeing, School of Medicine, Swansea University, Swansea, United Kingdom.

* Corresponding author

Email: Amy Locke 968097@swansea.a.c.uk

Abstract

Background: School feeding programmes (SFP) are one of the world's most extensive interventions to alleviate child poverty, food insecurity and malnutrition. Schools are ideal settings to promote healthy eating habits early in life since children have access to at least one main meal per day at school. However, there is a lack of clear evidence for the effectiveness of school feeding programmes on child health outcomes. Additionally, there is considerable debate on whether feeding programmes should be universal or targeted with countries taking different approaches. This review will therefore explore global research on school feeding programmes and consider different health outcomes through targeted or universal policies.

Methods: We conducted a search across four electronic databases. These studies investigated the impacts of school feeding programmes on children's health outcomes. 207 papers were identified with 76 studies subjected to full text screening.

Results: A total of 42 papers were included in the final review. All papers were published within the past fifteen years (2009-2024) and included thirteen countries in total. SFP were associated with child weight being more in the healthy range. Targeted SFP were negatively associated with stigma.

Conclusions: Universal SFP were effective at improving children's health outcomes such as healthy weight, improved behaviour and social support. Overall, both targeted and universal SFP positively impact children's health outcomes and address health disparities.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- The review mapped both universal and targeted provisions, providing a broad overview of the existing literature.
- The review included an appraisal of the methodological quality of the included studies. Only English studies were included.
- Only studies published in English were included, potentially excluding relevant research in other languages.
- The study did not perform a formal assessment of publication bias, which could impact the interpretation of the results.

Introduction

School feeding programmes (SFP) are one of the world's most extensive interventions to alleviate child poverty, food insecurity and malnutrition [1]. In 2022, 418 million children worldwide received a free or subsidised school meal as part of a SFP [2]. In recent years, growing concerns have arisen regarding the nutritional health of children [3]. With increased sugar intake, increases in childhood obesity and socio-economic factors affecting millions of families worldwide, there is emphasis on the need for interventions to reduce negative health outcomes [4]. Schools present an opportunity to promote healthy eating habits among children and can serve as a preventative measure to negative health implications [5].

From a public health perspective, schools are ideal settings to promote healthy eating behaviours early in life since children have access to at least one main meal per day at school in most schools [6]. There is a large body of literature dedicated to exploring the impacts of SFP on educational outcomes, noting positive outcomes, specifically in relation to key stage attainment and attendance [7, 8]. However, literature relating to the impacts of SFP on children's health outcomes are limited, and even less research considers SFP as a preventative tool for public health. Research by Chaudhary et al [9] demonstrated that initiatives targeting food and nutrition within schools could enhance dietary habits, promote healthy eating, and impact body measurements positively. Noting that interventions with a focus on meal provision, healthy eating promotion and food literacy are most effective in improving children's health outcomes.

It is well documented that good nutrition plays a vital role in maintaining good health throughout the life course, preventing malnutrition in all forms, and decreasing the risk of non-communicable diseases, such as cardiovascular disease, diabetes, and some cancers [10]. Increases in processed foods and changes in lifestyles have led to a shift in dietary patterns, with individuals opting for cheaper food options and convenience foods [11]. With the rise in

poor dietary habits, there is a negative impact on health outcomes. This has become a public health concern, specifically in relation to rising childhood obesity. Figures from 2022 indicate childhood obesity has reached an all-time high with an estimated 390 million children and adolescents aged 5–19 years being overweight or living with obesity. Of this, 37 million children under age five were classified as overweight [12, 13]. To date, no country is on track to curb the obesity crisis [14]. Moreover, in 2019, over 161 million children under five years of age were reported as having a nutritional deficiency globally from undernourishment [15]. Nutritional deficiencies have been attributed to childhood wasting, stunting, infections, as well as cognitive and behavioural disorders [16, 17, 18].

Health disparities due to poor nutritional intake is largely seen in poor economic groups [19], and often arise in early childhood, resulting in on-going implications into adulthood [20]. According to the World Health Organisation (WHO) [5], childhood and adolescence is a critical period for promoting nutritional health and reducing the risk of negative health implications and calls for policy reform to SFP to address diet quality in children are becoming widespread [21].

Despite the large potential health benefits of SFP, they have proven difficult to evaluate given the diversity in implementation policies across nations. For example, Europe alone has a range of SFP policies from universal to targeted approaches [22]. Universal systems refer to a provision open to all students, regardless of socio-economic status [23]. This system is well established in countries such as Sweden, where all children in both primary and secondary education (ages 5 - 16) have access to at least one free school meal (FSM) a day and is regarded as a symbol of national welfare [24]. However, there is also diversity among policies relating to universal systems, whereby countries offer universal school meals to children in particular year groups. For instance, Latvia offers free meals to children in grades

one to four (ages 7 -9), and Lithuania provides FSM from preschool to first grade (up to age 7) [22].

Targeted programmes on the other hand, target children from lower socio-economic backgrounds, usually using a means-tested eligibility system which is based on parental income [25]. This approach is implemented in Poland, Slovenia, and parts of the United Kingdom [26]. This approach, whilst beneficial to some families, has its own limitations, e.g. a threshold cut off means families slightly above the eligibility criteria may still be living in poverty but are able to access FSM and the stigma associated with being eligible may lead to a lack of uptake [23].

There is considerable debate among government policymakers as to whether SFP should be targeted (benefit eligibility depends on family income) or universal (benefits are provided to all students with no eligibility criteria) [27, 28]. However, given the notable gap in health-related literature, it is difficult to establish which system, if any, is most effective at addressing health disparities.

To our knowledge, no study has systematically synthesised information relating to the effectiveness of SFP on children's health outcomes. Therefore, this review aims to address the following questions; 1. To what extent does the existing literature examine the impacts of school feeding programmes on the physical, emotional, psychological, and social health of school children globally and 2. What challenges, including factors influencing uptake, are reported in the literature regarding the implementation of free school meal provision and 3. Does the effectiveness of SFP on children's health outcomes vary depending on the SFP policy (targeted or universal).

For the purpose of this scoping review, children's health and wellbeing will be categorised as physical health, emotional and psychological health, and social health.

Methods

Review Design

We conducted a scoping review of both qualitative and quantitative research to establish evidence of the effectiveness of school feeding programmes in improving children's health outcomes globally. This scoping review was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) guidelines [29]. The protocol for the review was registered on the Open Science Framework (OSF) [30]. A preliminary search of MEDLINE, the Cochrane Database of Systematic Reviews was conducted, and no current or underway systematic reviews or scoping reviews on the topic were identified. The study findings were summarised through a narrative synthesis.

Researcher (AL) extracted study data from full text papers. Each extraction was reviewed by a second reviewer. Any conflicts were presented to and resolved through group discussion.

Search Strategy

Four electronic databases – Medline, PubMed, Web of Science and Google Scholar (including Science Direct, Web of Science, ProQuest Central, EBSCO, CINAHL) – were searched to identify relevant research papers in English. Studies published from 2009 were included, with most papers from 2020 to 2023. The literature search was conducted by lead author (AL) in December 2023 with screening taking place in January and February 2024.

Additionally, the reference lists of all included sources were also screened for additional studies. Reports and case studies were also sourced and included in the final screening.

Search terms included: Universal Free School Meals OR School Feeding Programmes OR Free School Meals and Children's Health OR Free School Meals and Children's Wellbeing OR School Feeding Programmes OR Challenges with School Feeding Programmes OR School Meal Provision. Additionally, each string was searched using the phrase 'Impacts of.'

Eligibility

Inclusion Criteria

Studies exploring school feeding programmes were included if they met the following outcomes: (1) Children’s physical health – Illness, physical activity, and/or diet dietary intake. (2) Children’s emotional and psychological health– emotion regulation, resilience, self-esteem, and behaviour. (3) Children’s social health – social engagement/participation, building and maintaining relationships. (4) Studies published within the past fifteen years to ensure research is up-to-date and relevant. (5) Primary and secondary school aged children (ages 5 to 16). (6) Studies investigating the implementation of universal feeding programmes and its potential challenges. Both experimental and quasi-experimental study designs including randomised controlled trials, non-randomised controlled trials, before and after studies and interrupted time-series studies. Descriptive observational study designs including case series, individual case reports and descriptive cross-sectional studies were also included. Qualitative studies were also included that focused on children’s self-assessed health and wellbeing and perceptions of school feeding programmes.

Exclusion Criteria

Studies were excluded if they met the following criteria: (1) Children below or above school age (below age 5 or above 18). (2) Studies with a focus on SFP and academic outcomes – performance, attainment, or attendance without a health element. (3) Studies that only looked at parental funded meals/packed lunches. (4) Studies not in English and, (5) Studies published prior to 2009. Additionally, any studies that did not focus on the impacts of school meal provisions on children’s health, unless considering factors influencing uptake, were excluded.

Evidence Selection

Selected papers were uploaded to Covidence systematic review software [31], by the lead author (AL). After removing 30 duplicates, the remaining titles and abstracts (177) were screened against the selection criteria by authors (AL, MJ, HJ, RD, SB). Remaining studies undertook full-text screening by reviewers (AL, MJ, HJ, RD, FW). Any conflicts during the review process were resolved through group discussion. One paper was removed during data extraction as it became apparent that it was an abstract for a conference presentation and no full text was available.

Quality Assessment

Methodological quality of included studies was assessed using the Mixed Methods Appraisal Tool (MMAT) [32]. The MMAT is a quality appraisal tool developed to evaluate the methodological quality of empirical studies. The MMAT was first published in 2009. Since then, it has been validated in several studies testing its interrater reliability, usability, and content validity [33]. The tool can appraise five different categories of study designs: qualitative, randomised controlled trial, non-randomised, quantitative descriptive and mixed methods studies. As our review includes numerous study designs, this tool was deemed as appropriate. Each section is assessed based on five criteria for methodological quality, offering three responses: "Yes" (indicating the criterion is satisfied), "No" (indicating the criterion is not satisfied), and "Can't tell" (suggesting insufficient information to assess). Studies are categorised as low quality if they receive a "Yes" response for two or less questions, moderate quality for 3 questions, and high quality for four or more questions.

Four papers included in the review were impact reports and opinion pieces, and did not fit under an appraisal category within the MMAT, therefore the modified version of the Joanna Briggs Institute (JBI) critical appraisal checklist for text and opinion was used to

quality assess these papers [34]. Appraisal is subjected to the same response criteria as above; "Yes" (indicating the criterion is satisfied), "No" (indicating the criterion is not satisfied), and "Can't tell" (suggesting insufficient information to assess). No papers were excluded based on the results of the quality assessment. Quality assessment was conducted by one reviewer (AL) and checked by a second reviewer(s).

Data Extraction and Analysis

Relevant information from selected studies was extracted using a template (this is provided in the Supplementary Information) developed by lead author (AL). The data collection template included information on; author(s), year of publication, Country, type of meal provision, participants, study design, and key findings, gaps/limitations identified and recommendations for future research. The development of the template was guided by the JBI manual for evidence synthesis [35] and piloted on Covidence using two papers.

Data analysis consisted of a narrative synthesis due to heterogeneity of both study designs and outcomes among included studies, making statistical meta-analysis impractical. Analysis and synthesis were conducted following recommendations from the Guidance of the Conduct of Narrative Synthesis in Systematic Reviews [36]. Specifically, a grouping strategy was utilised to arrange findings into similar theme groups.

Results

Overview of studies

The literature search yielded a total of 207 titles, and these were imported to Covidence for title and abstract screening. After duplicates were removed (n= 30), 177 titles and abstracts remained for screening. Of these, 101 were excluded for not meeting the inclusion criteria, i.e., wrong outcomes measured. A further 34 studies did not meet the eligibility criteria and were excluded at full text screening phase, with a majority vote for wrong outcomes studied. A total of 42 papers were included in the final review. Figure 1 illustrates the screening process.




Fig 1. PRISMA Diagram

Quality of Included Papers

The quality assessment scores of included studies are shown in tables 1 and 2. Studies assessed with Mixed Methods Appraisal Tool (MMAT) [32] were categorised as high overall. 23 studies were categorised as having high methodological quality, six were classed as moderate and nine as being of low quality.

Study	Overall Score
Altindag et al. [57]	Moderate
Batista et al. [37]	Moderate
Bethmann and Cho [48]	High
Cardoso et al. [62]	High
Jessiman et al [65]	Low
Chambers et al. [63]	High
Chambers et al. [74]	High
Colombo et al. [38]	High
Davis et al. [57]	Low
Evans [39]	High
Garton et al. [73]	High
Goel et al. [55]	Moderate
Goodchild et al. [40]	Moderate
Guio [22]	High
Hecht [59]	High
Holford and Rabe [49]	Low

Holford and Rabe [50]	High
Horta et al. [41]	High
Illøkken et al. [64]	Low
James [42]	Low
Carlisle et al. [58]	High
Long et al. [43]	High
Lundborg et al. [20]	High
Mauer et al. [66]	High
McKelvie-Sebileau 1 et al. [67]	High
Meier et al. [54]	Low
Neervoort et al. [52]	Low
Parnham et al. [51]	High
Parnham et al. [44]	Low
Rahim et al. [71]	High
Sahota et al. [68]	Moderate
Spence et al. [45]	Moderate
Taylor et al. [60]	Low

Vik et al. [6]	High
Vik et al. [53]	High
Yamaguchi et al. [61]	High
Yang et al. [46]	High
Zailani et al. [47]	High

Table 1: MMAT Quality Assessment results

Using the modified version of the Joanna Briggs Institute (JBI) critical appraisal checklist for text and opinion [34], three papers were categorised as high, and the other was categorised as moderate. The three papers categorised as high [24, 69, 72], used a variety of methodologies or a strong evidence base to support their findings/opinion. The paper rated moderate [70] included less peer reviewed evidence to support their findings but methodological quality was satisfactory.

Paper	Overall score
Harper and Wood [69]	High
Persson and Fjellström [24]	High
Oostindjer et al. [72]	High

Kitchen et al. [70]	Moderate
---------------------	----------

Table 2: JBI critical appraisal checklist for text and opinion results.

Characteristics of Included Studies

Characteristics of the included studies are summarised in table 3. 11 studies were cross-sectional [37-47]. Five studies had a difference-in-difference design [20, 49-51]. Three were non-randomised control trials [6, 52, 53]. Four were quantitative descriptive [54-57]. Four were mixed methods [58-61]. Six were qualitative [62, 64-68]. Six were reports and/or opinion pieces [22, 24, 70-72]. One was a rapid narrative review [73] and two were case studies [63, 74].

Twenty-three papers were based on universal school feeding programmes [20, 24, 38, 40, 43, 45, 48, 49, 50, 51, 52, 53, 56, 57, 59, 60, 61, 63, 65, 66, 67, 70, 73]. Fifteen had targeted SFP systems [6, 37, 41, 42, 44, 46, 47, 54, 62, 64, 66, 68, 69, 72, 74]. Four papers were categorised as ‘other’. Papers were categorised as other if they looked at breakfast [55], compared both universal and targeted systems [71], compared packed lunch to school meals [39] or were not specified [22].

Nine papers included children in all school ages (primary and secondary) [24, 43, 44, 46, 56, 57, 59, 68, 70], Twelve papers included children just in primary schools [39, 40, 41, 45, 49, 50, 51, 52, 55, 61, 69, 71], Ten papers included children just in secondary schools [6, 38, 42, 53, 62, 64, 65, 66, 67, 74]. Two papers had an overlap of different age groups (children aged 4-14 [37] and children aged 9-16 [48]). One study looked at the impacts of SFP across the life course (from birth to 50+) [20]. One paper included school staff [60]. Five papers did not explicitly state age groups [22, 54, 63, 65, 72].

Seventeen papers were from UK (England n=11 [39, 40, 42, 45, 49, 50, 58, 65, 68, 70, 71], Scotland n= 2 [63, 74], England and Scotland n=1 [51], UK wide n=3 [44, 46, 69]. Six papers were from the USA [43, 54, 55, 56, 59, 60], three papers were from Sweden [20, 24, 38], four from Norway [6, 53, 64, 65], one from Portugal [62] one was Europe wide [22]. Two were from New Zealand [67, 73], two from Brazil [37, 41], two from South Korea [47, 48], two from Africa (Kenya [52] and Nigeria [47]), one from Japan [71] and one was cross national [72]. Figure 2 demonstrates the countries of the included papers.

BRAZIL
 JAPAN
 KENYA
 NEW ZEALAND
 NIGERIA
 NORWAY
 PORTUGAL

 REPUBLIC OF KOREA
 SWEDEN
 UNITED KINGDOM
 UNITED STATES OF AMERICA




Fig 2. Map showing countries included in review.

Author	Country	Provision Type	Participant School year	Design
Altindag et al. [57]	South Korea	Universal	All	Empirical analysis

Batista et al. [37]	Brazil	Targeted	Aged 4-14	Cross-Sectional
Bethmann and Cho [48]	South Korea	Universal	Aged 9-16	Difference-in-Difference
Cardoso et al. [62]	Portugal	Targeted	Secondary	Qualitative
Jessiman et al [65]	London	Universal	Secondary	Mixed Methods
Chambers et al. [63]	Scotland	Universal	N/A	Qualitative Case Study
Chambers et al. [74]	Scotland	Targeted	Secondary	Case Study
Colombo et al. [38]	Sweden	Universal	Secondary	Cross-sectional
Davis et al. [56]	Georgia-USA	Universal	All	Descriptive
Evans [39]	England	Other	Primary	Cross-sectional - Observational
Garton et al. [73]	New Zealand	Universal	Not specified	Rapid Narrative Review
Goel et al. [55]	USA -	Other	Primary	Observational

	Virginia			
Goodchild et al. [40]	England	Universal	Primary	Cross-sectional - Descriptive
Guio [22]	Europe	Other	N/A	Policy Recommendation
Harper and Wood [69]	UK Wide	Targeted	Primary	Report
Hecht [59]	USA	Universal	All	Mixed methods
Holford and Rabe [49]	England	Universal	Primary	Difference-in-Difference
Holford and Rabe [50]	England	Universal	Primary	Difference-in-Difference
Horta et al. [41]	Brazil	Targeted	Primary	Cross-sectional – Analytical
Illøkken et al. [64]	Norway	Targeted	Secondary	Qualitative
James [42]	England	Targeted	Secondary	Cross-sectional - Descriptive
Carlisle et al. [58]	London	Universal	Not specified	Qualitative
Kitchen, et al.	England	Universal	All	Impact report

[70]				
Long et al. [43]	USA	Universal	All	Cross-sectional
Lundborg et al. [20]	Sweden	Universal	Birth to 50+	Difference-in-Difference
Mauer et al. [66]	Norway	Targeted	Secondary	Qualitative
McKelvie-Sebileau et al. [67]	New Zealand	Universal	Secondary	Qualitative
Meier et al. [54]	USA	Targeted	N/A	Quantitative Exploratory
Neervoort et al. [52]	Kenya	Universal	Primary	Non-randomised Control Trial
Oostindjer et al. [72]	Cross-National	Other	N/A	Discussion/Essay
Parnham et al. [51]	England and Scotland	Universal	Primary	Difference-in-Difference
Parnham et al. [44]	UK	Targeted	All	Cross-sectional
Persson & Fjellström [25]	Sweden	Universal	All	Article piece

Rahim et al. [71]	England	Universal & Targeted	Primary	Government Report
Sahota et al. [68]	England	Targeted	All	Qualitative
Spence et al. [45]	England	Universal (Infant)	Primary	Cross-sectional
Taylor et al. [60]	Vermont, USA	Universal	Staff	Mixed methods
Vik et al. [6]	Norway	Targeted	Secondary	Non-randomised Control Trial
Vik et al. [53]	Norway	Universal	Secondary	Non-randomised Control Trial
Yamaguchi et al. [61]	Japan	Universal	Primary	Mixed methods
Yang et al. [46]	UK	Targeted	All	Cross-sectional
Zailani et al. [47]	Nigeria	Targeted	Primary	Cross-sectional

Table 3: Characteristics of the included studies.

Overall studies that investigated universal feeding programmes appear to have the most significant positive outcomes on children’s health, specifically in relation to behaviour, health, and socio-economic outcomes. Whilst targeted systems adequately address food

insecurity and child hunger, children eligible for targeted provision had a higher probability of experiencing poor mental health and stigma. Findings are discussed in more detail below.

Physical Health: BMI and Body Weight Outcomes

Five papers investigated the impacts of SFP on children's BMI and weight outcomes. Two with targeted provision [6, 37] and three with a universal system [48, 50, 56]. Studies had contradicting results.

Batista et al, [37], investigated the prevention of overweight children in Brazil through targeted SFP. The study which utilised a cross-sectional design and included 7,017 participants from 21 schools found that being overweight was prevalent in 30.6% of children and being underweight was present in 1.9%. This study highlighted that a large proportion of the foods served were ultra-processed, recommending a re-evaluation of food standards within schools. Researchers noted an inability to observe extracurricular activities relating to food consumption and physical activity which may also influence children being overweight.

Bethman and Cho [48], investigated the impacts of reintroducing universal free school meals (UFSM) in South Korea after the policy had been abolished a year prior. Findings indicated the removal of the policy had adverse effects on children's BMI outcomes. The reintroduction of the policy on the other hand reversed these impacts and saw more children with healthier weights. However, the sample size of the study is not explicitly stated and may not represent the intended population.

Holford and Rabe [50], suggested that longer exposure to a universal infant SFP positively influenced children's weight outcomes. The study included 153,522 primary school children aged between four and seven and used a difference-indifference analysis to measure children's weight outcomes over time. Findings concluded that children are more likely to be

a healthy weight after exposure to the provision over the course of the first year of school.

Though participants were very young, it is recommended that participants are tracked over a longer time period to see if this is a lasting effect.

A study by Vik et al, [6] conducted in a secondary school in Norway, investigated changes in children's dietary habits, BMI and waist circumference modulated by socio economic status. The study conducted a non-randomised control trial consisting of two groups, non-exposure to SFP and exposure to SFP under a targeted system. Findings suggested free school meals (FSM) increased the intake of healthy foods, particularly among children with lower socio-economic status, leading to an increase in the Healthy Food score of the intervention group between baseline and follow-up. However, children in the intervention group had a significant increase in BMI, though it is not explicitly stated if the children were underweighting prior to the trial. The study concluded that interventions to promote healthy eating are more effective in lower socio-economic groups if it is free or at reduced price.

Davis et al. [56] saw similar results when they examined the effects of a universal SFP in a deprivation area in the USA. Reporting that BMI increased in provision schools versus non provision schools. They further stated that children in provision schools had a higher BMI on average.

In conclusion, there is no evidence that SFP lower BMI, but children are more likely to be a healthy weight. Factors such as the prevalence of ultra-processed foods, socio-economic status, and policy reintroduction played significant roles in these outcomes. Longer exposure to SFP, particularly universal programs, was associated with healthier weight outcomes in younger children, though lasting effects require further study.

Nutritional Intake

Ten papers [24, 53, 51, 39, 45, 61, 52, 47, 38] explored the impacts of SFP on nutritional intake. Outcomes explored were diverse. An Article piece [24] claimed that Sweden was one of the countries with the highest vegetable intake, with a substantial intake at lunch. Sweden has the longest running universal SFP in the world and supplies food to all school aged children. The article notes that school meal provision in Sweden is seen as a universal welfare service and a part of public health work.

Vik et al. [53] used a non-randomised control trial including 164 secondary school children to investigate whether serving a free healthy school meal for one year resulted in a higher intake of fruit and vegetables and a lower intake of unhealthy snacks. Vik et al. [53] reported that a free healthy school meal for one year was associated with higher weekly intake of vegetables on sandwiches in the intervention group compared to the control group. However, it was not associated with a lower weekly intake of unhealthy snacks in the intervention group compared to the control group. It is important to note that the study had a small sample size and relied on self-reported data.

Parnham et al. [51] examined the impact of a universal infant SFP policy on dietary intake in primary school children in England and Scotland. The implementation of the policy led to more children participating in school meals. No impact was seen on fruit and vegetable intake but there was evidence that the policy lowered consumption of foods associated with packed lunches, such as crisps, and some nutrients, such as total fat and sodium.

A further study by Evans [39] also reported that school meals were a healthier option than packed lunches. The study utilised a cross-sectional design and included 2709 primary school

children in England. Findings suggested children having a packed lunch consumed on average 11g more total sugars than those who had a school meal whereas, children having a school meal were more likely to consume different types of vegetables and drink water. Overall, children with a packed lunch consumed a lower-quality diet over the whole school day. The study did not look directly at school meal provision and so it is difficult to note whether those on provision would yield differing results.

Spence et al. [45] explored the effect of pre- and post-Universal Infant Free School Meals (UIFSM) and school on pupil's dietary intakes using a cross-sectional research design. At lunchtime, there was a statistically significant decrease in students' non-milk extrinsic sugars intake pre implementation and increases of intakes of cake's pudding post implementation. However, only two schools were included, thus limiting the generalisability of results. It is recommended that schools consider healthier policies when implementing SFP.

Yamaguchi et al. [61] indicated that universal SFP were beneficial to individuals with low socio-economic backgrounds. Their study which used a mixed methods approach and included 719 primary school children in Japan, saw that children whose mothers were less educated had greater reliance on school lunch for their vegetable intake. Children with lower household income had more contribution from school lunch to their fruit intake. Household income was not explicitly considered in the study and mothers' education was used to determine socio-economic status, so it is unclear whether families were from lower socio-economic backgrounds. Further studies are needed to conclude the correlation between vegetable intake and household income in Japanese children.

Four papers, Neervoort et al. [52], Zailani et al. [47], Colombo et al. [38] and Goel [55]

considered the impacts of SFP on vitamin intake as well as nutritional deficiencies.

Neervoort et al. [52] conducted a non-randomised control trial including sixty-seven primary school children to investigate the impacts of a universal SFP on anaemia, stunting, wasting and malnutrition. Findings indicated the programme reduced anaemia and malnutrition, and improved child growth in the study group. Whilst improvements were seen, the study had several limitations. Such as a very small sample size and participants were from only one school. The study also lacked information on the content and nutritional value of provided meals, which could influence program effectiveness.

Zailania et al. [47] investigated portion sizes of targeted school meals as well as nutrient intake and found that the meals served through the SFP contributed at least 33% for energy, protein, iron, calcium, sodium, vitamin A, and zinc intake. However, there was no consumption of fruits, meat, poultry, and fish within the programme. The study did not consider the contribution of other food sources, including snacks and homemade meals which may account for nutritional values.

Colombo et al. [38] used a cross-sectional design to explore School Lunch Dietary intake under a universal SFP in Sweden. Results found that a quarter of the overall energy intake; between 22% and 30 % of selected nutrient intakes; almost half of vegetable intakes; roughly two-thirds of fish intakes; and around a third of red/processed meat intakes. These findings imply that school meals make an important contribution to children's diets on weekdays. Researchers also reported school meals as more nutritious than meals consumed outside of school, though meals consumed outside of school meals were not measured.

Goel et al. [55] examined total sugar in free breakfasts served in elementary (primary) schools in Virginia, USA through an observational study. Findings suggest that meals offered might contribute to excessive overall sugar availability for children. Nutritional information for added sugars was not provided within the study, however, the findings may explain contradictions in BMI above. Goel et al. [55] states districts and policymakers should collaborate to implement more effective guidelines concerning sugar availability in breakfast items to optimise children's dietary intake.

In summary, the evidence for improved nutritional intake is limited. However, school meals are an important contribution to diets during weekdays. Consuming school meals alleviated dietary discrepancies related to social inequalities.

Psychological and Emotional Health Outcomes

The impacts of SFP on children's psychological and emotional outcomes were investigated in four papers [57, 60, 48, 46].

A randomised control trial by Altindag et al. [57] evaluated the effects of targeted school meals on students' behaviour, bullying, and violence in schools. Behavioural incidents reduced by 35% since the implementation of a universal system in 2010 (particularly physical fights among students). It is believed that this reduction is the result of reduced stigma among students as socio-economic status cannot be identified via the system. Provision did not impact students' nutritional intake but reduced the stigma of receiving free meals through a targeted system by mitigating the possibility of identifying peers' socioeconomic statuses.

Taylor et al. [60] included 116 school staff in the USA to explore the impact of a universal

SFP on school climate, behaviour, and attainment. Taking a mixed-methods approach, over 725 of school staff surveyed reported that serving universal school meals has improved social climate as well as reductions in student stress, family financial stress and school administrator stress. This is thought to be a result of income differences being less visible, and the school community feeling more inclusive. It is recommended that further research include interviews with students and families to gain a better understanding.

As previously mentioned Bethman and Cho [48] saw positive results when looking at children's BMI. Similarly, the reintroduction of a universal policy also saw mental health improvements among students. Proclaiming that free school lunches help to improve health and benefit student welfare. However, mental health was assessed using a 'crying without any reason' measure. The measure used is not reported so validity and reliability cannot be determined. No other Mental Health outcomes were assessed.

A further paper by Yang et al. [46] assessed targeted school meals in the UK on children's mental health outcomes. The study used a cross-sectional design and included 2166 primary and secondary children. Yang et al [46] reported that poor mental health was observed in food insecure children receiving provision but also food insecure children not receiving provision. Interestingly, food insecure (i.e., not having access to sufficient food) children receiving provision had a higher probability of poor mental health than those who were food insecure and not receiving provision. It is thought that this is because targeted approaches are accosted with stigma.

In conclusion, universal approaches were associated with improved mental health outcomes, such as reduced behavioural incidents in school and an improved social climate. This was not

observed with targeted approaches.

Social Health Outcomes

Five papers [58, 65, 64, 66, 68] investigated the social impacts of SFP. The impacts of SFP are multifaceted with outcomes seen with social interactions, reduced stigma and narrowing inequalities. Findings are discussed further below.

A mixed method, quasi-experimental evaluation exploring the impact of universal provisions in two London secondary schools by Carlisle et al [58] saw improved social skills as children were eating with peers. Improved behaviours, less stigma as well as students eating more varied foods. These findings were as a result of paired student interviews, this may have caused socially desirable responses.

A further study by Jessiman et al. [65] which also included a sample of secondary school students in London also reported social health benefits, Students perceived feeling equal under a universal system whereas they reported feeling 'weak' under the targeted system.

Illøkken et al. [64] conducted a qualitative study in a Norwegian secondary school, involving 13 students and five teachers to investigate the effects of a targeted school feeding program on social inequalities. Participants viewed school meals as a social event where students made new friends and learned new skills. It was also reported that social equality among students increased. However, it is unclear whether this was the result of the feeding programme or school meals in general.

Another qualitative study conducted in a Norwegian secondary school by Mauer et al. [66],

also reported positive social outcomes. Students expressed that social time while eating school meals was important to them. This study also noted that popularity of the food was also important for attracting students to school meals.

Sahota et al. [68] investigated factors influencing take-up of targeted school meals through focus groups with children in both primary and secondary schools in England. Utilising focus groups to gain children's perspectives. Food choice, queuing, and the social aspects of lunch time, such as eating with friends was a major influence in uptake. Recognition of the importance of the social aspects of dining for pupils and facilitation of social interactions through the spatial (including flexible locations, e.g. outside) and temporal organisation of lunchtimes. It is reported that the schools involved in the study had a high level of targeted school meal entitlement which may have resulted in the normalisation of school meal uptake. It is recommended that social aspects of school meal provision be further investigated in schools with lower levels of eligibility to see if results differ.

Reducing Food Insecurity and Child Hunger

Three papers [73, 59, 41] looked at the impacts of SFP on food insecurity and child hunger. Most papers in this review touched on food insecurity, however, these three papers specifically looked at impacts of food insecurity outcome. A rapid narrative review by Garton et al. [73] concluded that SFP in New Zealand significantly reduced hunger and food insecurity in primary schools. Hecht [59] suggested that access to free meals through a universal provision in a high deprivation area in the USA, reduced food insecurity among children as well as decreased child hunger. Lastly, a cross-sectional analysis by Horta [40], investigated impacts of SFP on Vulnerability risk. Findings saw positive impacts from consuming school meals on children's diets, particularly among children living in high/very

high social vulnerability risk areas.

Factors Influencing Uptake

Ten papers explicitly investigated SFP participation and factors associated with non-take-up among students. Two studies gained perspectives through qualitative methods [67, 62].

A paper by McKelvie-Sebileau et al. [67], involving universal provision, found that lack of knowledge of the programme and loss of agency over meal choices were major drivers of non-take-up but, positive influences such as participants' food security, better nutritional knowledge and improved wellbeing were also perceived. Cardoso et al. [62] who evaluated a targeted approach in Portugal noted that quality of food was a concern when participating in school meals. Children who were not entitled to free or discounted meals reported eating at school less often than children with free or discounted lunches.

Three reports, Kitchen et al. [70], Rahim et al. [71] and Harper and Wood [69], all reported increased take-up under a universal policy, for both previously non eligible children and eligible children. It is suggested that improvements were seen as a result of reduced stigma and familiarising parents with school meals. Participants believed the pilot increased the range of food that pupils would eat, built their social skills at mealtimes and, for some pupils, resulted in health benefits associated with having a balanced meal, such as more energy, concentration and alertness and improved complexion. take-up was higher among primary school children than secondary school children. However, Rahim et al. [71] in particular, found schools experienced difficulty predicting the appropriate quantities of food required for each menu option, which meant that some meals either ran out early or were wasted.

A study by Holford and Rabe [49] also found that take-up of school meals in England by

non-eligible children rose from a consistent 30-35% in the eight years preceding the policy to approximately 85% in the universal period.

Additionally, two case studies [74, 63], one exploring a targeted system and the other a universal system saw similar results. Under the targeted provision [74] stigma, food quality and social aspects played a vital role in take-up. Results specifically noted that, being separated from friends decreased take-up. Implementation of a universal system [63] however, yielded more positive results such as, families who were previously above the eligibility threshold now being able to access provision. It was also documented that school-provided meals were of a higher nutritional quality than those from home, exposing children to a greater variety of foods and establishing healthier eating habits.

Finally, a cross-national comparison by Oostindjer [72] indicated that removing popular foods such as meat from school menus significantly reduced take-up of school meals. This was particularly evident in Finland where meat and fish were removed from the school menu on some days reducing participation rate, consequently producing up to 60% plate waste. Please note that this paper did not explicitly mention SFP but instead, school menu choices and take-up therefore, more research is needed on food choices under SFP to evaluate if there is an effect on take-up.

Wider Impacts

The remaining seven papers did not fit into a thematic category and will therefore be summarised individually below.

A UK wide study by Parnham et al. [44] exploring access to targeted SFP, reported that receiving an FSM was associated with increased odds of recently using a food bank [75] but

not reporting feeling hungry. In the month after the COVID-19 lockdown, 49% of eligible children did not receive any form of FSM. In this study the sample size was small, and this may lead to chance findings or in ability to detect differences.

A policy recommendation report by Guio [22], explored evidence of the short and longer-term benefits of school meal provisions in EU countries. The report found that health outcomes are dependent on the programme take-up and quality of food. Additionally, while eating formally, pupils learn to be sociable and develop interaction skills. There are some findings on health outcomes associated with school meal provision, but these are more difficult to establish. School-based food and nutrition interventions were able to improve dietary behaviour, healthy eating, and anthropometry, but the design of the intervention affects the magnitude of the effect.

James [42], used a cross-sectional design to investigate stigma associated with targeted provision and how this influences the peer effect. The study included 21,000 secondary school students in England. According to James [42], the presence of stigma dampens the peer effect and so students may not be benefiting from provision as much as possible.

However, information about the provision has the opposite impact on the peer effect. These findings suggest that information is a more important part of the peer effect for those living in areas of greater deprivation and stigma is more important for those in the least deprived regions.

Long et al. [43] assessed the impacts of a universal system on meal cost and food quality in 508 US schools. This study finds that participation in the provision was associated with lower per-meal full cost with no differences in dietary quality. This indicates that in the US, universal SFP can provide nutritious meals to more students without a financial disadvantage.

Meier et al. [54] conducted a quantitative exploratory study including 576 parents. Completed in the USA, the study aimed to gain parents overall perceptions of the targeted school meals program. This study found that parents of enrolled children were more likely to report positive perceptions of the school meals' program. Whereas parents of children not receiving the program were less likely to perceive the school meals program in a positive light. Two limitations were noted that may have influenced results. Firstly, Parents of children not receiving provision made up much of the sample and second, parents volunteered to participate in this survey, therefore participation bias could have resulted from those who are more involved or have strong opinions about school lunch.

A longitudinal study, utilising a difference-in-difference design by Lundborg et al. [20] evaluated the long-term impacts of the universal program on children's economic, educational, and health outcomes throughout the life course. The study involved 1,529,760 participants from birth to age 50+. The study noted several findings, for instance, children exposed to the program during their entire primary school period have 3% higher lifetime income. This effect was greater for pupils that were exposed at earlier ages and for pupils from poor households, suggesting that the program reduced socioeconomic inequalities in adulthood. Additionally, a year of exposure to UFSM increased height growth and nine years of school lunch exposure increase the likelihood of being of near perfect health. Exposure to school lunches also decreases the probability of being diagnosed with any health condition.

The final study by Goodchild et al. [40] investigated factors associated with a universal infant free school meal take-up and refusal in a multicultural urban community involving 676 parents of nursery aged children (4-7 years). The study explored two groups; non-provision, n

= 159 or took provision, n = 517. The non-provision group were more likely to be White-British, have higher socio-economic status, have English as a first language, and involve their child in the decision over whether to take the school meal, compared to the provision group. It should be noted that the area the study was conducted is multicultural and school meals must cater for children from a variety of cultural backgrounds. Parents who did not complete questionnaires correctly were more likely to have lower socio-economic status, be non-white British and have English as an additional language, meaning some groups were under-represented.

Discussion

A scoping review methodology in accordance with the PRISMA guidelines [29], was used to explore the global impacts of SFP on children's health and wellbeing outcomes. To our knowledge, this is the first scoping review to comprehensively appraise SFP depending on SFP type (targeted or universal). From a health perspective, SFP positively influenced children's physical health in terms of their social, emotional, and psychological well-being.

Research on the effectiveness of FSM on BMI, weight, and nutritional intake outcomes is limited but does indicate some positive impacts. Both targeted and universal approaches play a crucial role in addressing inequalities among disadvantaged children by providing access to food; however, evidence supporting one as more effective in impacting health outcomes is limited.

In terms of physical health, most included papers investigated BMI, with some conflicting findings. For example, most papers indicated positive impacts, however two papers [6, 57] observed increased BMI under SFP. This discrepancy may be due to pre-existing malnutrition being present prior to provision or the foods served. For example, Goel et al. [55] found an increased sugar intake in free school breakfast. Though, nutritional quality of foods served differs between countries, making this difficult to assess.

SFP appears to be a positive step forward in promoting positive psychological and emotional outcomes. However, studies in this area are limited and have several limitations. It is clear that more research is needed to evaluate the effectiveness of SFP on children's psychological and emotional outcomes.

SFP, whether universal or targeted, seems to ease food insecurity. Individuals from low socio-economic backgrounds benefited more from SFP, regardless of whether they were targeted or universal. However, targeted programs were associated with stigma, resulting in lower take-up and poorer mental health outcomes. Emphasising the importance of implementing policies to mitigate stigma. Universal systems eliminated stigma and significantly increased uptake. Nonetheless, issues such as loss of agency and food quality persisted in influencing take-up rates.

Implications for Research

As previously mentioned, heterogeneity in research designs makes evaluating SFP difficult. Most papers included in this review were conducted over a short time span. To fully understand the impacts of SFP on children's health outcomes, longer term research designs are essential. Additionally, sociality was deemed an important aspect of free school meals uptake, whereas stigma was associated with lower uptake. Therefore, future research should consider the implications of stigma on children's health and how this can be alleviated. Only three studies investigated SFP as a means of addressing nutritional deficiencies among children. Future research should seek to explore these in more detail as well as other health concerns, such as respiratory illnesses and other common childhood illnesses associated with poor dietary intake.

Implications for Practice

There is not a definitive global recommendation for SFP implementation. Based on available evidence, both targeted and universal provision appears to be effective at improving children's health outcomes to some degree, however there are considerable limitations in current research. There is a notable need to address factors influencing uptake, particularly food quality and choice. Information was deemed important to parents when registering for SFP. Therefore, educational settings should ensure parents have sufficient information relating to SFP. Going forward, best practice and nutritional quality should be considered.

Conclusions

Our review of 44 articles provides mixed findings on the effectiveness of SFP in improving children's health outcomes. SFP can contribute to healthy weight outcomes, although the nutritional benefits depend on the composition of the food offered. Universal provision reduced stigma and may lead to a more positive social environment. Overall, both targeted and universal approaches were effective in addressing inequalities and food insecurity.

Acknowledgements

Funding

This research was supported by the National Centre for Population Health and Wellbeing Research. Funding was provided by Public Health Wales and the Economic and Social Research Council (ESRC) through their support of a PhD studentship.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Author Contributions

Amy Locke led the review, was responsible for managing the project, analysing data and preparing an original draft.

Michaela James provided supervision, contributed to abstract and full text screening, quality assessment and editing.

Hope Jones contributed to abstract and full text screening, and quality assessment, editing.

Rachel Davis contributed to abstract and full text screening and quality assessment, editing.

Francesca Williams contributed to full text screening and quality assessment, editing.

Sinead Brophy provided supervision and advice for the scoping review and methodology and reviewed the final report.

References

- [1] World Food Programme. 2022. State of School Feeding Worldwide 2022. Rome, World Food Programme. ISBN 978-92-95050-16-7.
- [2] Watkins K. School meal programmes: A missing link in food systems reform Prepared by the Sustainable Financing Initiative (SFI) for School Health and Nutrition [Internet]. 2023 [cited 2024 Feb 21]. Available from: <https://www.edc.org/sites/default/files/School-meals-Food-Systems.pdf>
- [3] Jansen PW, Roza SJ, Jaddoe VW, Mackenbach JD, Raat H, Hofman A, Verhulst FC, Tiemeier H. Children's eating behavior, feeding practices of parents and weight problems in early childhood: results from the population-based Generation R Study.

- International Journal of Behavioral Nutrition and Physical Activity. 2012 Dec;9(1):1-1.
- [4] Hunter D, Giyose B, PoloGalante A, Tartanac F, Bundy D, Mitchell A, Moleah T, Friedrich J, Alderman A, Drake L, Kupka R. Schools as a system to improve nutrition: A new statement for school-based food and nutrition interventions.
- [5] World Health Organization. Food and nutrition policy for schools: a tool for the development of school nutrition programmes in the European Region. Copenhagen: WHO Regional Office for Europe; 2006.
- [5] Vik, F.N., Van Lippevelde, W. & Øverby, N.C. Free school meals as an approach to reduce health inequalities among 10–12- year-old Norwegian children. *BMC Public Health* 19, 951 (2019). <https://doi.org/10.1186/s12889-019-7286-z>
- [7] Cohen JF, Hecht AA, McLoughlin GM, Turner L, Schwartz MB. Universal school meals and associations with student participation, attendance, academic performance, diet quality, food security, and body mass index: A systematic review. *Nutrients*. 2021 Mar 11;13(3):911.
- [8] Destaw Z, Wencheke E, Kidane S, Endale M, Challa Y, Tiruneh M, Tamrat M, Samson H, Shaleka D, Ashenafi M. Impact of school meals on educational outcomes in Addis Ababa, Ethiopia. *Public Health Nutrition*. 2022 Sep;25(9):2614-24.
- [9] Chaudhary A, Sudzina F, Mikkelsen BE. Promoting healthy eating among young people—a review of the evidence of the impact of school-based interventions. *Nutrients*. 2020 Sep 22;12(9):2894.
- [10] Khodae GH, Emami Moghadam Z, Khademi G, Saeidi M. Healthy diet in children: Facts and keys. *International Journal of Pediatrics*. 2015 Dec 1;3(6.2):1183-94.
- [11] Ayton A, Ibrahim A. The dramatic rise of ultra-processed foods. *bmj*. 2019 Aug 5;366:14970.
- [12] Okunogbe A, Nugent R, Spencer G, Powis J, Ralston J, Wilding J. Economic impacts of overweight and obesity: current and future estimates for 161 countries. *BMJ global health*. 2022 Sep 1;7(9):e009773.
- [13] World Health Organization □ Obesity and Overweight.(2024) Available from <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
- [14] World Health Organization. WHO acceleration plan to stop obesity. Available from <https://iris.who.int/bitstream/handle/10665/370281/9789240075634-eng.pdf?sequence=1>
- [15] Yue T, Zhang Q, Li G, Qin H. Global burden of nutritional deficiencies among children under 5 Years of age from 2010 to 2019. *Nutrients*. 2022 Jun 28;14(13):2685.
- [16] Bringas Vega ML, Guo Y, Tang Q, Razzaq FA, Calzada Reyes A, Ren P, Paz Linares D, Galan Garcia L, Rabinowitz AG, Galler JR, Bosch-Bayard J. An age-adjusted EEG source classifier accurately detects school-aged barbadian children

that had protein energy malnutrition in the first year of life. *Frontiers in neuroscience*. 2019 Nov 29;13:476889.

- [17] Liu J, Qi X, Wang X, Qin Y, Jiang S, Han L, Kang Z, Shan L, Liang L, Wu Q. Evolving patterns of nutritional deficiencies burden in low-and middle-income countries: findings from the 2019 Global Burden of Disease Study. *Nutrients*. 2022 Feb 22;14(5):931.
- [18] Pasricha SR, Tye-Din J, Muckenthaler MU, Swinkels DW. Iron deficiency. *The Lancet*. 2021 Jan 16;397(10270):233-48.
- [19] Gupta S, Jawanda MK. The impacts of COVID-19 on children. *Acta Paediatrica* (Oslo, Norway: 1992). 2020 Nov;109(11):2181.
- [20] Lundborg P, Rooth DO, Alex-Petersen J. Long-term effects of childhood nutrition: evidence from a school lunch reform. *The Review of Economic Studies*. 2022 Mar 1;89(2):876-908.
- [21] Kupka R, Siekmans K, Beal T. The diets of children: Overview of available data for children and adolescents. *Global Food Security*. 2020 Dec 1;27:100442.
- [22] Guio AC. Free school meals for all poor children in Europe: An important and affordable target?.
- [23] Guio AC, Frazer H, Marlier E. Study on the economic implementing framework of a possible EU Child Guarantee scheme including its financial foundation: Second phase of the Feasibility Study for a Child Guarantee (FSCG2).
- [24] Persson Osowski C, Fjellström C. Understanding the ideology of the Swedish tax-paid school meal. *Health Education Journal*. 2019 Jun;78(4):388-98.
- [25] Hobbs G, Vignoles A. Is children's free school meal 'eligibility' a good proxy for family income?. *British Educational Research Journal*. 2010 Aug 1;36(4):673-90.
- [26] Lucas PJ, Patterson E, Sacks G, Billich N, Evans CE. Preschool and school meal policies: an overview of what we know about regulation, implementation, and impact on diet in the UK, Sweden, and Australia. *Nutrients*. 2017 Jul 11;9(7):736.
- [27] Morelli CJ, Seaman PT. Universal versus targeted benefits: the distributional effects of free school meals. *Environment and Planning C: Government and Policy*. 2005 Aug;23(4):583-98.
- [28] Taylor A. England's free school meals scheme should be expanded to help more children living in poverty. *BMJ*. 2022 Nov 2;379.
- [29] Moher D, Liberati A, Tetzlaff J, Altman DG, Altman D, Antes G, et al. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med* [Internet]. 2009 Jul 1 [cited 2024 Feb 07];6(7):e1000097. Available from: <https://journals.plos.org/plosmedicine/article?id=10.1371/journal.pmed.1000097>

- [30] Locke A, James M, Jones H, Davies R, Williams F, Brophy S. Impact of Global School Feeding Programmes on Children's Health and Wellbeing Outcomes: A Scoping Review Protocol 2024. doi:10.17605/OSF.IO/UZRWD.
- [31] Covidence systematic review software, Veritas Health Innovation, Melbourne, Australia. Available at www.covidence.org.
- [32] Hong QN, Fàbregues S, Bartlett G, Boardman F, Cargo M, Dagenais P, Gagnon MP, Griffiths F, Nicolau B, O'Cathain A, Rousseau MC. The Mixed Methods Appraisal Tool (MMAT) version 2018 for information professionals and researchers. *Education for information*. 2018 Jan 1;34(4):285-91.
- [33] Hong QN, Gonzalez-Reyes A, Pluye P. Improving the usefulness of a tool for appraising the quality of qualitative, quantitative and mixed methods studies, the Mixed Methods Appraisal Tool (MMAT). *Journal of evaluation in clinical practice*. 2018 Jun;24(3):459-67.
- [34] McArthur A, Klugarova J, Yan H, Florescu S. Innovations in the systematic review of text and opinion. *Int J Evid Based Healthcare*. 2015;13(3):188–195
- [35] Munn Z, Tufanaru C, Aromataris E. JBI's systematic reviews: data extraction and synthesis. *AJN The American Journal of Nursing*. 2014 Jul 1;114(7):49-54.
- [36] Popay J, Roberts H, Sowden A, Petticrew M, Arai L, Rodgers M, Britten N, Roen K, Duffy S. Guidance on the conduct of narrative synthesis in systematic reviews. A product from the ESRC methods programme Version. 2006 Apr 1;1(1):b92.
- [37] Batista MD, Mondini L, Jaime PC. Actions of the School Health Program and school meals in the prevention of childhood overweight: experience in the municipality of Itapevi, São Paulo State, Brazil, 2014. *Epidemiologia e Serviços de Saúde*. 2017 Jul;26:569-78.
- [38] Colombo PE, Patterson E, Elinder LS, Lindroos AK. The importance of school lunches to the overall dietary intake of children in Sweden: a nationally representative study. *Public health nutrition*. 2020 Jul;23(10):1705-15.
- [39] Evans CE, Mandl V, Christian MS, Cade JE. Impact of school lunch type on nutritional quality of English children's diets. *Public Health Nutrition*. 2016 Jan;19(1):36-45.
- [40] Goodchild GA, Faulks J, Swift JA, Mhesuria J, Jethwa P, Pearce J. Factors associated with universal infant free school meal take up and refusal in a multicultural urban community. *Journal of human nutrition and dietetics*. 2017 Aug;30(4):417-28.
- [41] Horta PM, do Carmo AS, Junior EV, Dos Santos LC. Consuming school meals improves Brazilian children's diets according to their social vulnerability risk. *Public Health Nutrition*. 2019 Oct;22(14):2714-9.
- [42] James J. Peer Effects in Free School Meals: Information or Stigma?.

- [43] Long MW, Marple K, Andreyeva T. Universal free meals associated with lower meal costs while maintaining nutritional quality. *Nutrients*. 2021 Feb 19;13(2):670.
- [44] Parnham JC, Lavery AA, Majeed A, Vamos EP. Half of children entitled to free school meals did not have access to the scheme during COVID-19 lockdown in the UK. *Public health*. 2020 Oct 1;187:161-4.
- [45] Spence S, Matthews JN, McSweeney L, Rowland MK, Orango P, Adamson AJ. Implementation of Universal Infant Free School Meals: A pilot study in NE England exploring the impact on Key Stage 1 pupil's dietary intake. *Public Health Nutrition*. 2021 Jul;24(10):3167-75.
- [46] Yang TC, Power M, Moss RH, Lockyer B, Burton W, Doherty B, Bryant M. Are free school meals failing families? exploring the relationship between child food insecurity, child mental health and free school meal status during COVID-19: national cross-sectional surveys. *BMJ open*. 2022 Jun 1;12(6):e059047.
- [47] Zailani H, Owolabi OA, Sallau AB. Contribution of school meals to the recommended nutrient and energy intake of children enrolled in the National Homegrown School Feeding Program in Zaria, Nigeria. *Archives de Pédiatrie*. 2023 Oct 1;30(7):471-6.
- [48] Bethmann D, Cho JI. The impacts of free school lunch policies on adolescent BMI and mental health: Evidence from a natural experiment in South Korea. *SSM-Population Health*. 2022 Jun 1;18:101072.
- [49] Holford A, Rabe B. Impact of the universal infant free school meal policy. Colchester: Institute for Social and Economic Research. 2020.
- [50] Holford A, Rabe B. Going universal. The impact of free school lunches on child body weight outcomes. *Journal of Public Economics Plus*. 2022 Jan 1;3:100016.
- [51] Parnham JC, Chang K, Millett C, Lavery AA, von Hinke S, Pearson-Stuttard J, de Vocht F, White M, Vamos EP. The impact of the Universal Infant Free School Meal policy on dietary quality in English and Scottish primary school children: evaluation of a natural experiment. *Nutrients*. 2022 Apr 12;14(8):1602.
- [52] Neervoort F, von Rosenstiel I, Bongers K, Demetriades M, Shacola M, Wolffers I. Effect of a school feeding programme on nutritional status and anaemia in an urban slum: a preliminary evaluation in Kenya. *Journal of tropical pediatrics*. 2013 Jun 1;59(3):165-74.
- [53] Vik FN, Heslien KE, Van Lippevelde W, Øverby NC. Effect of a free healthy school meal on fruit, vegetables and unhealthy snacks intake in Norwegian 10-to 12-year-old children. *BMC Public Health*. 2020 Dec;20:1-8.
- [54] Meier CL, Brady P, Askelson N, Ryan G, Delger P, Scheidel C. What do parents think about school meals? An exploratory study of rural middle school parents' perceptions. *The Journal of school nursing*. 2022 Jun;38(3):226-32.

- [55] Goel NJ, Caccavale LJ, Mazzeo SE, Raynor HA, Bean MK. Total sugar in free breakfasts served in Virginia elementary schools. *Health Behavior and Policy Review*. 2019 Sep 1;6(5):455-62.
- [56] Davis W, Kreisman D, Musaddiq T. The effect of universal free school meals on child BMI. *Education Finance and Policy*. 2023 Sep 15:1-31.
- [57] Altindag DT, Baek D, Lee H, Merkle J. Free lunch for all? The impact of universal school lunch on student misbehavior. *Economics of Education Review*. 2020 Feb 1;74:101945.
- [58] Carlisle VR, Jessiman PE, Breheny K, Campbell R, Jago R, Leonard N, Robinson M, Strong S, Kidger J. A Mixed Methods, Quasi-Experimental Evaluation Exploring the Impact of a Secondary School Universal Free School Meals Intervention Pilot. *International Journal of Environmental Research and Public Health*. 2023 Mar 22;20(6):5216.
- [59] Hecht AA. Universal free school meals: Implementation of the community eligibility provision and impacts on student nutrition, behavior and academic performance (Doctoral dissertation, Johns Hopkins University).
- [60] Taylor J, Garnett B, Horton MA, Farineau G. Universal free school meal programs in Vermont show multi-domain benefits. *Journal of hunger & environmental nutrition*. 2020 Nov 1;15(6):753-66.
- [61] Yamaguchi M, Kondo N, Hashimoto H. Universal school lunch programme closes a socioeconomic gap in fruit and vegetable intakes among school children in Japan. *The European Journal of Public Health*. 2018 Aug 1;28(4):636-41.
- [62] Cardoso SG, Truninger M, Ramos V, Augusto FR. School meals and food poverty: Children's views, parents' perspectives and the role of school. *Children & Society*. 2019 Nov;33(6):572-86.
- [63] Chambers S, Boydell N, Ford A, Eadie D. Learning from the implementation of Universal Free School Meals in Scotland using Normalisation Process Theory: Lessons for policymakers to engage multiple stakeholders. *Food Policy*. 2020 Aug 1;95:101936.
- [64] Illøkken KE, Johannessen B, Barker ME, Hardy-Johnson P, Øverby NC, Vik FN. Free school meals as an opportunity to target social equality, healthy eating, and school functioning: Experiences from students and teachers in Norway. *Food & nutrition research*. 2021;65.
- [65] Jessiman PE, Carlisle VR, Breheny K, Campbell R, Jago R, Robinson M, Strong S, Kidger J. A qualitative process evaluation of universal free school meal provision in two London secondary schools. *BMC public health*. 2023 Feb 9;23(1):300.
- [66] Mauer S, Torheim LE, Terragni L. Children's participation in free school meals: a qualitative study among pupils, parents, and teachers. *Nutrients*. 2022 Mar 18;14(6):1282.

- [67] McKelvie-Sebileau P, Swinburn B, Glassey R, Tipene-Leach D, Gerritsen S. Health, wellbeing and nutritional impacts after 2 years of free school meals in New Zealand. *Health Promotion International*. 2023 Aug 1;38(4):daad093.
- [68] Sahota P, Woodward J, Molinari R, Pike J. Factors influencing take-up of free school meals in primary-and secondary-school children in England. *Public health nutrition*. 2014 Jun;17(6):1271-9.
- [69] Harper C, Wood L. Please Sir? Can we have some more?—lessons from Free School Meal initiatives.
- [70] Kitchen S, Tanner E, Brown V, Payne C, Crawford C, Dearden L, Greaves E, Purdon S. Evaluation of the free school meals pilot. Department for Education: London, UK. 2013.
- [71] Rahim N, Kotecha M, Callanan M, White C, Tanner E. Implementing the free school meals pilot. Department for Education; 2012.
- [72] Oostindjer M, Aschemann-Witzel J, Wang Q, Skuland SE, Egelanddal B, Amdam GV, Schjøll A, Pachucki MC, Rozin P, Stein J, Lengard Almli V. Are school meals a viable and sustainable tool to improve the healthiness and sustainability of children's diet and food consumption? A cross-national comparative perspective. *Critical reviews in food science and nutrition*. 2017 Dec 12;57(18):3942-58.
- [73] Garton K, Riddell C, McKelvie-Sebileau P, Glassey R, Leech DT, Rees D, Swinburn B. Not Just a Free Lunch a logic model and evidence review for the Ka Ora, Ka Ako| Healthy School Lunch programme. *Policy Quarterly*. 2023 May 31;19(2):75-87.
- [74] Chambers S, Dundas R, Torsney B. School and local authority characteristics associated with take-up of free school meals in Scottish secondary schools, 2014. *In Civic Engagement 2017 Sep 19* (pp. 52-63). Routledge.
- [75] The Trussell Trust. (2024). Emergency Food- The Trussell Trust. <https://www.trusselltrust.org/get-help/emergency-food/>