medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .

Title: A Sex-specific Mendelian Randomization-Phenome-Wide Association Study of Body Mass Index

Short title: Phenome-Wide Study of BMI

Authors: Zhu Liduzi Jiesisibieke[1], Io Ieong Chan[2], Jack Chun Man Ng[1], C Mary Schooling[1,3]

[1] School of Public Health, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong SAR, China

[2] Department of Public Health and Medicinal Administration, Faculty of Health Sciences, University of Macau, Macao SAR, China

[3] Graduate School of Public Health and Health Policy, City University of New York, New York, United States

[*]**Corresponding Author:**

C Mary Schooling

Graduate School of Public Health and Health Policy, City University of New York, New York, United States

Email address: mary.schooling@sph.cuny.edu

Abstract word count: 203

Text word count: 2,604

Figures: 4

ORCID: C Mary Schooling (0000-0001-9933-5887)

Zhu Liduzi Jiesisibieke (0000-0002-4986-653X)

Jack Chun Man Ng (0000-0002-3084-236X)

Grants or fellowships supporting the writing of the paper: The authors declare that there is no funding support regarding the publication of this paper.

Disclosure summary: The authors have nothing to disclose.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .

Impact statement: BMI may affect a wide range of health-related attributes and there are notable sex differences in its impact, including opposite associations for certain attributes, such as ApoB and neuroticism. These findings emphasize the importance of maintaining a healthy BMI.

Keywords: BMI, Mendelian randomization, PheWAS, sex difference, UK Biobank

medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .

ABSTRACT

Background: Trials of incretins are making it increasingly clear that body mass index (BMI) is linked to several diseases throughout life, but trials cannot provide a comprehensive assessment of the role of BMI in health-related attributes for men and women. To systematically investigate the role of BMI, we conducted a sex-specific Mendelian randomization-phenome-wide association study.

Methods: We comprehensively examined the associations of genetically predicted BMI in women (n: 194,174) and men (n: 167,020) with health-related attributes from the UK Biobank with inverse variance weighting and sensitivity analysis.

Results: BMI impacted 232 of 776 traits considered in women and 204 of 681 traits in men, after adjusting for false discovery; differences by sex were found for 105 traits. BMI was more strongly positively associated with heart disease, heart failure and hypertensive heart disease in men than women. BMI was more strongly positively associated with apolipoprotein B (ApoB), diastolic blood pressure, neuroticism, arthritis and triglycerides in women than men.

Conclusion: Our study revealed that BMI might affect a wide range of health-related attributes and highlights notable sex differences in its impact, including opposite associations for certain attributes, such as ApoB and neuroticism. These findings emphasize the importance of maintaining a healthy BMI.

Funding: None

(Word count: 200)

medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .

Introduction

Global obesity prevalence more than doubled from 1980 to 2008 (Finucane et al., 2011). In 2022, one in eight people were obese worldwide (WHO, 2024). Body mass index (BMI) is a longstanding measure of obesity, despite its high specificity but low sensitivity (Chooi et al., 2019; Okorodudu et al., 2010). Many observational studies have found BMI associated with disease risk factors and lifespan (Collaboration, 2009; Kahn et al., 2006; Khan et al., 2018), generally, a higher BMI is detrimental to health. However, BMI does not account well for fat distribution. For men and women with the same BMI, women tend to store more fat (Power & Schulkin, 2008). Furthermore, men tend to store fat around their organs, while women are more likely to store it subcutaneously (Power & Schulkin, 2008), potentially leading to different health impacts by sex (Costanzo et al., 2022; Power & Schulkin, 2008). As such, the effect of BMI on health may vary between men and women and may be a modifiable factor contributing to differences in lifespan by sex.

Few previous observational studies have systematically assessed the role of BMI in health and disease, and are limited by their susceptibility to confounding (Davey Smith & Ebrahim, 2003). Mendelian randomization (MR) studies reduce confounding by using genetic proxies, such as single nucleotide polymorphisms (SNPs), for exposures. Previous phenome-wide association studies using MR (MR-PheWASs) have identified impacts of BMI on endocrine disorders, circulatory diseases, inflammatory and dermatological conditions, and some biomarkers (Hyppönen et al., 2019; Millard et al., 2015), but were conducted before the advent of large-scale genome-wide association studies (GWASs) and were not sex-specific. Previous MR studies and recent trials of incretins have expanded our knowledge about a broad range of effects of BMI (Larsson et al., 2020; Marso et al., 2016). To our knowledge, no sex-specific PheWAS has investigated the effects of BMI on health outcomes. To address this gap, we conducted a sex-specific PheWAS, utilizing the largest available sex-specific GWAS, to explore the impact of BMI on health-related attributes.

medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .

Methods

Data sources

Exposure: Body mass index

To obtain genetic information for BMI (inverse rank normalized) sex-specifically, we used BMI from two sources, the UK Biobank, a prospective cohort study of half a million adults, and the Genetic Investigation of ANthropometric Traits (GIANT) consortium (Locke et al., 2015), a GWAS meta-analysis of 339,224 participants mostly of European ancestry. The UK Biobank has the advantage of a larger sample size and denser genotyping but is the only large-scale sex-specific source for many outcomes. The sex-specific UK Biobank BMI GWAS conducted by Neale Lab (women: 194,174; men: 167,020) was adjusted for age, age squared and the first 20 principal components (https://www.nealelab.is/uk-biobank/faq). The GIANT GWAS has the advantage of using a different sample from the UK Biobank but is smaller with less dense genotyping (women: 171,977; men: 152,893). The sex-specific GIANT GWAS was adjusted for age, age squared, and study-specific covariates (Locke et al., 2015). We also considered overall BMI from the GIANT (n: 681,275) which includes the UK Biobank participants (approximately 64%), and was adjusted for age, age squared, principal components and study-specific covariates (Yengo et al., 2018).

Outcomes

The UK Biobank is currently the largest and most comprehensive source for sex-specific GWAS, provided by Neale Lab (women: 194,174; men: 167,020), including many disease outcomes and physiological attributes. The average age at recruitment of UK Biobank participants was 57 years, as described previously (Collins, 2012).

Outcomes: Inclusion and exclusion criteria

For continuous outcomes only rank normalized health attributes with at least 1,000 participants (Verma et al., 2018) were included to maintain statistical power. For

medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .

binary attributes only those with at least 200 cases were included; duplicated phenotypes were excluded. We further excluded very similar attributes. Where an attribute has a standard measure, such as forced expiratory volume in 1 second, we used that in preference to similar measures. We also excluded attributes only assessed in selected subgroups of the UK Biobank participants, such as an electrocardiogram, which was only conducted in healthy people (Ramírez et al., 2021). Some attributes were available from both self-report and doctor diagnosis, in which case we used self-report because of the higher case number and hence greater statistical power. Finally, attributes with known measurement issues, such as estrogen and immature reticulocyte fraction, were excluded (Newman & Handelsman, 2014; Piva et al., 2015). Detailed exclusion criteria are listed in **Figure 1**.

Outcomes: Attribute categorization

We categorized attributes as age at recruitment, physical measures, lifestyle and environmental, medical conditions, operations, physiological factors, cognitive function, health and medical history, sex-specific factors, blood assays and urine assays, as recommended by the UK Biobank. Binary attributes with an International Classification of Diseases-10 code were categorized by International Classification of Diseases-10 chapter, as (I) infectious diseases, (II) neoplasms, (III) diseases of the hematopoietic system and blood disorders, (IV) endocrine and metabolic diseases, (V) psychological disorders, (VI) diseases of the nervous system, (VII and VIII) diseases of the sensory system (eyes and ears), (IX) diseases of the circulatory system, (X) diseases of the respiratory system, (XI) diseases of the digestive system, (XII) diseases of skin and subcutaneous tissue, (XIII) diseases of the musculoskeletal system and connective tissue, (XIV) diseases of the genitourinary system, (XV) pregnancy, childbirth, and the puerperium, and (XVIII) symptoms.

Statistical analysis

MR was used to assess sex-specific effects of genetically predicted BMI on each attribute considered. Specifically, inverse-variance weighted estimates were used

medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .

initially, i.e., meta-analysis of Wald estimates (SNP on outcome divided by SNP on exposure), and then sensitivity analysis (MR-Egger) was used for any associations found. The significance level was adjusted for the false discovery rate to account for multiple comparisons (Benjamini & Hochberg, 1995). Specifically, we ranked the p-values for men and women respectively, calculated the Benjamini-Hochberg (BH) value, and identified the significant attributes influenced by BMI using the BH value. We used a z-test to evaluate differences by sex (Altman & Bland, 2003). We used the R packages "TwoSampleMR" (version: 0.6.1), "Mendelian Randomization" (version: 0.10.0) and "metafor" (version: 4.6-0) to assess sex-differences.

Results

Initial analysis using sex-specific BMI from the GIANT yielded similar estimates to using sex-specific BMI from the UK Biobank but had fewer single nucleotide polymorphisms resulting in wider confidence intervals (**S Table 1 and S Table 2**). We presented the results obtained using sex-specific BMI from the UK Biobank.

Sex-specific estimates

In men, BMI was associated with 204 of the 681 health-related attributes considered using the false discovery rate (Figure 2, **S Table 3**). As expected, BMI was positively associated with heart disease, heart failure, hypertensive heart disease, diabetes, hypertension, sleep apnoea, daytime dozing, triglycerides, urinary potassium, urinary sodium and major surgeries. BMI was also inversely associated with age at recruitment, osteoporosis, hay fever, high density lipoprotein cholesterol (HDL-c), cholesterol, low density lipoprotein (LDL), sex hormone binding globulin (SHBG) and total testosterone. Positive associations of BMI with higher risk of digestive system cancers, lymphomas, primary lymphoid and hematopoietic malignancies, and esophageal cancer were also found in men. Furthermore, we found higher BMI associated with accelerated facial aging and baldness.

In women, BMI was associated with 232 of 776 attributes considered using the false

medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .

discovery rate (Figure 3, **S Table 3)**. As expected, BMI was positively associated with heart disease, heart failure, hypertensive heart disease, diabetes, hypertension, sleep apnoea, daytime dozing, apolipoprotein B (ApoB), triglycerides, total testosterone and urinary potassium and sodium and major surgeries. BMI was also inversely associated with osteoporosis, HDL-c, cholesterol, LDL and SHBG. BMI was positively associated with risk of uterine cancer, cancer of the bronchus and lung and intrathoracic organs, as well as cancers affecting the skin and female reproductive organs. Higher BMI in women was also associated with accelerated facial aging.

Differences by sex

We found significant differences by sex in the associations of BMI with 105 health-related attributes. Most differences (74) were in magnitude but not direction, directionally different associations were found for 31 attributes. BMI was inversely associated with ApoB, psoriatic arthropathy, iron deficiency anemia, hernia, hand grip strength and total testosterone in men, while positively associated with these traits in women (**Figure 4**). BMI was positively associated with the risk of malignant neoplasm of the urinary tract and colon in men but was inversely associated in women. BMI was positively associated with hernia, cervicalgia, mononeuropathies of lower limb and degeneration of macula and posterior pole in women, whilst inversely associated in men. BMI was inversely associated with anemias in men, but positively associated in women. BMI was inversely associated with neuroticism scores, sensitivity to hurt feelings, and frequency of seeking of medical advice for nerves, anxiety, tension, or depression in men. However, BMI was positively with neuroticism scores and seeking medical advice for these issues in women.

Several notable differences in magnitude by sex were also found. BMI was more strongly and inversely associated with osteoporosis SHBG, albumin and bilirubin in women than in men. BMI was more strongly and inversely associated with LDL and hay fever in men than women. BMI was more strongly and positively associated with diastolic blood pressure, depression, arthritis, triglycerides and hypothyroidism in

medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .

women than men. BMI was more strongly and positively associated with heart disease, heart failure, hypertensive heart disease, lymphomas, facial aging, and sleep apnoea in men than women.

Discussion

Consistent with previous studies BMI was positively associated with many health-related attributes, such as heart disease, heart failure, hypertensive heart disease, diabetes, hypertension and sleep apnoea. Our study adds by showing sex-differences in some traits related to cancer and psychological disorders as well as ApoB and showing stronger associations in men than women for common cardiovascular diseases.

Comparison with previous studies

Consistent with previous MR studies, BMI was positively associated with the risk of heart disease, heart failure, hypertensive heart disease (Riaz et al., 2018), hypertension (Riaz et al., 2018) and diabetes (Larsson & Burgess, 2021) in both men and women as would be expected. We also found BMI was inversely associated with osteoporosis (Larsson & Burgess, 2021), with a stronger association in women than men. A previous MR study found height and fat-free mass positively associated with follicular lymphoma (Zhou et al., 2024), however, we found BMI positively associated with lymphomas in men, but not women (although directionally consistent but not significant after considering multiple comparisons). Another MR study found BMI positively associated with asthma and poorer lung function, but not with hay fever (Skaaby et al., 2018), when we found BMI inversely associated with hay fever in men, but not in women. Previous studies have shown BMI inversely associated with Na/K (Zanetti et al., 2020), while we found BMI positively associated with urinary potassium and sodium. Additionally, we found a positive association of BMI with major surgeries. Consistent with the previous MR study (Ardissino et al., 2022), we also found BMI positively associated with sleep apnoea we also found positive associations with daytime dozing in both sexes, which may indicate effects of BMI on

medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .

quality of life. Higher BMI was associated with a younger age at recruitment among men, suggesting poorer survival to recruitment in men given the GWAS adjusted for age and the short recruitment window makes period effects unlikely. However, the association was less evident in women, with no sex difference. Notably, higher BMI may be more detrimental to lifespan in men than women, especially at younger ages (Fontaine et al., 2003).

Differences in the associations of body mass index with health-related attributes in men and women

BMI was more strongly associated with heart disease in men than women. BMI was positively associated with ApoB in women, whereas in men, the association was inverse (but not significant after correction for multiple comparisons); however, the sex difference was significant. This pattern contrasts with earlier MR studies that reported an inverse association of BMI with ApoB in the general population (Bell et al., 2022). BMI being more strongly associated with heart disease in men than women while also being protective for the key heart disease risk factor of ApoB requires some explanation. BMI is a complex phenotype that may represent different attributes or have different consequences in men and women. Fat mass storage tends to differ between men and women (visceral versus subcutaneous) (Power & Schulkin, 2008). The causes of BMI may also differ between men and women, due to occupational roles, gender-based food preferences or cultural norms (Kanter & Caballero, 2012). Alternatively, unknown factors affected by BMI could contribute to heart disease specifically to men. In terms of psychological disorders, women being more affected psychologically by higher BMI is consistent with sociocultural pressures and norms surrounding women's rather than men's bodies (Esnaola et al., 2010; Schwartz & Brownell, 2004). We also found that BMI was associated with balding in men but not in women. Moreover, BMI was positively associated with facial aging in both men and women, with larger effects in men.

Strengths and limitations

medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .

A major strength of this study was the focus on sex differences, which have not been explicitly explored in previous PheWAS of BMI, despite differences in lifespan by sex Despite the large sample size and the broad range of outcomes considered in this study, limitations and concerns exist. First, MR has stringent assumptions, i.e., relevance, independence and exclusion restriction. The F-statistics were above 10, which addresses relevance. MR is largely free from confounding by design. For a few attributes (37 out of 1,457), the MR-Egger intercept was significant while the IVW estimate was not (**S Table 4**), potentially indicating horizontal pleiotropic effects or the play of chance, which requires further investigation. Second, MR assesses BMI lifetime effects of BMI, which may differ from those of short-term weight changes or weight cycling. Third, not all attributes were included. For example, heart rate from ECG and age at asthma diagnosis were excluded, because such information was limited to selected subgroups. Fourth, given the study's exploratory nature, we allowed for multiple comparisons to minimize chance findings, which means some BMI effects may not be captured. However, such comprehensive consideration may identify previously unknown or ignored impacts, such as facial aging and balding. Fifth, while the UK Biobank does not fully represent the UK population, representativeness is not crucial for causal inference , as long as the sample is not selected on exposure and outcome (Greenland, 2003). Sixth, focusing on a European population may limit generalizability, but helps reduce bias from population stratification. Seventh, we excluded replicated or very similar attributes, potentially missing some attributes. However, this reduces false negatives when adjusting for multiple comparisons. Eighth, this study is open to selection bias because genetic endowment is lifelong but the UK Biobank participants are middle-aged or older, so potential recruits who did not live to recruitment because of their BMI are missing, which may bias estimates towards the null or inverse for harmful binary outcomes (Thompson et al., 2013), but less likely affects continuous outcomes (Smit et al., 2019). Ninth, we focused on BMI, because it is well accepted and easy to measure, although waist-to-hip ratio may be a better marker for mortality. Tenth, although this study primarily utilized sex-specific BMI, we also conducted analyses using overall

medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .

BMI from GIANT including the UK Biobank, which gave a generally similar interpretation (**S Table 5**). Using a sex-specific BMI may lead to lower statistical power than using overall population BMI, but allows for the detection of traits that are affected differently by BMI by sex. Including findings from the overall population BMI makes the results more comparable to previous similar studies. Eleventh, our study did not find associations of BMI with some early-onset cancers possibly because early-onset cancers tend to be of germline origin (Qing et al., 2020). Twelfth, we used information from Neale Lab, which removed participants whose self-reported sex differed from biological sex in the quality check (*Whole genome analysis pipeline*), so our findings only relate to cis men and women. Thirteenth, while overlapping samples for BMI and the outcomes pose a potential issue in the main analysis, we also replicated the analysis using sex-specific instruments for BMI independent of the UK Biobank, and results were similar, suggesting bias from overlapping samples is minimal. Consistently, overlapping samples have been shown to make most difference for relatively small samples and for MR-Egger estimates, particularly when the I^2_{GX} is low, leading to confounded estimates. (Minelli et al., 2021). Lastly, we used linear MR, so we cannot exclude the possibility of a "J" shaped association, however trials of incretins suggest a "J" curve may be a manifestation of bias (Wilding et al., 2021).

Public health implications

Trials of incretins have clearly indicated many benefits of weight loss particularly for older people (Leiter et al., 2019; Lincoff et al., 2023). Our study showing how BMI might increase risk of chronic diseases, reduce quality of life, and depress mental health in women, underscores the importance of addressing the obesity epidemic equitably in men and women, given differences by sex in some consequences of adiposity, such as psychological disorders, lipids and cancer. Overweight and obesity is more common in men than women in developed countries (Kanter & Caballero, 2012; Maruyama & Nakamura, 2018). With ongoing global economic development, the same pattern is likely to become more common. Currently women are more likely

medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .

to seek medical intervention for weight management, such as semaglutide, than men (Luthra, 2023). To promote population health, it is important to address the unique challenges faced by overweight men and women. Men are at higher risk of heart disease and premature death due to high BMI, so targeted public health interventions, such as sex-specific weight recommendations or guidelines, could perhaps be considered.

Conclusion

Our contemporary systematic examination found BMI associated with a broad range of health-related attributes. We also found significant sex differences in many traits, underscoring the importance of addressing higher BMI in both men and women possibly as means of redressing differences in life expectancy. Ultimately, our study emphasizes the harmful effects of obesity and the importance of maintaining a healthy BMI. Whether BMI recommendations should differ by sex might be considered.

medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .

Abbreviations

ApoB: apolipoprotein B; BMI: body mass index; GIANT: Genetic Investigation of ANthropometric Traits; GWAS: genome-wide association study; HDL-c: high density lipoprotein cholesterol; LDL: low density lipoprotein; MR: Mendelian Randomization; SHBG: sex hormone binding globulin

Acknowledgements

We would like to thank the Neale Lab for providing GWAS summary-level statistics.

Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Consent for publication

All participants were consent for publication.

Ethical approval

The study protocol was not pre-registered. We used only publicly available summary-level data and did not collect any original data in this study. Ethics approval and consent from individual participants can be found in the original publications.

Availability of data and material

This study used data from the MR-base platform (https://www.mrbase.org/), UK Biobank (http://www.nealelab.is/uk-biobank/).

Credit author statement

Conceptualization: Zhu Liduzi Jiesisibieke, C Mary Schooling

Data curation: Zhu Liduzi Jiesisibieke, C Mary Schooling, Io Ieong Chan, Jack Chun Man Ng

Formal analysis: Zhu Liduzi Jiesisibieke, C Mary Schooling

Methodology: Zhu Liduzi Jiesisibieke, C Mary Schooling, Io Ieong Chan, Jack Chun Man Ng

Project administration: C Mary Schooling

Resources: C Mary Schooling

medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .

Software: Zhu Liduzi Jiesisibieke, C Mary Schooling

Supervision: C Mary Schooling

Visualization: Zhu Liduzi Jiesisibieke, C Mary Schooling

Writing – original draft: Zhu Liduzi Jiesisibieke, C Mary Schooling

Writing – review & editing: Zhu Liduzi Jiesisibieke, C Mary Schooling

medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .

References

Altman, D. G., & Bland, J. M. (2003). Interaction revisited: the difference between two estimates. *bmj*, *326*(7382), 219. https://doi.org/10.1136/bmj.326.7382.219

Ardissino, M., Reddy, R. K., Slob, E. A., Patel, K. H., Ryan, D. K., Gill, D., & Ng, F. S. (2022). Sleep disordered breathing, obesity and atrial fibrillation: A Mendelian randomisation study. *Genes*, *13*(1), 104.

Bell, J. A., Richardson, T. G., Wang, Q., Sanderson, E., Palmer, T., Walker, V., O'Keeffe, L. M., Timpson, N. J., Cichonska, A., & Julkunen, H. (2022). Effects of general and central adiposity on circulating lipoprotein, lipid, and metabolite levels in UK Biobank: a multivariable Mendelian randomization study. *The Lancet Regional Health–Europe*, *21*.

Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)*, *57*(1), 289-300.

Chooi, Y. C., Ding, C., & Magkos, F. (2019). The epidemiology of obesity. *Metabolism*, *92*, 6-10.

Collaboration, P. S. (2009). Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *The Lancet*, *373*(9669), 1083-1096.

Collins, R. (2012). What makes UK Biobank special? *The Lancet*, *379*(9822), 1173-1174.

Costanzo, M., Caterino, M., Sotgiu, G., Ruoppolo, M., Franconi, F., & Campesi, I. (2022). Sex differences in the human metabolome. *Biology of sex differences*, *13*(1), 30.

Davey Smith, G., & Ebrahim, S. (2003). 'Mendelian randomization': can genetic epidemiology contribute to understanding environmental determinants of disease? *International journal of epidemiology*, *32*(1), 1-22.

Esnaola, I., Rodríguez, A., & Goñi, A. (2010). Body dissatisfaction and perceived sociocultural pressures: Gender and age differences. *Salud mental*, *33*(1), 21-29.

Finucane, M. M., Stevens, G. A., Cowan, M. J., Danaei, G., Lin, J. K., Paciorek, C. J., Singh, G. M., Gutierrez, H. R., Lu, Y., & Bahalim, A. N. (2011). National, regional, and global trends in body-mass index since 1980: systematic analysis of health examination surveys and epidemiological studies with 960 country-years and 9· 1 million participants. *The Lancet*, *377*(9765), 557-567.

Fontaine, K. R., Redden, D. T., Wang, C., Westfall, A. O., & Allison, D. B. (2003). Years of life lost due to obesity. *Jama*, *289*(2), 187-193.

Greenland, S. (2003). Quantifying biases in causal models: classical confounding vs collider-stratification bias. *Epidemiology*, *14*(3), 300-306.

Hyppönen, E., Mulugeta, A., Zhou, A., & Santhanakrishnan, V. K. (2019). A data-driven approach for studying the role of body mass in multiple diseases: a phenome-wide registry-based case-control study in the UK Biobank. *The Lancet Digital Health*, *1*(3), e116-e126.

Kahn, S. E., Hull, R. L., & Utzschneider, K. M. (2006). Mechanisms linking obesity to insulin resistance and type 2 diabetes. *Nature*, *444*(7121), 840-846.

Kanter, R., & Caballero, B. (2012). Global gender disparities in obesity: a review. *Advances in Nutrition*, *3*(4), 491-498.

Khan, S. S., Ning, H., Wilkins, J. T., Allen, N., Carnethon, M., Berry, J. D., Sweis, R. N., & Lloyd-Jones, D. M. (2018). Association of body mass index with lifetime risk of cardiovascular disease and compression of morbidity. *JAMA cardiology*, *3*(4), 280-287.

medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .

Larsson, S. C., Bäck, M., Rees, J. M., Mason, A. M., & Burgess, S. (2020). Body mass index and body composition in relation to 14 cardiovascular conditions in UK Biobank: a Mendelian randomization study. *European heart journal*, *41*(2), 221-226.

Larsson, S. C., & Burgess, S. (2021). Causal role of high body mass index in multiple chronic diseases: a systematic review and meta-analysis of Mendelian randomization studies. *BMC medicine*, *19*, 1-10.

Leiter, L. A., Bain, S. C., Hramiak, I., Jódar, E., Madsbad, S., Gondolf, T., Hansen, T., Holst, I., & Lingvay, I. (2019). Cardiovascular risk reduction with once-weekly semaglutide in subjects with type 2 diabetes: a post hoc analysis of gender, age, and baseline CV risk profile in the SUSTAIN 6 trial. *Cardiovascular Diabetology*, *18*, 1-12.

Lincoff, A. M., Brown-Frandsen, K., Colhoun, H. M., Deanfield, J., Emerson, S. S., Esbjerg, S., Hardt-Lindberg, S., Hovingh, G. K., Kahn, S. E., & Kushner, R. F. (2023). Semaglutide and cardiovascular outcomes in obesity without diabetes. *New England Journal of Medicine*, *389*(24), 2221-2232.

Locke, A. E., Kahali, B., Berndt, S. I., Justice, A. E., Pers, T. H., Day, F. R., Powell, C., Vedantam, S., Buchkovich, M. L., & Yang, J. (2015). Genetic studies of body mass index yield new insights for obesity biology. *Nature*, *518*(7538), 197-206.

Luthra, S. (2023). Weight-loss drugs like Ozempic are more likely to interest women than men, poll finds. *The 19th News*. https://19thnews.org/2023/08/weight-loss-drugs-ozempic-women-poll/

Marso, S. P., Bain, S. C., Consoli, A., Eliaschewitz, F. G., Jódar, E., Leiter, L. A., Lingvay, I., Rosenstock, J., Seufert, J., & Warren, M. L. (2016). Semaglutide and cardiovascular outcomes in patients with type 2 diabetes. *New England Journal of Medicine*, *375*(19), 1834-1844.

Maruyama, S., & Nakamura, S. (2018). Why are women slimmer than men in developed countries? *Economics & Human Biology*, *30*, 1-13.

Millard, L. A., Davies, N. M., Timpson, N. J., Tilling, K., Flach, P. A., & Smith, G. D. (2015). MR-PheWAS: hypothesis prioritization among potential causal effects of body mass index on many outcomes, using Mendelian randomization. *Scientific reports*, *5*(1), 16645.

Minelli, C., Del Greco M, F., van der Plaat, D. A., Bowden, J., Sheehan, N. A., & Thompson, J. (2021). The use of two-sample methods for Mendelian randomization analyses on single large datasets. *International journal of epidemiology*, *50*(5), 1651-1659.

Newman, J. D., & Handelsman, D. J. (2014). Challenges to the measurement of oestradiol: comments on an endocrine society position statement. *The Clinical Biochemist Reviews*, *35*(2), 75.

Okorodudu, D. O., Jumean, M., Montori, V. M., Romero-Corral, A., Somers, V. K., Erwin, P. J., & Lopez-Jimenez, F. (2010). Diagnostic performance of body mass index to identify obesity as defined by body adiposity: a systematic review and meta-analysis. *International journal of obesity*, *34*(5), 791-799.

Piva, E., Brugnara, C., Spolaore, F., & Plebani, M. (2015). Clinical utility of reticulocyte parameters. *Clinics in Laboratory Medicine*, *35*(1), 133-163.

Power, M. L., & Schulkin, J. (2008). Sex differences in fat storage, fat metabolism, and the health risks from obesity: possible evolutionary origins. *British journal of nutrition*, *99*(5), 931-940.

Qing, T., Mohsen, H., Marczyk, M., Ye, Y., O'Meara, T., Zhao, H., Townsend, J. P., Gerstein, M., Hatzis, C., & Kluger, Y. (2020). Germline variant burden in cancer genes correlates with age at diagnosis and somatic mutation burden. *Nature communications*, *11*(1), 2438.

medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .

Ramírez, J., van Duijvenboden, S., Young, W. J., Orini, M., Jones, A. R., Lambiase, P. D., Munroe, P. B., & Tinker, A. (2021). Analysing electrocardiographic traits and predicting cardiac risk in UK biobank. *JRSM Cardiovasc Dis*, *10*, 20480040211023664. https://doi.org/10.1177/20480040211023664

Riaz, H., Khan, M. S., Siddiqi, T. J., Usman, M. S., Shah, N., Goyal, A., Khan, S. S., Mookadam, F., Krasuski, R. A., & Ahmed, H. (2018). Association between obesity and cardiovascular outcomes: a systematic review and meta-analysis of Mendelian randomization studies. *JAMA network open*, *1*(7), e183788-e183788.

Schwartz, M. B., & Brownell, K. D. (2004). Obesity and body image. *Body image*, *1*(1), 43-56.

Skaaby, T., Taylor, A. E., Thuesen, B. H., Jacobsen, R. K., Friedrich, N., Møllehave, L. T., Hansen, S., Larsen, S. C., Völker, U., & Nauck, M. (2018). Estimating the causal effect of body mass index on hay fever, asthma and lung function using Mendelian randomization. *Allergy*, *73*(1), 153-164.

Smit, R. A., Trompet, S., Dekkers, O. M., Jukema, J. W., & Le Cessie, S. (2019). Survival bias in Mendelian randomization studies: a threat to causal inference. *Epidemiology*, *30*(6), 813-816.

Thompson, C. A., Zhang, Z.-F., & Arah, O. A. (2013). Competing risk bias to explain the inverse relationship between smoking and malignant melanoma. *European journal of epidemiology*, *28*, 557-567.

Verma, A., Bradford, Y., Dudek, S., Lucas, A. M., Verma, S. S., Pendergrass, S. A., & Ritchie, M. D. (2018). A simulation study investigating power estimates in phenome-wide association studies. *BMC bioinformatics*, *19*, 1-8.

WHO. (2024). *Obesity and Overweight*. https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight

Whole genome analysis pipeline. Neale lab. Retrieved June 10 from https://github.com/Nealelab/whole_genome_analysis_pipeline/blob/main/README.md#sex-check-sample-qc

Wilding, J. P., Batterham, R. L., Calanna, S., Davies, M., Van Gaal, L. F., Lingvay, I., McGowan, B. M., Rosenstock, J., Tran, M. T., & Wadden, T. A. (2021). Once-weekly semaglutide in adults with overweight or obesity. *New England Journal of Medicine*, *384*(11), 989-1002.

Yengo, L., Sidorenko, J., Kemper, K. E., Zheng, Z., Wood, A. R., Weedon, M. N., Frayling, T. M., Hirschhorn, J., Yang, J., & Visscher, P. M. (2018). Meta-analysis of genome-wide association studies for height and body mass index in∼ 700000 individuals of European ancestry. *Human molecular genetics*, *27*(20), 3641-3649.

Zanetti, D., Bergman, H., Burgess, S., Assimes, T. L., Bhalla, V., & Ingelsson, E. (2020). Urinary albumin, sodium, and potassium and cardiovascular outcomes in the UK Biobank: observational and Mendelian randomization analyses. *Hypertension*, *75*(3), 714-722.

Zhou, Y., Zhang, X., Li, X., Zhu, G., Gao, T., Deng, Y., Huang, L., & Liu, Z. (2024). Anthropometric indicators may explain the high incidence of follicular lymphoma in Europeans: Results from a bidirectional two-sample two-step Mendelian randomisation. *Gene*, *911*, 148320.

medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .

Figure legends

Figure 1. Selection criteria for the health-related attributes

Figure 2. Manhattan plot of body mass index on health-related attributes in men (n: 167,020) from the UK Biobank

Figure 3. Manhattan plot of body mass index on health-related attributes in women (n: 194,174) from the UK Biobank

Figure 4. Sex differences of body mass index on health-related attributes in men (n: 167,020) and women (n: 194,174)

Supplementary tables

S Table 1. Phenome-wide association analysis of causal estimates for body mass index (women: 194,174; men: 167,020) on attributes (women: 194,174; men: 167,020) from the UK Biobank based on inverse variance weighting

S Table 2. Phenome-wide association analysis of causal estimates for body mass index (women: 171,977, men: 152,983) from the Genetic Investigation of ANthropometric Traits (GIANT) on attributes (women: 194,174; men: 167,020) from the UK Biobank based on inverse variance weighting

S Table 3. Number of significant associations of body mass index (women: 194,174; men: 167,020) with traits (women: 194,174; men: 167,020) by category

S Table 4. MR-Egger sensitivity analysis of phenome-wide association causal estimates for body mass index on various attributes in women (n:94,174) and men (n:167,020) from the UK Biobank

S Table 5. Phenome-wide association analysis of causal estimates for body mass index (n: 681,275) from the Genetic Investigation of ANthropometric Traits (GIANT) on attributes (women: 194,174; men: 167,020) from the UK Biobank based on inverse variance weighting

Abbreviations used in supplementary tables:

Irnt: inverse rank normalized; SHBG: sex hormone binding globulin; GIANT: Genetic Investigation of ANthropometric Traits; GV: genetic variant; GWAS: genome-wide association study; ICD: international classification of diseases.



medRxiv preprint doi: https://doi.org/10.1101/2024.09.12.24313524; this version posted September 13, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a CC-BY 4.0 International license .


Men
-log10(p_BH)
4.20
4.00
3.50
3.00
2.50
2.00
1.35
Basal metabolic rate
HDL cholesterol
Diabetes diagnosed by doctor
High blood pressure
Testosterone
Carpal tunnel syndrome
Ischaemic heart disease
Facial ageing
Cholesterol
Major coronary heart disease event
FEV1
SHBG
LDL direct
Sleep apnoea
Triglycerides
Heart failure
Malignant neoplasm of digestive organs
Hypertensive heart and/or renal disease
Primary_lymphoid and hematopoietic malignant neoplasms
Malignant neoplasm of oesophagus
Hair/balding pattern: Pattern 4
Age at recruitment
Blood assays
Age at recruitment
Cognitive function
Diseases of the circulatory system
Diseases of the digestive system
Diseases of the genitourinary system
Diseases of the hematopoietic system and blood disorders
Diseases of the musculoskeletal system and connective tissue
Diseases of the nervous system
Diseases of the respiratory system
Diseases of the sensory system
Diseases of the skin and subcutaneous tissue
Endocrine and metabolic diseases
Health and medical history
Infectious diseases
Lifestyle and environmental
Medical conditions
Neoplasms
Operations
Physical measures
Physiological factors
Psychological disorders
Sex-specific factors
Symptoms
Urine assays
Direction
Inverse association
Positive association


Women
-log10(p_BH)
4.20
4.00
3.50
3.00
2.50
2.00
1.35
Basal metabolic rate
Diabetes diagnosed by doctor
High blood pressure
HDL cholesterol
Triglycerides
SHBG
Carpal tunnel syndrome
FVC
FEV1
Ischaemic heart disease
Major coronary heart disease event
hypothyroidism/myxoedema
Sleep apnoea
Cholesterol
Heart failure
depression
Apolipoprotein B
Facial ageing
uterine/endometrial cancer
Malignant neoplasm of bronchus and lung
malignant neoplasm of female genital organs
Urine assays
Symptoms
Sex-specific factors
Psychological disorders
Pregnancy, childbirth, and the puerperium
Physiological factors
Physical measures
Operations
Neoplasms
Medical conditions
Lifestyle and environmental
Infectious diseases
Health and medical history
Endocrine and metabolic diseases
Diseases of the skin and subcutaneous tissue
Diseases of the sensory system
Diseases of the respiratory system
Diseases of the nervous system
Diseases of the musculoskeletal system and connective
Diseases of the hematopoietic system and blood disorde
Diseases of the genitourinary system
Diseases of the digestive system
Diseases of the circulatory system
Age at recruitment
Cognitive function
Blood assays

Health-related traits	Category	Men	Women	Directionally discord between sexes	Absolute estimations
SHBG	Blood assays	Inverse	Inverse	No	Greater absolute estimations in women
Albumin	Blood assays	Inverse	Inverse	No	Greater absolute estimations in women
Vitamin D	Blood assays	Inverse	Inverse	No	Greater absolute estimations in women
Cholesterol	Blood assays	Inverse	Inverse	No	Greater absolute estimations in men
Total bilirubin	Blood assays	Inverse	Inverse	No	Greater absolute estimations in women
LDL direct	Blood assays	Inverse	Inverse	No	Greater absolute estimations in men
Direct bilirubin	Blood assays	Inverse	Inverse	No	Greater absolute estimations in women
Platelet count	Blood assays	Inverse	Positive	Yes	Greater absolute estimations in men
Testosterone	Blood assays	Inverse	Positive	Yes	Greater absolute estimations in men
Apoliprotein B	Blood assays	Inverse	Positive	Yes	Greater absolute estimations in women
Platelet crit	Blood assays	Inverse	Positive	Yes	Greater absolute estimations in women
Aspartate aminotransferase	Blood assays	Positive	Positive	No	Greater absolute estimations in men
Alkaline phosphatase	Blood assays	Positive	Positive	No	Greater absolute estimations in women
Neutrophill count	Blood assays	Positive	Positive	No	Greater absolute estimations in women
Red blood cell (erythrocyte) distribution width	Blood assays	Positive	Positive	No	Greater absolute estimations in women
Triglycerides	Blood assays	Positive	Positive	No	Greater absolute estimations in women
Urate	Blood assays	Positive	Positive	No	Greater absolute estimations in women
C-reactive protein	Blood assays	Positive	Positive	No	Greater absolute estimations in women
Diagnoses - main ICD10: I35 Nonrheumatic aortic valve disorders	Diseases of the circulatory system	Positive	Positive	No	Greater absolute estimations in men
Peripheral artery disease	Diseases of the circulatory system	Positive	Positive	No	Greater absolute estimations in men
Death due to cardiac causes	Diseases of the circulatory system	Positive	Positive	No	Greater absolute estimations in men
Heart failure,strict	Diseases of the circulatory system	Positive	Positive	No	Greater absolute estimations in men
Diagnoses - main ICD10: I26 Pulmonary embolism	Diseases of the circulatory system	Positive	Positive	No	Greater absolute estimations in men
DVT of lower extremities and pulmonary embolism	Diseases of the circulatory system	Positive	Positive	No	Greater absolute estimations in men
Venous thromboembolism	Diseases of the circulatory system	Positive	Positive	No	Greater absolute estimations in men
Myocardial infarction, strict	Diseases of the circulatory system	Positive	Positive	No	Greater absolute estimations in men
Major coronary heart disease event	Diseases of the circulatory system	Positive	Positive	No	Greater absolute estimations in men
Major coronary heart disease event excluding revascularizations	Diseases of the circulatory system	Positive	Positive	No	Greater absolute estimations in men
Coronary atherosclerosis	Diseases of the circulatory system	Positive	Positive	No	Greater absolute estimations in men
Ischaemic heart disease, wide definition	Diseases of the circulatory system	Positive	Positive	No	Greater absolute estimations in men
Diseases of the circulatory system	Diseases of the circulatory system	Positive	Positive	No	Greater absolute estimations in men
Diagnoses - main ICD10: K60 Fissure and fistula of anal and rectal regions	Diseases of the digestive system	Positive	Positive	No	Greater absolute estimations in men
Diagnoses - main ICD10: K81 Cholecystitis	Diseases of the digestive system	Positive	Positive	No	Greater absolute estimations in women
Hernia	Diseases of the digestive system	Inverse	Positive	Yes	Greater absolute estimations in women
Diagnoses - main ICD10: K80 Cholelithiasis	Diseases of the digestive system	Positive	Positive	No	Greater absolute estimations in women
Disorders of gallbladder, biliary tract and pancreas	Diseases of the digestive system	Positive	Positive	No	Greater absolute estimations in women
Diseases of the digestive system	Diseases of the digestive system	Positive	Positive	No	Greater absolute estimations in women
Diagnoses - main ICD10: N17 Acute renal failure	Diseases of the genitourinary system	Positive	Positive	No	Greater absolute estimations in men
Diagnoses - main ICD10: N39 Other disorders of urinary system	Diseases of the genitourinary system	Positive	Positive	No	Greater absolute estimations in women
Other anaemias	Diseases of the hematopoietic system and blood disorders	Inverse	Positive	Yes	Greater absolute estimations in women
Other and unspecified anaemias	Diseases of the hematopoietic system and blood disorders	Inverse	Positive	Yes	Greater absolute estimations in women
Iron deficiency anaemia	Diseases of the hematopoietic system and blood disorders	Inverse	Positive	Yes	Greater absolute estimations in women
Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	Diseases of the hematopoietic system and blood disorders	Positive	Positive	No	Greater absolute estimations in women
Cervicalgia	Diseases of the musculoskeletal system and connective tissue	Inverse	Positive	Yes	Greater absolute estimations in women
Soft tissue disorders related to use, overuse and pressure	Diseases of the musculoskeletal system and connective tissue	Positive	Positive	No	Greater absolute estimations in men
Impingement syndrome of shoulder	Diseases of the musculoskeletal system and connective tissue	Positive	Positive	No	Greater absolute estimations in women
Pain in limb	Diseases of the musculoskeletal system and connective tissue	Positive	Positive	No	Greater absolute estimations in women
Other soft tissue disorders, not elsewhere classified	Diseases of the musculoskeletal system and connective tissue	Positive	Positive	No	Greater absolute estimations in women
Diagnoses - main ICD10: M79 Other soft tissue disorders, not elsewhere classified	Diseases of the musculoskeletal system and connective tissue	Positive	Positive	No	Greater absolute estimations in women
Diagnoses - main ICD10: G57 Mononeuropathies of lower limb	Diseases of the nervous system	Inverse	Positive	Yes	Greater absolute estimations in women
Sleep apnoea	Diseases of the nervous system	Positive	Positive	No	Greater absolute estimations in men
Diagnoses - main ICD10: G47 Sleep disorders	Diseases of the nervous system	Positive	Positive	No	Greater absolute estimations in men
Carpal tunnel syndrome	Diseases of the nervous system	Positive	Positive	No	Greater absolute estimations in women
Nerve, nerve root and plexus disorders	Diseases of the nervous system	Positive	Positive	No	Greater absolute estimations in women
Diagnoses - main ICD10: J18 Pneumonia, organism unspecified	Diseases of the respiratory system	Positive	Positive	No	Greater absolute estimations in men
Diseases of the respiratory system	Diseases of the respiratory system	Positive	Positive	No	Greater absolute estimations in men
Diagnoses - main ICD10: H61 Other disorders of external ear	Diseases of the sensory system	Positive	Inverse	Yes	Greater absolute estimations in men
Degeneration of macula and posterior pole	Diseases of the sensory system	Inverse	Positive	Yes	Greater absolute estimations in women
Diagnoses - main ICD10: L03 Cellulitis	Diseases of the skin and subcutaneous tissue	Positive	Positive	No	Greater absolute estimations in men
Diabetes diagnosed by doctor	Endocrine and metabolic diseases	Positive	Positive	No	Greater absolute estimations in men
Blood clot, DVT, bronchitis, emphysema, asthma, rhinitis, eczema, allergy diagnosed by doctor: Hayfever, allergic rhinitis or eczema	Health and medical history	Inverse	Inverse	No	Greater absolute estimations in men
Blood clot, DVT, bronchitis, emphysema, asthma, rhinitis, eczema, allergy diagnosed by doctor: None of the above	Health and medical history	Positive	Inverse	Yes	Greater absolute estimations in women
Vascular/heart problems diagnosed by doctor: Heart attack	Health and medical history	Positive	Positive	No	Greater absolute estimations in men
Chest pain or discomfort	Health and medical history	Positive	Positive	No	Greater absolute estimations in women
Mouth/teeth dental problems: Dentures	Health and medical history	Positive	Positive	No	Greater absolute estimations in women
Morning/evening person (chronotype)	Lifestyle and environmental	Inverse	Inverse	No	Greater absolute estimations in men
Facial ageing	Lifestyle and environmental	Positive	Positive	No	Greater absolute estimations in men
Non-cancer illness code, self-reported: osteoporosis	Medical conditions	Inverse	Inverse	No	Greater absolute estimations in women
Non-cancer illness code, self-reported: headaches (not migraine)	Medical conditions	Inverse	Positive	Yes	Greater absolute estimations in men
Non-cancer illness code, self-reported: pernicious anaemia	Medical conditions	Inverse	Positive	Yes	Greater absolute estimations in women
Non-cancer illness code, self-reported: psoriatic arthropathy	Medical conditions	Inverse	Positive	Yes	Greater absolute estimations in women
Non-cancer illness code, self-reported: eye/eyelid problem	Medical conditions	Inverse	Positive	Yes	Greater absolute estimations in men
Non-cancer illness code, self-reported: allergy/hypersensitivity/anaphylaxis	Medical conditions	Inverse	Positive	Yes	Greater absolute estimations in women
Non-cancer illness code, self-reported: ulcerative colitis	Medical conditions	Positive	Inverse	Yes	Greater absolute estimations in men
Non-cancer illness code, self-reported: bladder problem (not cancer)	Medical conditions	Inverse	Positive	Yes	Greater absolute estimations in women
Non-cancer illness code, self-reported: helicobacter pylori	Medical conditions	Positive	Positive	No	Greater absolute estimations in women
Non-cancer illness code, self-reported: muscle or soft tissue injuries	Medical conditions	Positive	Inverse	Yes	Greater absolute estimations in men
Non-cancer illness code, self-reported: muscle/soft tissue problem	Medical conditions	Inverse	Positive	Yes	Greater absolute estimations in women
Non-cancer illness code, self-reported: gout	Medical conditions	Positive	Positive	No	Greater absolute estimations in men
Non-cancer illness code, self-reported: cervical spondylosis	Medical conditions	Positive	Positive	No	Greater absolute estimations in women
Non-cancer illness code, self-reported: arthritis (nos)	Medical conditions	Positive	Positive	No	Greater absolute estimations in women
Non-cancer illness code, self-reported: unclassifiable	Medical conditions	Inverse	Positive	Yes	Greater absolute estimations in women
Non-cancer illness code, self-reported: cholelithiasis/gall stones	Medical conditions	Positive	Positive	No	Greater absolute estimations in women
Non-cancer illness code, self-reported: depression	Medical conditions	Positive	Positive	No	Greater absolute estimations in women
Non-cancer illness code, self-reported: hypothyroidism/myxoedema	Medical conditions	Positive	Positive	No	Greater absolute estimations in women
Non-cancer illness code, self-reported: gastro-oesophageal reflux (gord) / gastric reflux	Medical conditions	Positive	Positive	No	Greater absolute estimations in women
Non-cancer illness code, self-reported: angina	Medical conditions	Positive	Positive	No	Greater absolute estimations in men
Non-cancer illness code, self-reported: asthma	Medical conditions	Positive	Positive	No	Greater absolute estimations in women
Non-cancer illness code, self-reported: osteoarthritis	Medical conditions	Positive	Positive	No	Greater absolute estimations in women
Malignant neoplasm of urinary organs	Neoplasms	Positive	Inverse	Yes	Greater absolute estimations in men
Diagnoses - main ICD10: D35 Benign neoplasm of other and unspecified endocrine glands	Neoplasms	Positive	Positive	No	Greater absolute estimations in women
Malignant neoplasm of colon	Neoplasms	Positive	Inverse	Yes	Greater absolute estimations in men
Lymphomas	Neoplasms	Positive	Positive	No	Greater absolute estimations in men
Malignant neoplasm of digestive organs	Neoplasms	Positive	Positive	No	Greater absolute estimations in men
Hand grip strength (right)	Physical measures	Positive	Inverse	Yes	Greater absolute estimations in women
Diastolic blood pressure, automated reading	Physical measures	Positive	Positive	No	Greater absolute estimations in women
Sensitivity / hurt feelings	Physiological factors	Inverse	Positive	Yes	Greater absolute estimations in women
Neuroticism score	Physiological factors	Inverse	Positive	Yes	Greater absolute estimations in men
Seen doctor (GP) for nerves, anxiety, tension or depression	Physiological factors	Inverse	Positive	Yes	Greater absolute estimations in women
Miserableness	Physiological factors	Positive	Positive	No	Greater absolute estimations in women
Loneliness, isolation	Physiological factors	Positive	Positive	No	Greater absolute estimations in women
Fed-up feelings	Physiological factors	Positive	Positive	No	Greater absolute estimations in women
Diagnoses - main ICD10: R04 Haemorrhage from respiratory passages	Symptoms	Positive	Inverse	Yes	Greater absolute estimations in men
Diagnoses - main ICD10: R31 Unspecified haematuria	Symptoms	Positive	Inverse	Yes	Greater absolute estimations in men
Potassium in urine	Urine assays	Positive	Positive	No	Greater absolute estimations in women