

1 **Analysis of IL-4 Levels, IL-1 α , Cortisol Hormone, and Blood Cell Number**
2 **in Parkinson's Disease Patients Undergoing Dance-based Movement**
3 **Therapy: A Preliminary Study**

4 Ethical committee of the Federal University of Pará (CEP: 3.965.311/ CAAE:
5 27811119.4.0000.0018) and the Brazilian Registry of Clinical Trials (ReBEC RBR-
6 3bhbrb5, UTN code: U1111-1275-6679)

7 Karen Adriana Pastana Marques¹, Gabriel Mesquita da Conceição Bahia¹, Juliana dos
8 Santos Duarte¹, Denise da Silva Pinto², Arnaldo Jorge Martins Filho³, Evander de Jesus
9 Oliveira Batista⁴, Anderson Manoel Herculano Oliveira da Silva⁵, Karen Renata Herculano
10 Matos Oliveira⁵, Lane Viana Krejcová⁶, and Carlomagno Pacheco Bahia^{1*}.

11
12 **Affiliations**

13 ¹Neuroplasticity Lab, Institute of Health Sciences, Federal University of Pará, Belém,
14 Pará, Brazil.

15 ²Functional Rehab Laboratory, Institute of Health Sciences, Federal University of Pará,
16 Belém, Pará, Brazil.

17 ³Evandro Chagas Institute, Belém, Pará, Brazil.

18 ⁴Protozoology Laboratory, Tropical Medicine Nucleus, Federal University of Pará, Belém,
19 Pará, Brazil.

20 ⁵Experimental Neuropharmacology Lab, Institute of Biologic Science, Federal University
21 of Pará, Belém, Pará, Brazil.

22 ⁶College of Dance, Institute of Art Sciences, Federal University of Pará, Belém, Pará,
23 Brazil.

24
25 * **Corresponding author:** Carlomagno Pacheco Bahia. Email address:
26 carlomagnobahia@gmail.com carlomagno@ufpa.br (CPB)

27

28 **Abstract**

29 **Introduction:** The interplay between neuroinflammation and Parkinson's disease (PD) is
30 still unclear, although several studies suggest that neuroinflammation plays an
31 important role in the pathophysiological pathways of neurodegenerative disorders.
32 Dance-based movement therapy has been studied as an adjuvant treatment for PD
33 symptoms by modulating some inflammatory interleukins and improving the overall
34 effect of pharmacological treatments.

35 **Objective:** To evaluate the levels of Interleukin 4, Interleukin 1 α , cortisol hormone, and
36 blood cell total numbers in PD subjects undergoing dance-based movement therapy.

37 **Methods:** Twenty-five participants were selected for this study. Seventeen subjects
38 diagnosed with PD were divided into two groups: PD group (n=8) and PD+dance group
39 (n=9). Another group was composed of age-matched healthy controls (n=8). The dance-
40 based movement therapy was performed twice a week for 50-60 minutes/session over
41 six months.

42 **Results:** The serum IL-4 levels were significantly lower in the PD group when compared
43 to the healthy group, while the PD+Dance group showed similar levels to the healthy
44 group. The levels of IL-1 α , cortisol, and blood cell count showed no statistical differences
45 between groups.

46 **Conclusion:** Our results suggest that regular practice of dance-based movement therapy
47 may induce a neuroprotective effect by modulating the levels of IL-4 in people with PD
48 similarly to healthy participants.

49

50 **Keywords:** Parkinson's Disease; Dance-based Movement Therapy; Interleukins; Cortisol;
51 Blood Cells.

52

53 **Trial registration:** The study was registered in the Brazilian registry of clinical trials: RBR-
54 3bhbrb5.

55

56 **Short title:** Dance-based movement therapy as adjuvant treatment for Parkinson's
57 disease

58

59 Introduction

60 The interplay between inflammation and Parkinson’s disease (PD) is still unclear,
61 although inflammation plays an important role in neurodegenerative disorders[1] and
62 may help promote neuronal death and involve blood cell abnormalities[2].

63 The effects of physical exercise on motor and non-motor symptoms in people
64 with PD effectively modulate the inflammatory process[3], promoting neuronal survival
65 by increasing the production of neurotrophic factors, releasing neurotransmitters, and
66 improving neurogenesis, angiogenesis, modulating inflammatory cytokines, and cortisol
67 hormone expression[3,4].

68 Thus, therapy based on dance movements may be a physical activity alternative
69 to improve PD treatment [5], once dance involves motor skills such as motion planning,
70 rhythm, motor memory, proprioception, imitation[5–7], emotional aspects, and
71 visuospatial information, in addition to increasing body balance, integrating brain areas,
72 decreasing cortisol hormone, and improving neuroplasticity [5,6,8,9].

73 To this date, no study has analyzed the serum levels of inflammatory cytokines,
74 the stress hormone (cortisol), and the blood cell number in people with PD undergoing
75 dance-based movement therapy. Therefore, the present study evaluated the serum
76 levels of Interleukin 4 (IL-4), Interleukin 1 α (IL-1 α), cortisol hormone, and total blood cell
77 numbers in people with Parkinson's disease undergoing physical activity based on dance
78 movements.

79

80 Methods

81 Study design and Participants

82 We performed a longitudinal study aiming to analyze the effects of physical
83 activity based on dance movements using the “Baila Parkinson” method as a
84 complementary therapy for Parkinson’s disease [10].

85 We submitted this study to the ethical committee of the Federal University of
86 Pará (CEP: 3.965.311/ CAAE: 27811119.4.0000.0018) and registered it in the Brazilian
87 Registry of Clinical Trials (ReBEC RBR-3bhbrb5, UTN code: U1111-1275-6679). The
88 research was developed from July to December 2019, the participants were recruited
89 through social media announcements, the follow-up was until march 2020, and the
90 inclusion criteria were: 1) a diagnosis of PD according to the UK Parkinson’s Disease
91 Society Brain Bank (UKPDSBB); 2) Hoehn and Yahr stages I to III under pharmacological
92 treatment for at least 3 years; and 3) the ability to walk independently. Exclusion criteria:
93 1) presence of other neurologic or neuropsychiatric conditions; 2) comorbid
94 inflammatory disease; or 3) severe cardiopulmonary disease. This study was conducted
95 in accordance with the Declaration of the World Medical Association and all participants
96 provided written informed consent.

97 The study included 25 subjects. Seventeen (n=17) subjects with Parkinson's
98 disease, and eight (n=8) healthy controls (Healthy Group - HG) were selected.
99 Demographic and clinical parameters were documented. Among the subjects with PD,
100 eight received their PD medical treatment as usual (PD Group, n=8), whereas the other
101 nine subjects with PD received medical treatment associated to dance therapy
102 (PD+Dance, n=9). All PD subjects were submitted to evaluation of disease progression
103 through the Unified Parkinson Disease Rating Scale (UPDRS), and fulfilled UKPDSBB
104 clinical criteria for idiopathic PD (H&Y Scale $1,88 \pm 0,33$; UPDRS III $30,55 \pm 13,60$; UPDRS
105 Total $48,77 \pm 19,63$).

106 **Intervention: Dance-based movement therapy**

107 The dance-based movement therapy was performed twice a week, 50-60
108 minutes per session for six months following a protocol based on the “Baila Parkinson”
109 method [10].

110

111 **Data Collection and Statistical Analysis**

112 We collected the peripheral blood of all participants (10 mL through peripheral
113 vein puncture; at 8-9 am after 8h of fasting) and centrifuged it at 3500 g for 10 min. The
114 serum was removed and stored at -80 °C (Sanyo VIP Plus -80°C Freezer) until analysis.

115 The levels of IL-4, IL-1 α , and cortisol were measured using the Enzyme-Linked
116 Immunosorbent Assay - ELISA (Invitrogen®) method. The blood cell number was
117 calculated using impedance and colorimetric measurements (Medical Laboratory
118 Equipment Blood Test CBC). The results were analyzed using GraphPad Prism® 8.0
119 Software. We tested the data for normality (Kolmogorov-Smirnov) and homogeneity
120 (Levine test). The data were analyzed using a one-way analysis of variance (ANOVA)
121 corrected by the Tukey intergroup post-test or Kruskal-Wallis test for non-parametric
122 data. Statistical significance was $p < 0.05$, and all data are presented as mean \pm standard
123 deviation (M \pm SD).

124

125 **Results**

126 The present study aimed to the serum levels of Interleukin 4 (IL-4), Interleukin
127 1 α (IL-1 α), cortisol hormone, and total blood cell numbers in people with Parkinson's

128 disease undergoing physical activity based on dance movements. Peripheral blood was
129 collected once and IL-1 α , IL-4, cortisol levels and blood cell count were measured.

130 Twenty-five individuals enrolled in this study which eight were healthy and
131 seventeen were with idiopathic PD. Nine individuals with idiopathic PD were selected to
132 attend the therapeutic sessions (Fig 1). were selected to attend the therapeutic sessions
133 (Fig 1). The demographic and clinical parameters of the three subjects' groups are shown
134 in Table 1. There were no statistically significant differences in age or gender among the
135 three groups.

136 >>> INSERT FIGURE 1 HERE

137

138 **Figure 1. Flowchart of the participant's recruitment for the study.**

139

140 **Table 1. Demographic and clinical features of the three groups.**

Demographic and Clinical Features	Parkinson's Group with Dance (PD + Dance) N=9	Parkinson's Without Dance Group (PD) N=8	Healthy Group (HG) N=8
Sex			
Male	4	4	3
Female	5	4	5
Age (years)	63,22 \pm 9,40	67,25 \pm 13,41	59,12 \pm 10,73
Time since Diagnosis (Years)	4,55 \pm 2,50	7,22 \pm 5,80	-
H&Y Scale	1,88 \pm 0,33	-	-
UPDRS III	30,55 \pm 13,60	-	-
UPDRS Total	48,77 \pm 19,63	-	-

141 UPDRS: Unified Parkinson's disease rating scale; H&Y Scale: Hoehn and Yahr Scale

142 **Pro and Anti-Inflammatory Cytokine Levels**

143 Serum blood levels of IL-4 in the PD group were significantly lower when
144 compared to the healthy group (HG) while the PD+Dance group had IL-4 serum levels
145 similar to the healthy group (HG: 0.37 ± 0.05 ; PD: 0.1280 ± 0.02 ; PD+Dance: 0.2300 ± 0.08 ;
146 $p=0,0079$). Regarding the levels of IL-1 α , there was no statistically significant difference
147 between the groups, but the concentration of IL-1 α in the PD+Dance subjects was lower
148 in relation to the PD group (HG: 0.008 ± 0.004 ; PD group: 0.022 ± 0.02 ; PD+Dance group:
149 0.01 ± 0.01 ; $p=0,5051$), see Figure 2 A and B.

150

151 **Serum Cortisol Levels**

152 The serum blood level of cortisol hormone did not show statistically significant
153 differences among the three groups analyzed (HG: 12.40 ± 3.86 ; PD Group: 11.57 ± 5.31 ;
154 PD + Dance Group: 10.32 ± 2.75 ; ANOVA $p=0,56$), see Figure 2C.

155

156 >>> INSERT FIGURE 2 HERE

157

158 **Figure 2. The blood levels of IL-4, IL-1 α , and cortisol hormone analysis ($\mu\text{g/dL}$).**
159 Statistical difference assessed using one-way ANOVA test ($p < 0.05$). A) Serum levels of
160 interleukin 4: HG $0.37 (\pm 0.05)$; PD $0.1280 (\pm 0.02)$; PD + Dance $0.2300 (\pm 0.08)$; $p=0,0079$.
161 B) Serum levels of interleukin 1 α : HG $0.008 (\pm 0.004)$; PD $0.022 (\pm 0.02)$ and PD + Dance
162 $0.01 (\pm 0.01)$; $p=0,5051$. C) Serum levels of cortisol: HG 12.40 ± 3.86 ; PD Group
163 11.57 ± 5.31 ; PD + Dance Group 10.32 ± 2.75 ; ANOVA $p=0,56$.

164

165 **Blood Cells Count**

166 The blood cell counts did not show statistical differences among the three groups
 167 ($p > 0.05$), see Table 2. However, two participants of the PD group that underwent dance-
 168 based movement therapy (PD+Dance) had changes in red cell numbers, which may
 169 indicate anemia and positive microcytosis (see Table 2). There was also one case with
 170 erythrocytosis, one with hypochromic, another with thrombocytopenia, and the last
 171 with atypical lymphocytes (Table S1).

172

173 **Table 2. Blood cell total numbers of the three groups (Data shown as mean \pm standard**
 174 **deviation and one-way ANOVA test).**

Blood Cell Count	Parkinson's Group with Dance (PD + Dance) N=9	Parkinson's Without Dance Group (PD) N=8	Healthy Group (HG) N=8	p Value
Red Blood Cells ($10^6/\text{mm}^3$)	4848888,88 \pm 589246,22	4878750 \pm 817879,44	5120000 \pm 571564,26	>0,05
Hemoglobin (g/dl)	14,28 \pm 2,16	14,60 \pm 2,99	14,77 \pm 1,40	>0,05
Hematocrit (%)	42,98 \pm 6,24	44,48 \pm 8,67	44,37 \pm 4,13	>0,05
MCV (μm^3)	88,56 \pm 6,49	90,92 \pm 4,07	87,03 \pm 3,33	>0,05
MCH (pg)	29,42 \pm 2,42	29,81 \pm 1,97	28,96 \pm 1,32	>0,05
MCHC (g/dl)	33,21 \pm 0,72	32,78 \pm 0,96	33,77 \pm 1,41	>0,05
White Blood Cells ($10^3/\text{mm}^3$)	6503,33 \pm 1320,70	7070 \pm 1417,58	6343,75 \pm 1079,53	>0,05
Lymphocytes (p/mm^3)	2033,22 \pm 646,03	2173,25 \pm 461,21	2333,50 \pm 567,02	>0,05
Monocytes (p/mm^3)	607204,23 \pm 136,84	400,50 \pm 87,34	438,75 \pm 130,26	>0,05
Platelets ($10^3/\text{mm}^3$)	233111,11 \pm 36285,13	205250 \pm 46092,29	222875 \pm 32148,92	>0,05
MPV (μm^3)	8,48 \pm 1,24	10,93 \pm 3,82	8,81 \pm 0,96	>0,05

175 MCV: Mean Corpuscular Volume; MCH: Mean Corpuscular Hemoglobin; MCHC: Mean
 176 Corpuscular Hemoglobin Concentration; MPV: Mean Platelet Volume.

177

178 **Discussion**

179 The participants in the PD group presented lower blood levels of IL-4 when
180 compared to the HG group. The PD+Dance group had IL-4 concentration levels similar
181 to those of the HG. This finding is relevant given the importance of IL-4 as a regulator
182 for the inflammation process and its relationship with normal neuronal function which
183 is damaged in neurodegenerative diseases[11]. Once physical activity may regulate
184 different kinds of inflammatory cytokines, inhibiting neuroinflammatory responses via
185 the upregulation of interleukins [12], our results suggest that dance-based movement
186 therapy modulated neuroprotective mechanisms across the increasing IL-4 blood serum
187 concentrations in the PD+Dance group (Figure 2A). However, it is necessary to evaluate
188 the cellular and molecular mechanisms involved in the effects of regular therapy based
189 on dance movements.

190 There was no difference among the three groups for IL-1 α serum blood levels.
191 However, the IL-1 α blood serum levels of the PD+Dance group showed lower absolute
192 values compared to the PD group (Figure 2B). Once the cell death is directly related to
193 the regulation of IL-1 α processing and release, conceivably the neurodegenerative and
194 chronic character of Parkinson's disease must be inducing cell death across the constant
195 release of IL-1 α [1,12].

196 There was no difference among the three participants' groups regarding serum
197 cortisol blood levels (Figure 2C). Although our results were not statistically significant,
198 our work demonstrated that therapy based on dance movements was able to conserve
199 cortisol levels in people with PD.

200 There was no statistically significant difference among the three experimental
201 groups regarding the total number of blood cells. For the parameters of blood cell
202 counting, see Table 2. However, the proportion of red blood cells and the platelet count
203 of both the PD+Dance Group and the PD Group showed alterations in some cases, such
204 as anemia and microcytosis (Table S1). The number of red blood cells, leukocytes, and
205 hematocrit values were not affected by the experimental conditions and were similar to
206 a previous study that supports the association between low hemoglobin levels and PD
207 [13] [14].

208

209 **Conclusion**

210 Dance-based movement therapy reduced blood serum levels of IL-4, suggesting
211 that its regular practice may modulate neuroprotective effective action to delay the
212 progression of Parkinson's disease.

213

214 **Acknowledgment**

215 We thank the Health Science Institute and the Federal University of Para for their
216 support and technical assistance.

217

218

219 References

- 220 1. Marogianni C, Sokratous M, Dardiotis E, Hadjigeorgiou GM, Bogdanos D,
221 Xiromerisiou G. Neurodegeneration and inflammation—an interesting interplay
222 in parkinson’s disease. *Int J Mol Sci.* 2020;21: 1–15. doi:10.3390/ijms21228421
223
- 224 2. Bosman GJCGM. Disturbed red blood cell structure and function: An exploration
225 of the role of red blood cells in neurodegeneration. *Front Med (Lausanne).*
226 2018;5. doi:10.3389/fmed.2018.00198
227
- 228 3. Almikhlaifi MA. The role of exercise in Parkinson’s Disease. *Neurosciences.*
229 2023;28: 4–12. doi:10.17712/nsj.2023.1.20220105
230
- 231 4. Ernst M, Folkerts AK, Gollan R, Lieker E, Caro-Valenzuela J, Adams A, et al.
232 Physical exercise for people with Parkinson’s disease: a systematic review and
233 network meta-analysis. *Cochrane Database of Systematic Reviews.* 2023;2023.
234 doi:10.1002/14651858.CD013856.pub2
235
- 236 5. Cheng-Cheng Wu, Huan-Yu Xiong, Jie-Jiao Zheng, Xue-Qiang Wang. Dance
237 movement therapy for neurodegenerative diseases: A systematic review. *Front*
238 *Aging Neurosci.* 2022;14. doi:10.3389/fnagi.2022.975711
239
- 240 6. Lopez-Nieves I, Jakobsche CE. Biomolecular Effects of Dance and
241 Dance/Movement Therapy: A Review. *Am J Dance Ther.* 2022;44: 241–263.
242 doi:10.1007/s10465-022-09368-z
243
- 244 7. Burzynska AZ, Finc K, Taylor BK, Knecht AM, Kramer AF. The dancing brain:
245 Structural and functional signatures of expert dance training. *Front Hum*
246 *Neurosci.* 2017;11. doi:10.3389/fnhum.2017.00566
247
- 248 8. Smyth N, Rossi E, Wood C. Effectiveness of stress-relieving strategies in
249 regulating patterns of cortisol secretion and promoting brain health.
250 *International Review of Neurobiology.* Academic Press Inc.; 2020. pp. 219–246.
251 doi:10.1016/bs.irn.2020.01.003
252
- 253 9. Teixeira-Machado L, Arida RM, de Jesus Mari J. Dance for neuroplasticity: A
254 descriptive systematic review. *Neurosci Biobehav Rev.* 2019;96: 232–240.
255 doi:10.1016/j.neubiorev.2018.12.010
256
- 257 10. dos Santos Duarte J, Alcantara WA, Brito JS, Barbosa LCS, Machado IPR, Furtado
258 VKT, et al. Physical activity based on dance movements as complementary
259 therapy for Parkinson’s disease: Effects on movement, executive functions,
260 depressive symptoms, and quality of life. *PLoS One.* 2023;18.
261 doi:10.1371/journal.pone.0281204
262

- 263 11. Amanollahi M, Jameie M, Heidari A, Rezaei N. The Dialogue Between
264 Neuroinflammation and Adult Neurogenesis: Mechanisms Involved and
265 Alterations in Neurological Diseases. *Molecular Neurobiology*. Springer; 2023.
266 pp. 923–959. doi:10.1007/s12035-022-03102-z
267
- 268 12. Mee-Inta O, Zhao ZW, Kuo YM. Physical exercise inhibits inflammation and
269 microglial activation. *Cells*. 2019;8. doi:10.3390/cells8070691
270
- 271 13. Deng Q, Zhou X, Chen J, Pan M, Gao H, Zhou J, et al. Lower hemoglobin levels in
272 patients with parkinson’s disease are associated with disease severity and iron
273 metabolism. *Brain Res*. 2017;1655: 145–151.
274 doi:10.1016/j.brainres.2016.11.007
275
- 276 14. Filar-Mierzwa K, Marchewka A, Dąbrowski Z, Bac A, Marchewka J. Effects of
277 dance movement therapy on the rheological properties of blood in elderly
278 women. *Clin Hemorheol Microcirc*. 2019;72: 211–219. doi:10.3233/CH-180470
279




Figure 1




Figure 2