Novel antibiotic resistance genes from the hospital effluent 1

are disseminated into the marine environment in Norway 2

3	
4	Vera Radisic ^{1,2†} , Manish, P. Victor ^{1†} , Didrik H. Grevskott ¹ , Nachiket P. Marathe ^{1#}
5	
6	¹ Department of Contaminants and Biohazards, Institute of Marine Research (IMR), Bergen,
7	Norway
8	² Department of Biological Sciences, University of Bergen (UiB), Bergen, Norway
9	Vera Radisic and Manish, P. Victor contributed equally to this work.
10	
11	[#] Correspondence can be addressed to Nachiket P. Marathe, <u>nachiket.marathe@hi.no</u>
12	
13	
14	
15	
16	
17	
18	
19	
20 21 22 23 24 25 26	
27 28	
29	nois in preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

30 Abstract

31 Hospital effluent comprises feces of many individuals, including patients undergoing 32 antibiotic treatment. Although, hospital effluent is an important source for contamination of 33 the environment with antibiotic resistance genes (ARGs) and pathogens, how hospital 34 effluent in a low resistance setting contributes to antimicrobial resistance (AMR) in the 35 environment is largely understudied. The aim of our study was to understand the microbiota 36 and resistome of hospital effluent, and its role in the spread of AMR in the marine 37 environment in Norway. We further aimed at describing/characterizing novel resistance 38 factors from hospital effluent and the receiving sewage treatment plant (STP). 24-hour 39 composite samples of the hospital effluent and the influent and effluent of the receiving STP 40 were collected at two sampling time-points (February and April 2023) in Bergen city, 41 Norway. Isolation of Escherichia coli and Klebsiella spp. was performed, using ECC and SCAI plates with cefotaxime, tigecycline or meropenem, followed by antibiotic susceptibility 42 43 testing, using EUVSEC3 plates. Whole-genome sequencing of selected strains (n=36) and 44 shotgun metagenomics of sewage samples (n=6) were performed, using Illumina NovaSeq. 45 ARGs were identified with USEARCH, and known and novel ARGs were assembled with 46 fARGene. All E. coli strains (n=66) were multidrug-resistant (MDR), while 92.3% of the 47 Klebsiella spp. strains (n=55) showed MDR phenotype. The sequenced strains carried 48 multiple clinically important ARGs, including carbapenemases such as NDM-5 (n=3) and 49 KPC-3 (n=3). We obtained 238 Gigabases of sequence data from which we identified 676 50 unique ARGs with >200 ARGs shared across samples. We assembled 1,205 ARGs using 51 fARGene, 365 gene sequences represented novel ARGs (< 90% amino acid (aa) identity). 52 Both known and novel ARGs (n=54) were shared between the hospital effluent and the 53 treated effluent of the receiving STP. We show that hospital effluent in Norway has a high 54 diversity of both known and novel ARGs. Our study demonstrates that hospital effluent is a 55 source of clinically relevant pathogens, as well as known and novel ARGs, reaching the 56 marine environment in Norway through treated sewage. 57 58 59 60 61 62 63 64 65 66 67 Keywords: Norway; Hospital effluent; Antibiotic resistance genes; fARGene; Novel ARGs; 68 Last resort antibiotics 69 70

- 71
- 72
- 73

74

75 1. Introduction

76 Antimicrobial resistance (AMR) is an emerging global health problem (WHO, 2017). 77 Pathogens of particular concern to the public health include extended spectrum β-lactamases 78 (ESBL)- and carbapenemase-producing Enterobacterales (WHO, 2017). Carbapenems are 79 last resort antibiotics used for treatment of severe infections cause by multidrug-resistant 80 (MDR) pathogens (Sheu et al., 2019; Zhu et al., 2020). Resistance to carbapenems is largely 81 due to the presence of carbapenem hydrolyzing β -lactamases or carbapenemases (Sheu et al., 82 2019; Tamma & Simner, 2018). The five clinically most relevant carbapenemases are KPC, 83 IMP, VIM, NDM and OXA-48 variants (Bonnin et al., 2021). KPC belongs to Ambler class 84 A, while IMP, VIM and NDM belongs to class B (metallo- β -lactamases (MBLs)) and OXA-85 48 variants belong to class D of β-lactamases (van Duin & Doi, 2017). MBLs are of 86 particular concern as they hydrolyze most β-lactam antibiotics, and have successfully being 87 disseminated globally (Boyd et al., 2020; Logan & Bonomo, 2016). Although, the use of 88 carbapenems in clinics is low in Norway, the prevalence of carbapenemase-producing clinical 89 strains has increased over the past few years (NORM/NORM-VET, 2023).

90

91 Pathogens, antibiotic resistance genes (ARGs) and residues of antibiotics may be introduced

92 into the environment through different routes, such as pollution with untreated sewage from 93 the population, industries and/or hospitals (Stalder et al., 2012). The hospital effluent, in 94 particular, is an important source of clinically relevant pathogens, ARGs and residues of 95 antibiotics reaching the receiving environment (Karkman et al., 2019; Khan et al., 2021; 96 Marathe et al., 2019). Hospital effluent contains fecal material from many individuals 97 undergoing antibiotic treatment, and invariably contains antibiotic resistant bacteria, resistant 98 genes, as well as residues of different antimicrobial compounds (Deguenon et al., 2022; Orias 99 & Perrodin, 2013). Hospital effluents are treated on-site in some countries, but not in Norway 100 (VKM, 2020). How untreated hospital effluent contributes to the spread of AMR in the

101 receiving STP and eventually to the marine environment in Norway is largely unknown.

102

103 The aim our study was to understand the microbiota and resistome of hospital effluent and its 104 role in the spread of AMR in the marine environment in Norway. We further aimed at 105 describing/characterizing novel resistance genes from hospital effluent and the receiving 106 sewage treatment plant (STP).

107

108 2. Materials and Methods

109 2.1 Collection of sewage samples

Composite samples, representing a 24-hour period, from both raw sewage (influent) and 110 treated sewage (effluent) were collected on two occasions (February 7th and April 12th, 2023) 111 112 from the second largest sewage treatment plant (STP), Holen, in Bergen, Norway. This 113 treatment plant, serving 132,000 inhabitants, receives sewage from two hospitals: a 933-bed 114 and a 170-bed hospital. Moreover, composite sewage samples were also collected directly 115 from the sewer line from the Haukeland University hospital in Bergen, on the same two occasions. The sewage samples were transported to the laboratory in sterile containers at 4°C, 116117 and analyses were initiated within 6 h after collection.

118

119 2.2 Isolation and identification of pathogens

The samples were serially diluted ten-fold with sterile saline (0.85% NaCl) before plating on 120 121 ECC (CHROMagarTM, France) chromogenic media and on Simmons Citrate Agar with myo-122 Inositol (SCAI) (Sigma-Aldrich, Germany) media supplemented with three different 123 antibiotics: 2 mg/L cefotaxime (Sigma-Aldrich, Germany), 0.5 mg/L tigecycline (Glentham 124 Life Sciences, UK), or 0.125 mg/L meropenem (Sigma-Aldrich, Germany). The ECC plates were incubated at 37°C for 20-24 h, while the SCAI plates were incubated 37°C for 24-48 h. 125 126 E. coli CCUG 17620 and K. pneumoniae CCUG 225T were included as negative controls for 127 verifying the ECC and SCAI plates, respectively. From each sample, 10-15 isolated colonies 128 were picked depending on growth, restreaked and incubated at 37°C overnight. Identification 129 of strains was performed, using matrix-assisted laser desorption ionization-time of flight mass 130 spectrometry (MALDI-TOF MS) (Bruker Daltonics, Germany) at the Institute of Marine 131 Research (IMR, Norway). Subsequently, confirmed strains were stored at -80°C in Mueller-132 Hinton (MH) broth (Oxoid, UK) with respective antibiotics and 20% glycerol until further 133 use.

134

135 2.3 Antibiotic sensitivity testing

136 The resistance profile of 66 E. coli and 55 Klebsiella spp. strains were determined against 15 137 antibiotics, using a broth microdilution assay with Sensititre® EUVSEC3 plates (Thermo 138 Scientific, USA), following manufacturer's protocol. The plates were incubated at 37°C for 139 20-22 h. The strains were defined as 'susceptible' or 'resistant', according to the EUCAST 140 clinical breakpoints tables v.13.1 (EUCAST, 2023). For certain antibiotics, like azithromycin, 141 nalidixic acid, sulfamethoxazole and tetracyclines, not having defined breakpoints in 142 EUCAST, resistance was determined if growth was observed in the highest antibiotic 143 concentration tested. E. coli CCUG 17620 was included as negative controls, while E. coli 144 CCUG 73937 was included as positive controls (Grevskott et al., 2020).

145

146 2.4 DNA extraction and sequencing

147 Based on phenotypic resistance profiles, 19 presumptive E. coli and 17 presumptive K. 148 pneumoniae strains were selected for whole genome sequencing (WGS). These isolates were 149 grown overnight on MH agar (Oxoid, UK) containing 2 mg/L cefotaxime (Sigma-Aldrich, 150 Germany), 0.5 mg/L tigecycline (Glentham Life Sciences, UK), or 0.125 mg/L meropenem 151 (Sigma-Aldrich, Germany), based on the resistance profiles, at 37°C. Genomic DNA was 152 extracted from the strains, using the DNeasy Blood and Tissue kit (Qiagen, Germany) 153 following the manufacturer's protocol. The extracted DNA was quantified, using QubitTM 2.0 154 2000 Fluorometer with the dsDNA BR (Broad-Range) kit and NanoDrop[™] 155 Spectrophotometer (Thermo Scientific, USA) assay. Sequencing libraries was prepared, using 156 Nextera DNA Flex Library Prep kit (Illumina, USA). Sequencing was performed, using 157 Illumina NovaSeq platform (Illumina, USA), with 2×150 bp on a S4 flow cell, at Eurofins

- 158 Genomics A/S Germany.
- 159

160 2.5 Genome assembly and sequencing analysis

- 161 Obtained raw reads from the Illumina NovaSeq sequencing were quality trimmed and 162 assembled as previously described (Radisic et al., 2020), follow by annotation using National 163 Center for Biotechnology Information (NCBI) Prokaryotic Genomes Automatic Annotation 164 Pipeline (PGAAP) (Tatusova et al., 2016). Antibiotic resistance genes (ARGs) and sequence
- 165 types (STs) of *Klebsiella* spp. were screened, using Kleborate v.2.1.0 (Lam et al., 2021),
- 166 while the PubMLST database (https://pubmlst.org/bigsdb?db=pubmlst escherichia seqdef) 167 was used for identification of STs among E. coli strains. ResFinder v.4.1 (Bortolaia et al.,
- 168 2020) and the comprehensive antibiotic resistance database (CARD) v.3.1.4 (Alcock et al.,
- 169 2020) were used for further identification of ARGs among strains of *Klebsiella* spp. and *E*.
- 170 coli. PlasmidFinder v.2.1 was used for identification of plasmid replicons (Carattoli et al.,
- 171 2014).
- 172

173 2.6 Sampling, DNA extraction and shotgun sequencing

174 From each sample, 30-50 mL was vacuum-filtered through S-Pak filters (pore size 0.22 µm) 175 (Millipore, USA), after which the filters were transferred to 50 mL Falcon tubes and stored at 176 -20°C until further use. Genomic DNA was extracted from the filters, using DNeasy 177 PowerWater kit (Qiagen, Germany) following the manufacturer's protocol. The extracted 178 DNA was quantified, using dsDNA BR (Broad-Range) assay kit on the Qubit[™] 2.0 179 Fluorometer and NanoDrop[™] 2000 Spectrophotometer (Thermo Scientific, USA) assay. 180 Sequencing libraries for the shotgun metagenomics were prepared, using Nextera DNA Flex 181 Library Prep kit (Illumina, USA) and sequencing was performed, using Illumina NovaSeq 182 platform (Illumina, USA), with 2×150 bp chemistry on a S4 flow cell, at the Norwegian Sequencing Centre (Oslo University Hospital, Ullevål, Oslo, Norway). 183

184

185 2.7 Sequence analysis of shotgun metagenomic data

186 The sequences quality-trimmed with Trim Galore v.0.6.10 were 187 (http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/), using the default settings; 188 trim galore --quality 30 --phred33 --trim n --cores 50 --paired seqForward.fg seqReverse.fg 189 with a phred quality score of 33 and a maximum error rate of 0.5. The quality-processed 190 reads from the metagenomes were mapped against protein sequences from a high-quality and 191 manually curated database of ARGs, mobile genetic elements (MGEs), biocide/metal 192 resistance virulence USEARCH genes (BMRGs) and genes. using 193 v.usearch11.0.667 i86linux64 with the default settings; usearch -usearch global reads.fastg -194 db DB seq.fasta -id 0.9 -query cov 1.0 -blast6out hits.b6 -strand plus -maxaccepts 1. The 195 database for BMRGs were obtained from Pal et al. (2014), virulence factors (experimentally 196 verified) from Chen et al. (2016), MGEs from Pärnänen et al. (2018), while for the ARGs 197 database we used а combination of the **NCBI** AMR database 198 https://www.ncbi.nlm.nih.gov/pathogens/antimicrobial-resistance/, CARD (Alcock et al., 199 2020), and Antibiotic Resistance Gene-ANNOTation (ARG-ANNOT) (Gupta et al., 2014). 200 For quantitative taxonomic distribution, quality-filtered shotgun reads were used as input to 201 extract the reads corresponding to small subunit (SSU) 16S bacterial ribosomal RNA (rRNA) 202 genes from the metagenomes and assigned them to different taxonomic groups, using Metaxa2 v.2.2.3 (Bengtsson-Palme et al., 2015) with the default settings; metaxa2 203 -0 204 metaxa out -1 (sequence1) -2 (sequence2) -f fastq -g ssu --mode metagenome --usearch

205 v11.0.667 i86linux64 --usearch bin usearch11 --cpu 25 --table T --reltax T and metaxa2 ttt 206 -i (taxonomy text file) -o metaxa ttt out. fARGene was used to predict probable ARGs for 207 β -lactamases (class A, subclass B1-B3, class C and D1-2), quinolones, tetracyclines, 208 macrolides, and aminoglycosides. Known and novel ARGs were assembled from shotgun 209 sequence data, using fARGene as previously described by Berglund et al. (2019) with the 210 default settings; Fargene --infiles (sequence1) (sequence2) --hmm-model (model name) --211 meta --output (folder out) --force --processes (cpu) --rerun --amino-dir class a out tmpdir --212 orf-finder'. ARGs in different samples were clustered at 100/95/90% amino acid (aa) 213 sequence similarity together with previously characterized ARGs with USEARCH. The 214 query (obtained fARGenes) was blasted against nucleotide non-redundant (nr) database 215 (https://ftp.ncbi.nlm.nih.gov/blast/db/FASTA/) with query coverage of > 90% and E-value = 216 10⁻⁵, using diamond v.2.1.9 (https://github.com/bbuchfink/diamond). Novel ARGs were 217 identified, using cutoff < 90% aa identity (Inda-Díaz et al., 2023). The number of shared 218 ARGs, both known and novel, across samples were clustered at 100% identity for each 219 sample and presented Venn diagram. then in a using pyvenn 220 (https://github.com/tctianchi/pyvenn).

221

222 3. Results

223 3.1 Strains obtained from media with antibiotics

224 Most *Klebsiella* spp. isolates and *E. coli* isolates were obtained from the hospital effluent 225 samples (n=31; n=31), followed by STP influent samples (n=19; n=21) and STP treated 226 effluent samples (n=10; n=14). In total, 130 isolates were obtained on both ECC plates with 227 antibiotics (n=66) and SCAI plates with antibiotics (n=64) and identified with MALDI-TOF 228 MS. Most of the obtained isolates on the SCAI plates (n=64) were identified as K. 229 pneumoniae (n=34, 53.1%), followed by K. oxytoca (n=21, 32.8%), Raoultella 230 ornithinolytica (n=5, 7.8%), Acinetobacter bereziniae (n=2, 3.1%), A. baumannii (n=1, 1.6%) 231 and Citrobacter freundii (n=1, 1.6%). Isolates identified as other genera than K. pneumoniae, 232 K. oxytoca and E. coli were not included in further analysis.

233

234 3.2 Resistance rates in *E. coli* and *Klebsiella* spp. strains

235 We obtained a total of 64 E. coli isolates on cefotaxime-containing plates and two E. coli 236 isolates on meropenem-containing plates. All E. coli isolates (n=64) on cefotaxime-237 containing plates were multidrug-resistant (MDR), showing resistance against at least three 238 different classes of antibiotics (Supplementary Table S1). E. coli isolates obtained on 239 cefotaxime-containing plates from the hospital effluent samples (n=31) showed highest 240 resistance against ampicillin (100%), ceftazidime (100%), nalidixic acid (71%) and 241 ciprofloxacin (71%). Colistin resistance was detected in one E. coli isolate (3.2%). Further, 242 E. coli isolates (n=20) obtained on cefotaxime-containing plates from STP influent samples 243 showed highest resistance against ampicillin (100%), trimethoprim (81%) and 244 sulfamethoxazole (76%). Highest resistance against ampicillin (100%), nalidixic acid (79%), 245 ceftazidime (45%) and tetracycline (45%) was observed in E. coli isolates (n=13) on 246 cefotaxime-containing plates from STP effluent samples. E. coli isolates (n=2), obtained from 247 both STP influent (n=1) and STP effluent samples (n=1) on meropenem-containing plates, 248 were MDR (Supplementary Table S1). Several E. coli isolates from hospital effluent

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

(February: n=3; April: n=5) and STP influent (February: n=4) samples on cefotaximecontaining plates displaying reduced susceptibility against meropenem. No resistance was
observed against tigecycline in these isolates.

252

253 On the cefotaxime-containing plates we obtained a total of 31 Klebsiella spp. isolates (n=21 hospital effluent; n=10 STP influent; n=10 STP effluent). All Klebsiella spp. isolates on 254 255 cefotaxime-resistant were MDR (Supplementary Table S1). Klebsiella spp. isolates (n=21) 256 obtained on cefotaxime-containing plates from the hospital effluent samples showed highest 257 resistance against ampicillin (100%), cefotaxime (100%), nalidixic acid (100%) and 258 ciprofloxacin (100%). Resistance against colistin and tigecycline was observed in 9.5% and 259 57.1% of the isolates, respectively. Further, *Klebsiella* spp. isolates (n=10) obtained on 260 cefotaxime-containing plates from STP influent samples showed highest resistance against 261 ampicillin (100%), cefotaxime (100%), ceftazidime (100%) and trimethoprim (100%). 262 Highest resistance against ampicillin (100%), cefotaxime (100%), ceftazidime (100%), and 263 sulfamethoxazole (100%) was observed in *Klebsiella* spp. isolates (n=10) on the cefotaxime-264 containing plates from STP effluent samples. During the two-sampling time-points no 265 resistance against meropenem or amikacin was detected among these *Klebsiella* spp. isolates.

266

267 We further detected 11 K. pneumoniae isolates (n=7 hospital effluent; n=4 STP influent) on 268 tigecycline-containing plates and three K. oxytoca isolates (n=3 hospital effluent) on 269 meropenem-containing plates. Most tigecycline-resistant K. pneumoniae isolates (91.7%) 270 were MDR and showed resistance against third generation cephalosporins, such as 271 cefotaxime (63.6%) and ceftazidime (63.6%) (Supplementary Table S1). None of the 272 tigecycline-resistant K. pneumoniae isolates were resistant to meropenem, amikacin or 273 colistin. Three presumptive K. oxytoca isolates obtained on meropenem-containing plates 274 showed intermediate resistance for meropenem (Supplementary Table S1).

275

276 3.3 Genome sequence and STs of sequenced strains of *E. coli* and *Klebsiella* spp.

- 277 A complete overview of the genome sequence statistics is presented in Supplementary Table 278 S2. The total length ranged from 5,857,362 bp - 6,384,740 bp and 4,590,194 bp - 5,361,939 279 bp for *Klebsiella* spp. strains and *E. coli* strains, respectively. Six *Klebsiella* spp. strains 280 which were initially identified as K. oxytoca by MALDI TOF MS were after sequencing re-281 identified as either K. michiganensis (n=3) (K11-303, K11-314 and K11-325) or K. 282 pneumoniae (n=3) (K10-M302, K10-M304 and K10-M306) with Kleborate v.2.1.0. 283 Additionally, four K. pneumoniae strains (K10-T302, K10-T303, K10-T306 and K10-321) 284 were re-identified as K. michiganensis. A high ST diversity was observed among the 285 sequenced E. coli strains, including pathogenic ST131 and ST38. The most common ST 286 among the sequenced K. pneumoniae strains was ST307 (n=6), followed by ST2947 (n=4) 287 (Supplementary Table S2).
- 288

289 3.4 Clinically important ARGs detected in E. coli and Klebsiella spp. strains

Acquired ARGs conferring resistance against tetracyclines, aminoglycosides, sulfonamides and quinolones were detected in sequenced *Klebsiella* spp. and *E. coli* strains (Supplementary

292 Table S3). In addition, different β -lactamase genes with bla_{OXA-1} (52.65%) as most dominant,

293 as well as carbapenemases, such as NDM-5 (n=3) and KPC-3 (n=3), were detected among 294 the sequenced strains. Four K. pneumoniae strains (K11-304, K11-305, K11-327 and K11-295 329) with MIC of $\geq 2\mu g/mL$ for colistin carried mutations in the mgrB gene, contributing to a

- 296 truncated protein of only 24 aa instead of 47 aa long protein.
- 297

298 3.5 Shotgun sequencing

- 299 In total, 238 Gigabases of sequence data was generated from the shotgun metagenomic
- 300 sequencing of the hospital effluent samples and STP influent and STP effluent samples, 301 representing 2.1×10^{11} quality filtered reads from six samples. In the hospital effluent samples, 302 109.987 reads (0.000053% of the total reads from hospital effluent samples) matched to 303 ARGs, while 39,737 reads (0.000019% of the total reads from STP influent samples) and 304 45,069 reads (0.000022% of the total reads from STP effluent samples) matched to ARGs in 305 the STP influent and STP effluent samples, respectively. All together, these accounted for 306 between 0.064 (STP effluent from February) to 0.376 (hospital effluent from April) copies of 307 ARGs per 16S rRNA gene. We detected 676 unique ARGs in hospital effluent samples, as
- 308 well as 517 unique ARGs in STP influent and 567 unique ARGs in STP effluent samples,
- 309 respectively (Supplementary Table S4). These ARGs conferred resistance to multiple classes
- 310 of antibiotics, including glycylcyclines, polymyxins, glycopeptides and oxazolidinones.
- 311

312 **3.6 Bacterial diversity**

313 A total of 28 bacterial phyla were detected across all samples, with *Pseudomonadota* as most

- 314 dominant followed by Bacillota and Bacteroidota for most of the samples (Figure 1). Within
- 315 Pseudomonadota, β -proteobacteria was most dominant class followed by γ -proteobacteria
- 316 for most samples. At genus level, *Acinetobacter* was found to be the dominant in the majority
- 317 of the samples. For STP influent samples collected in February, Bacillota dominated
- 318 followed by *Pseudomonadota* and *Bacteroidota*. Within *Pseudomonadota*, β -proteobacteria
- 319 was most dominant class followed by *y-protebacteria*. At genus level, *Lachnospiraceae* was
- 320 found to be most dominant. For STP effluent samples collected in February, Within
- 321 Pseudomonadota dominated followed Bacteroidota and Bacillota. by
- 322 *Pseudomonadota*, β -proteobacteria was most dominant class followed by γ -protebacteria.

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .



323

Figure 1: Relative abundance of bacterial diversity detected in sewage from two sampling timepoints
 February (Feb) and April (Apr), including hospital effluent (eff), STP influent (inf) and STP treated
 effluent (eff) samples. Only *Pseudomonadota* are represented at class level.

327 3.7 Abundance of ARGs, MGEs, BMRGs and virulence genes

328 In total, 676 unique ARGs were detected across samples. The macrolide resistance genes 329 msr(E) and mph(E) were the most abundant ARGs in hospital effluent collected in February 330 and April, and in STP influent and STP effluent samples collected in April. For STP influent 331 and STP effluent collected in February, the tetracycline resistance genes tet(Q) and tet(W)332 were the most abundant ARGs (Figure 2). For majority of the samples, we detected genes 333 conferring resistance to macrolides as most abundant, followed by genes conferring 334 resistance to β -lactams and tetracyclines. Genes conferring resistance against last resort 335 antibiotics, such as tigecycline (tet(X)), colistin (mcr), vancomycin (van type genes), and 336 linezolid (cfr(C)) were detected across all samples (Supplementary Table S4).

337

Overall, there was a higher abundance of integron-associated integrases (*intl*) and ISCR
transposases in hospital effluent samples compared to STP influent and STP effluent samples.
Among the studied MGEs, class 1 integrase was the most abundant followed by class 2
integrase and ISCR2 for most of the samples (Supplementary Table S5).

342

343 Several BMRGs conferring resistance to a range of compounds, such as arsenic, cobalt,344 copper, lead, mercury, nickel, silver, and zinc compounds were detected across samples and

345 found in higher abundance in hospital effluent compared to STP influent and STP effluent.

346 For all samples, the relative abundance of genes conferring resistance to mercury (*mer* genes)

347 was higher compared to other BMRGs. A complete overview of the BMRGs is presented in

348 Supplementary Table S6.

349

Virulence genes involved in adhesion factors, capsule formation, pilus formation, siderophore production, and toxin productions were also detected in all samples. Important virulence genes, such as aerobactin, enterobactin, yersiniabactin, salmochelin, cytotoxic necrotizing factor 1, colibactin, and hemolysin, were detected across all samples (Supplementary Table S7). The important virulence factor, yersiniabactin, was also detected in five (K11-302, K11-304, K11-305, K11-327 and K11-329) out of 17 sequenced K. pneumoniae strains.





3.8 Shared ARGs across samples

More than 200 ARGs were shared among the samples, including genes conferring resistance against last resort antibiotics, such as colistin, carbapenems, tigecycline, linezolid and vancomycin. A total of 1,205 predicted gene sequences were assembled, using fARGene. Several ARGs conferring resistance to aminoglycosides, macrolides, tetracyclines and quinolones were shared across samples (Supplementary Table S8). Among all assembled ARGs, 365 represented novel ARGs using cutoff < 90% aa identity (Table 1). The majority of the predicted novel ARGs (n=244) were β -lactamases, followed by aminoglycoside resistance genes (n=98) (Table 1). Among novel β -lactamases, we detected 32 novel MBLs across all samples, where most of the novel MBLs were detected in the STP treated effluent samples.

Table 1: Number of antibiotic resistance genes (ARGs) assembled using fARGene and the antibiotic classes they confer resistant against.

Antibiotic class	Aminoglycosides			β-lactams			Macrolides			Quinolones			Tetracyclines		
Sample	UA	KA	NA	UA	KA	NA	UA	KA	NA	UA	KA	NA	UA	KA	NA
Hosp-eff-F	55	35	20	136	105	31	23	20	3	6	5	1	17	17	0
Hosp-eff-A	39	12	27	92	78	14	16	15	1	5	4	1	12	11	1
STP-inf-F	34	24	10	111	79	32	23	21	2	5	4	1	17	17	0
STP-inf-A	38	24	14	94	65	29	14	13	1	2	1	1	12	12	0
STP-T-eff-F	25	19	6	120	65	56	11	11	0	2	1	1	11	11	0
STP-T-eff-A	49	28	21	187	104	83	23	18	5	3	2	1	22	19	3

Legend: UA (unique ARGs); KA (known ARGs); NA (novel ARGs); Hosp (hospital); eff (effluent);

F (February); A (April); STP (sewage treatment plant); inf (influent); T (treated). Novel genes <90% aa identity.

In total, 40 unique assembled ARGs were shared across all samples (Figure 3), representing both novel (n=1) (belonging to class A β -lactamase) and known ARGs (n=39). A total of 54 unique ARGs were shared between hospital effluent and STP effluent, including three novel ARGs (two β-lactamases and one ribosomal RNA methyltransferase Erm).

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .



406

407 Figure 3: Venn diagram showing the shared (46, 40, 14 and 15) and exclusive (190, 274 and

408 259) antibiotic resistance genes (ARGs), including both known and novel, across the three

409 samples (hospital effluent, sewage treatment plant (STP) influent and STP treated effluent).

410

411 4. Discussion

412 To the best of our knowledge this is the first comprehensive study, investigating the 413 microbiota and resistome of hospital effluent and receiving STP in Norway. Although 414 resistance is low in Norway, we show high diversity of ARGs in hospital effluent and the 415 receiving STP, including genes for resistance against last resort antibiotics. We further show 416 that both novel genes and pathogenic strains are shared between the hospital effluent and STP 417 effluent (Grevskott et al., 2024). Thus, suggesting that hospital effluent is contributing to the

- 418 spread of AMR even in a low prevalence setting like Norway.
- 419

420 We detected resistance against meropenem (with a MIC of > $16\mu g/mL$) in three out of 66 421 MDR *E. coli* isolates (4.5%) from STP influent samples (n=2) and STP effluent (n=1) 422 samples. All sequenced meropenem-resistant *E. coli* strains (11-330, 11-M301 and 11-M302) 423 carried NDM-5 and belonged to ST11681 (Supplementary Table S3). The *bla*_{NDM-5} gene was

- 424 located on a highly conserved region (IS 30Δ -bla_{NDM}-ble-trpT-IS26), which has previously
- 425 been observed in NDM-carrying *E. coli* strains in Norway and in Europe (Chakraborty et al.,
- 426 2021; Grevskott et al., 2023; Turton et al., 2022). In the clinics in Norway, the occurrence of
- 427 NDM-5-producing *E. coli* strains has increased during the last decade (NORM/NORM-VET,
- 428 2023). The spread of NDM-5 carrying *E. coli* strains in the environment through sewage may
- 429 have contributed to this increased detection in clinics in Norway.
- 430

431 The bla_{KPC} genes, initially described in *K. pneumoniae*, represent globally the most 432 disseminated carbapenemase genes in pathogens like *Enterobacterales* (Hansen, 2021; 433 Munoz-Price et al., 2013), *Acinetobacter* spp. and *Pseudomonas* spp. (Robledo et al., 2011).

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

434 Among the KPC variants, KPC-2 and KPC-3 are responsible for most epidemic outbreaks 435 worldwide (Chen et al., 2011). We detected three bla_{KPC-3} carrying K. oxytoca isolates, (K10-436 M302, K10-M304 and K10-M306) which were later reidentified as K. pneumoniae, using 437 Kleborate. These stains belonged to ST307 which represents a globally disseminated high 438 risk clone (Fostervold et al., 2022; NORM/NORM-VET, 2023; Peirano et al., 2020). This ST 439 is also the most common among K. pneumoniae strains in clinics in Norway and is often 440 associated with carbapenemases and ESBLs (NORM/NORM-VET, 2023; Peirano et al., 2020). KPC-3-producing K. pneumoniae belonging to ST307 have also been reported from 441 442 clinics in Norway (NORM/NORM-VET, 2023).

443

444 We assembled 365 novel ARGs across samples, representing 32 novel MBLs. The MBLs are 445 enzymes that confer resistance against nearly all β-lactams, including carbapenems, thus 446 limiting available treatment options (Yang et al., 2023). MBLs, such as IMP, VIM, and NDM 447 have been successfully spread among pathogens (Bakthavatchalam et al., 2016; Cornaglia et 448 al., 2011). MBLs can be divided into subclass B1 and B3 (two zinc ions) or subclass B2 (one 449 zinc ion) (Berglund et al., 2017; Palzkill, 2013). Most of the novel MBLs in our study 450 belonged to subclass B1, which also include NDM, VIM and IMP (Palzkill, 2013). These 451 were detected in the STP effluent, suggesting that these ARGs are reaching the receiving 452 environment. Presence of novel MBLs along with clinical strains may provide an opportunity 453 for pathogens to acquire these genes and this may eventually become a problem in clinics 454 (Berglund et al., 2017).

455

456 Hospital effluent has been shown to contribute to dissemination of carbapenem-resistant 457 pathogens and genes in the environment (Lamba et al., 2017). In a previous longitudinal 458 temporal study, we showed the presence of carbapenem-resistant E. coli strains only in the 459 STP

460 (Holen) receiving hospital sewage on different time-points (Grevskott et al., 2021; Grevskott 461 et al., 2024; Marathe et al., 2023). Further, we showed that OXA-244 carbapenemase-462 producing E. coli ST38 from raw and treated sewage from STP receiving hospital effluent 463 were identical to OXA-244 carbapenemase-producing E. coli ST38 from clinics in Norway 464 (Grevskott et al., 2024). Although, the prevalence of resistance in clinics in Norway is low, 465 we detected a high diversity of ARGs in our study. A total of 54 unique ARGs, including both known (n=51) and novel (n=3), were present in the hospital effluent and STP effluent, 466 467 suggesting that the genes from the hospital effluent are introduced in the environment 468 through treated sewage. Put together, our studies strongly suggest that the hospital effluent 469 contributes to the spread of epidemic-causing pathogens, carbapenemaseses as well as novel 470 ARGs into the receiving marine environment in Norway (Grevskott et al., 2024; Radisic et 471 al., 2023).

472

473 **5.** Conclusions

474 Our study shows that hospital effluent is a rich source of both known and novel ARGs,

- 475 although Norway has low prevalence of AMR and low usage of antibiotics. We detected E.
- 476 coli and Klebsiella spp. strains resistant to last resort antibiotics in hospital effluent as well as
- 477 in treated

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

- 478 effluent of the receiving plant. We further show that clinical strains carrying carbapenemases,
- 479 such as NDM-5 and KPC-3, are reaching the marine environment through treated sewage, in
- 480 Norway. We further show that the hospital effluent contributes to the spread of both known
- 481 and novel ARGs into the receiving environment in Norway. Our study suggests that
- 482 mitigation strategies, such as on-site treatment of hospital effluent or bacterial removal from
- 483 sewage, are needed in order to limit the ongoing dissemination of AMR into the receiving
- 484 environment in Norway.
- 485

486 **Author contributions**

487 D.H.G.: Methodology, Investigation, Formal analysis, Visualization, Writing – original draft, 488 Writing - review & editing. V.R.: Methodology, Investigation, Formal analysis, 489 Visualization, Writing - original draft, Writing - review & editing. M.P.V.: Methodology, 490 Investigation, Formal analysis, Data curation, Writing - original draft. N.P.M.: 491 Conceptualization, Methodology, Investigation, Validation, Writing - review & editing, 492 Supervision, Project administration, Funding acquisition.

493

494 Funding

- 495 The Research Council of Norway under the ResMarine project (grant number 315266) 496 funded this work.
- 497

498 Acknowledgements

499 The authors would like to thank Kristine S. Akervold (Bergen municipality), Ingvild Fenne

500 (Bergen municipality) for permission and support as well as Ole Gunnar (Bergen Vann) and

- 501 the people working at the STP for sewage sample collection.
- 502

503 **Conflicts of Interest**

504 None.

505

506 References

- 507 Alcock, B. P., Raphenya, A. R., Lau, T. T., Tsang, K. K., Bouchard, M., Edalatmand, A., Huynh, 508 W., Nguyen, A.-L. V., Cheng, A. A., Liu, S., Min, S. Y., Miroshnichenko, A., Tran, H.-509 K., Werfalli, R. E., Nasir, J. A., Oloni, M., Speicher, D. J., Florescu, A., Singh, B.,... 510 McArthur, A. G. (2020). CARD 2020: antibiotic resistome surveillance with the
- 511 comprehensive antibiotic resistance database. Nucleic acids research, 48(D1), D517-D525. 512 https://doi.org/10.1093/nar/gkz935
- 513 Bakthavatchalam, Y. D., Anandan, S., & Veeraraghavan, B. (2016). Laboratory detection and clinical 514 implication of oxacillinase-48 like carbapenemase: the hidden threat. Journal of global 515 infectious diseases, 8(1), 41. https://doi.org/10.4103/0974-777X.176149
- 516 Bengtsson-Palme, J., Hartmann, M., Eriksson, K. M., Pal, C., Thorell, K., Larsson, D. G. J., & 517 Nilsson, R. H. (2015). METAXA2: improved identification and taxonomic classification of 518 small and large subunit rRNA in metagenomic data. Molecular ecology resources, 15(6), 519 1403-1414. https://doi.org/10.1111/1755-0998.12399
- 520 Berglund, F., Marathe, N. P., Österlund, T., Bengtsson-Palme, J., Kotsakis, S., Flach, C.-F., Larsson, 521 D. J., & Kristiansson, E. (2017). Identification of 76 novel B1 metallo-β-lactamases through 522 large-scale screening of genomic and metagenomic data. Microbiome, 5, 1-13. 523 https://doi.org/10.1186/s40168-017-0353-8

- Berglund, F., Österlund, T., Boulund, F., Marathe, N. P., Larsson, D. J., & Kristiansson, E. (2019).
 Identification and reconstruction of novel antibiotic resistance genes from metagenomes. *Microbiome*, 7, 1-14. https://doi.org/10.1186/s40168-019-0670-1
- Bonnin, R. A., Jousset, A. B., Emeraud, C., Oueslati, S., Dortet, L., & Naas, T. (2021). Genetic diversity, biochemical properties, and detection methods of minor carbapenemases in *Enterobacterales.* Frontiers in Medicine, 7, 616490.
 https://doi.org/10.3389/fmed.2020.616490
- Bortolaia, V., Kaas, R. S., Ruppe, E., Roberts, M. C., Schwarz, S., Cattoir, V., Philippon, A., Allesoe,
 R. L., Rebelo, A. R., Florensa, A. F., Fagelhauer, L., Chakraborty, T., Neumann, B., Werner,
 G., Bender, J. K., Stingl, K., Nguyen, M., Coppens, J., Xavier, B. B.,... Aarestrup, F. M.
 (2020). ResFinder 4.0 for predictions of phenotypes from genotypes. *Journal of Antimicrobial Chemotherapy*, 75(12), 3491-3500. https://doi.org/10.1093/jac/dkaa345
- Boyd, S. E., Livermore, D. M., Hooper, D. C., & Hope, W. W. (2020). Metallo-β-lactamases:
 structure, function, epidemiology, treatment options, and the development pipeline. *Antimicrobial Agents and Chemotherapy*, 64(10), 00397-00320.
 https://doi.org/10.1128/AAC.00397-20
- 540 Carattoli, A., Zankari, E., García-Fernández, A., Voldby Larsen, M., Lund, O., Villa, L., Møller
 541 Aarestrup, F., & Hasman, H. (2014). In silico detection and typing of plasmids using
 542 PlasmidFinder and plasmid multilocus sequence typing. *Antimicrobial Agents and*543 *Chemotherapy*, 58(7), 3895-3903. https://doi.org/10.1128/aac.02412-14
- 544 Chakraborty, T., Sadek, M., Yao, Y., Imirzalioglu, C., Stephan, R., Poirel, L., & Nordmann, P. (2021).
 545 Cross-border emergence of *Escherichia coli* producing the carbapenemase NDM-5 in
 546 Switzerland and Germany. *Journal of Clinical Microbiology*, *59*(3), 02238-02220.
 547 https://doi.org/10.1128/jcm.02238-20
- 548 Chen, L., Mediavilla, J. R., Endimiani, A., Rosenthal, M. E., Zhao, Y., Bonomo, R. A., & Kreiswirth,
 549 B. N. (2011). Multiplex real-time PCR assay for detection and classification of *Klebsiella*550 *pneumoniae* carbapenemase gene (*bla*_{KPC}) variants. *Journal of Clinical Microbiology*, 49(2),
 551 579-585. https://doi.org/10.1128/jcm.01588-10
- 552 Chen, L., Zheng, D., Liu, B., Yang, J., & Jin, Q. (2016). VFDB 2016: hierarchical and refined dataset
 553 for big data analysis 10 years on. *Nucleic acids research*, 44(D1), D694-D697.
 554 https://doi.org/10.1093/nar/gkv1239
- 555 Cornaglia, G., Giamarellou, H., & Rossolini, G. M. (2011). Metallo-β-lactamases: a last frontier for β-lactams? *The Lancet infectious diseases*, 11(5), 381-393. https://doi.org/10.1016/S1473-3099(11)70056-1
- Deguenon, E., Dougnon, V., Houssou, V. M. C., Gbotche, E., Ahoyo, R. A., Fabiyi, K., Agbankpe, J.,
 Mousse, W., Lougbegnon, C., & Klotoe, J. R. (2022). Hospital effluents as sources of
 antibiotics residues, resistant bacteria and heavy metals in Benin. *SN Applied Sciences*, 4(8),
 206. https://doi.org/10.1007/s42452-022-05095-9
- 562 EUCAST. (2023). Breakpoint tables for interpretation of MICs and zone diameters Version 13.1.
 563 European Committee on Antimicrobial Suceptibility Testing (EUCAST).
 564 https://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/
 565 v_13.1_Breakpoint_Tables.pdf
- Fostervold, A., Hetland, M. A., Bakksjø, R., Bernhoff, E., Holt, K. E., Samuelsen, Ø., Simonsen, G.
 S., Sundsfjord, A., Wyres, K. L., & Löhr, I. H. (2022). A nationwide genomic study of clinical *Klebsiella pneumoniae* in Norway 2001–15: introduction and spread of ESBLs facilitated by clonal groups CG15 and CG307. *Journal of Antimicrobial Chemotherapy*, 77(3), 665-674. https://doi.org/10.1093/jac/dkab463
- 571 Grevskott, D. H., Ghavidel, F. Z., Svanevik, C. S., & Marathe, N. P. (2021). Resistance profiles and diversity of β-lactamases in *Escherichia coli* strains isolated from city-scale sewage
 573 surveillance in Bergen, Norway mimic clinical prevalence. *Ecotoxicology and environmental* safety, 226, 112788. https://doi.org/10.1016/j.ecoenv.2021.112788
- 575 Grevskott, D. H., Radisic, V., Salvà-Serra, F., Moore, E. R., Akervold, K. S., Victor, M. P., & Marathe,
 576 N. P. (2024). Emergence and dissemination of epidemic-causing OXA-244 carbapenemase577 producing *Escherichia coli* ST38 through hospital sewage in Norway, 2020-2022. *Journal of*578 *Hospital infection*. https://doi.org/10.1016/j.jhin.2023.12.020

- 579 Grevskott, D. H., Salvà-Serra, F., Moore, E. R., & Marathe, N. P. (2020). Nanopore sequencing
 580 reveals genomic map of CTX-M-type extended-spectrum β-lactamases carried by
 581 *Escherichia coli* strains isolated from blue mussels (*Mytilus edulis*) in Norway. *BMC*582 *Microbiology*, 20(1), 1-10. https://doi.org/10.1186/s12866-020-01821-8
- Grevskott, D. H., Salvà-Serra, F., Moore, E. R., & Marathe, N. P. (2023). Complete sequence of a new conjugative multidrug-resistance encoding IncFII/IncFIA/IncFIB plasmid carrying NDM-6 metallo-β-lactamase from pathogenic *Escherichia coli* sequence type 167 isolated from sewage in Norway. *Journal of Global Antimicrobial Resistance, 33*, 291-293. https://doi.org/10.1016/j.jgar.2023.04.005
- 588 Gupta, S. K., Padmanabhan, B. R., Diene, S. M., Lopez-Rojas, R., Kempf, M., Landraud, L., &
 589 Rolain, J.-M. (2014). ARG-ANNOT, a new bioinformatic tool to discover antibiotic
 590 resistance genes in bacterial genomes. *Antimicrobial Agents and Chemotherapy*, 58(1), 212591 220. https://doi.org/10.1128/aac.01310-13
- Hansen, G. T. (2021). Continuous evolution: perspective on the epidemiology of carbapenemase
 resistance among *Enterobacterales* and other Gram-negative bacteria. *Infectious diseases and therapy*, 10, 75-92. https://doi.org/10.1007/s40121-020-00395-2
- Inda-Díaz, J. S., Lund, D., Parras-Moltó, M., Johnning, A., Bengtsson-Palme, J., & Kristiansson, E. (2023). Latent antibiotic resistance genes are abundant, diverse, and mobile in human, animal, and environmental microbiomes. *Microbiome*, 11(1), 1-16. https://doi.org/10.1186/s40168-023-01479-0
- Karkman, A., Pärnänen, K., & Larsson, D. J. (2019). Fecal pollution can explain antibiotic resistance
 gene abundances in anthropogenically impacted environments. *Nature communications*, 10(1), 80. https://doi.org/10.1038/s41467-018-07992-3
- Khan, M. T., Shah, I. A., Ihsanullah, I., Naushad, M., Ali, S., Shah, S. H. A., & Mohammad, A. W.
 (2021). Hospital wastewater as a source of environmental contamination: An overview of management practices, environmental risks, and treatment processes. *Journal of Water Process Engineering*, 41, 101990. https://doi.org/10.1016/j.jwpe.2021.101990
- Lam, M. M., Wick, R. R., Watts, S. C., Cerdeira, L. T., Wyres, K. L., & Holt, K. E. (2021). A genomic surveillance framework and genotyping tool for *Klebsiella pneumoniae* and its related species complex. *Nature communications, 12*(1), 4188. https://doi.org/10.1038/s41467-021-24448-3
- Lamba, M., Graham, D. W., & Ahammad, S. (2017). Hospital wastewater releases of carbapenemresistance pathogens and genes in urban India. *Environmental Science & Technology*, 51(23), 13906-13912. https://doi.org/10.1021/acs.est.7b03380
- 612 Logan, L. K., & Bonomo, R. A. (2016). Metallo-β-lactamase (MBL)-producing *Enterobacteriaceae* in
 613 United States children. *Open Forum Infectious Diseases*, 3(2), ofw090.
 614 https://doi.org/10.1093/ofid/ofw090
- Marathe, N. P., Berglund, F., Razavi, M., Pal, C., Dröge, J., Samant, S., Kristiansson, E., & Larsson,
 D. J. (2019). Sewage effluent from an Indian hospital harbors novel carbapenemases and
 integron-borne antibiotic resistance genes. *Microbiome*, 7, 1-11.
 https://doi.org/10.1186/s40168-019-0710-x
- 619 Marathe, N. P., Radisic, V., Salvà-Serra, F., Moore, E. R., & Grevskott, D. H. (2023). Emergence of 620 new IncHI2 multidrug-resistance plasmids carrying VIM-1 metallo-β-lactamase in 621 Escherichia coli in Norway. The Lancet Microbe, 4(9), E663-E664. 622 https://doi.org/10.1016/S2666-5247(22)00389-5
- Munoz-Price, L. S., Poirel, L., Bonomo, R. A., Schwaber, M. J., Daikos, G. L., Cormican, M.,
 Cornaglia, G., Garau, J., Gniadkowski, M., & Hayden, M. K. (2013). Clinical epidemiology
 of the global expansion of *Klebsiella pneumoniae* carbapenemases. *The Lancet infectious diseases, 13*(9), 785-796. https://doi.org/10.1016/S1473-3099(13)70190-7
- NORM/NORM-VET. (2023). Usage of Antimicrobial Agents and Occurrence of Antimicrobial Resistance in Norway 2022. Norwegian monitoring programme on antimicrobial resistance in bacteria from humans, food, feed and animals (NORM/NORM-VET).
 https://www.vetinst.no/en/surveillance-programmes/norm-norm-vet-report
- 631 Orias, F., & Perrodin, Y. (2013). Characterisation of the ecotoxicity of hospital effluents: a review.
 632 Science of the Total Environment, 454, 250-276.
 633 https://doi.org/10.1016/j.scitotenv.2013.02.064

- Pal, C., Bengtsson-Palme, J., Rensing, C., Kristiansson, E., & Larsson, D. J. (2014). BacMet: antibacterial biocide and metal resistance genes database. *Nucleic acids research*, 42(D1), D737-D743. https://doi.org/10.1093/nar/gkt1252
- 637 Palzkill, T. (2013). Metallo-β-lactamase structure and function. *Annals of the New York Academy of* 638 *Sciences*, 1277(1), 91-104. https://doi.org/10.1111/j.1749-6632.2012.06796.x
- Pärnänen, K., Karkman, A., Hultman, J., Lyra, C., Bengtsson-Palme, J., Larsson, D. J., Rautava, S.,
 Isolauri, E., Salminen, S., & Kumar, H. (2018). Maternal gut and breast milk microbiota affect infant gut antibiotic resistome and mobile genetic elements. *Nature communications*, 9(1), 3891. https://doi.org/10.1038/s41467-018-06393-w
- Peirano, G., Chen, L., Kreiswirth, B. N., & Pitout, J. D. (2020). Emerging antimicrobial-resistant
 high-risk *Klebsiella pneumoniae* clones ST307 and ST147. *Antimicrobial Agents and Chemotherapy*, 64(10), 01148-01120. https://doi.org/10.1128/aac.01148-20
- Radisic, V., Grevskott, D. H., Lunestad, B. T., Øvreås, L., & Marathe, N. P. (2023). Sewage-based
 surveillance shows presence of *Klebsiella pneumoniae* resistant against last resort antibiotics
 in the population in Bergen, Norway. *International Journal of Hygiene and Environmental Health, 248*, 114075. https://doi.org/10.1016/j.ijheh.2022.114075
- Radisic, V., Nimje, P. S., Bienfait, A. M., & Marathe, N. P. (2020). Marine plastics from Norwegian
 west coast carry potentially virulent fish pathogens and opportunistic human pathogens
 harboring new variants of antibiotic resistance genes. *Microorganisms, 8*(8), 1200.
 https://doi.org/10.3390/microorganisms8081200
- Robledo, I. E., Aquino, E. E., & Vázquez, G. J. (2011). Detection of the KPC gene in *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, and *Acinetobacter baumannii* during
 a PCR-based nosocomial surveillance study in Puerto Rico. *Antimicrobial Agents and Chemotherapy*, 55(6), 2968-2970. https://doi.org/10.1128/AAC.01633-10
- 658
 658
 659
 659
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
 660
- Stalder, T., Barraud, O., Casellas, M., Dagot, C., & Ploy, M.-C. (2012). Integron involvement in environmental spread of antibiotic resistance. *Frontiers in Microbiology*, *3*, 119. https://doi.org/10.3389/fmicb.2012.00119
- Tamma, P. D., & Simner, P. J. (2018). Phenotypic detection of carbapenemase-producing organisms
 from clinical isolates. *Journal of Clinical Microbiology*, 56(11), 01140-01118.
 https://doi.org/10.1128/jcm.01140-18
- Tatusova, T., DiCuccio, M., Badretdin, A., Chetvernin, V., Nawrocki, E. P., Zaslavsky, L., Lomsadze,
 A., Pruitt, K. D., Borodovsky, M., & Ostell, J. (2016). NCBI prokaryotic genome annotation
 pipeline. *Nucleic acids research*, 44(14), 6614-6624. https://doi.org/10.1093/nar/gkw569
- Turton, J. F., Pike, R., Perry, C., Jenkins, C., Turton, J. A., Meunier, D., & Hopkins, K. L. (2022).
 Wide distribution of *Escherichia coli* carrying IncF plasmids containing bla NDM-5 and *rmtB* resistance genes from hospitalized patients in England. *Journal of medical microbiology*, 71(8), 001569. https://doi.org/10.1099/jmm.0.001569
- 674 van Duin, D., & Doi, Y. (2017). The global epidemiology of carbapenemase-producing
 675 *Enterobacteriaceae*. *Virulence*, 8(4), 460-469.
 676 https://doi.org/10.1080/21505594.2016.1222343
- 677 VKM. (2020). Assessment of the impact of wastewater and sewage sludge treatment methods on antimicrobial resistance. Scientific opinion of the Panel on Microbial Ecology of the Norwegian Scientific Committee for Food and Environment. (VKM 2020: 08). Norwegian Scientific Committee for Food and Environment. (VKM 2020: 08). Norwegian Scientific Committee for Food and Environment (VKM) https://nmbu.brage.unit.no/nmbuxmlui/bitstream/handle/11250/2829018/
- Assessment+of+the+impact+of+wastewater+and+sewage+sludge+treatment+methods+on+an
 timicrobial+resistance+-01.10.2020.pdf?sequence=2
- WHO. (2017, February 27). WHO publishes list of bacteria for which new antibiotics are urgently
 needed. World Health Organization (WHO). https://www.who.int/news/item/27-02-2017 who-publishes-list-of-bacteria-for-which-new-antibiotics-are-urgently-needed

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

- Yang, Y., Yan, Y.-H., Schofield, C. J., McNally, A., Zong, Z., & Li, G.-B. (2023). Metallo-β-lactamase-mediated antimicrobial resistance and progress in inhibitor discovery. *Trends in microbiology*, *31*(7), 735-748. https://doi.org/10.1016/j.tim.2023.01.013
- Zhu, J., Li, Q., Li, X., Kang, J., Song, Y., Song, J., Yin, D., & Duan, J. (2020). Successful control of
 the first carbapenem-resistant *Klebsiella pneumoniae* outbreak in a Chinese hospital 2017–
 2019. Antimicrobial Resistance & Infection Control, 9(1), 91. https://doi.org/10.1186/s13756020-00757-y
- 694
- 695
- 696

697 Supplementary Materials

698 Supplementary Table S1: Minimum inhibitory concentrations (MICs) of *Escherichia coli*699 (n= 66) and *Klebsiella* spp. (n=55) strains in this study.

- **Supplementary Table S2:** Genome assembly statistics of sequenced strains (n=36) in thisstudy.
- 702 Supplementary Table S3: Detected antibiotic resistance genes (ARGs) and sequence types
- 703 (STs) in sequenced strains (n=36).
- 704 Supplementary Table S4: Detected antibiotic resistance genes (ARGs) across six samples.
- 705 Supplementary Table S5: Relative abundance of most detected mobile genetic elements
- 706 (MGEs) across samples.
- 707 Supplementary Table S6: Detected biocide/metal resistance genes (BMRGs) across all708 samples.
- 709 Supplementary Table S7: Detected virulence genes across all samples.
- 710 Supplementary Table S8: Assembled known and novel antibiotic resistance genes (ARGs)
- 711 across all samples.
- 712