

1 Characterizing Highly Fermentable Carbohydrate Foods in the Diets of Children with
2 Disorders of Gut-Brain Interaction and Healthy Children

3

4 Vishnu Narayana^{1¶}, Jocelyn Chang^{1,2¶}, Ann R. McMeans^{1,2}, Rona L. Levy³, Robert J.
5 Shulman^{1,2&}, Bruno P. Chumpitazi^{4,5&*}

6 ¹ Department of Pediatrics, Baylor College of Medicine, Houston, TX, USA

7 ² Children's Nutrition Research Center, Houston, TX, USA

8 ³ School of Social Work, University of Washington, Seattle, WA, USA

9 ⁴ Duke University School of Medicine, Department of Pediatrics, Durham, NC USA

10 ⁵ Duke Clinical Research Institute, Durham, NC USA

11

12 * Corresponding Author:

13 Bruno P. Chumpitazi

14 Email: Bruno.chumpitazi@duke.edu (BPC)

15

16 ¶ These authors contributed equally to the work

17 & These authors also contributed equally to the work

18 **Abstract**

19 **Objectives**

20 Restricting fermentable oligosaccharides, disaccharides, monosaccharides, and polyols
21 (**FODMAPs**) intake can alleviate symptoms in children with gut-brain interaction disorders
22 (**DGBI**). Due to the restrictive nature of the low FODMAP diet (**LFD**), the less restrictive
23 FODMAP Gentle diet (**FGD**) has been suggested. However, the types of high FODMAP
24 foods and carbohydrates commonly consumed by US children are unknown, as is the
25 impact of the FGD on a typical diet. This project aimed to identify the high FODMAP foods
26 and proportions of FODMAP carbohydrates consumed by children with DGBI and healthy
27 children (**HC**), and to determine which usually ingested FODMAPs would be restricted on
28 the FGD.

29 **Methods**

30 Three-day diet records from both HC and children with DGBI were analyzed to assess
31 the type of high FODMAP foods and carbohydrates ingested. Results were compared
32 between the groups. The ingested FODMAPs that would be restricted on the FGD was
33 determined.

34 **Results**

35 The number of foods ingested daily was similar between children with DGBI and HC (12.3
36 ± 4.2 vs 12.9 ± 3.4 , respectively); high FODMAP foods comprised most foods eaten in
37 both groups. Children with DGBI (vs HC) ate less high FODMAP foods per day (6.5 ± 2.3
38 vs 8.7 ± 2.4 , $P < 0.0001$, respectively). Fructans were the most consumed FODMAP
39 carbohydrate in both groups and children with DGBI (vs HC) consumed fewer fructans,

40 lactose, fructose, and polyols (all $P < 0.0001$). The top 3 food categories consumed in both
41 groups were wheat-containing foods, dairy, and fruits and 100% fruit juices. In children
42 with DGBI, 80.9% of the high FODMAP foods consumed would be limited on the FGD.

43 **Discussion**

44 Children with DGBI consume significantly fewer high FODMAP foods and
45 carbohydrates than HC. In both groups, the top consumed FODMAP carbohydrates are
46 fructans, lactose, and fructose. A FODMAP Gentle diet would restrict a large majority of
47 high FODMAP foods consumed by children with DGBI.

48 Introduction

49 Abdominal pain-related disorders of gut-brain interaction (**DGBI**) such as irritable
50 bowel syndrome are highly prevalent. Contributors to symptoms include visceral
51 hypersensitivity, psychosocial distress, gut dysmotility, abnormal gut immune function,
52 abnormal gut microbiota composition, and diet.(1) Dietary culprits include highly
53 fermentable oligosaccharides, disaccharides, monosaccharides, and polyols
54 (**FODMAPs**) which are carbohydrates that are poorly absorbed in the small intestine.(2)
55 After reaching the colon, FODMAPs are rapidly fermented, producing gas and other
56 metabolites.(2-4) The gas and metabolite production, in addition to osmotically-driven
57 water secretion into the lumen of the colon can contribute to DGBI symptom
58 generation.(3) Restricting FODMAP dietary intake through a low FODMAP diet (**LFD**)
59 may ameliorate pediatric and adult DGBI symptoms, including pain, bloating, flatus,
60 diarrhea, nausea, and heartburn.(5-7)

61 Unfortunately, there also can be challenges with implementing a LFD, including
62 the need for appropriate educational resources (e.g., dietitian knowledgeable about the
63 LFD) and patient adherence.(8, 9) Some patients have reported cost of the diet to be a
64 barrier to adherence.(10) In addition, given the restrictive nature of a LFD, there is
65 concern for potential adverse effects such as decreases in bacteria associated with health
66 (e.g., *Bifidobacteria*), the potential for unintentional weight loss, and risk of developing
67 disordered eating. Due to these possible negative impacts, some experts do not currently
68 recommend a LFD for children.(11-14) This has led to proposals for diets that restrict
69 fewer fermentable carbohydrates (e.g., the “FODMAP Gentle” diet, **FGD**) in an attempt to
70 be more pediatric-friendly.(14, 15) However, determining potentially effective and less

71 restrictive diets such as the FGD for children is hampered by lack of knowledge related
72 to which high FODMAP foods normally are consumed by children and whether the
73 FODMAP intake of healthy children (**HC**) differs from that of children with DGBI. This
74 information is needed to determine the degree to which a less restrictive diet (e.g., FGD)
75 may impact FODMAP intake. Therefore, we sought to identify the high FODMAP foods
76 and proportions of FODMAP carbohydrates consumed by children with DGBI and HC,
77 and to determine which usually ingested FODMAPs would be restricted on the FGD.

78 **Materials and methods**

79 **Study Design**

80 The data analyzed in this study were dietary intake from a large trial of children
81 with DGBI (NCT03823742) recruited in the study between February 22, 2019 through
82 May 1, 2024, and the dietary intake of HC from a separate study which recruited subjects
83 between May 1, 2013 and October 29, 2018. The Baylor College of Medicine Institutional
84 Review Board for Human Subjects approved both studies. Written informed consent was
85 obtained from study participants' parents (or guardians) and assent from study
86 participants. Prior to any kind of intervention, parents of participants in both studies
87 completed a 3-day baseline food record of their child's habitual diet after receiving
88 detailed information on how to complete the diet records.

89 **Study Population**

90 Participants were children 7-12 years of age from the greater Houston, Texas
91 metropolitan area. Exclusion criteria for both children with DGBI and HC included:
92 children following specific diet alterations (e.g., LFD); those with eating disorders; any
93 recent weight loss ($\geq 5\%$ of body weight); or organic gastrointestinal or systemic diseases
94 (e.g., celiac disease, Crohn's disease, cancer). Children with DGBI met pediatric Rome
95 IV criteria for an abdominal pain-related DGBI.(16) HC could not have had any healthcare
96 visits related to abdominal pain.

97 **Data Collection**

98 Following enrollment, 3-day food records of foods and drinks consumed by the
99 children were completed by the parents. Details included the name of the food, amount

100 eaten, and parts of the food if the food had multiple components (e.g., a sandwich had all
101 components listed). Diet records then were analyzed by two research dietitians well-
102 versed in DGBI and FODMAPs and in providing LFD education. Data were entered into
103 the University of Minnesota Nutrition Data System for Research software version
104 2019.(17) Both children with DGBI and HC kept 2-week pain and stooling diaries as
105 previously reported to insure they met entry criteria.(18, 19)

106 **Analysis and Classification of Foods**

107 High FODMAP foods are identified using guidelines derived from the Monash
108 University Low FODMAP dietary food guide and smartphone app.(20) High FODMAP
109 foods are further characterized by the type of FODMAP carbohydrate within the food, the
110 food categories to which the foods belong, and whether the foods contained high fructose
111 corn syrup as a major ingredient. Statistical comparisons between the dietary intake of
112 the two groups were made using both chi-square for categorical data and t-testing for
113 continuous data using IBM SPSS Statistics for Windows (IBM Corp. Released 2023. IBM
114 SPSS Statistics for Windows, Version 29.0.2.0 Armonk, NY: IBM Corp). In addition, based
115 on published guidelines related to the FGD, we determined whether the high FODMAP
116 foods consumed by children with DGBI would have been restricted.(14, 15)

117 Results

118 At the time of analysis, 77 children with DGBI had completed baseline diet records
119 for the ongoing study. From the previous study, data from 64 HC were available. Self-
120 reported demographic data were similar between groups (Table 1).

121

122 **Table 1. Demographics of Children with DGBI versus Healthy Children**

	DGBI (n=77)	HC (n=64)	P-value
Age (years \pm SD)	10.0 \pm 1.7	9.7 \pm 1.5	0.25
Sex			0.68
Female, n (%)	46 (59.7)	36 (56.3)	
Male, n (%)	31 (40.3)	28 (43.8)	
Race, ethnicity , n (%)			0.07
White, non-Hispanic	22 (28.6)	32 (50.0)	
White, Hispanic	16 (20.8)	14 (21.9)	
Black, non-Hispanic	23 (29.9)	10 (15.6)	
Black, Hispanic	4 (5.2)	2 (3.1)	
Asian, non-Hispanic	2 (2.6)	0 (0.0)	
American Indian/ Hawaiian, non-Hispanic	0 (0.0)	0 (0.0)	
American Indian/ Hawaiian, Hispanic	4 (5.2)	1 (1.6)	
More than one race, non-Hispanic	1 (1.3)	3 (4.7)	
More than one race, Hispanic	1 (1.3)	2 (3.1)	
Black, no ethnicity identified	1 (1.3)	0 (0.0)	
Hispanic, no race identified	3 (3.9)	0 (0.0)	

123

124 Out of a total of 4734 foods consumed in both groups, 2792 (59.0%) were high
125 FODMAP. Children with DGBI consumed fewer high FODMAP foods daily per child than
126 HC (Table 2). High FODMAP foods comprised 1481 (52.5%) and 1311 (68.6%) of total
127 foods consumed by children with DGBI and HC, respectively.

128

129 **Table 2. Daily Food Consumption and High FODMAP Intake in Children with DGBI**
 130 **versus Healthy Children**

	DGBI (n=77)	HC (n=64)	P-value
Total Foods Consumed	2822 ¹	1912 ¹	
Total Foods per subject per day (mean ± SD)	12.3 ± 4.2	12.9 ± 3.4	0.36
High FODMAP Foods per subject per day (mean ± SD)	6.5 ± 2.3	8.7 ± 2.4	<0.001
FODMAP Sugar, n (%)²			
Fructans ³	850 (30)	748 (39)	<0.0001
Lactose ⁴	552 (20)	527 (28)	<0.0001
Fructose ⁵	412 (15)	570 (30)	<0.0001
Polyols ⁶	82 (3)	100 (5)	<0.0001
Galacto-oligosaccharides ⁷	82 (3)	53 (3)	0.79
HFCS ⁸	248 (9)	388 (20)	<0.0001
Food Category, n (%)²			
Wheat	557 (20)	534 (28)	<0.0001
Dairy	344 (12)	342 (18)	<0.0001
Fruits and 100% Fruit Juices	168 (6)	194 (10)	<0.0001
Beverages (e.g., sodas, juice w/ high fructose corn syrup)	144 (5)	93 (5)	0.71
Condiments (e.g., salad dressing, mayonnaise, ketchup, jelly, jam)	125 (4)	74 (4)	0.35
Sweet Snacks (e.g., candy, cookies, brownies)	125 (4)	143 (7)	<0.0001
Vegetables	111 (4)	80 (4)	0.67
Savory Snacks (e.g., chips)	84 (3)	90 (5)	0.002
Sweeteners (e.g., fructose, honey, added polyols)	64 (2)	35 (2)	0.30
Legumes	42 (1)	16 (1)	0.06
Nuts, seeds, nut/seed butters	8 (<1)	13 (1)	0.04
Top High FODMAP Foods Consumed, n (%)²			
Milk	146 (5.2)	121 (6.3)	.09
Sodas ⁹	69 (2.4)	31 (1.6)	.05
Fructose-sweetened drinks ⁹	67 (2.4)	45 (2.4)	0.96
Breaded Animal Protein ¹⁰	65 (2.3)	34 (1.8)	0.22
Fresh fruit	62 (2.2)	74 (3.9)	<0.001
Loaf bread ¹¹	61 (2.2)	63 (3.3)	0.02
Savory condiments ¹²	58 (2.1)	18 (0.9)	<0.005
100% Fruit juice	44 (1.6)	46 (2.4)	0.04
Sweet/enriched bread ¹¹	43 (1.5)	51 (2.8)	<0.01
Pizza ¹⁰	42 (1.5)	34 (1.8)	0.44
Mixed pasta ¹⁰	35 (1.2)	35 (1.8)	0.10
Cookies ¹¹	34 (1.2)	35 (1.8)	0.08
Seasonings ¹³	23 (0.8)	41 (2.1)	<0.001

Crackers ¹¹	20 (0.7)	40 (2.1)	<0.0001
Fruit snacks	20 (0.7)	35 (1.8)	<0.0005

131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155

¹Total count of all foods consumed

²Percent out of total foods consumed

³Fructans are found in wheat, garlic, onion, legumes, and inulin and chicory root which are types of fiber and the main ingredient in fiber supplements or added to baked goods to increase fiber content.

⁴Lactose is found in dairy products and may be added to some food products such as chocolate bars, flavored chips, and baked sweets.

⁵Fructose is found in many fruits and some vegetables such as apples, pears, and asparagus; it is found in honey and agave syrup and may be added as fructose or high fructose corn syrup to provide a sweet flavor in different food products, such as sodas, jams, ketchup, and baked sweets.

⁶Polyols can be found in different fruits such as peaches, plums, and avocado and may be added in different forms (i.e., sorbitol, mannitol, xylitol) to impart a sweet flavor to different food products, most notably sugar free candies and sugar free gums.

⁷Galacto-oligosaccharides can be found in beans and nuts.

⁸High fructose corn syrup; a syrup of fructose and glucose, usually found in a 55%/45% ratio in food products

⁹Contains high fructose corn syrup

¹⁰Contains wheat, and onion and/or garlic seasoning

¹¹Contains wheat and may contain high fructose corn syrup

¹²Contains onion, garlic, and/or high fructose corn syrup

¹³Contains dried or powdered onion and/or garlic added during at home food preparation process. Does not include packaged foods that already contain spices.

156 Based on the FODMAP carbohydrate, fructan-containing foods were the most
157 numerous in the diet of both groups, followed by foods containing lactose and fructose
158 (Table 2). Individual foods containing more than one type of FODMAP contributed 501
159 (33.8%) and 638 (48.7%) of the high FODMAP foods consumed by children with DGBI
160 and HC, respectively. Children with DGBI (vs HC) consumed significantly less fructans,
161 lactose, fructose, and polyols but not galactans.

162 Based on food category, wheat-containing foods were the most prevalent, followed
163 by dairy and fruits/100% fruit juices (Table 2); children with DGBI consumed significantly
164 fewer of these foods compared to HC. Children with DGBI (vs HC) consumed less
165 FODMAP-containing sweet and savory snacks. There was no significant difference in
166 intake of beverages, condiments, vegetables, and sweeteners.

167 Based on the high FODMAP-containing foods, children with DGBI consumed most
168 frequently milk, sodas, and fructose-sweetened drinks (e.g., fruit-flavored drinks
169 containing high fructose corn syrup) whereas HC consumed milk, fresh fruit, and wheat-
170 containing loaf bread (Table 2). High FODMAP foods containing high fructose corn syrup
171 as a major ingredient were consumed by 16.7% (248) of children with DGBI and 29.6%
172 (388) of HC.

173 In children with DGBI, 1198 (80.9%) of the consumed high FODMAP foods would
174 be restricted on a FGD. Of the 283 high FODMAP foods permitted on a FGD, 139 (49.1%)
175 contained high fructose corn syrup as a major ingredient. Of these, beverages and
176 condiments (e.g., sodas, jam/jelly, ready-to-drink lemonade, pancake syrup, and flavored
177 drinks) comprised a significant proportion.

178 **Discussion**

179 Determining the type of high FODMAP foods and FODMAP carbohydrates
180 consumed by children with DGBI and HC children is important for tailoring educational
181 content and aid in the creation of diets that restrict FODMAP intake. We determined that
182 high FODMAP foods comprise a majority of the foods eaten by both children with DGBI
183 and HC. Children with DGBI (vs HC) consumed fewer high FODMAP foods and several
184 FODMAP carbohydrates. We also identified that a large majority of FODMAP foods
185 normally eaten by children with DGBI would be restricted on a FGD.

186 We hypothesize the less frequent ingestion of high FODMAP foods by children
187 with DGBI may relate to self-restriction (either intentional or due to unconscious behavior)
188 to avoid worsening their GI symptoms. Supporting the former, a recent study of school
189 children reporting bothersome GI symptoms identified an association between high
190 FODMAP foods and worse GI symptoms.(21) Furthermore, children with IBS self-restrict
191 more foods than healthy children.(22) Additional studies are needed to determine the
192 reasons why children with DGBI eat fewer FODMAPs than HC.

193 To our knowledge, this is the first study to describe FODMAP intake in children
194 with DGBI and HC under normal dietary circumstances. Fructans were the FODMAP
195 carbohydrate ingested most commonly in both groups. A U.S. Department of Agriculture
196 survey from 1994-1996 estimated that 75% of fructan intake in individuals was from wheat
197 and 25% from onions.(23) Our study findings are in agreement with the survey data, as
198 wheat-containing products, particularly breaded meat products such as chicken nuggets
199 and wheat bread, were the primary contributors to fructan intake for both groups. Onion
200 and garlic powders via seasoning contributed to fructan intake to a lesser degree. Some

201 of the consumed foods containing onion and garlic powder seasonings included flavored
202 chips, processed and seasoned meats, and seasoned beans (i.e., refried beans).
203 Fructans, which arrive essentially intact into the colon after consumption, are often a focus
204 of a low FODMAP diet as fructans have been found to induce more DGBI symptoms than
205 other FODMAPs.(24) Given our finding that wheat (a FODMAP) is the most commonly
206 ingested category of food, particular attention to restriction of wheat products should be
207 a focus of a reduced FODMAP diet.

208 After fructans, lactose and fructose were the most commonly ingested FODMAPs.
209 Lactose malabsorption is not more common in those with DGBI than HC but lactose
210 intolerance appears to be; likely related to visceral hyperalgesia commonly found in those
211 with DGBI.(25) Similar to education regarding wheat ingestion, our data would suggest
212 emphasis also should be given to a reduction in fructose-sweetened drinks including
213 sodas and 100% fruit juice in children with DGBI. In contrast, a global reduction in fresh
214 fruit might not be warranted given its value as an important component of a healthy diet.

215 We found that the vast majority of high FODMAP foods commonly consumed by
216 the children with DGBI would be restricted on a FGD which has been proposed as an
217 alternative, pediatric-friendly and less restrictive version to the traditional LFD.(14, 15)
218 The United Kingdom's National Institute of Clinical Excellence (NICE) dietary guidelines
219 that reduce FODMAP intake also has been recommended but data are conflicting as to
220 whether it has similar (worse or better) efficacy as the traditional LFD in adults and
221 children.(14) One particular advantage of the FGD may be its relative ease related to
222 education and adherence as only 15 different food and/or food types (e.g., wheat) are
223 restricted. However, its efficacy in children with DGBI remains to be determined.

224 Of the high FODMAP foods consumed by the children in our study that would be
225 allowed on a FGD, about half contained high fructose corn syrup as a major ingredient-
226 not surprising given its prevalence in many foods in the US. Important to note is that the
227 FGD was developed in Australia where high fructose corn syrup is not added to food as
228 frequently as in the US.(15) Thus, in the US, restriction of foods containing this FODMAP
229 (e.g., sodas, jams) is likely needed when children are placed on a FGD.

230 There are some limitations to our study. Parents may not have accurately recorded
231 food intake of their children with the use of food records. However, 3-day food records
232 are considered a valid method to record food consumption and are considered a better
233 method of food capture compared to other approaches such as 24-hour recall.(26)
234 Children were recruited from a single geographic area and diets may differ based on
235 geographic location. On the other hand, our study population included a racially and
236 ethnically diverse population which may increase the generalizability of our findings.
237 Further studies in different geographic areas can be conducted to confirm our findings

238 A major strength of our study is that, to our knowledge, it is the first to characterize
239 and compare the FODMAP intake of children with DGBI and HC. Also, the two
240 populations were well-matched in terms of demographics, increasing the rigor of the
241 findings. Additionally, the dietitians participating in the study had experience with
242 analyzing food records for high FODMAP foods and maintained high quality in their
243 review, including follow-up with families, if needed, regarding specific food record entries.

244 In summary, both children with DGBI and HC consume a significant amount of high
245 FODMAP foods, though children with DGBI consume relatively less.. Fructans, lactose,
246 and fructose were the most consumed FODMAP carbohydrates via wheat-containing

247 foods, dairy, and fruits/100% fruit juices. Targeted and less restrictive diets such as the
248 FGD, with modification based on our findings, may effectively restrict a large majority of
249 the high FODMAP foods consumed by children with DGBI.

250 REFERENCES

- 251 1. Thapar N, Benninga MA, Crowell MD, Di Lorenzo C, Mack I, Nurko S, et al. Paediatric
252 functional abdominal pain disorders. *Nat Rev Dis Primers*. 2020;6(1):89.
- 253 2. Algera J, Colomier E, Simren M. The Dietary Management of Patients with Irritable
254 Bowel Syndrome: A Narrative Review of the Existing and Emerging Evidence.
255 *Nutrients*. 2019;11(9).
- 256 3. Barrett JS, Geary RB, Muir JG, Irving PM, Rose R, Rosella O, et al. Dietary poorly
257 absorbed, short-chain carbohydrates increase delivery of water and fermentable
258 substrates to the proximal colon. *Aliment Pharmacol Ther*. 2010;31(8):874-82.
- 259 4. Shepherd SJ, Lomer MC, Gibson PR. Short-chain carbohydrates and functional
260 gastrointestinal disorders. *Am J Gastroenterol*. 2013;108(5):707-17.
- 261 5. Chumpitazi BP, Cope JL, Hollister EB, Tsai CM, McMeans AR, Luna RA, et al.
262 Randomised clinical trial: gut microbiome biomarkers are associated with clinical
263 response to a low FODMAP diet in children with the irritable bowel syndrome. *Aliment*
264 *Pharmacol Ther*. 2015;42(4):418-27.
- 265 6. Khalighi Sikaroudi M, Soltani S, Ghoreishy SM, Ebrahimi Z, Shidfar F, Dehnad A.
266 Effects of a low FODMAP diet on the symptom management of patients with irritable
267 bowel syndrome: a systematic umbrella review with the meta-analysis of clinical
268 trials. *Food Funct*. 2024;15(10):5195-208.
- 269 7. Jent S, Bez NS, Haddad J, Catalano L, Egger KS, Raia M, et al. The efficacy and
270 real-world effectiveness of a diet low in fermentable oligo-, di-, monosaccharides and
271 polyols in irritable bowel syndrome: A systematic review and meta-analysis. *Clinical*
272 *nutrition*. 2024;43(6):1551-62.

- 273 8. Narayana V, McMeans AR, Levy RL, Shulman RJ, Chumpitazi BP. Children with
274 functional abdominal pain disorders successfully decrease FODMAP food intake on
275 a low FODMAP diet with modest improvements in nutritional intake and diet quality.
276 *Neurogastroenterol Motil.* 2022;34(10):e14392.
- 277 9. Tenenbaum RB, Czyzewski D, McMeans A, Narayana V, Chumpitazi BP, Levy RL,
278 et al. Factors Associated With Adherence to a Low Fermentable Carbohydrate Diet
279 in Children With Functional Gastrointestinal Disorders. *Journal of the Academy of*
280 *Nutrition and Dietetics.* 2024;124(6):757-62.
- 281 10. Maagaard L, Ankersen DV, Vegh Z, Burisch J, Jensen L, Pedersen N, et al. Follow-
282 up of patients with functional bowel symptoms treated with a low FODMAP diet. *World*
283 *J Gastroenterol.* 2016;22(15):4009-19.
- 284 11. Frieling T, Heise J, Krummen B, Hundorf C, Kalde S. Tolerability of FODMAP -
285 reduced diet in irritable bowel syndrome - efficacy, adherence, and body weight
286 course. *Z Gastroenterol.* 2019;57(6):740-4.
- 287 12. Halmos EP, Christophersen CT, Bird AR, Shepherd SJ, Gibson PR, Muir JG. Diets
288 that differ in their FODMAP content alter the colonic luminal microenvironment. *Gut.*
289 2015;64(1):93-100.
- 290 13. Mitchell H, Porter J, Gibson PR, Barrett J, Garg M. Review article: implementation of
291 a diet low in FODMAPs for patients with irritable bowel syndrome-directions for future
292 research. *Aliment Pharmacol Ther.* 2019;49(2):124-39.
- 293 14. Rhys-Jones D, Varney JE, Muir JG, Gibson PR, Halmos EP. Application of The
294 FODMAP Diet in a Paediatric Setting. *Nutrients.* 2022;14(20).

- 295 15. Halmos EP, Gibson PR. Controversies and reality of the FODMAP diet for patients
296 with irritable bowel syndrome. *J Gastroenterol Hepatol*. 2019;34(7):1134-42.
- 297 16. Hyams JS, Di Lorenzo C, Saps M, Shulman RJ, Staiano A, van Tilburg M. Functional
298 Disorders: Children and Adolescents. *Gastroenterology*. 2016;150:1456–68.
- 299 17. Feskanich D, Sielaff BH, Chong K, Buzzard IM. Computerized collection and analysis
300 of dietary intake information. *Comput Methods Programs Biomed*. 1989;30(1):47-57.
- 301 18. Czyzewski DI, Lane MM, Weidler EM, Williams AE, Swank PR, Shulman RJ. The
302 interpretation of Rome III criteria and method of assessment affect the irritable bowel
303 syndrome classification of children. *Aliment Pharmacol Ther*. 2011;33(3):403-11.
- 304 19. Self MM, Czyzewski DI, Chumpitazi BP, Weidler EM, Shulman RJ. Subtypes of
305 irritable bowel syndrome in children and adolescents. *Clin Gastroenterol Hepatol*.
306 2014;12(9):1468-73.
- 307 20. [Available from: [https://www.monashfodmap.com/ibs-central/i-have-ibs/get-the-](https://www.monashfodmap.com/ibs-central/i-have-ibs/get-the-app/)
308 [app/](https://www.monashfodmap.com/ibs-central/i-have-ibs/get-the-app/)].
- 309 21. Saps M, Velasco-Benitez CA, Velasco-Suarez DA, Alvarez-Baumgartner M, Balda
310 AN, Arrizabalo S. Gastrointestinal Symptoms and Fermentable Oligosaccharides,
311 Disaccharides, Monosaccharides, and Polyols Sources in Schoolchildren-A Pilot
312 Study. *Children (Basel)*. 2024;11(6).
- 313 22. Chumpitazi BP, Weidler EM, Lu DY, Tsai CM, Shulman RJ. Self-Perceived Food
314 Intolerances Are Common and Associated with Clinical Severity in Childhood Irritable
315 Bowel Syndrome. *Journal of the Academy of Nutrition and Dietetics*.
316 2016;116(9):1458-64.

- 317 23. Moshfegh AJ, Friday JE, Goldman JP, Ahuja JK. Presence of inulin and oligofructose
318 in the diets of Americans. *J Nutr.* 1999;129(7 Suppl):1407S-11S.
- 319 24. Shepherd SJ, Parker FC, Muir JG, Gibson PR. Dietary triggers of abdominal
320 symptoms in patients with irritable bowel syndrome: randomized placebo-controlled
321 evidence. *Clin Gastroenterol Hepatol.* 2008;6(7):765-71.
- 322 25. Yang J, Deng Y, Chu H, Cong Y, Zhao J, Pohl D, et al. Prevalence and presentation
323 of lactose intolerance and effects on dairy product intake in healthy subjects and
324 patients with irritable bowel syndrome. *Clin Gastroenterol Hepatol.* 2013;11(3):262-8
325 e1.
- 326 26. Crawford PB, Obarzanek E, Morrison J, Sabry ZI. Comparative advantage of 3-day
327 food records over 24-hour recall and 5-day food frequency validated by observation
328 of 9- and 10-year-old girls. *J Am Diet Assoc.* 1994;94(6):626-30.