It is made available under a CC-BY 4.0 International license .

1	Characterizing Highly Fermentable Carbohydrate Foods in the Diets of Children with
2	Disorders of Gut-Brain Interaction and Healthy Children
3	
4	Vishnu Narayana ^{1¶} , Jocelyn Chang ^{1,2¶} , Ann R. McMeans ^{1,2} , Rona L. Levy ³ , Robert J.
5	Shulman ^{1,2&} , Bruno P. Chumpitazi ^{4,5&*}
6	¹ Department of Pediatrics, Baylor College of Medicine, Houston, TX, USA
7	² Children's Nutrition Research Center, Houston, TX, USA
8	³ School of Social Work, University of Washington, Seattle, WA, USA
9	⁴ Duke University School of Medicine, Department of Pediatrics, Durham, NC USA
10	⁵ Duke Clinical Research Institute, Durham, NC USA
11	
12	* Corresponding Author:
13	Bruno P. Chumpitazi
14	Email: Bruno.chumpitazi@duke.edu (BPC)
15	
16	[¶] These authors contributed equally to the work
17	^{&} These authors also contributed equally to the work

It is made available under a CC-BY 4.0 International license .

Abstract

19 **Objectives**

Restricting fermentable oligosaccharides, disaccharides, monosaccharides, and polyols 20 (FODMAPs) intake can alleviate symptoms in children with gut-brain interaction disorders 21 (DGBI). Due to the restrictive nature of the low FODMAP diet (LFD), the less restrictive 22 FODMAP Gentle diet (FGD) has been suggested. However, the types of high FODMAP 23 foods and carbohydrates commonly consumed by US children are unknown, as is the 24 impact of the FGD on a typical diet. This project aimed to identify the high FODMAP foods 25 and proportions of FODMAP carbohydrates consumed by children with DGBI and healthy 26 children (**HC**), and to determine which usually ingested FODMAPs would be restricted on 27 28 the FGD.

29 Methods

Three-day diet records from both HC and children with DGBI were analyzed to assess the type of high FODMAP foods and carbohydrates ingested. Results were compared between the groups. The ingested FODMAPs that would be restricted on the FGD was determined.

34 **Results**

The number of foods ingested daily was similar between children with DGBI and HC (12.3 ± 4.2 vs 12.9 ± 3.4, respectively); high FODMAP foods comprised most foods eaten in both groups. Children with DGBI (vs HC) ate less high FODMAP foods per day (6.5 ± 2.3 vs 8.7 ± 2.4, P<0.0001, respectively). Fructans were the most consumed FODMAP carbohydrate in both groups and children with DGBI (vs HC) consumed fewer fructans,

40 lactose, fructose, and polyols (all P<0.0001). The top 3 food categories consumed in both

groups were wheat-containing foods, dairy, and fruits and 100% fruit juices. In children

42 with DGBI, 80.9% of the high FODMAP foods consumed would be limited on the FGD.

43 **Discussion**

Children with DGBI consume significantly fewer high FODMAP foods and carbohydrates than HC. In both groups, the top consumed FODMAP carbohydrates are fructans, lactose, and fructose. A FODMAP Gentle diet would restrict a large majority of high FODMAP foods consumed by children with DGBI.

It is made available under a CC-BY 4.0 International license .

48 Introduction

Abdominal pain-related disorders of gut-brain interaction (DGBI) such as irritable 49 bowel syndrome are highly prevalent. Contributors to symptoms include visceral 50 hypersensitivity, psychosocial distress, gut dysmotility, abnormal gut immune function, 51 abnormal gut microbiota composition, and diet.(1) Dietary culprits include highly 52 fermentable oligosaccharides, disaccharides, monosaccharides, 53 and polyols 54 (FODMAPs) which are carbohydrates that are poorly absorbed in the small intestine.(2) After reaching the colon, FODMAPs are rapidly fermented, producing gas and other 55 metabolites.(2-4) The gas and metabolite production, in addition to osmotically-driven 56 water secretion into the lumen of the colon can contribute to DGBI symptom 57 generation.(3) Restricting FODMAP dietary intake through a low FODMAP diet (LFD) 58 may ameliorate pediatric and adult DGBI symptoms, including pain, bloating, flatus, 59 diarrhea, nausea, and heartburn.(5-7) 60

Unfortunately, there also can be challenges with implementing a LFD, including 61 the need for appropriate educational resources (e.g., dietitian knowledgeable about the 62 LFD) and patient adherence. (8, 9) Some patients have reported cost of the diet to be a 63 barrier to adherence.(10) In addition, given the restrictive nature of a LFD, there is 64 concern for potential adverse effects such as decreases in bacteria associated with health 65 (e.g., *Bifidobacteria*), the potential for unintentional weight loss, and risk of developing 66 disordered eating. Due to these possible negative impacts, some experts do not currently 67 recommend a LFD for children.(11-14) This has led to proposals for diets that restrict 68 fewer fermentable carbohydrates (e.g., the "FODMAP Gentle" diet, FGD) in an attempt to 69 be more pediatric-friendly. (14, 15) However, determining potentially effective and less 70

It is made available under a CC-BY 4.0 International license .

restrictive diets such as the FGD for children is hampered by lack of knowledge related to which high FODMAP foods normally are consumed by children and whether the FODMAP intake of healthy children (**HC**) differs from that of children with DGBI. This information is needed to determine the degree to which a less restrictive diet (e.g., FGD) may impact FODMAP intake. Therefore, we sought to identify the high FODMAP foods and proportions of FODMAP carbohydrates consumed by children with DGBI and HC, and to determine which usually ingested FODMAPs would be restricted on the FGD.

78 Materials and methods

79 Study Design

The data analyzed in this study were dietary intake from a large trial of children 80 with DGBI (NCT03823742) recruited in the study between February 22, 2019 through 81 May 1, 2024, and the dietary intake of HC from a separate study which recruited subjects 82 between May 1, 2013 and October 29, 2018. The Baylor College of Medicine Institutional 83 Review Board for Human Subjects approved both studies. Written informed consent was 84 obtained from study participants' parents (or guardians) and assent from study 85 participants. Prior to any kind of intervention, parents of participants in both studies 86 completed a 3-day baseline food record of their child's habitual diet after receiving 87 88 detailed information on how to complete the diet records.

89 Study Population

Participants were children 7-12 years of age from the greater Houston, Texas metropolitan area. Exclusion criteria for both children with DGBI and HC included: children following specific diet alterations (e.g., LFD); those with eating disorders; any recent weight loss (\geq 5 % of body weight); or organic gastrointestinal or systemic diseases (e.g., celiac disease, Crohn's disease, cancer). Children with DGBI met pediatric Rome IV criteria for an abdominal pain-related DGBI.(16) HC could not have had any healthcare visits related to abdominal pain.

97 Data Collection

Following enrollment, 3-day food records of foods and drinks consumed by the children were completed by the parents. Details included the name of the food, amount

eaten, and parts of the food if the food had multiple components (e.g., a sandwich had all
 components listed). Diet records then were analyzed by two research dietitians well versed in DGBI and FODMAPs and in providing LFD education. Data were entered into
 the University of Minnesota Nutrition Data System for Research software version
 2019.(17) Both children with DGBI and HC kept 2-week pain and stooling diaries as
 previously reported to insure they met entry criteria.(18, 19)

Analysis and Classification of Foods

High FODMAP foods are identified using guidelines derived from the Monash 107 University Low FODMAP dietary food guide and smartphone app.(20) High FODMAP 108 foods are further characterized by the type of FODMAP carbohydrate within the food, the 109 food categories to which the foods belong, and whether the foods contained high fructose 110 corn syrup as a major ingredient. Statistical comparisons between the dietary intake of 111 112 the two groups were made using both chi-square for categorical data and t-testing for continuous data using IBM SPSS Statistics for Windows (IBM Corp. Released 2023. IBM 113 SPSS Statistics for Windows, Version 29.0.2.0 Armonk, NY: IBM Corp). In addition, based 114 115 on published guidelines related to the FGD, we determined whether the high FODMAP foods consumed by children with DGBI would have been restricted.(14, 15) 116

It is made available under a CC-BY 4.0 International license .

117 **Results**

118 At the time of analysis, 77 children with DGBI had completed baseline diet records

119 for the ongoing study. From the previous study, data from 64 HC were available. Self-

- reported demographic data were similar between groups (Table 1).
- 121
- 122

Table 1. Demographics of Children with DGBI versus Healthy Children

	DGBI (n=77)	HC (n=64)	P-value
Age (years ± SD)	10.0 ± 1.7	9.7 ± 1.5	0.25
Sex			0.68
Female, n (%)	46 (59.7)	36 (56.3)	
Male, n (%)	31 (40.3)	28 (43.8)	
Race, ethnicity, n (%)			0.07
White, non-Hispanic	22 (28.6)	32 (50.0)	
White, Hispanic	16 (20.8)	14 (21.9)	
Black, non-Hispanic	23 (29.9)	10 (15.6)	
Black, Hispanic	4 (5.2)	2 (3.1)	
Asian, non-Hispanic	2 (2.6)	0 (0.0)	
American Indian/ Hawaiian, non-	0 (0.0)	0 (0.0)	
Hispanic			
American Indian/ Hawaiian, Hispanic	4 (5.2)	1 (1.6)	
More than one race, non-Hispanic	1 (1.3)	3 (4.7)	
More than one race, Hispanic	1 (1.3)	2 (3.1)	
Black, no ethnicity identified	1 (1.3)	0 (0.0)	
Hispanic, no race identified	3 (3.9)	0 (0.0)	

123

Out of a total of 4734 foods consumed in both groups, 2792 (59.0%) were high FODMAP. Children with DGBI consumed fewer high FODMAP foods daily per child than HC (Table 2). High FODMAP foods comprised 1481 (52.5%) and 1311 (68.6%) of total foods consumed by children with DGBI and HC, respectively.

It is made available under a CC-BY 4.0 International license .

Table 2. Daily Food Consumption and High FODMAP Intake in Children with DGBI versus Healthy Children

	DGBI	HC	P-value
	(n=77)	(n=64)	
Total Foods Consumed	2822 ¹	1912 ¹	
Total Foods per subject per day	12.3 ± 4.2	12.9 ± 3.4	0.36
(mean ± SD)			
High FODMAP Foods per subject per day	6.5 ± 2.3	8.7 ± 2.4	<0.001
(mean ± SD)			
FODMAP Sugar, n (%) ²			
Fructans ³	850 (30)	748 (39)	< 0.0001
Lactose ⁴	552 (20)	527 (28)	< 0.0001
Fructose ⁵	412 (15)	570 (30)	< 0.0001
Polyols ⁶	82 (3)	100 (5)	<0.0001
Galacto-oligosaccharides ⁷	82 (3)	53 (3)	0.79
HFCS [®]	248 (9)	388 (20)	<0.0001
Food Category n (%) ²			
Wheat	557 (20)	534 (28)	<0.0001
Dairy	344 (12)	342 (18)	<0.0001
Fruits and 100% Fruit Juices	168 (6)	194 (10)	<0.0001
Beverages (e.g. sodas juice w/ high fructose	144 (5)	93 (5)	0.71
corn syrup)	111(0)		0.71
Condiments (e.g., salad dressing, mayonnaise,	125 (4)	74 (4)	0.35
Ketchup, jelly, jam) Sweet Spacks (e.g., candy, cookies, brownies)	125 (4)	1/13 (7)	<0.0001
Vegetables	111 (4)	80 (4)	0.67
Savory Spacks (e.g., chins)	84 (3)	90 (5)	0.07
Sweeteners (e.g., tructose honey added	64 (2)	35 (2)	0.002
polyols)	04 (2)	00 (2)	0.00
Legumes	42 (1)	16 (1)	0.06
Nuts, seeds, nut/seed butters	8 (<1)	13 (1)	0.04
Top High FODMAP Foods Consumed n (%) ²			
Milk	146 (5.2)	121 (6.3)	.09
Sodas ⁹	69 (2.4)	31 (1.6)	.05
Fructose-sweetened drinks ⁹	67 (2.4)	45 (2.4)	0.96
Breaded Animal Protein ¹⁰	65 (2.3)	34 (1.8)	0.22
Fresh fruit	62 (2.2)	74 (3.9)	< 0.001
Loaf bread ¹¹	61 (2.2)	63 (3.3)	0.02
Savory condiments ¹²	58 (2.1)	18 (0.9)	< 0.005
100% Fruit juice	44 (1.6)	46 (2.4)	0.04
Sweet/enriched bread ¹¹	43 (1.5)	51 (2.8)	<0.01
Pizza ¹⁰	42 (1.5)	34 (1.8)	0.44
Mixed pasta ¹⁰	35 (1.2)	35 (1.8)	0.10
Cookies ¹¹	34 (1.2)	35 (1.8)	0.08
Seasonings ¹³	23 (0.8)	41 (2.1)	<0.001

It is made available under a CC-BY 4.0 International license .

Crackers ¹¹	20 (0.7)	40 (2.1)	<0.0001
Fruit snacks	20 (0.7)	35 (1.8)	<0.0005

131

- 132 ¹Total count of all foods consumed
- 133 ²Percent out of total foods consumed
- ¹³⁴³Fructans are found in wheat, garlic, onion, legumes, and inulin and chicory root which are types
- of fiber and the main ingredient in fiber supplements or added to baked goods to increase fiber
- 136 content.
- ¹³⁷ ⁴Lactose is found in dairy products and may be added to some food products such as chocolate
- 138 bars, flavored chips, and baked sweets.
- ¹³⁹ ⁵Fructose is found in many fruits and some vegetables such as apples, pears, and asparagus; it
- is found in honey and agave syrup and may be added as fructose or high fructose corn syrup to
- 141 provide a sweet flavor in different food products, such as sodas, jams, ketchup, and baked
- sweets.
- ⁶Polyols can be found in different fruits such as peaches, plums, and avocado and may be
- added in different forms (i.e., sorbitol, mannitol, xylitol) to impart a sweet flavor to different food
- 145 products, most notably sugar free candies and sugar free gums.
- ⁷Galacto-oligosaccharides can be found in beans and nuts.
- ¹⁴⁷ ⁸High fructose corn syrup; a syrup of fructose and glucose, usually found in a 55%/45% ratio in
- 148 food products
- ⁹Contains high fructose corn syrup
- ¹⁰Contains wheat, and onion and/or garlic seasoning
- ¹⁵¹¹¹Contains wheat and may contain high fructose corn syrup
- ¹²Contains onion, garlic, and/or high fructose corn syrup
- ¹⁵³¹³Contains dried or powdered onion and/or garlic added during at home food preparation
- 154 process. Does not include packaged foods that already contain spices.

155

Based on the FODMAP carbohydrate, fructan-containing foods were the most numerous in the diet of both groups, followed by foods containing lactose and fructose (Table 2). Individual foods containing more than one type of FODMAP contributed 501 (33.8%) and 638 (48.7%) of the high FODMAP foods consumed by children with DGBI and HC, respectively. Children with DGBI (vs HC) consumed significantly less fructans, lactose, fructose, and polyols but not galactans.

Based on food category, wheat-containing foods were the most prevalent, followed by dairy and fruits/100% fruit juices (Table 2); children with DGBI consumed significantly fewer of these foods compared to HC. Children with DGBI (vs HC) consumed less FODMAP-containing sweet and savory snacks. There was no significant difference in intake of beverages, condiments, vegetables, and sweeteners.

Based on the high FODMAP-containing foods, children with DGBI consumed most frequently milk, sodas, and fructose-sweetened drinks (e.g., fruit-flavored drinks containing high fructose corn syrup) whereas HC consumed milk, fresh fruit, and wheatcontaining loaf bread (Table 2). High FODMAP foods containing high fructose corn syrup as a major ingredient were consumed by 16.7% (248) of children with DGBI and 29.6% (388) of HC.

In children with DGBI, 1198 (80.9%) of the consumed high FODMAP foods would be restricted on a FGD. Of the 283 high FODMAP foods permitted on a FGD, 139 (49.1%) contained high fructose corn syrup as a major ingredient. Of these, beverages and condiments (e.g., sodas, jam/jelly, ready-to-drink lemonade, pancake syrup, and flavored drinks) comprised a significant proportion.

It is made available under a CC-BY 4.0 International license .

178 **Discussion**

Determining the type of high FODMAP foods and FODMAP carbohydrates consumed by children with DGBI and HC children is important for tailoring educational content and aid in the creation of diets that restrict FODMAP intake. We determined that high FODMAP foods comprise a majority of the foods eaten by both children with DGBI and HC. Children with DGBI (vs HC) consumed fewer high FODMAP foods and several FODMAP carbohydrates. We also identified that a large majority of FODMAP foods normally eaten by children with DGBI would be restricted on a FGD.

We hypothesize the less frequent ingestion of high FODMAP foods by children with DGBI may relate to self-restriction (either intentional or due to unconscious behavior) to avoid worsening their GI symptoms. Supporting the former, a recent study of school children reporting bothersome GI symptoms identified an association between high FODMAP foods and worse GI symptoms.(21) Furthermore, children with IBS self-restrict more foods than healthy children.(22) Additional studies are needed to determine the reasons why children with DGBI eat fewer FODMAPs than HC.

To our knowledge, this is the first study to describe FODMAP intake in children 193 with DGBI and HC under normal dietary circumstances. Fructans were the FODMAP 194 195 carbohydrate ingested most commonly in both groups. A U.S. Department of Agriculture survey from 1994-1996 estimated that 75% of fructan intake in individuals was from wheat 196 and 25% from onions.(23) Our study findings are in agreement with the survey data, as 197 198 wheat-containing products, particularly breaded meat products such as chicken nuggets and wheat bread, were the primary contributors to fructan intake for both groups. Onion 199 and garlic powders via seasoning contributed to fructan intake to a lesser degree. Some 200

It is made available under a CC-BY 4.0 International license .

201 of the consumed foods containing onion and garlic powder seasonings included flavored 202 chips, processed and seasoned meats, and seasoned beans (i.e., refried beans). 203 Fructans, which arrive essentially intact into the colon after consumption, are often a focus 204 of a low FODMAP diet as fructans have been found to induce more DGBI symptoms than 205 other FODMAPs.(24) Given our finding that wheat (a FODMAP) is the most commonly 206 ingested category of food, particular attention to restriction of wheat products should be 207 a focus of a reduced FODMAP diet.

After fructans, lactose and fructose were the most commonly ingested FODMAPs. Lactose malabsorption is not more common in those with DGBI than HC but lactose intolerance appears to be; likely related to visceral hyperalgesia commonly found in those with DGBI.(25) Similar to education regarding wheat ingestion, our data would suggest emphasis also should be given to a reduction in fructose-sweetened drinks including sodas and 100% fruit juice in children with DGBI. In contrast, a global reduction in fresh fruit might not be warranted given its value as an important component of a healthy diet.

We found that the vast majority of high FODMAP foods commonly consumed by 215 the children with DGBI would be restricted on a FGD which has been proposed as an 216 217 alternative, pediatric-friendly and less restrictive version to the traditional LFD.(14, 15) The United Kingdom's National Institute of Clinical Excellence (NICE) dietary guidelines 218 that reduce FODMAP intake also has been recommended but data are conflicting as to 219 220 whether it has similar (worse or better) efficacy as the traditional LFD in adults and children.(14) One particular advantage of the FGD may be its relative ease related to 221 222 education and adherence as only 15 different food and/or food types (e.g., wheat) are 223 restricted. However, its efficacy in children with DGBI remains to be determined.

It is made available under a CC-BY 4.0 International license .

Of the high FODMAP foods consumed by the children in our study that would be allowed on a FGD, about half contained high fructose corn syrup as a major ingredientnot surprising given its prevalence in many foods in the US. Important to note is that the FGD was developed in Australia where high fructose corn syrup is not added to food as frequently as in the US.(15) Thus, in the US, restriction of foods containing this FODMAP (e.g., sodas, jams) is likely needed when children are placed on a FGD.

There are some limitations to our study. Parents may not have accurately recorded 230 food intake of their children with the use of food records. However, 3-day food records 231 are considered a valid method to record food consumption and are considered a better 232 method of food capture compared to other approaches such as 24-hour recall.(26) 233 Children were recruited from a single geographic area and diets may differ based on 234 geographic location. On the other hand, our study population included a racially and 235 ethnically diverse population which may increase the generalizability of our findings. 236 Further studies in different geographic areas can be conducted to confirm our findings 237

A major strength of our study is that, to our knowledge, it is the first to characterize and compare the FODMAP intake of children with DGBI and HC. Also, the two populations were well-matched in terms of demographics, increasing the rigor of the findings. Additionally, the dietitians participating in the study had experience with analyzing food records for high FODMAP foods and maintained high quality in their review, including follow-up with families, if needed, regarding specific food record entries. In summary, both children with DGBI and HC consume a significant amount of high

FODMAP foods, though children with DGBI consume relatively less.. Fructans, lactose, and fructose were the most consumed FODMAP carbohydrates via wheat-containing

- foods, dairy, and fruits/100% fruit juices. Targeted and less restrictive diets such as the
- FGD, with modification based on our findings, may effectively restrict a large majority of
- the high FODMAP foods consumed by children with DGBI.

It is made available under a CC-BY 4.0 International license .

250 **REFERENCES**

- Thapar N, Benninga MA, Crowell MD, Di Lorenzo C, Mack I, Nurko S, et al. Paediatric
 functional abdominal pain disorders. Nat Rev Dis Primers. 2020;6(1):89.
- 253 2. Algera J, Colomier E, Simren M. The Dietary Management of Patients with Irritable
- Bowel Syndrome: A Narrative Review of the Existing and Emerging Evidence.
 Nutrients. 2019;11(9).
- Barrett JS, Gearry RB, Muir JG, Irving PM, Rose R, Rosella O, et al. Dietary poorly
 absorbed, short-chain carbohydrates increase delivery of water and fermentable
 substrates to the proximal colon. Aliment Pharmacol Ther. 2010;31(8):874-82.
- Shepherd SJ, Lomer MC, Gibson PR. Short-chain carbohydrates and functional
 gastrointestinal disorders. Am J Gastroenterol. 2013;108(5):707-17.
- Chumpitazi BP, Cope JL, Hollister EB, Tsai CM, McMeans AR, Luna RA, et al.
 Randomised clinical trial: gut microbiome biomarkers are associated with clinical
 response to a low FODMAP diet in children with the irritable bowel syndrome. Aliment
 Pharmacol Ther. 2015;42(4):418-27.
- Khalighi Sikaroudi M, Soltani S, Ghoreishy SM, Ebrahimi Z, Shidfar F, Dehnad A.
 Effects of a low FODMAP diet on the symptom management of patients with irritable
 bowel syndrome: a systematic umbrella review with the meta-analysis of clinical
 trials. Food Funct. 2024;15(10):5195-208.
- Jent S, Bez NS, Haddad J, Catalano L, Egger KS, Raia M, et al. The efficacy and real-world effectiveness of a diet low in fermentable oligo-, di-, monosaccharides and polyols in irritable bowel syndrome: A systematic review and meta-analysis. Clinical nutrition. 2024;43(6):1551-62.

It is made available under a CC-BY 4.0 International license .

- 8. Narayana V, McMeans AR, Levy RL, Shulman RJ, Chumpitazi BP. Children with
- functional abdominal pain disorders successfully decrease FODMAP food intake on
- a low FODMAP diet with modest improvements in nutritional intake and diet quality.
- 276 Neurogastroenterol Motil. 2022;34(10):e14392.
- 9. Tenenbaum RB, Czyzewski D, McMeans A, Narayana V, Chumpitazi BP, Levy RL,
- et al. Factors Associated With Adherence to a Low Fermentable Carbohydrate Diet
- in Children With Functional Gastrointestinal Disorders. Journal of the Academy of
 Nutrition and Dietetics. 2024;124(6):757-62.
- 10. Maagaard L, Ankersen DV, Vegh Z, Burisch J, Jensen L, Pedersen N, et al. Follow-
- up of patients with functional bowel symptoms treated with a low FODMAP diet. World
 J Gastroenterol. 2016;22(15):4009-19.
- 11. Frieling T, Heise J, Krummen B, Hundorf C, Kalde S. Tolerability of FODMAP reduced diet in irritable bowel syndrome efficacy, adherence, and body weight
 course. Z Gastroenterol. 2019;57(6):740-4.
- Halmos EP, Christophersen CT, Bird AR, Shepherd SJ, Gibson PR, Muir JG. Diets
 that differ in their FODMAP content alter the colonic luminal microenvironment. Gut.
 2015;64(1):93-100.
- Mitchell H, Porter J, Gibson PR, Barrett J, Garg M. Review article: implementation of
 a diet low in FODMAPs for patients with irritable bowel syndrome-directions for future
 research. Aliment Pharmacol Ther. 2019;49(2):124-39.
- 14. Rhys-Jones D, Varney JE, Muir JG, Gibson PR, Halmos EP. Application of The
 FODMAP Diet in a Paediatric Setting. Nutrients. 2022;14(20).

It is made available under a CC-BY 4.0 International license .

- 15. Halmos EP, Gibson PR. Controversies and reality of the FODMAP diet for patients
 with irritable bowel syndrome. J Gastroenterol Hepatol. 2019;34(7):1134-42.
- 16. Hyams JS, Di Lorenzo C, Saps M, Shulman RJ, Staiano A, van Tilburg M. Functional
- 298 Disorders: Children and Adolescents. Gastroenterology. 2016;150:1456–68.
- 17. Feskanich D, Sielaff BH, Chong K, Buzzard IM. Computerized collection and analysis
- of dietary intake information. Comput Methods Programs Biomed. 1989;30(1):47-57.
- 18. Czyzewski DI, Lane MM, Weidler EM, Williams AE, Swank PR, Shulman RJ. The
 interpretation of Rome III criteria and method of assessment affect the irritable bowel
- 303 syndrome classification of children. Aliment Pharmacol Ther. 2011;33(3):403-11.
- 304 19. Self MM, Czyzewski DI, Chumpitazi BP, Weidler EM, Shulman RJ. Subtypes of
 305 irritable bowel syndrome in children and adolescents. Clin Gastroenterol Hepatol.
 306 2014;12(9):1468-73.
- 307 20. [Available from: <u>https://www.monashfodmap.com/ibs-central/i-have-ibs/get-the-</u> 308 app/.
- Saps M, Velasco-Benitez CA, Velasco-Suarez DA, Alvarez-Baumgartner M, Balda
 AN, Arrizabalo S. Gastrointestinal Symptoms and Fermentable Oligosaccharides,
 Disaccharides, Monosaccharides, and Polyols Sources in Schoolchildren-A Pilot
 Study. Children (Basel). 2024;11(6).
- 22. Chumpitazi BP, Weidler EM, Lu DY, Tsai CM, Shulman RJ. Self-Perceived Food
 Intolerances Are Common and Associated with Clinical Severity in Childhood Irritable
 Bowel Syndrome. Journal of the Academy of Nutrition and Dietetics.
 2016;116(9):1458-64.

It is made available under a CC-BY 4.0 International license .

- Moshfegh AJ, Friday JE, Goldman JP, Ahuja JK. Presence of inulin and oligofructose
 in the diets of Americans. J Nutr. 1999;129(7 Suppl):1407S-11S.
- Shepherd SJ, Parker FC, Muir JG, Gibson PR. Dietary triggers of abdominal
 symptoms in patients with irritable bowel syndrome: randomized placebo-controlled
- evidence. Clin Gastroenterol Hepatol. 2008;6(7):765-71.
- 25. Yang J, Deng Y, Chu H, Cong Y, Zhao J, Pohl D, et al. Prevalence and presentation
- 323 of lactose intolerance and effects on dairy product intake in healthy subjects and
- patients with irritable bowel syndrome. Clin Gastroenterol Hepatol. 2013;11(3):262-8
- 325 e1.
- 26. Crawford PB, Obarzanek E, Morrison J, Sabry ZI. Comparative advantage of 3-day
- food records over 24-hour recall and 5-day food frequency validated by observation
- of 9- and 10-year-old girls. J Am Diet Assoc. 1994;94(6):626-30.