1	Dominant harmonic pattern as an ictal marker of the epileptogenic zone in				
2	focal neocortical epilepsy				
3	Lingqi Ye <sup>1, 2, 3</sup> , Lingli Hu <sup>1</sup> , Hongyi Ye <sup>1</sup> , Yihe Chen <sup>1</sup> , Junming Zhu <sup>2, 4</sup> , Zhe Zhe				
4	Hongjie Jiang <sup>4</sup> , Dongping Yang <sup>5</sup> , Cong Chen <sup>1</sup> , Shan Wang <sup>1</sup> , Zhongjin Wang <sup>1</sup> , We				
5	Ming <sup>1</sup> , Yi Wang <sup>6</sup> , Cenglin Xu <sup>6</sup> , Jin Wang <sup>3</sup> , Meiping Ding <sup>1</sup> , Shuang Wang <sup>1, 2</sup> *				
6					
7	1. Department of Neurology and Epilepsy Center, Second Affiliated Hospital, School				
8	of Medicine, Zhejiang University, Hangzhou, China				
9	2. Nanhu Brain-computer Interface Institute, Hangzhou, China				
10	3. Department of Neurology, Sir Run Run Shaw Hospital, School of Medicine,				
11	Zhejiang University, Hangzhou, China				
12	4. Department of Neurosurgery, Second Affiliated Hospital, School of Medicine,				
13	Zhejiang University, Hangzhou, China				
14	5. Research Center for Augmented Intelligence, Research Institute of Artificial				
15	Intelligence, Zhejiang Lab, Hangzhou, China				
16	6. Key Laboratory of Neuropharmacology and Translational Medicine of Zhejiang				
17	Province, School of Pharmaceutical Sciences, Zhejiang Chinese Medical				
18	University, Hangzhou, China				
19					
20	*Correspondence to :				
21	Shuang Wang <sup>*</sup> , Department of Neurology, Second Affiliated Hospital, School of				
22	Medicine, Zhejiang University, Hangzhou 310009, China; Tel: +86-571-87783872;				
23	E-mail: wangs77@zju.edu.cn				

### 24 Abstract

**Objective:** The ictal *Harmonic* pattern (*H* pattern), produced by the non-linear characteristics of EEG waveforms, may hold significant potential for localizing the epileptogenic zone (EZ) in focal epilepsy. However, further validation is needed to establish the *H* pattern's effectiveness as a biomarker for measuring the EZ.

29 Methods: We retrospectively enrolled 131 patients diagnosed with drug-resistant 30 focal epilepsy, all of whom had complete stereo-electroencephalographic (SEEG) data. 31 From this cohort, we selected 85 patients for outcome analysis. We analyzed the 32 morphological and time-frequency (TF) features of the H pattern using TF plots. A 33 third quartile (Q3) threshold was applied to classify channels expressing either 34 dominant (*Channel<sub>dH pattern</sub>*) or non-dominant H patterns (*Channel<sub>non-dH pattern</sub>*). We 35 then examined associations between the morphological features of the H pattern and patients' clinical characteristics, as well as the correlations between the extent of 36 37 channel removal and seizure outcomes.

38 **Results:** We found no significant correlations between the morphological features of the ictal H pattern and clinical factors, including lesional MRI findings, epileptic 39 40 onset patterns, epilepsy type, pathology, or surgical outcomes. The non-localizable H41 pattern appeared exclusively in patients with non-focal onset patterns. Notably, the 42 proportion of *Channel<sub>dH pattern</sub>* was higher in the seizure-onset zone (SOZ) compared 43 to the early propagation zone. The seizure-free group demonstrated significantly higher removal proportions of  $Channel_{dH pattern}$ , both within and outside the SOZ (p =44 45 0.014; p = 0.036), with AUCs of 0.606 and 0.660, respectively, in a seizure freedom 46 prediction model. Survival analysis confirmed that complete removal of these regions 47 correlated with long-term seizure freedom (p = 0.008; p = 0.028). Further subgroup 48 analysis showed a significant correlation in neocortical epilepsy (p = 0.0004; p =0.011), but not in mesial temporal lobe epilepsy. Additionally, multivariate analysis 49 50 identified the complete removal of *Channel*<sub>dH pattern</sub> as the only independent predictor for seizure freedom (*p* = 0.022; OR 6.035, 95% CI 1.291-28.211). 51

52 **Conclusions:** Our study supports the notion that the dominance of the ictal *H* pattern,

- 53 regardless of its morphology, serves as a novel biomarker for the EZ in focal epilepsy.
- 54 The non-linearity in EEG waveforms provides new insights into understanding ictal
- 55 spreading propagation and offers potential improvements for surgical planning in
- 56 neocortical epilepsy.
- 57
- 58 Keywords: harmonic pattern; focal epilepsy; epileptogenic zone; epilepsy surgery;
- 59 seizure outcome

### 60 Introduction

61 Resective surgery currently remains the most effective treatment for drug-resistant 62 epilepsy (DRE). However, only 50% of patients achieve prolonged seizure freedom 63 following surgery. Despite advancements in intracranial electroencephalography 64 (iEEG) technology, it has yet to surpass a 70% ceiling in achieving favorable surgical outcomes <sup>1-3</sup>. The success of the surgery largely depends on the precise localization 65 and complete removal of the epileptogenic zone (EZ), which is defined as the brain 66 area essential for generating seizures <sup>4</sup>. Postsurgical seizure freedom indicates that the 67 EZ was likely successfully removed. Although the seizure-onset zone (SOZ) does not 68 necessarily equate to the EZ<sup>5,6</sup>, experienced clinicians' visual identification of the 69 70 SOZ is a crucial step in the process of localizing the EZ.

71 Many studies have aimed to identify ictal EEG biomarkers capable of quantitatively 72 delineating the EZ. Established semi-quantitative algorithms, such as the epileptogenic index (EI)  $^{7}$  and epileptogenic map  $^{8}$ , have been shown to be effective in 73 74 cases where ictal onset patterns include low-voltage fast activity. Additionally, the 75 removal of areas exhibiting ictal high-frequency oscillations (HFOs) has been reported as a prognostic indicator of favorable seizure outcomes <sup>9-11</sup>. The concept of 76 77 the EEG *fingerprint*, proposed as an ictal biomarker of the EZ, suggests a possible synchronous activation of GABAergic neurons at seizure onset <sup>12</sup>. Similarly, brain 78 *chirp*<sup>13</sup> highlights the importance of fast EEG activity in defining the EZ. 79

80 In our previous study, we identified the *Harmonic* pattern (*H* pattern), a spectral 81 feature of specific ictal EEG components that contains valuable information for 82 seizure propagation and EZ localization. The H pattern is characterized by an 83 equidistant spectral band distribution of varying frequencies on time-frequency (TF) 84 plots. Generated by the specific non-sinusoidal waveforms, the ictal H pattern may 85 represent periodically synchronized neural firings, rather than a methodological artifact. Furthermore, the H pattern is frequently observed in seizures with various 86 onset patterns and across different seizure stages  $^{14}$ . The dominance of the ictal H 87

pattern in different brain regions reflects higher non-linearity and has proven useful inlocalizing the EZ.

90 Despite its potential, the full spectrum of the ictal H pattern remains unclear. The H91 pattern shows significant variability in morphology across patients; however, within 92 an individual patient, its morphology and spatial localization tend to remain consistent 93 across seizures, emphasizing the personalized nature of ictal neural dynamics. In this 94 expanded cohort, we first investigated clinical factors related to the morphological 95 properties of the *H* pattern, specifically focusing on its clarity and shape. High clarity 96 indicates a narrower frequency band with reduced background noise, while the shape 97 of the spectral bands reflects changes in the frequency of ictal neural oscillations at a 98 macroscopic level. This study aims to further support the hypothesis that the 99 dominance of the ictal H pattern could serve as a novel biomarker for the EZ. 100 Additionally, we analyzed the potential impact of the H pattern's morphological 101 features.

102

### 103 Methods

### 104 Selection of patient cohort

105 We retrospectively reviewed 133 patients with drug-resistant epilepsy who had 106 completed stereo-electroencephalography (SEEG) at the Second Affiliated Hospital 107 Zhejiang University School of Medicine (SAHZU) between April 2013 and October 108 2022. We excluded two patients diagnosed with hypothalamic hamartoma from the 109 study. Therefore, we performed H pattern analysis on the remaining 131 cases. From 110 this cohort, we selected 85 patients for outcome analysis based on the following 111 inclusion criteria (Figure S1): (1) a focal-onset pattern on SEEG with a clearly 112 defined SOZ localized to a discrete cortical region; (2) undergoing either resective 113 surgery or thermocoagulation treatment; (3) recording at least two habitual clinical 114 seizures during SEEG monitoring; (4) identifying a discernible and localizable ictal H115 pattern (as described in the *Identification of ictal H pattern* section); (5) a follow-up

period of at least 12 months after surgery. The Medical Ethics Committee of SAHZU
approved this study (Study No. 2023-1260, 2021-0585). We obtained written
informed consent from each participant.

119

### 120 SEEG evaluation

121 All patients were recommended for SEEG evaluation during a multidisciplinary 122 patient management conference. The non-invasive presurgical evaluation included a 123 detailed medical history, neuropsychological assessment, scalp video-EEG monitoring, 3.0-Tesla cerebral MRI, and interictal <sup>18</sup>F-fluorodeoxyglucose positron 124 emission tomography (<sup>18</sup>F-FDG-PET). The procedure for SEEG electrode 125 implantation has been previously described <sup>15</sup>. Based on the working hypothesis, 126 127 multiple-contact intracerebral electrodes (8-16 contacts, each 2 mm in length, 0.8 mm 128 in diameter, and spaced 1.5 mm apart) were implanted using a robotic arm-assisted 129 system. SEEG signals were recorded at sampling rates of either 1,000 or 2,000 Hz 130 using EEG systems with 256 channels (Nihon Kohden) or 128 channels (Xltek, 131 Natus). The anatomical locations of each contact along the electrode trajectory in all 132 subjects were confirmed by co-registering post-implantation CT scans with pre-implantation MRI using Sinovation 2.0<sup>16</sup>. A three-dimensional schematic 133 134 diagram of the reconstruction is provided in the flowchart (Figure 1C).

135

### 136 Surgical outcome

Following SEEG evaluation, patients underwent either resective surgery or laser ablation. The planning of these surgical interventions was conducted independently of this study. Each contact was classified as either *removed* or *non-removed* based on postsurgical MRI evaluations. Surgical outcomes were assessed at the most recent follow-up and patients were categorized as either seizure-free (SF, Engel class Ia) or not seizure-free (NSF, >Engel class Ia) <sup>17</sup>.

143

### 144 Visual EEG analysis

Two experienced and board-certified neurophysiologists reviewed the SEEG data to identify the SOZ and the rapid propagation zone (PZ) using our previously established methods <sup>15, 18</sup>. The SOZ is defined as the brain area that shows the earliest changes on SEEG. The PZ is characterized by the rapid spread of ictal activity within 3-5 seconds after seizure onset. Areas outside these defined zones were classified as 'other regions.' Both the PZ and these 'other regions' were collectively categorized as non-SOZ.

152

## 153 Identification of ictal H pattern

154 We selected only contacts within cortical gray matter for quantitative analysis. We 155 analyzed all EEG data using a bipolar montage. For each identified seizure, we 156 extracted a 110-second segment of SEEG data: 10 seconds before and 100 seconds 157 after the EEG onset. Additionally, we collected a segment of preictal EEG data 158 ranging from -230 to -120 seconds before EEG onset as baseline data for 159 normalization. To enhance the reliability of our results, we applied two analytical methods: the Morlet wavelet transform<sup>19</sup> and the multitaper method <sup>20, 21</sup> in the TF 160 161 analysis.

162 We identified the ictal H pattern from TF maps based on the following criteria: the 163 presence of two or more distinct high-intensity bands with varying frequencies 164 occurring at any point during a seizure, with equal frequency intervals between 165 adjacent bands (Figure 1A). For each patient, we analyzed only the initially occurring 166 H pattern. We estimated various features of the ictal H pattern, including its shape, 167 clarity, onset and end times, fundamental and maximal frequencies, frequency 168 intervals, and the number of bands. If the H pattern was consistently stereotyped 169 across different seizures for a single patient, we included only one data set in the 170 analysis. To determine the dominance of the H pattern, we established a threshold

171 known as Q3, which represents the upper quartile (75th percentile) of the data 172 distribution for the maximal count of frequency bands per patient. We calculated Q3173 using the following formula:

174 Q3 = (n+1) \* 3/4,

where *n* denotes the highest count of frequency bands observed across all channels. If 175 176 the number of bands in a channel showing the ictal H pattern (*Channel<sub>H pattern</sub>*) 177 exceeds the Q3 threshold, we marked it as exhibiting a dominant H pattern 178 (Channel<sub>dH pattern</sub>). Conversely, a channel not meeting this threshold was classified as 179 expressing a non-dominant H pattern (Channel<sub>non-dH pattern</sub>) (Figure 1D). A localizable 180 H pattern should be consistent across seizures and present in specific regions. If no 181 gradient in band numbers was observed across all channels, the H pattern was 182 classified as non-localizable.

183

### 184 Morphology of Ictal Harmonic Pattern

185 We classified the clarity of the *H* pattern into two levels. Grade A indicated a highly 186 clear frequency band with distinct borders against the background activity. Grade B 187 referred to a less clear frequency band with fuzzier borders. We identified three 188 distinct shapes of the *H* pattern: Shape 1 resembled a parabola, where the fundamental 189 frequency initially increases and then decreases; Shape 2 appeared as a declining line, 190 representing a decrease in the frequency harmonics of the spectral band; and Shape 3 191 took the form of an ascending line, indicating an increase in the frequency harmonics 192 of the spectral band. This resulted in a total of six categories: A1, A2, A3, B1, B2, and 193 B3 (Figure S2). Initially, two investigators (Ye L.Q. and Hu L.L.) performed these 194 classifications. In cases of disagreement, a third epileptologist (Wang S.) made the 195 final decision.

196

### 197 Prediction of Surgical Outcome

198 For each patient, we categorized channels as either SOZ or non-SOZ based on visual

199 analysis, and as *Channel*<sub>dH pattern</sub> or *Channel*<sub>non-dH pattern</sub> based on TF analysis. We

200 determined the ratio of removed channels using the following formula:

 $Removal ratio(\%) = \frac{\#Channel_{removed}}{\#Channel_{removed}} \times 100\%.$ 201 To estimate the correlation between the removal of *Channel\_{non-removed*} and seizure
202 freedom, we used the *Chi-Squared* test and receiver operating characteristic (ROC)
203 curve analysis, from which we derived the area under the curve (AUC). We then
204 generated Kaplan-Meier (KM) curves to depict the cumulative probability of
205 achieving seizure freedom. We compared the retention of seizure freedom between
206 groups using the log-rank test, including patients with complete (100%) removal and
207 those with incomplete or no removal (0-100%).

208

### 209 Statistical Analysis

We used MATLAB 2021a (The MathWorks, Inc.) and SPSS 26.0 for statistical 210 211 analysis. We evaluated the normality of the distribution of all continuous variables 212 using the Kolmogorov-Smirnov test. Normally distributed variables were presented as 213 mean  $\pm$  standard deviation (SD), while non-normally distributed variables were 214 reported as median (interquartile range, IQR). We expressed categorical variables as 215 frequencies. We applied an *unpaired t*-test or a nonparametric *Mann-Whitney U* test 216 to analyze continuous variables, and we assessed categorical variables using the 217 Chi-Squared test. We also conducted a binary regression analysis to identify 218 predictive factors affecting surgical outcomes. For multiple comparisons, we 219 employed the Bonferroni correction method. p < 0.05 represents statistically 220 significant.

221

222 **Results** 

### 223 *Patient Demographics*

In the entire cohort, 75.6% (99 patients) exhibited a focal-onset pattern, while the

225 remaining 24.4% (32 patients) had a non-focal onset pattern (**Table S1**). Among those 226 with a focal-onset pattern, 92.9% (92 patients) displayed an ictal H pattern, all of 227 which were localizable. In contrast, 93.8% (30 patients) of those with a non-focal 228 onset pattern exhibited at least one ictal H pattern. Of these, 80.0% (24 patients) were 229 localizable, and 20.0% (6 patients) were non-localizable. Most patients presented with 230 only one type of H pattern. However, two different types of H pattern were observed 231 exclusively in approximately 20% of patients with a non-focal onset pattern (Table 232 **S1**).

233 Of the 85 patients included in the outcome analysis, 38.9% (33 patients) had participated in our previous H pattern study (**Table 1**)<sup>14</sup>. In the current cohort, 61.2% 234 235 (52 patients) achieved seizure freedom after an average follow-up period of 47.8 236 months. We found no significant differences between the SF and NSF groups 237 regarding age at disease onset, age at SEEG recording, or the 238 lateralization/localization of seizures. The incidence of detectable abnormalities on 239 MRI was similar between the two groups (75.0% in the SF group and 72.7% in the 240 NSF group, p = 0.816). However, the NSF group had a significantly higher number of 241 implanted electrodes (p = 0.025). Focal cortical dysplasia (FCD)  $\Box$  was the most 242 common pathological finding, occurring in 34.1% of patients. Among those with 243 malformations of cortical development (MCD) other than FCD  $\square$  (19.5% of patients), 244 11 had FCD I, 3 had tuberous sclerosis complex, 1 had FCD  $\Box$  c, and 1 had a mild 245 malformation of cortical development with oligodendroglial hyperplasia and epilepsy. 246 Patients with FCD  $\Box$  a were included in the hippocampal sclerosis (HS) group. We 247 observed a significant difference in the pathological findings between the SF and NSF 248 groups (p = 0.014), particularly between FCD  $\square$  and gliosis/non-specific findings.

249

### 250 Morphology, Distribution and Parameters of Ictal Harmonic Pattern

We summarized the morphological characteristics of the ictal *H* pattern in patients who exhibited only one type of *H* pattern in **Table S2**. The most frequently observed

shapes were the declining line (57.3%) and the parabola (40.2%) (**Figure S2**). Our analysis revealed no correlation between the clarity or shape of the *H* pattern and clinical factors, including the presence of lesional MRI findings, focal-onset pattern, neocortical epilepsy, pathology, or surgical outcome.

257 Out of 85 patients with focal-onset patterns, we identified both the SOZ and the PZ in 258 65 patients. Within these zones, the proportion of channels exhibiting the ictal H259 pattern was significantly higher in the SOZ (100.0%, IQR 57.3-100%) compared to 260 the PZ (26.3%, IQR 0-50.0%; p < 0.0001) (Figure 2A). Similarly, the ictal dH pattern 261 appeared more frequently in the SOZ (38.5%, IQR 16.0-80.0%) than in the PZ (0%, 262 IQR 0-5.8%; p < 0.0001) (Figure 2A). The ictal H pattern was less common in other 263 regions, and we did not observe a significant difference in the distribution of the 264 non-dH pattern between the SOZ and PZ. Among the 65 patients, 37 exhibited the ictal H pattern simultaneously in both the SOZ and PZ. The number of frequency 265 266 bands of the ictal H pattern was notably higher in the SOZ (3.1, IQR 2.2-6.2)267 compared to the PZ (2.5, IQR 1.9-4.2; p < 0.0001), as was the maximal frequency 268 (SOZ, 118.8, IQR 92.5-333.0 Hz; PZ, 90.0, IQR 66.3-112.6 Hz; p < 0.0001). The 269 onset time of the ictal H pattern did not differ between the SOZ and PZ, indicating 270 almost simultaneous initiation across these regions. Furthermore, we did not observe 271 differences in the termination, fundamental frequency, or frequency interval of the 272 ictal *H* pattern between the SOZ and PZ (Figure 2B-D). These parameters of the ictal 273 *H* pattern also did not differ between the SF and NSF groups (Figure 3B-E). In the 274 remaining 20 patients without a PZ, the dH pattern was predominantly expressed in 275 88.9% of channels within the SOZ (IQR, 54.6-88.9%) and was rarely observed in 276 other regions.

277

### 278 Ictal Dominant Harmonic Pattern

Figure 3A shows that a higher removal ratio of the *Channel*<sub>dH pattern</sub> significantly correlates with achieving seizure freedom (p = 0.014). The SF group demonstrated a

higher removal ratio of the *Channel*<sub>dH pattern</sub> within non-SOZ (100.0, IQR 100-100%)

282 compared to the NSF group (100.0, IQR 0-100%; p = 0.036).

283

### 284 Predictive Value of Removing Ictal Dominant Harmonic Pattern

285 Regression analysis identified several independent factors significantly associated with achieving seizure freedom. These factors include the complete removal of the 286 287 *Channel*<sub>dH pattern</sub> (p = 0.022; OR 6.035, 95% CI 1.291-28.211), FCD  $\Box$  or other MCD 288 types (as opposed to gliosis or other non-specific pathologies), and a smaller number 289 of implanted electrodes (Figure 4). Additionally, the association between the complete removal of the Channel<sub>dH pattern</sub> and seizure freedom was supported by an 290 291 analysis showing statistical significance (p = 0.031, AUC = 0.606) (Figure 5). The 292 highest AUC value of 0.660 was recorded when considering the complete removal of 293 the Channel<sub>dH pattern</sub> within non-SOZ, highlighting the importance of the ictal dH294 pattern for localizing the EZ (Figure S3).

295

### 296 Survival Analysis

297 Figure 6 and Figure S4 illustrate the 10-year prognosis of seizure freedom for 298 patients who underwent complete removal (100%) compared to those with incomplete 299 or no removal (0~100%). Among the 85 patients in the cohort, 58.2% achieved 300 seizure freedom at the 10-year mark. Notably, patients who underwent complete 301 removal of the *Channel<sub>dH pattern</sub>* had a significantly higher probability of attaining 302 long-term seizure freedom (p = 0.008). Specifically, among the 37 patients with 303 complete removal of the *Channel*<sub>dH pattern</sub> within non-SOZ, there was also an increased 304 likelihood of achieving seizure freedom (p = 0.028). In cases of neocortical epilepsy, 305 the complete removal of the Channel<sub>dH pattern</sub> was strongly associated with long-term 306 seizure freedom (p=0.0004), regardless of whether the removal occurred within the 307 SOZ(p = 0.013) or non-SOZ (p = 0.011) (Figure S5). In contrast, for mesial temporal 308 lobe epilepsy (mTLE), complete removal of the SOZ, rather than the *Channel<sub>dH pattern</sub>*,

309 was significantly associated with better long-term seizure freedom outcomes (p =310

0.019).

311

### 312 Discussion

313 The ictal *H* pattern, which reflects the non-linear features of ictal SEEG signals on TF 314 maps, may serve as an innovative ictal biomarker and offer a new approach for 315 delineating the boundary of the EZ. Our key findings reveal: (1) In patients with focal 316 onset patterns, the morphology of the ictal H pattern is consistent within individuals 317 but heterogeneous across the cohort; (2) the morphological features of the ictal H318 pattern do not correlate with clinical variables; (3) the dominance of the ictal H319 pattern indicates high epileptogenicity, and removing regions that exhibit this pattern 320 is crucial for achieving seizure freedom in focal epilepsy, particularly in neocortical 321 epilepsy.

322 Previous studies have introduced various quantitative biomarkers for measuring spectral components of SEEG signals, including EI<sup>7, 22</sup>, HFOs<sup>23</sup>, and ictal non-linear 323 metrics such as the  $h^2$  coefficient <sup>24, 25</sup>, coherence analysis <sup>26, 27</sup>, entropy <sup>28, 29</sup>, and 324 infraslow activity <sup>30-32</sup>. These computational EEG markers aim to enhance the 325 326 precision of EZ delineation. Our findings suggest that the ictal dH pattern could be a 327 valuable biomarker for localizing the core epileptogenic cortex across different brain 328 regions. Visual inspection showed that the distribution of the ictal H pattern aligned 329 with the ictal onset zone, regardless of whether it was focal-, multifocal-, or 330 diffuse-onset. In individuals with focal-onset patterns, the ictal H pattern typically 331 exhibited a consistent spatial distribution across various seizures. In contrast, 332 individuals with non-focal onset patterns displayed variable types of the ictal H333 pattern, indicating an unstable seizure propagation pattern. The near-simultaneous 334 onset of the ictal H pattern across different regions, along with a uniform fundamental 335 frequency, suggests clear inter-regional synchronization during this period. Notably, 336 our findings indicate that regions showing the dH pattern can extend into the PZ 337 beyond the boundary of the SOZ. As our previous research indicated, the H pattern

reflects spectral skewness or asymmetry in EEG waveforms, and the dH pattern represents stronger inter-regional synchronization during seizure evolution, particularly in focal epilepsy. These findings further support the utility of the dHpattern as a non-linear feature in delineating the EZ. Future clinical practices in defining the EZ would benefit from incorporating an integrated EEG biomarker that includes analysis of the dH pattern.

344 We analyzed the morphological features of the *H* pattern in terms of shape and clarity. 345 The shape of the spectral bands in the *H* pattern reflects variations in the frequency of 346 synchronized neural oscillations. The most frequently observed shape is a declining line, which closely resembles a spectral signature known as *chirps* <sup>33-35</sup>. This shape 347 348 may indicate a decline in neural firing, possibly due to increased inhibition. In our 349 cohort, nearly half of the SOPs exhibiting low-frequency activity (LVFA) expressed *chirp*<sup>13</sup> patterns, with proportions similar to those of the parabolic shape. We propose 350 351 that some H patterns with a parabolic shape and very short upward lines might be 352 classified as chirps due to the limitations of visual inspection. This potential 353 misclassification could narrow the scope of the ictal H pattern and limit its 354 applicability. A corticothalamic computational model has similarly elucidated how 355 changes in physiological parameters affect the dynamics of tonic-clonic seizures, 356 similar to the H pattern. This model revealed that increasing the connection strength enhances the power, count, and duration of harmonic bands <sup>36</sup>. Conversely, the clarity 357 358 of the H pattern signifies the distinctness of the harmonic signals against the 359 background. A highly clear H pattern is characterized by a narrow frequency band 360 with minimal noise. Interestingly, these morphological properties do not correlate 361 with clinical factors such as seizure-onset pattern, seizure origin, MRI findings, 362 pathological findings, or seizure outcome. The primary clinically relevant aspect of 363 the *H* pattern is its degree of non-linearity. Regardless of its shape, the prominence of 364 the *H* pattern indicates epileptogenicity, and its complete removal has been linked to 365 the potential achievement of seizure freedom.

366 We observed a stronger association between the dH pattern and long-term seizure

367 freedom in neocortical epilepsy compared to mTLE. Previous studies have reported that patients with mTLE generally experience more favorable surgical outcomes than 368 those with neocortical epilepsy <sup>37</sup>. Additionally, the classification of seizure outcomes 369 370 can vary among patients. Therefore, we used Engel class Ia as the criterion for a 371 favorable outcome to evaluate the efficacy of the dH pattern in localizing the EZ. 372 Earlier iEEG studies have indicated that the extent of SOZ removal does not always correlate with seizure freedom across all patients <sup>38, 39</sup>. In contrast, we found that, for 373 374 mTLE, the complete removal of the SOZ had a stronger correlation with long-term 375 seizure freedom than the dH pattern. Furthermore, our current mTLE cohort included more MRI-negative cases compared to our previous cohort <sup>14</sup>. Surgical outcomes in 376 MRI-negative cases are usually less favorable compared to those in TLE-HS<sup>40, 41</sup>, 377 378 suggesting a more extensive epileptogenic network in MRI-negative TLE. Previous 379 research has highlighted differences in the involved neural networks between mTLE and neocortical epilepsy. For instance, an <sup>18</sup>F-FDG-PET study revealed distinct 380 381 metabolic profiles between these two types of epilepsy, suggesting variations in seizure propagation <sup>42</sup>. Additionally, an iEEG study showed that the interictal 382 383 epileptogenic network in mTLE exhibits a more regular pattern of graph connectivity compared to neocortical epilepsy <sup>43</sup>. It is well-recognized that epileptic networks 384 385 differ between mTLE and neocortical temporal lobe epilepsy, as well as between TLE and extra-TLE<sup>44, 45</sup>. A longer follow-up period is necessary to assess the final 386 387 prognosis in patients with mTLE and to further estimate the value of the H pattern in 388 this population.

Nevertheless, our study has several limitations. First, we based the determination of the *H* pattern's dominance solely on visual interpretation throughout the study. Future investigations should focus on optimizing the definition of the d*H* pattern and developing a more automated algorithm to quantify the *H* pattern. This would enhance the precision and objectivity of the analysis. We plan to create a quantitative index for the *H* pattern to address this issue. Additionally, we did not analyze the occurrence of the second *H* pattern during the later ictal stage of one seizure in this

396 study, which warrants further investigation.

397

### 398 Conclusions

In focal epilepsy, the dominance of the ictal *H* pattern offers new insights for localizing the EZ, regardless of its morphology. The distinct non-linear feature of the H pattern has significant implications for presurgical assessments and surgical planning. Applying this biomarker in neocortical epilepsy may offer greater value in achieving long-term favorable outcomes compared to mTLE.

404

## 405 Data availability

406 The that support the findings of this study are available upon reasonable request from407 the corresponding author.

## 408 Acknowledgements

This study received support from the National Natural Science Foundation of China
(grant number: 82171437, 82301636, 82471469) and the Major Program of the
National Natural Science Foundation of Zhejiang Province (grant number:
LD24H090003).

### 413 Disclosures

414 The authors have no conflicts of interest to disclose related to this study.

## 415 Supplementary material

416 Supplementary material is available online.

### 417 **References**

- 418 1. Cardinale F, Rizzi M, Vignati E, et al. Stereoelectroencephalography:
  419 retrospective analysis of 742 procedures in a single centre. Brain
  420 2019;142:2688-2704.
- 421 2. Jehi L, Morita-Sherman M, Love TE, et al. Comparative Effectiveness of
  422 Stereotactic Electroencephalography Versus Subdural Grids in Epilepsy Surgery. Ann
  423 Neurol 2021;90:927-939.
- 3. Sivaraju A, Hirsch L, Gaspard N, et al. Factors Predicting Outcome After
  Intracranial EEG Evaluation in Patients With Medically Refractory Epilepsy.
  Neurology 2022;99:e1-e10.
- 427 4. Vakharia VN, Duncan JS, Witt JA, Elger CE, Staba R, Engel J, Jr. Getting the
  428 best outcomes from epilepsy surgery. Ann Neurol 2018;83:676-690.
- 429 5. Rosenow F, Lüders H. Presurgical evaluation of epilepsy. Brain430 2001;124:1683-1700.
- 431 6. Widjaja E, Li B, Schinkel CD, et al. Cost-effectiveness of pediatric epilepsy
  432 surgery compared to medical treatment in children with intractable epilepsy. Epilepsy
  433 Res 2011;94:61-68.
- 434 7. Bartolomei F, Chauvel P, Wendling F. Epileptogenicity of brain structures in
  435 human temporal lobe epilepsy: a quantified study from intracerebral EEG. Brain
  436 2008;131:1818-1830.
- 437 8. David O, Blauwblomme T, Job AS, et al. Imaging the seizure onset zone with
  438 stereo-electroencephalography. Brain 2011;134:2898-2911.
- 9. Ochi A, Otsubo H, Donner EJ, et al. Dynamic changes of ictal high-frequency
  oscillations in neocortical epilepsy: using multiple band frequency analysis. Epilepsia
  2007;48:286-296.
- 442 10. Modur PN, Zhang S, Vitaz TW. Ictal high-frequency oscillations in neocortical
  443 epilepsy: implications for seizure localization and surgical resection. Epilepsia
  444 2011;52:1792-1801.
- 445 11. Fujiwara H, Greiner HM, Lee KH, et al. Resection of ictal high-frequency

- 446 oscillations leads to favorable surgical outcome in pediatric epilepsy. Epilepsia447 2012;53:1607-1617.
- 448 12. Grinenko O, Li J, Mosher JC, et al. A fingerprint of the epileptogenic zone in449 human epilepsies. Brain 2018;141:117-131.
- 450 13. Di Giacomo R, Burini A, Chiarello D, et al. Ictal fast activity chirps as markers of
- 451 the epileptogenic zone. Epilepsia 2024;65:e97-e103.
- 452 14. Lingli H, Lingqi Y, Hongyi Y, et al. Harmonic patterns embedding ictal EEG
- 453 signals in focal epilepsy: a new insight into the epileptogenic zone. medRxiv
  454 2024:2023.2012.2020.23300274.
- 455 15. Hu L, Xiong K, Ye L, et al. Ictal EEG desynchronization during low-voltage fast
- 456 activity for prediction of surgical outcomes in focal epilepsy. J Neurosurg 2022:1-10.
- 457 16. Tadel F, Baillet S, Mosher JC, Pantazis D, Leahy RM. Brainstorm: a user-friendly
- 458 application for MEG/EEG analysis. Comput Intell Neurosci 2011;2011:879716.
- 459 17. Engel J. Outcome with respect to epileptic seizures. 1993.
- 460 18. Wang S, So NK, Jin B, et al. Interictal ripples nested in epileptiform discharge
- 461 help to identify the epileptogenic zone in neocortical epilepsy. Clin Neurophysiol462 2017;128:945-951.
- 463 19. Bernardino A, Santos-Victor J. A Real-Time Gabor Primal Sketch for Visual Attention.
- 464 2005; Berlin, Heidelberg: Springer Berlin Heidelberg: 335-342.
- 20. Thomson DJ. Spectrum estimation and harmonic analysis. Proceedings of theIEEE 1982;70:1055-1096.
- 467 21. Babadi B, Brown EN. A review of multitaper spectral analysis. IEEE Trans
  468 Biomed Eng 2014;61:1555-1564.
- 469 22. Aubert S, Wendling F, Regis J, et al. Local and remote epileptogenicity in focal
- 470 cortical dysplasias and neurodevelopmental tumours. Brain 2009;132:3072-3086.
- 471 23. González Otárula KA, von Ellenrieder N, Cuello-Oderiz C, Dubeau F, Gotman J.
- 472 High-Frequency Oscillation Networks and Surgical Outcome in Adult Focal Epilepsy.
- 473 Ann Neurol 2019;85:485-494.
- 474 24. Bartolomei F, Lagarde S, Wendling F, et al. Defining epileptogenic networks:

475 Contribution of SEEG and signal analysis. Epilepsia 2017;58:1131-1147.

476 25. Hu L, Xiong K, Ye L, et al. Ictal EEG desynchronization during low-voltage fast

- 477 activity for prediction of surgical outcomes in focal epilepsy. J Neurosurg478 2023;139:238-247.
- 479 26. Li YH, Ye XL, Liu QQ, et al. Localization of epileptogenic zone based on graph
  480 analysis of stereo-EEG. Epilepsy Res 2016;128:149-157.
- 27. Ren Y, Cong F, Ristaniemi T, Wang Y, Li X, Zhang R. Transient seizure onset
  network for localization of epileptogenic zone: effective connectivity and graph
  theory-based analyses of ECoG data in temporal lobe epilepsy. J Neurol
  2019;266:844-859.
- 28. Travnicek V, Klimes P, Cimbalnik J, et al. Relative entropy is an easy-to-use
  invasive electroencephalographic biomarker of the epileptogenic zone. Epilepsia
  2023;64:962-972.
- 488 29. Bratu IF, Makhalova J, Garnier E, et al. Permutation entropy-derived parameters
  489 to estimate the epileptogenic zone network. Epilepsia 2023.
- 30. Rodin E, Modur P. Ictal intracranial infraslow EEG activity. Clin Neurophysiol2008;119:2188-2200.
- 492 31. Kanazawa K, Matsumoto R, Imamura H, et al. Intracranially recorded ictal direct
  493 current shifts may precede high frequency oscillations in human epilepsy. Clin
  494 Neurophysiol 2015;126:47-59.
- 32. Nakatani M, Inouchi M, Daifu-Kobayashi M, et al. Ictal direct current shifts
  contribute to defining the core ictal focus in epilepsy surgery. Brain Commun
  2022;4:fcac222.
- 33. Schiff SJ, Colella D, Jacyna GM, et al. Brain chirps: spectrographic signatures of
  epileptic seizures. Clin Neurophysiol 2000;111:953-958.
- 34. Breakspear M, Roberts JA, Terry JR, Rodrigues S, Mahant N, Robinson PA. A
  unifying explanation of primary generalized seizures through nonlinear brain
  modeling and bifurcation analysis. Cereb Cortex 2006;16:1296-1313.
- 503 35. Dutta S, Iyer KK, Vanhatalo S, Breakspear M, Roberts JA. Mechanisms
  504 underlying pathological cortical bursts during metabolic depletion. Nat Commun

505 2023;14:4792.

- 506 36. Deeba F, Sanz-Leon P, Robinson PA. Effects of physiological parameter
- 507 evolution on the dynamics of tonic-clonic seizures. PLoS One 2020;15:e0230510.
- 508 37. Spencer S, Huh L. Outcomes of epilepsy surgery in adults and children. Lancet
- 509 Neurol 2008;7:525-537.
- 510 38. Khan M, Chari A, Seunarine K, et al. Proportion of resected seizure onset zone
- 511 contacts in pediatric stereo-EEG-guided resective surgery does not correlate with
- 512 outcome. Clin Neurophysiol 2022;138:18-24.
- 513 39. Thomas J, Kahane P, Abdallah C, et al. A Subpopulation of Spikes Predicts
  514 Successful Epilepsy Surgery Outcome. Ann Neurol 2023;93:522-535.
- 40. Fong JS, Jehi L, Najm I, Prayson RA, Busch R, Bingaman W. Seizure outcome
- and its predictors after temporal lobe epilepsy surgery in patients with normal MRI.
- 517 Epilepsia 2011;52:1393-1401.
- 41. McIntosh AM, Wilson SJ, Berkovic SF. Seizure outcome after temporal
  lobectomy: current research practice and findings. Epilepsia 2001;42:1288-1307.
- 520 42. Zhu HY, Tang YX, Xiao L, et al. Metabolic profiles and correlation with surgical
- 521 outcomes in mesial versus neocortical temporal lobe epilepsy. CNS Neurosci Ther 522 2023;29:2656-2665.
- 43. Bartolomei F, Bettus G, Stam CJ, Guye M. Interictal network properties in mesial
- temporal lobe epilepsy: a graph theoretical study from intracerebral recordings. Clin
  Neurophysiol 2013;124:2345-2353.
- 44. González Otárula KA, Schuele S. Networks in Temporal Lobe Epilepsy.
  Neurosurg Clin N Am 2020;31:309-317.
- 45. Hall GR, Hutchings F, Horsley J, et al. Epileptogenic networks in extra temporal
  lobe epilepsy. Netw Neurosci 2023;7:1351-1362.
- 530

## 531 Figure Legends

### 532 Figure 1. Examples and characteristics of the *H* pattern (data from patient #057).

- 533 (A) Ictal SEEG recordings from both SOZ and PZ of a patient with right parietal focal
- 534 cortical dysplasia showing bursts of high-amplitude poly-spikes followed by
- 535 low-voltage fast activity at seizure onset. Below the SEEG recordings are
- 536 corresponding time-frequency plots of the two channels. An ictal *H* pattern could be
- 537 identified in both plots: two or more unique high-intensity spectral bands with varying
- 538 frequency starting at 4 s (red arrow) after 0 s (seizure onset), and ending at 19 s (red
- arrow). The frequencies of the bottom and top spectral bands are defined as the
- 540 *fundamental frequency* and *maximal frequency*, respectively (yellow arrow).
- 541 Importantly, the ictal *H* pattern exhibits a distinctive feature of equidistant distribution
- on the TF plots. For instance, at 6.3 s (long blue dotted arrows), the frequency interval
- 543 between adjacent spectral bands is nearly uniform, at approximately 26-28 Hz, as
- observed in the power spectral density diagrams. The ictal *H* pattern shown in the two
- 545 channels are classified as a 'dominant H pattern (dH pattern)' and a 'non-dominant H
- 546 pattern (non-dH pattern)', respectively. (B) 3D representation of an example SEEG
- 547 exploration. Electrode contacts are categorized into three regions: the SOZ (red dots),
- the PZ (orange dots), and the other regions (blue dots), according to a patient's ictal
- 549 SEEG recordings. (C) Total number of visually determined spectral bands on the TF
- 550 plot of each channel each time a seizure is counted. Channels are then ranked based
- on the number of bands, and if a channel's count exceeds the threshold of Q3, it is
- 552 marked as expressing the ictal dominant *H* pattern. SEEG,
- stereo-electroencephalography; SOZ, seizure-onset zone; PZ, early propagation zone;
- 554 *H* pattern, *Harmonic* pattern; *Q3*, the 75th percentile of data distribution.
- 555

### 556 Figure 2. Distribution and time-frequency features of ictal *H* patterns in SOZ

and PZ. (A) Proportions of channels expressing the ictal *H* pattern, the ictal dominant

- 558 *H* pattern, and the ictal non-dominant *H* pattern between SOZ (red dots) and PZ
- (orange dots) in 65 patients who had both SOZ and PZ. (B-F) Onset and end time,

- 560 fundamental frequency, maximal frequency, frequency interval, and number of
- spectral bands of the ictal *H* pattern between SOZ and PZ in 37 patients that
- simultaneously expressed the ictal *H* pattern in both SOZ and PZ.
- 563 \*\*\*\*: *p* < 0.0001, nonparametric Mann-Whitney U test; *H* pattern, *Harmonic* pattern;
- 564 SOZ, seizure-onset zone; PZ, early propagation zone.
- 565
- 566 Figure 3. Associations between proportions of removal, time-frequency features
- 567 of the ictal *H* patterns, and seizure outcomes. (A) Violin diagrams comparing
- removed ratios in channels of SOZ, channels expressing the ictal *H* pattern, channels
- 569 expressing the ictal dominant *H* pattern, and channels expressing the ictal
- 570 non-dominant *H* pattern between seizure-free (blue hollow circles) and not
- 571 seizure-free (red hollow triangles) patients. (B-E) Onset time, end time, fundamental
- 572 frequency, maximal frequency, frequency interval, and the number of spectral bands
- 573 of the ictal *H* pattern between seizure-free (blue dots) and not seizure-free (red dots)
- 574 patients. *H pattern*, *Harmonic* pattern; SOZ, seizure-onset zone.
- 575

576 **Figure 4. Binary regression for classification of seizure outcomes.** Seven variables 577 were recruited into the equation: EEG pattern harboring the *H* pattern, lesional MRI

578 findings, seizure localization (mesial temporal lobe or neocortex), pathological

579 findings, removed ratios of channels of the seizure-onset zone, channels expressing

the ictal H pattern, and channels expressing the ictal dominant H pattern. The forest

- 581 plot shows complete removal of channels expressing the ictal dominant H pattern,
- 582 FCD type-II, or other MCD types (compared to gliosis/other non-specific pathology).
- 583 Less implanted electrodes were correlated with seizure freedom. EEG,

584 electroencephalography; *H* pattern, *Harmonic* pattern; MRI, magnetic resonance

- 585 imaging; FCD, focal cortical dysplasia; MCD, malformation of cortical development;
- 586 OR, odds ratio; CI, confidence interval.
- 587

588 **Figure 5. Predictive values for seizure outcomes.** (A-D) Four-fold tables and

589 receiver operator characteristic curve (ROC) for the removed ratios of channels within

590 the seizure-onset zone (dark red), channels expressing the ictal *H* pattern (orange),

- 591 channels expressing the ictal dominant *H* patten (scarlet), and channels expressing the
- 592 ictal non-dominant H pattern (yellow) as predictors of post-surgical outcomes. In the
- 593 four-fold tables, the numbers of seizure-free patients and not seizure-free patients with
- all the corresponding channels removed ('Yes', upper left of the table), or
- incompletely/none removed ('No', bottom left of the table) are listed. p-values for each
- 596 *Chi-Squared* test are provided above each table. In the ROC curves, each colored dot
- 597 represents a combined value of the false positive rate (X-axis) and the true positive
- 598 rate (Y-axis). The areas under curves (AUCs) are respectively listed on the bottom
- 599 right corner.
- 600
- **Figure 6. Survival analysis.** Kaplan-Meier (KM) plots of cumulative probability of
- 602 continuous seizure freedom separated by complete (100%) removal (green continuous
- line) and incomplete/none (0~100%) removal (blue continuous line) of the whole
- 604 cohort, channels of seizure-onset zone, channels expressing the ictal *H* pattern,
- 605 channels expressing the ictal dominant *H* patten, and channels expressing the ictal
- 606 non-dominant *H* pattern. The 'number at risk (number censored)' of each year is
- annotated below the KM plot for every category.

	Tatal salasut	Seizure-free	Non-seizure free		
	Iotal conort	(Engel class Ia)	(> Engel class Ia)	<i>p</i> value	
Number of patients, n	85	52	33	-	
Gender, M, n (%)	47 (55.3%)	29 (55.8%)	18 (54.5%)	0.912 <sup>a</sup>	
Age at onset, y	10.0 (6.0-15.4)	8.5 (5.8-18.8)	12.0 (6.5-15.4)	0.338 <sup>c</sup>	
Age at SEEG recording, y	$23.3\pm10.4$	$23.0\pm11.1$	$23.7\pm9.4$	0.753 <sup>b</sup>	
EEG pattern at <i>H</i> pattern, n (%)					
Irregular spiking (IS)	29 (34.1%)	17 (32.7%)	12 (36.4%)	0 728 <sup>a</sup>	
Fast activity (FA)	56 (65.9%)	35 (67.3%)	21 (63.6%)	0.728	
Lateralization, n (%)					
Left	36 (42.4%)	21 (40.4%)	18 (54.5%)	0 <i>6 1 5 <sup>a</sup></i>	
Right	49 (57.6%)	31 (59.6%)	15 (45.5%)	0.043	
Localization, n (%)					
Mesial temporal lobe	28 (32.9%)	15 (28.8%)	13 (39.4%)	0 212 <sup>a</sup>	
Neocortex	57 (67.1%)	37 (71.2%)	20 (60.6%)	0.515	
Lesional MRI, n (%)	63 (74.1%)	39 (75.0%)	24 (72.7%)	0.816 <sup>a</sup>	
Number of Electrodes, n	10.0 (8.0-11.5)	9.0 (7.0-11.0)	10.0 (9.0-12.5)	<b>0.025</b> <sup>c</sup>	
Number of Contacts, n	112.0 (88.5-127.0)	112.5 (81.3-127.8)	112.0 (96.0-126.5)	0.433 <sup>c</sup>	
Bilateral implantation, n (%)	27 (31.8%)	16 (30.8%)	11 (33.3%)	$0.874^{a}$	
Pathology, n (%)	82	52	30		
Hippocampal sclerosis	13 (15.9%)	8 (15.4%)	5 (16.7%)		
FCD type-II	28 (34.1%)	22 (42.3%) <sup>e</sup>	$6(20.0\%)^{e}$		
Other MCD types	16 (19.5%)	11 (21.2%)	5 (16.7%)	<b>0.014</b> <sup>d</sup>	
Tumor	3 (3.7%)	3 (5.8%)	0		
Gliosis/other non-specific	22 (26.8%)	8 (15.4%) <sup>f</sup>	14 (46.7%) <sup>f</sup>		

## Table 1. Patients' demographics and surgical outcome

Normal variables were presented as mean  $\pm$  standard deviation (SD). Non-normal variables were reported as median (interquartile range, IQR). a. *Pearson's chi-square* test. b. *Unpaired t-test*. c. *Mann-Whitney U* test. d. *Fisher exact* test. e & f. *Bonferroni* correction: In the SF and NSF groups, there was a discrepancy in pathological finding (p = 0.014). A significant difference was found between FCD type-II and gliosis/non-specific findings.

SEEG, stereo-electroencephalography, H pattern, Harmonic pattern, MRI, magnetic resonance imaging, FCD, focal cortical dysplasia, MCD, malformation of cortical development

Notion: The category *Other MCD types* consists of FCD type-I, tuberous sclerosis complex, heterotopia, and mild malformation of cortical development with oligodendroglial hyperplasia in epilepsy.









# Pathology-

(Other MCD types vs. Gliosis/non-specific)

# Pathology-

0

- (FCD type-II vs. Gliosis/non-specific)
- Number of implanted electrodes

Resected ratio of Channel<sub>dH pattern</sub> - (100% vs. 0-100%)



# OR (95% CI) *P* 5.606 (1.174-26.773) 0.031

## 7.387 (1.824-29.906) 0.005

## 0.746 (0.595-0.936) 0.011

# 6.035 (1.291-28.211) 0.022





# Figure 6