

²⁴**Abstract**

²⁵**Objective:** The ictal *Harmonic* pattern (*H* pattern), produced by the non-linear 26 characteristics of EEG waveforms, may hold significant potential for localizing the 27 epileptogenic zone (EZ) in focal epilepsy. However, further validation is needed to 28 establish the *H* pattern's effectiveness as a biomarker for measuring the EZ.

²⁹**Methods:** We retrospectively enrolled 131 patients diagnosed with drug-resistant 30 focal epilepsy, all of whom had complete stereo-electroencephalographic (SEEG) data. ³¹From this cohort, we selected 85 patients for outcome analysis. We analyzed the ³²morphological and time-frequency (TF) features of the *H* pattern using TF plots. A 33 third quartile (Q3) threshold was applied to classify channels expressing either ³⁴dominant (*ChanneldH pattern*) or non-dominant *H* patterns (*Channelnon-dH pattern*). We 35 then examined associations between the morphological features of the *H* pattern and 36 patients' clinical characteristics, as well as the correlations between the extent of 37 channel removal and seizure outcomes.

³⁸**Results:** We found no significant correlations between the morphological features of 39 the ictal *H* pattern and clinical factors, including lesional MRI findings, epileptic 40 onset patterns, epilepsy type, pathology, or surgical outcomes. The non-localizable *H* 41 pattern appeared exclusively in patients with non-focal onset patterns. Notably, the 42 proportion of *Channel_{dH pattern*} was higher in the seizure-onset zone (SOZ) compared 43 to the early propagation zone. The seizure-free group demonstrated significantly 44 higher removal proportions of *Channel_{dH pattern*}, both within and outside the SOZ ($p =$ 45 0.014 ; $p = 0.036$), with AUCs of 0.606 and 0.660, respectively, in a seizure freedom 46 prediction model. Survival analysis confirmed that complete removal of these regions 47 correlated with long-term seizure freedom ($p = 0.008$; $p = 0.028$). Further subgroup 48 analysis showed a significant correlation in neocortical epilepsy ($p = 0.0004$; $p =$ ⁴⁹0.011), but not in mesial temporal lobe epilepsy. Additionally, multivariate analysis 50 identified the complete removal of *Channel_{dH pattern*} as the only independent predictor 51 for seizure freedom ($p = 0.022$; OR 6.035, 95% CI 1.291-28.211).

⁵²**Conclusions:** Our study supports the notion that the dominance of the ictal *H* pattern,

- 53 regardless of its morphology, serves as a novel biomarker for the EZ in focal epilepsy.
- 54 The non-linearity in EEG waveforms provides new insights into understanding ictal
- 55 spreading propagation and offers potential improvements for surgical planning in
- 56 neocortical epilepsy.
-
- ⁵⁸**Keywords:** harmonic pattern; focal epilepsy; epileptogenic zone; epilepsy surgery;
- 59 seizure outcome

⁶⁰**Introduction**

61 Resective surgery currently remains the most effective treatment for drug-resistant 62 epilepsy (DRE). However, only 50% of patients achieve prolonged seizure freedom 63 following surgery. Despite advancements in intracranial electroencephalography ⁶⁴(iEEG) technology, it has yet to surpass a 70% ceiling in achieving favorable surgical 65. outcomes $1-3$. The success of the surgery largely depends on the precise localization 66 and complete removal of the epileptogenic zone (EZ), which is defined as the brain 67 area essential for generating seizures⁴. Postsurgical seizure freedom indicates that the ⁶⁸EZ was likely successfully removed. Although the seizure-onset zone (SOZ) does not 69 necessarily equate to the $EZ^{5, 6}$, experienced clinicians' visual identification of the 70 SOZ is a crucial step in the process of localizing the EZ.

⁷¹Many studies have aimed to identify ictal EEG biomarkers capable of quantitatively ⁷²delineating the EZ. Established semi-quantitative algorithms, such as the 73 epileptogenic index (EI)⁷ and epileptogenic map⁸, have been shown to be effective in 74 cases where ictal onset patterns include low-voltage fast activity. Additionally, the 75 removal of areas exhibiting ictal high-frequency oscillations (HFOs) has been 76 reported as a prognostic indicator of favorable seizure outcomes $9-11$. The concept of 77 the EEG *fingerprint*, proposed as an ictal biomarker of the EZ, suggests a possible synchronous activation of GABAergic neurons at seizure onset ¹² ⁷⁸. Similarly, *brain z*9 *chirp* ¹³ highlights the importance of fast EEG activity in defining the EZ.

⁸⁰In our previous study, we identified the *Harmonic* pattern (*H* pattern), a spectral 81 feature of specific ictal EEG components that contains valuable information for 82 seizure propagation and EZ localization. The *H* pattern is characterized by an 83 equidistant spectral band distribution of varying frequencies on time-frequency (TF) 84 plots. Generated by the specific non-sinusoidal waveforms, the ictal H pattern may 85 represent periodically synchronized neural firings, rather than a methodological 86 artifact. Furthermore, the *H* pattern is frequently observed in seizures with various 87 onset patterns and across different seizure stages 14 . The dominance of the ictal *H*

88 pattern in different brain regions reflects higher non-linearity and has proven useful in 89 localizing the EZ.

90 Despite its potential, the full spectrum of the ictal *H* pattern remains unclear. The *H* 91 pattern shows significant variability in morphology across patients; however, within 92 an individual patient, its morphology and spatial localization tend to remain consistent 93 across seizures, emphasizing the personalized nature of ictal neural dynamics. In this 94 expanded cohort, we first investigated clinical factors related to the morphological 95 properties of the *H* pattern, specifically focusing on its clarity and shape. High clarity 96 indicates a narrower frequency band with reduced background noise, while the shape 97 of the spectral bands reflects changes in the frequency of ictal neural oscillations at a ⁹⁸macroscopic level. This study aims to further support the hypothesis that the 99 dominance of the ictal *H* pattern could serve as a novel biomarker for the EZ. ¹⁰⁰Additionally, we analyzed the potential impact of the *H* pattern's morphological 101 features.

¹⁰³**Methods**

¹⁰⁴*Selection of patient cohort*

¹⁰⁵We retrospectively reviewed 133 patients with drug-resistant epilepsy who had 106 completed stereo-electroencephalography (SEEG) at the Second Affiliated Hospital 107 Zhejiang University School of Medicine (SAHZU) between April 2013 and October ¹⁰⁸2022. We excluded two patients diagnosed with hypothalamic hamartoma from the 109 study. Therefore, we performed *H* pattern analysis on the remaining 131 cases. From 110 this cohort, we selected 85 patients for outcome analysis based on the following ¹¹¹inclusion criteria (**Figure S1**): (1) a focal-onset pattern on SEEG with a clearly 112 defined SOZ localized to a discrete cortical region; (2) undergoing either resective 113 surgery or thermocoagulation treatment; (3) recording at least two habitual clinical 114 seizures during SEEG monitoring; (4) identifying a discernible and localizable ictal *H* 115 pattern (as described in the *Identification of ictal H pattern* section); (5) a follow-up

116 period of at least 12 months after surgery. The Medical Ethics Committee of SAHZU 117 approved this study (Study No. 2023-1260, 2021-0585). We obtained written 118 informed consent from each participant.

119

¹²⁰*SEEG evaluation*

¹²¹All patients were recommended for SEEG evaluation during a multidisciplinary 122 patient management conference. The non-invasive presurgical evaluation included a ¹²³detailed medical history, neuropsychological assessment, scalp video-EEG 124 monitoring, 3.0-Tesla cerebral MRI, and interictal 18 F-fluorodeoxyglucose positron 125 emission tomography $(^{18}$ F-FDG-PET). The procedure for SEEG electrode 126 implantation has been previously described 15 . Based on the working hypothesis, ¹²⁷multiple-contact intracerebral electrodes (8-16 contacts, each 2 mm in length, 0.8 mm 128 in diameter, and spaced 1.5 mm apart) were implanted using a robotic arm-assisted 129 system. SEEG signals were recorded at sampling rates of either 1,000 or 2,000 Hz 130 using EEG systems with 256 channels (Nihon Kohden) or 128 channels (Xltek, 131 Natus). The anatomical locations of each contact along the electrode trajectory in all 132 subjects were confirmed by co-registering post-implantation CT scans with 133 pre-implantation MRI using *Sinovation 2.0*¹⁶. A three-dimensional schematic 134 diagram of the reconstruction is provided in the flowchart (**Figure 1C**).

¹³⁶*Surgical outcome*

137 Following SEEG evaluation, patients underwent either resective surgery or laser 138 ablation. The planning of these surgical interventions was conducted independently of 139 this study. Each contact was classified as either *removed* or *non-removed* based on 140 postsurgical MRI evaluations. Surgical outcomes were assessed at the most recent 141 follow-up and patients were categorized as either seizure-free (SF, Engel class Ia) or 142 not seizure-free (NSF, $>$ Engel class Ia)¹⁷.

¹⁴⁴*Visual EEG analysis*

¹⁴⁵Two experienced and board-certified neurophysiologists reviewed the SEEG data to 146 identify the SOZ and the rapid propagation zone (PZ) using our previously established 147 methods $15, 18$. The SOZ is defined as the brain area that shows the earliest changes on ¹⁴⁸SEEG. The PZ is characterized by the rapid spread of ictal activity within 3-5 seconds 149 after seizure onset. Areas outside these defined zones were classified as 'other 150 regions.' Both the PZ and these 'other regions' were collectively categorized as 151 non-SOZ.

¹⁵³*Identification of ictal H pattern*

¹⁵⁴We selected only contacts within cortical gray matter for quantitative analysis. We ¹⁵⁵analyzed all EEG data using a bipolar montage. For each identified seizure, we 156 extracted a 110-second segment of SEEG data: 10 seconds before and 100 seconds 157 after the EEG onset. Additionally, we collected a segment of preictal EEG data 158 ranging from -230 to -120 seconds before EEG onset as baseline data for 159 normalization. To enhance the reliability of our results, we applied two analytical 160 methods: the Morlet wavelet transform¹⁹ and the multitaper method $20, 21$ in the TF 161 analysis.

¹⁶²We identified the ictal *H* pattern from TF maps based on the following criteria: the 163 presence of two or more distinct high-intensity bands with varying frequencies 164 occurring at any point during a seizure, with equal frequency intervals between 165 adjacent bands (**Figure 1A**). For each patient, we analyzed only the initially occurring ¹⁶⁶*H* pattern. We estimated various features of the ictal *H* pattern, including its shape, 167 clarity, onset and end times, fundamental and maximal frequencies, frequency 168 intervals, and the number of bands. If the *H* pattern was consistently stereotyped 169 across different seizures for a single patient, we included only one data set in the 170 analysis. To determine the dominance of the *H* pattern, we established a threshold

171 known as *Q3*, which represents the upper quartile (75th percentile) of the data 172 distribution for the maximal count of frequency bands per patient. We calculated *Q3* 173 using the following formula:

 174 $03 = (n + 1) * 3/4$.

175 where *n* denotes the highest count of frequency bands observed across all channels. If 176 the number of bands in a channel showing the ictal *H* pattern (*Channel_{H pattern*)} ¹⁷⁷exceeds the *Q3* threshold*,* we marked it as exhibiting a dominant *H* pattern 178 (*Channel_{dH pattern*). Conversely, a channel not meeting this threshold was classified as} ¹⁷⁹expressing a non-dominant *H* pattern (*Channelnon-dH pattern*) (**Figure 1D**). A localizable ¹⁸⁰*H* pattern should be consistent across seizures and present in specific regions. If no 181 gradient in band numbers was observed across all channels, the *H* pattern was 182 classified as non-localizable.

183

¹⁸⁴*Morphology of Ictal Harmonic Pattern*

¹⁸⁵We classified the clarity of the *H* pattern into two levels. Grade A indicated a highly 186 clear frequency band with distinct borders against the background activity. Grade B 187 referred to a less clear frequency band with fuzzier borders. We identified three ¹⁸⁸distinct shapes of the *H* pattern: Shape 1 resembled a parabola, where the fundamental 189 frequency initially increases and then decreases; Shape 2 appeared as a declining line, 190 representing a decrease in the frequency harmonics of the spectral band; and Shape 3 191 took the form of an ascending line, indicating an increase in the frequency harmonics 192 of the spectral band. This resulted in a total of six categories: A1, A2, A3, B1, B2, and ¹⁹³B3 (**Figure S2**). Initially, two investigators (Ye L.Q. and Hu L.L.) performed these 194 classifications. In cases of disagreement, a third epileptologist (Wang S.) made the 195 final decision.

¹⁹⁷*Prediction of Surgical Outcome*

¹⁹⁸For each patient, we categorized channels as either SOZ or non-SOZ based on visual

199 analysis, and as *Channel_{dH pattern* or *Channel_{non-dH pattern* based on TF analysis. We}}

200 determined the ratio of removed channels using the following formula:

 $Removal ratio (\%) = \frac{$ $HChannel_{removed} +$ $#Channel_{removed} +$ H Channe v_{pemoved} \top H Channe $v_{\text{non-removed}}$ \longrightarrow 100% 201 To estimate the correlation between the removal of *Channel_{dH pattern* and seizure} 202 freedom, we used the *Chi-Squared* test and receiver operating characteristic (ROC) 203 curve analysis, from which we derived the area under the curve (AUC). We then 204 generated Kaplan-Meier (KM) curves to depict the cumulative probability of 205 achieving seizure freedom. We compared the retention of seizure freedom between 206 groups using the log-rank test, including patients with complete (100%) removal and 207 those with incomplete or no removal $(0-100\%)$.

208

²⁰⁹**Statistical Analysis**

²¹⁰We used MATLAB 2021a (The MathWorks, Inc.) and SPSS 26.0 for statistical 211 analysis. We evaluated the normality of the distribution of all continuous variables 212 using the *Kolmogorov-Smirnov* test. Normally distributed variables were presented as 213 mean \pm standard deviation (SD), while non-normally distributed variables were 214 reported as median (interquartile range, IQR). We expressed categorical variables as 215 frequencies. We applied an *unpaired t*-test or a nonparametric *Mann-Whitney U* test 216 to analyze continuous variables, and we assessed categorical variables using the ²¹⁷*Chi-Squared* test. We also conducted a binary regression analysis to identify 218 predictive factors affecting surgical outcomes. For multiple comparisons, we 219 employed the Bonferroni correction method. $p < 0.05$ represents statistically 220 significant.

²²²**Results**

²²³*Patient Demographics*

²²⁴In the entire cohort, 75.6% (99 patients) exhibited a focal-onset pattern, while the

²²⁵remaining 24.4% (32 patients) had a non-focal onset pattern (**Table S1**). Among those 226 with a focal-onset pattern, 92.9% (92 patients) displayed an ictal *H* pattern, all of 227 which were localizable. In contrast, 93.8% (30 patients) of those with a non-focal 228 onset pattern exhibited at least one ictal *H* pattern. Of these, 80.0% (24 patients) were 229 localizable, and 20.0% (6 patients) were non-localizable. Most patients presented with 230 only one type of *H* pattern. However, two different types of *H* pattern were observed 231 exclusively in approximately 20% of patients with a non-focal onset pattern (Table ²³²**S1**).

233 Of the 85 patients included in the outcome analysis, 38.9% (33 patients) had 234 participated in our previous *H* pattern study (**Table 1**)¹⁴. In the current cohort, 61.2% ²³⁵(52 patients) achieved seizure freedom after an average follow-up period of 47.8 236 months. We found no significant differences between the SF and NSF groups 237 regarding age at disease onset, age at SEEG recording, or the 238 lateralization/localization of seizures. The incidence of detectable abnormalities on ²³⁹MRI was similar between the two groups (75.0% in the SF group and 72.7% in the 240 NSF group, $p = 0.816$). However, the NSF group had a significantly higher number of 241 implanted electrodes ($p = 0.025$). Focal cortical dysplasia (FCD) \Box was the most 242 common pathological finding, occurring in 34.1% of patients. Among those with 243 malformations of cortical development (MCD) other than FCD \Box (19.5% of patients), 244 11 had FCD I, 3 had tuberous sclerosis complex, 1 had FCD \Box c, and 1 had a mild ²⁴⁵malformation of cortical development with oligodendroglial hyperplasia and epilepsy. 246 Patients with FCD \Box a were included in the hippocampal sclerosis (HS) group. We 247 observed a significant difference in the pathological findings between the SF and NSF 248 groups ($p = 0.014$), particularly between FCD \Box and gliosis/non-specific findings.

²⁵⁰*Morphology, Distribution and Parameters of Ictal Harmonic Pattern*

²⁵¹We summarized the morphological characteristics of the ictal *H* pattern in patients ²⁵²who exhibited only one type of *H* pattern in **Table S2**. The most frequently observed

²⁵³shapes were the declining line (57.3%) and the parabola (40.2%) (**Figure S2**). Our ²⁵⁴analysis revealed no correlation between the clarity or shape of the *H* pattern and 255 clinical factors, including the presence of lesional MRI findings, focal-onset pattern, 256 neocortical epilepsy, pathology, or surgical outcome.

257 Out of 85 patients with focal-onset patterns, we identified both the SOZ and the PZ in ²⁵⁸65 patients. Within these zones, the proportion of channels exhibiting the ictal *^H* 259 pattern was significantly higher in the SOZ (100.0%, IQR 57.3-100%) compared to 260 the PZ (26.3%, IQR 0-50.0%; $p < 0.0001$) (**Figure 2A**). Similarly, the ictal d*H* pattern 261 appeared more frequently in the SOZ $(38.5\%, IQR 16.0-80.0\%)$ than in the PZ $(0\%, IQR 16.0-80.0\%)$ 262 IQR 0-5.8%; $p < 0.0001$) (**Figure 2A**). The ictal *H* pattern was less common in other 263 regions, and we did not observe a significant difference in the distribution of the ²⁶⁴non-d*H* pattern between the SOZ and PZ. Among the 65 patients, 37 exhibited the 265 ictal *H* pattern simultaneously in both the SOZ and PZ. The number of frequency 266 bands of the ictal *H* pattern was notably higher in the SOZ $(3.1, IQR 2.2-6.2)$ 267 compared to the PZ (2.5, IQR 1.9-4.2; $p < 0.0001$), as was the maximal frequency ²⁶⁸(SOZ, 118.8, IQR 92.5-333.0 Hz; PZ, 90.0, IQR 66.3-112.6 Hz; *p* < 0.0001). The 269 onset time of the ictal *H* pattern did not differ between the SOZ and PZ, indicating 270 almost simultaneous initiation across these regions. Furthermore, we did not observe 271 differences in the termination, fundamental frequency, or frequency interval of the 272 ictal *H* pattern between the SOZ and PZ (**Figure 2B-D**). These parameters of the ictal ²⁷³*H* pattern also did not differ between the SF and NSF groups (**Figure 3B-E**). In the 274 remaining 20 patients without a PZ, the dH pattern was predominantly expressed in ²⁷⁵88.9% of channels within the SOZ (IQR, 54.6-88.9%) and was rarely observed in 276 other regions.

²⁷⁸*Ictal Dominant Harmonic Pattern*

279 Figure 3A shows that a higher removal ratio of the *Channel_{dH pattern*} significantly 280 correlates with achieving seizure freedom $(p = 0.014)$. The SF group demonstrated a

- 281 higher removal ratio of the *Channel_{dH pattern*} within non-SOZ (100.0, IQR 100-100%)
- 282 compared to the NSF group (100.0, IQR 0-100%; $p = 0.036$).
- 283

²⁸⁴*Predictive Value of Removing Ictal Dominant Harmonic Pattern*

²⁸⁵Regression analysis identified several independent factors significantly associated 286 with achieving seizure freedom. These factors include the complete removal of the ²⁸⁷*ChanneldH pattern* (*p* = 0.022; OR 6.035, 95% CI 1.291-28.211), FCD or other MCD 288 types (as opposed to gliosis or other non-specific pathologies), and a smaller number 289 of implanted electrodes (Figure 4). Additionally, the association between the 290 complete removal of the *Channel_{dH pattern*} and seizure freedom was supported by an 291 analysis showing statistical significance ($p = 0.031$, AUC = 0.606) (**Figure 5**). The 292 highest AUC value of 0.660 was recorded when considering the complete removal of 293 the *Channel_{dH pattern*} within non-SOZ, highlighting the importance of the ictal dH ²⁹⁴pattern for localizing the EZ (**Figure S3)**.

295

²⁹⁶*Survival Analysis*

²⁹⁷**Figure 6** and **Figure S4** illustrate the 10-year prognosis of seizure freedom for 298 patients who underwent *complete removal* (100%) compared to those with *incomplete* 299 or *no removal* $(0~100\%)$. Among the 85 patients in the cohort, 58.2% achieved 300 seizure freedom at the 10-year mark. Notably, patients who underwent complete 301 removal of the *Channel_{dH pattern*} had a significantly higher probability of attaining 302 long-term seizure freedom $(p = 0.008)$. Specifically, among the 37 patients with 303 complete removal of the *Channel_{dH pattern*} within non-SOZ, there was also an increased 304 likelihood of achieving seizure freedom $(p = 0.028)$. In cases of neocortical epilepsy, 305 the complete removal of the *Channel_{dH pattern* was strongly associated with long-term} 306 seizure freedom $(p=0.0004)$, regardless of whether the removal occurred within the 307 SOZ($p = 0.013$) or non-SOZ($p = 0.011$) (**Figure S5**). In contrast, for mesial temporal 308 lobe epilepsy (mTLE), complete removal of the SOZ, rather than the *Channel_{dH pattern*,}

309 was significantly associated with better long-term seizure freedom outcomes ($p =$

³¹⁰0.019).

³¹²**Discussion**

313 The ictal H pattern, which reflects the non-linear features of ictal SEEG signals on TF ³¹⁴maps, may serve as an innovative ictal biomarker and offer a new approach for 315 delineating the boundary of the EZ. Our key findings reveal: (1) In patients with focal 316 onset patterns, the morphology of the ictal H pattern is consistent within individuals 317 but heterogeneous across the cohort; (2) the morphological features of the ictal *H* 318 pattern do not correlate with clinical variables; (3) the dominance of the ictal *H* 319 pattern indicates high epileptogenicity, and removing regions that exhibit this pattern 320 is crucial for achieving seizure freedom in focal epilepsy, particularly in neocortical 321 epilepsy.

³²²Previous studies have introduced various quantitative biomarkers for measuring 323 spectral components of SEEG signals, including EI $^{7, 22}$, HFOs 23 , and ictal non-linear 324 metrics such as the h^2 coefficient ^{24, 25}, coherence analysis ^{26, 27}, entropy ^{28, 29}, and 325 infraslow activity $30-32$. These computational EEG markers aim to enhance the 326 precision of EZ delineation. Our findings suggest that the ictal dH pattern could be a 327 valuable biomarker for localizing the core epileptogenic cortex across different brain 328 regions. Visual inspection showed that the distribution of the ictal H pattern aligned 329 with the ictal onset zone, regardless of whether it was focal-, multifocal-, or 330 diffuse-onset. In individuals with focal-onset patterns, the ictal *H* pattern typically 331 exhibited a consistent spatial distribution across various seizures. In contrast, 332 individuals with non-focal onset patterns displayed variable types of the ictal *H* 333 pattern, indicating an unstable seizure propagation pattern. The near-simultaneous 334 onset of the ictal *H* pattern across different regions, along with a uniform fundamental 335 frequency, suggests clear inter-regional synchronization during this period. Notably, 336 our findings indicate that regions showing the dH pattern can extend into the PZ ³³⁷beyond the boundary of the SOZ. As our previous research indicated, the *H* pattern 338 reflects spectral skewness or asymmetry in EEG waveforms, and the dH pattern 339 represents stronger inter-regional synchronization during seizure evolution, 340 particularly in focal epilepsy. These findings further support the utility of the d*H* 341 pattern as a non-linear feature in delineating the EZ. Future clinical practices in 342 defining the EZ would benefit from incorporating an integrated EEG biomarker that 343 includes analysis of the dH pattern.

³⁴⁴We analyzed the morphological features of the *H* pattern in terms of shape and clarity. 345 The shape of the spectral bands in the *H* pattern reflects variations in the frequency of 346 synchronized neural oscillations. The most frequently observed shape is a declining 347 line, which closely resembles a spectral signature known as *chirps* ³³⁻³⁵. This shape ³⁴⁸may indicate a decline in neural firing, possibly due to increased inhibition. In our 349 cohort, nearly half of the SOPs exhibiting low-frequency activity (LVFA) expressed 350 *chirp*¹³ patterns, with proportions similar to those of the parabolic shape. We propose 351 that some H patterns with a parabolic shape and very short upward lines might be 352 classified as chirps due to the limitations of visual inspection. This potential 353 misclassification could narrow the scope of the ictal *H* pattern and limit its 354 applicability. A corticothalamic computational model has similarly elucidated how 355 changes in physiological parameters affect the dynamics of tonic-clonic seizures, 356 similar to the *H* pattern. This model revealed that increasing the connection strength 357 enhances the power, count, and duration of harmonic bands 36 . Conversely, the clarity 358 of the *H* pattern signifies the distinctness of the harmonic signals against the 359 background. A highly clear *H* pattern is characterized by a narrow frequency band 360 with minimal noise. Interestingly, these morphological properties do not correlate 361 with clinical factors such as seizure-onset pattern, seizure origin, MRI findings, 362 pathological findings, or seizure outcome. The primary clinically relevant aspect of 363 the *H* pattern is its degree of non-linearity. Regardless of its shape, the prominence of 364 the *H* pattern indicates epileptogenicity, and its complete removal has been linked to 365 the potential achievement of seizure freedom.

³⁶⁶We observed a stronger association between the d*H* pattern and long-term seizure

367 freedom in neocortical epilepsy compared to mTLE. Previous studies have reported 368 that patients with mTLE generally experience more favorable surgical outcomes than 369 those with neocortical epilepsy 3^7 . Additionally, the classification of seizure outcomes 370 can vary among patients. Therefore, we used Engel class Ia as the criterion for a 371 favorable outcome to evaluate the efficacy of the dH pattern in localizing the EZ. ³⁷²Earlier iEEG studies have indicated that the extent of SOZ removal does not always 373 correlate with seizure freedom across all patients $38, 39$. In contrast, we found that, for ³⁷⁴mTLE, the complete removal of the SOZ had a stronger correlation with long-term 375 seizure freedom than the dH pattern. Furthermore, our current mTLE cohort included 376 more MRI-negative cases compared to our previous cohort 14 . Surgical outcomes in 377 MRI-negative cases are usually less favorable compared to those in TLE-HS $40, 41$, 378 suggesting a more extensive epileptogenic network in MRI-negative TLE. Previous 379 research has highlighted differences in the involved neural networks between mTLE 380 and neocortical epilepsy. For instance, an 18 F-FDG-PET study revealed distinct 381 metabolic profiles between these two types of epilepsy, suggesting variations in 382 seizure propagation 42 . Additionally, an iEEG study showed that the interictal 383 epileptogenic network in mTLE exhibits a more regular pattern of graph connectivity 384 compared to neocortical epilepsy 43 . It is well-recognized that epileptic networks 385 differ between mTLE and neocortical temporal lobe epilepsy, as well as between TLE 386 and extra-TLE $^{44, 45}$. A longer follow-up period is necessary to assess the final 387 prognosis in patients with mTLE and to further estimate the value of the *H* pattern in 388 this population.

³⁸⁹Nevertheless, our study has several limitations. First, we based the determination of 390 the *H* pattern's dominance solely on visual interpretation throughout the study. Future 391 investigations should focus on optimizing the definition of the dH pattern and ³⁹²developing a more automated algorithm to quantify the *H* pattern. This would ³⁹³enhance the precision and objectivity of the analysis. We plan to create a quantitative 394 index for the *H* pattern to address this issue. Additionally, we did not analyze the 395 occurrence of the second H pattern during the later ictal stage of one seizure in this

396 study, which warrants further investigation.

³⁹⁸**Conclusions**

399 In focal epilepsy, the dominance of the ictal H pattern offers new insights for 400 localizing the EZ, regardless of its morphology. The distinct non-linear feature of the 401 H pattern has significant implications for presurgical assessments and surgical 402 planning. Applying this biomarker in neocortical epilepsy may offer greater value in 403 achieving long-term favorable outcomes compared to mTLE.

⁴⁰⁵**Data availability**

406 The that support the findings of this study are available upon reasonable request from 407 the corresponding author.

⁴⁰⁸**Acknowledgements**

409 This study received support from the National Natural Science Foundation of China ⁴¹⁰(grant number: 82171437, 82301636, 82471469) and the Major Program of the 411 National Natural Science Foundation of Zhejiang Province (grant number: 412 LD24H090003).

- ⁴¹³**Disclosures**
- 414 The authors have no conflicts of interest to disclose related to this study.

⁴¹⁵**Supplementary material**

416 Supplementary material is available online.

⁴¹⁷**References**

- ⁴¹⁸1. Cardinale F, Rizzi M, Vignati E, et al. Stereoelectroencephalography: 419 retrospective analysis of 742 procedures in a single centre. Brain ⁴²⁰2019;142:2688-2704.
- ⁴²¹2. Jehi L, Morita-Sherman M, Love TE, et al. Comparative Effectiveness of 422 Stereotactic Electroencephalography Versus Subdural Grids in Epilepsy Surgery. Ann 423 Neurol 2021;90:927-939.
- ⁴²⁴3. Sivaraju A, Hirsch L, Gaspard N, et al. Factors Predicting Outcome After ⁴²⁵Intracranial EEG Evaluation in Patients With Medically Refractory Epilepsy. 426 Neurology 2022;99:e1-e10.
- 427 4. Vakharia VN, Duncan JS, Witt JA, Elger CE, Staba R, Engel J, Jr. Getting the 428 best outcomes from epilepsy surgery. Ann Neurol 2018;83:676-690.
- 429 5. Rosenow F, Lüders H. Presurgical evaluation of epilepsy. Brain ⁴³⁰2001;124:1683-1700.
- ⁴³¹6. Widjaja E, Li B, Schinkel CD, et al. Cost-effectiveness of pediatric epilepsy 432 surgery compared to medical treatment in children with intractable epilepsy. Epilepsy 433 Res 2011;94:61-68.
- ⁴³⁴7. Bartolomei F, Chauvel P, Wendling F. Epileptogenicity of brain structures in 435 human temporal lobe epilepsy: a quantified study from intracerebral EEG. Brain ⁴³⁶2008;131:1818-1830.
- ⁴³⁷8. David O, Blauwblomme T, Job AS, et al. Imaging the seizure onset zone with 438 stereo-electroencephalography. Brain 2011;134:2898-2911.
- ⁴³⁹9. Ochi A, Otsubo H, Donner EJ, et al. Dynamic changes of ictal high-frequency 440 oscillations in neocortical epilepsy: using multiple band frequency analysis. Epilepsia ⁴⁴¹2007;48:286-296.
- ⁴⁴²10. Modur PN, Zhang S, Vitaz TW. Ictal high-frequency oscillations in neocortical 443 epilepsy: implications for seizure localization and surgical resection. Epilepsia ⁴⁴⁴2011;52:1792-1801.
- ⁴⁴⁵11. Fujiwara H, Greiner HM, Lee KH, et al. Resection of ictal high-frequency

- 446 oscillations leads to favorable surgical outcome in pediatric epilepsy. Epilepsia ⁴⁴⁷2012;53:1607-1617.
- ⁴⁴⁸12. Grinenko O, Li J, Mosher JC, et al. A fingerprint of the epileptogenic zone in 449 human epilepsies. Brain 2018;141:117-131.
- 450 13. Di Giacomo R, Burini A, Chiarello D, et al. Ictal fast activity chirps as markers of
- 451 the epileptogenic zone. Epilepsia 2024;65:e97-e103.
- ⁴⁵²14. Lingli H, Lingqi Y, Hongyi Y, et al. Harmonic patterns embedding ictal EEG
- ⁴⁵³signals in focal epilepsy: a new insight into the epileptogenic zone. medRxiv ⁴⁵⁴2024:2023.2012.2020.23300274.
- ⁴⁵⁵15. Hu L, Xiong K, Ye L, et al. Ictal EEG desynchronization during low-voltage fast
- 456 activity for prediction of surgical outcomes in focal epilepsy. J Neurosurg 2022:1-10.
- 457 16. Tadel F, Baillet S, Mosher JC, Pantazis D, Leahy RM. Brainstorm: a user-friendly
- 458 application for MEG/EEG analysis. Comput Intell Neurosci 2011;2011:879716.
- 459 17. Engel J. Outcome with respect to epileptic seizures. 1993.
- 460 18. Wang S, So NK, Jin B, et al. Interictal ripples nested in epileptiform discharge
- 461 help to identify the epileptogenic zone in neocortical epilepsy. Clin Neurophysiol ⁴⁶²2017;128:945-951.
- ⁴⁶³19. Bernardino A, Santos-Victor J. A Real-Time Gabor Primal Sketch for Visual Attention.
- 464 2005; Berlin, Heidelberg: Springer Berlin Heidelberg: 335-342.
- ⁴⁶⁵20. Thomson DJ. Spectrum estimation and harmonic analysis. Proceedings of the 466 IEEE 1982;70:1055-1096.
- ⁴⁶⁷21. Babadi B, Brown EN. A review of multitaper spectral analysis. IEEE Trans 468 Biomed Eng 2014;61:1555-1564.
- 469 22. Aubert S, Wendling F, Regis J, et al. Local and remote epileptogenicity in focal
- 470 cortical dysplasias and neurodevelopmental tumours. Brain 2009;132:3072-3086.
- ⁴⁷¹23. González Otárula KA, von Ellenrieder N, Cuello-Oderiz C, Dubeau F, Gotman J.
- ⁴⁷²High-Frequency Oscillation Networks and Surgical Outcome in Adult Focal Epilepsy.
- 473 Ann Neurol 2019;85:485-494.
- ⁴⁷⁴24. Bartolomei F, Lagarde S, Wendling F, et al. Defining epileptogenic networks:

475 Contribution of SEEG and signal analysis. Epilepsia 2017;58:1131-1147.

- 476 25. Hu L, Xiong K, Ye L, et al. Ictal EEG desynchronization during low-voltage fast
- ⁴⁷⁷activity for prediction of surgical outcomes in focal epilepsy. J Neurosurg ⁴⁷⁸2023;139:238-247.
- 479 26. Li YH, Ye XL, Liu QQ, et al. Localization of epileptogenic zone based on graph 480 analysis of stereo-EEG. Epilepsy Res 2016;128:149-157.
- ⁴⁸¹27. Ren Y, Cong F, Ristaniemi T, Wang Y, Li X, Zhang R. Transient seizure onset 482 network for localization of epileptogenic zone: effective connectivity and graph ⁴⁸³theory-based analyses of ECoG data in temporal lobe epilepsy. J Neurol ⁴⁸⁴2019;266:844-859.
- 485 28. Travnicek V, Klimes P, Cimbalnik J, et al. Relative entropy is an easy-to-use 486 invasive electroencephalographic biomarker of the epileptogenic zone. Epilepsia ⁴⁸⁷2023;64:962-972.
- 488 29. Bratu IF, Makhalova J, Garnier E, et al. Permutation entropy-derived parameters 489 to estimate the epileptogenic zone network. Epilepsia 2023.
- ⁴⁹⁰30. Rodin E, Modur P. Ictal intracranial infraslow EEG activity. Clin Neurophysiol ⁴⁹¹2008;119:2188-2200.
- 492 31. Kanazawa K, Matsumoto R, Imamura H, et al. Intracranially recorded ictal direct 493 current shifts may precede high frequency oscillations in human epilepsy. Clin 494 Neurophysiol 2015;126:47-59.
- ⁴⁹⁵32. Nakatani M, Inouchi M, Daifu-Kobayashi M, et al. Ictal direct current shifts 496 contribute to defining the core ictal focus in epilepsy surgery. Brain Commun 497 2022;4:fcac222.
- ⁴⁹⁸33. Schiff SJ, Colella D, Jacyna GM, et al. Brain chirps: spectrographic signatures of 499 epileptic seizures. Clin Neurophysiol 2000;111:953-958.
- ⁵⁰⁰34. Breakspear M, Roberts JA, Terry JR, Rodrigues S, Mahant N, Robinson PA. A ⁵⁰¹unifying explanation of primary generalized seizures through nonlinear brain ⁵⁰²modeling and bifurcation analysis. Cereb Cortex 2006;16:1296-1313.
- ⁵⁰³35. Dutta S, Iyer KK, Vanhatalo S, Breakspear M, Roberts JA. Mechanisms ⁵⁰⁴underlying pathological cortical bursts during metabolic depletion. Nat Commun

⁵⁰⁵2023;14:4792.

- ⁵⁰⁶36. Deeba F, Sanz-Leon P, Robinson PA. Effects of physiological parameter 507 evolution on the dynamics of tonic-clonic seizures. PLoS One 2020;15:e0230510.
-
- ⁵⁰⁸37. Spencer S, Huh L. Outcomes of epilepsy surgery in adults and children. Lancet
- 509 Neurol 2008;7:525-537.
- 510 38. Khan M, Chari A, Seunarine K, et al. Proportion of resected seizure onset zone
- 511 contacts in pediatric stereo-EEG-guided resective surgery does not correlate with
- 512 outcome. Clin Neurophysiol 2022;138:18-24.
- ⁵¹³39. Thomas J, Kahane P, Abdallah C, et al. A Subpopulation of Spikes Predicts 514 Successful Epilepsy Surgery Outcome. Ann Neurol 2023;93:522-535.
- ⁵¹⁵40. Fong JS, Jehi L, Najm I, Prayson RA, Busch R, Bingaman W. Seizure outcome
- 516 and its predictors after temporal lobe epilepsy surgery in patients with normal MRI.
- ⁵¹⁷Epilepsia 2011;52:1393-1401.
- ⁵¹⁸41. McIntosh AM, Wilson SJ, Berkovic SF. Seizure outcome after temporal 519 Iobectomy: current research practice and findings. Epilepsia 2001;42:1288-1307.
- ⁵²⁰42. Zhu HY, Tang YX, Xiao L, et al. Metabolic profiles and correlation with surgical
- 521 outcomes in mesial versus neocortical temporal lobe epilepsy. CNS Neurosci Ther ⁵²²2023;29:2656-2665.
- ⁵²³43. Bartolomei F, Bettus G, Stam CJ, Guye M. Interictal network properties in mesial
- 524 temporal lobe epilepsy: a graph theoretical study from intracerebral recordings. Clin 525 Neurophysiol 2013;124:2345-2353.
- ⁵²⁶44. González Otárula KA, Schuele S. Networks in Temporal Lobe Epilepsy. 527 Neurosurg Clin N Am 2020;31:309-317.
- ⁵²⁸45. Hall GR, Hutchings F, Horsley J, et al. Epileptogenic networks in extra temporal 529 lobe epilepsy. Netw Neurosci 2023;7:1351-1362.
-

⁵³¹**Figure Legends**

⁵³²**Figure 1. Examples and characteristics of the** *H* **pattern (data from patient #057).**

- ⁵³³(A) Ictal SEEG recordings from both SOZ and PZ of a patient with right parietal focal
- 534 cortical dysplasia showing bursts of high-amplitude poly-spikes followed by
- 535 low-voltage fast activity at seizure onset. Below the SEEG recordings are
- 536 corresponding time-frequency plots of the two channels. An ictal *H* pattern could be
- 537 identified in both plots: two or more unique high-intensity spectral bands with varying
- ⁵³⁸frequency starting at 4 s (red arrow) after 0 s (seizure onset), and ending at 19 s (red
- 539 arrow). The frequencies of the bottom and top spectral bands are defined as the
- ⁵⁴⁰*fundamental frequency* and *maximal frequency*, respectively (yellow arrow).
- ⁵⁴¹Importantly, the ictal *H* pattern exhibits a distinctive feature of equidistant distribution
- 542 on the TF plots. For instance, at 6.3 s (long blue dotted arrows), the frequency interval
- ⁵⁴³between adjacent spectral bands is nearly uniform, at approximately 26-28 Hz, as
- 544 observed in the power spectral density diagrams. The ictal *H* pattern shown in the two
- ⁵⁴⁵channels are classified as a 'dominant *H* pattern (d*H* pattern)' and a 'non-dominant *^H*
- 546 pattern (non-dH pattern)', respectively. (B) 3D representation of an example SEEG
- 547 exploration. Electrode contacts are categorized into three regions: the SOZ (red dots),
- 548 the PZ (orange dots), and the other regions (blue dots), according to a patient's ictal
- 549 SEEG recordings. (C) Total number of visually determined spectral bands on the TF
- 550 plot of each channel each time a seizure is counted. Channels are then ranked based
- 551 on the number of bands, and if a channel's count exceeds the threshold of $Q3$, it is
- 552 marked as expressing the ictal dominant *H* pattern. SEEG,
- 553 stereo-electroencephalography; SOZ, seizure-onset zone; PZ, early propagation zone;
- ⁵⁵⁴*H* pattern, *Harmonic* pattern; *Q3*, the 75th percentile of data distribution.
- 555

⁵⁵⁶**Figure 2. Distribution and time-frequency features of ictal** *H* **patterns in SOZ**

⁵⁵⁷**and PZ.** (A) Proportions of channels expressing the ictal *H* pattern, the ictal dominant

- ⁵⁵⁸*H* pattern, and the ictal non-dominant *H* pattern between SOZ (red dots) and PZ
- ⁵⁵⁹(orange dots) in 65 patients who had both SOZ and PZ. (B-F) Onset and end time,

- 560 fundamental frequency, maximal frequency, frequency interval, and number of
- 561 spectral bands of the ictal *H* pattern between SOZ and PZ in 37 patients that
- 562 simultaneously expressed the ictal *H* pattern in both SOZ and PZ.
- ⁵⁶³****: *p* < 0.0001, nonparametric Mann-Whitney U test; *H* pattern, *Harmonic* pattern;
- 564 SOZ, seizure-onset zone; PZ, early propagation zone.
-
- ⁵⁶⁶**Figure 3. Associations between proportions of removal, time-frequency features**
- ⁵⁶⁷**of the ictal** *H* **patterns, and seizure outcomes.** (A) Violin diagrams comparing
- 568 removed ratios in channels of SOZ, channels expressing the ictal *H* pattern, channels
- 569 expressing the ictal dominant *H* pattern, and channels expressing the ictal
- 570 non-dominant *H* pattern between seizure-free (blue hollow circles) and not
- 571 seizure-free (red hollow triangles) patients. (B-E) Onset time, end time, fundamental
- 572 frequency, maximal frequency, frequency interval, and the number of spectral bands
- 573 of the ictal *H* pattern between seizure-free (blue dots) and not seizure-free (red dots)
- 574 patients. *H pattern*, *Harmonic* pattern; SOZ, seizure-onset zone.
- 575 575

⁵⁷⁶**Figure 4. Binary regression for classification of seizure outcomes.** Seven variables

⁵⁷⁷were recruited into the equation: EEG pattern harboring the *H* pattern, lesional MRI

⁵⁷⁸findings, seizure localization (mesial temporal lobe or neocortex), pathological

- 579 findings, removed ratios of channels of the seizure-onset zone, channels expressing
- 580 the ictal *H* pattern, and channels expressing the ictal dominant *H* pattern. The forest
- 581 plot shows complete removal of channels expressing the ictal dominant *H* pattern,
- ⁵⁸²FCD type-II, or other MCD types (compared to gliosis/other non-specific pathology).
- 583 Less implanted electrodes were correlated with seizure freedom. EEG,
- ⁵⁸⁴electroencephalography; *H* pattern, *Harmonic* pattern; MRI, magnetic resonance
- ⁵⁸⁵imaging; FCD, focal cortical dysplasia; MCD, malformation of cortical development;
- 586 OR, odds ratio; CI, confidence interval.
- 587
- ⁵⁸⁸**Figure 5. Predictive values for seizure outcomes.** (A-D) Four-fold tables and
- 589 receiver operator characteristic curve (ROC) for the removed ratios of channels within

590 the seizure-onset zone (dark red), channels expressing the ictal *H* pattern (orange),

- 591 channels expressing the ictal dominant *H* patten (scarlet), and channels expressing the
- 592 ictal non-dominant *H* pattern (yellow) as predictors of post-surgical outcomes. In the
- ⁵⁹³four-fold tables, the numbers of seizure-free patients and not seizure-free patients with
- ⁵⁹⁴all the corresponding channels removed ('*Yes*', upper left of the table), or
- 595 incompletely/none removed (*'No'*, bottom left of the table) are listed. *p*-values for each
- ⁵⁹⁶*Chi-Squared* test are provided above each table. In the ROC curves, each colored dot
- 597 represents a combined value of the false positive rate (X-axis) and the true positive
- 598 rate (Y-axis). The areas under curves $(AUCs)$ are respectively listed on the bottom
- 599 right corner.
-
- ⁶⁰¹**Figure 6. Survival analysis.** Kaplan-Meier (KM) plots of cumulative probability of
- 602 continuous seizure freedom separated by complete (100%) removal (green continuous
- 603 line) and incomplete/none $(0~100\%)$ removal (blue continuous line) of the whole
- ⁶⁰⁴cohort, channels of seizure-onset zone, channels expressing the ictal *H* pattern,
- 605 channels expressing the ictal dominant *H* patten, and channels expressing the ictal
- 606 non-dominant *H* pattern. The 'number at risk (number censored)' of each year is
- 607 annotated below the KM plot for every category.

	Total cohort	Seizure-free	Non-seizure free	p value
		(Engel class Ia)	(> Engel class Ia)	
Number of patients, n	85	52	33	
Gender, M, n (%)	47 (55.3%)	29 (55.8%)	18 (54.5%)	0.912^a
Age at onset, y	$10.0(6.0-15.4)$	$8.5(5.8-18.8)$	$12.0(6.5-15.4)$	0.338 c
Age at SEEG recording, y	23.3 ± 10.4	23.0 ± 11.1	23.7 ± 9.4	0.753^b
EEG pattern at H pattern, n $(\%)$				
Irregular spiking (IS)	29 (34.1%)	17(32.7%)	12 (36.4%)	0.728^{a}
Fast activity (FA)	56 (65.9%)	35 (67.3%)	21 (63.6%)	
Lateralization, n (%)				
Left	36 (42.4%)	21 (40.4%)	18 (54.5%)	0.645^{a}
Right	49 (57.6%)	31 (59.6%)	15 (45.5%)	
Localization, n (%)				
Mesial temporal lobe	28 (32.9%)	15 (28.8%)	13 (39.4%)	0.313^{a}
Neocortex	$57(67.1\%)$	$37(71.2\%)$	$20(60.6\%)$	
Lesional MRI, n (%)	63 (74.1%)	39 (75.0%)	24 (72.7%)	0.816^{a}
Number of Electrodes, n	$10.0(8.0-11.5)$	$9.0(7.0-11.0)$	$10.0(9.0-12.5)$	0.025°
Number of Contacts, n	$112.0(88.5-127.0)$	$112.5(81.3-127.8)$	$112.0(96.0-126.5)$	0.433 ^c
Bilateral implantation, n (%)	27 (31.8%)	$16(30.8\%)$	11 (33.3%)	0.874^{a}
Pathology, n $(\%)$	82	52	30	
Hippocampal sclerosis	13 (15.9%)	8 (15.4%)	5(16.7%)	
FCD type-II	28 (34.1%)	22 $(42.3\%)^e$	6 $(20.0\%)^e$	
Other MCD types	16 (19.5%)	11(21.2%)	5(16.7%)	0.014^d
Tumor	3(3.7%)	$3(5.8\%)$	$\mathbf{0}$	
Gliosis/other non-specific	22 (26.8%)	$8(15.4\%)^f$	14 $(46.7\%)^f$	

Table 1. Patients' demographics and surgical outcome

Normal variables were presented as mean \pm standard deviation (SD). Non-normal variables were reported as median (interquartile range, IQR). a. *Pearson's chi-square* test. b. *Unpaired t-test*. c. *Mann-Whitney U* test. d. *Fisher exact* test. e & f. *Bonferroni* correction: In the SF and NSF groups, there was a discrepancy in pathological finding $(p = 0.014)$. A significant difference was found between FCD type-II and gliosis/non-specific findings.

SEEG, stereo-electroencephalography, H pattern, Harmonic pattern, MRI, magnetic resonance imaging, FCD, focal cortical dysplasia, MCD, malformation of cortical development

Notion: The category *Other MCD types* consists of FCD type-I, tuberous sclerosis complex, heterotopia, and mild malformation of cortical development with oligodendroglial hyperplasia in epilepsy.

Pathology-

(Other MCD types vs. Gliosis/non-specific)

Pathology-

 $\mathbf 0$

- (FCD type-II vs. Gliosis/non-specific)
- Number of implanted electrodes +

Resected ratio of ChanneldH pattern $(100\% \text{ vs. } 0-100\%)$

OR (95% CI) \boldsymbol{P} 5.606 (1.174-26.773) 0.031

7.387 (1.824-29.906) 0.005

$0.746(0.595-0.936)$ 0.011

6.035 (1.291-28.211) 0.022

Figure 6

100% resection $0~100\%$ resection Whole cohort

