Integrating health care and early years support for children and young people living in

deprivation: A cost effectiveness analysis of a novel integrated clinic in Birmingham, UK

Melyda Melyda,¹ Mark Monahan,¹ Chris Bird,² Tracy Roberts,¹ Lorraine Harper,³ Ian Litchfield,³

1. Health Economics Unit, Department of Applied Health Sciences, College of Medicine and Health,

University of Birmingham

2. Birmingham Women's and Children's NHS Foundation Trust

3. Birmingham Health Partners, Department of Applied Health Sciences, College of Medicine and

Health, University of Birmingham

Joint corresponding authors: Dr Mark Monahan (m.monahan.1@bham.ac.uk) and Dr Ian Litchfield

(i.litchfield@bham.ac.uk)

Abstract

Background: Increased use of emergency and secondary care by children and young people,

especially in deprived populations, drive increased costs in health and social care systems in rich

country settings, without necessarily delivering quality care.

Aim: To assess the potential cost-effectiveness of the Sparkbrook's Children Zone (SCZ), a pilot clinic

for children and young people which integrates health and early years support in a highly deprived

area of Birmingham, the UK's second city, compared with standard primary care.

Methods: A decision-analytic model taking healthcare and partial social care perspective was

developed using best available, though limited, evidence from aggregated data of an ongoing pilot,

published literature, expert opinions and assumptions. Effectiveness was measured as proportion of

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

patients attending the emergency department. Deterministic and probabilistic sensitivity analyses were performed to assess the impact of parameter uncertainties.

Results: The integrated SCZ clinic may potentially be cost-effective based on this preliminary modelbased analysis. The SCZ had lower proportion of patients attending emergency department, 0.017 compared with 0.029 for standard primary care, reducing proportion of emergency department visits by 0.012. The average cost of SCZ was £66.22 compared with £110.36 for standard primary care, leading to a cost-saving of £44.08 per patient. This potential reduction in total costs resulting from fewer referrals to children's social care and secondary medical services, including the emergency department. Extensive sensitivity analysis supported the indications that the intervention was likely to be cost-effective.

Conclusion: The SCZ pilot shows potential in integrating health and social care within a community setting, with Its early years services likely enhancing the cost-effectiveness of the SCZ compared to standard primary care. Further robust data and trial evaluation are essential to confirm these findings, ensuring the scalability and sustainability of such programs.

INTRODUCTION

The significant rise in emergency department and outpatient presentations to secondary care among children and young people (CYP) over the past decade has put a strain on the UK's healthcare system, especially in deprived areas with limited access to primary, community care services, and early interventions. And To tackle these challenges, there is growing support from National Health Service (NHS) England and the Royal College of Paediatrics and Child Health, and recent Health and Care Act 2022 to move towards integrating community-based health and social care for children. Evidence from various initiatives in the UK show that integrating children's services reduces outpatient and emergency department use, hospital admissions, general practitioner (GP) appointments, and inappropriate referrals to Child and Adolescent Mental Health Services, with emerging evidence of their long-term cost effectiveness.

The Sparkbrook Children's Zone (SCZ) is one of several place-based integrated care services being piloted by NHSE.⁵ It was established in March 2022 as a pilot integrated health and social care clinic through the Balsall Heath, Moseley, and Sparkhill Primary Care Network, with a list of 14,000 CYP. This site was selected due to the area's high levels of diversity and deprivation, ⁵infant mortality (8/1000), childhood obesity (Year 6 children 42% overweight), special educational needs and disabilities, poor oral health, and low immunisation uptake.¹⁰ These characteristics of the local population mean that SCZ aligns with NHS England's long-standing goal to reduce health inequalities, focusing on the CORE20PLUS population – targeting the most deprived 20% of the national population and other disadvantaged groups.¹¹ The area is also the source of a significant number of inappropriate emergency department attendances.⁵

The SCZ offers a comprehensive range of services from GP, family support workers, mental health outreach, and paediatricians, focusing on five key areas: (1) improved management of common

conditions (asthma, eczema, and constipation); (2) healthy weight; (3) immunisation; (4) oral health; and (5) Early Help Family Support. A key component of the SCZ offer is the co-located Early Help support service, aimed at addressing social determinants of health. Their advisors play a pivotal role in signposting and referring families to local support services to prevent costly late interventions.

If the SCZ effectively improves healthcare outcomes for CYP in this area, it could have significant economic implications for the healthcare sector. However, with tight healthcare budgets and competing demands on resources, justifying the additional costs associated with integrated services by demonstrating value for money, is crucial. We aimed to evaluate the cost-effectiveness of SCZ compared to usual primary care services, analysing how costs are allocated and where savings are generated. Given the limited evidence available on the benefits of integrating health and social support for underserved CYP populations, 13 our findings are intended to inform the development of similar initiatives in the West Midlands and beyond by identifying potentially cost-effective strategies.

METHODS

We developed a decision-analytic model in the form of decision tree model in Microsoft Excel 2024 (Microsoft Corporation, Redmond, Washington, USA) which was parameterized to reflect the patient pathway within the SCZ and standard primary care. Decision trees offer a step-by-step, visual representation, enhancing understanding, especially in complex scenarios. They are particularly useful for decision-making processes involving multiple potential outcomes, allowing decisionmakers to explore various pathways and weigh probabilities. ¹⁴ Given the short-term nature of patient turnover within the SCZ, the decision tree model was well-suited for our analysis. We evaluated the costs and benefits of the SCZ in comparison to the standard primary care from the healthcare and partial social care perspective.

MODEL STRUCTURE

The model structure was developed in consultation with clinical experts from the SCZ. The agreedupon model structure, depicted in Figure 1, shows pathways for patients attending the SCZ clinic

compared to the normal pathway provided in primary care.

In the SCZ arm, patients can either make appointments to visit the SCZ or directly visit the emergency department without any prior referral. Patients may either attend their appointment or not show up (Did Not Attend/DNA). Patients are offered combinations of services from not only GP, but also from a paediatrician, outreach nurse (who also delivers preventive health/mental health professional, and from the Early Help family support team. After receiving treatment or services, patients may be discharged or require follow-up care with their GP or the SCZ clinic. Patients who cannot be clinically managed within these settings will be referred to secondary care services, either to outpatient specialists or, to the emergency department. Patients who need social care support will receive Early Help services in the SCZ. A proportion of patients who receive this service may need to fill out a family support form to obtain more intensive support or be referred to children's social care services for more complex cases.

The standard primary care offer is similar to that of the SCZ but does not include access to a paediatrician, preventive health, an outreach nurse or Early Help services. Patients who require social care support are assumed to be directed to Children's Social Care instead of Early Help services. Once referred to Children's Social Care, children can receive a decision of no further action without further assessment, or they can be assessed and classified into categories: children not in

perpetuity.

It is made available under a CC-BY-NC-ND 4.0 International license .

need (minimal intervention/light touch/general guidance), children in need, children under a child in need plan, children under a child protection plan, and children in care/looked after.¹⁵




Figure 1. Model Structure

MODEL PARAMETERS

Probabilities

Probabilities represent the anticipated likelihood of patients progressing through various model pathways. ¹⁶ To ensure that the model accurately reflects the current conditions in the SCZ, we obtained probability estimates from the SCZ database. When data were unavailable, particularly concerning the comparator arm due to time constraints inherent to the study, we sought expert opinion, made informed assumptions, and referred to published literature to derive the necessary probabilities. All assumptions were confirmed and agreed upon before conducting any analysis and were not subsequently revised at any time.

Table 1. Model probabilities

Parameter	Value Used (SE)	Distribution	Distribution Parameter	Source
Sparkbrook's Children Zone	?			
Did not attend	0.146	Beta	α=5.19	SCZ database
	(0.029)		β=30.26	
Referrals from SCZ	0.039	Beta	α=5.96	SCZ database
Referrals from Sez	(800.0)	Deta	β=147.16	SCZ database
Referred to secondary	0.158		α=5.10	
care services from SCZ		Beta	β=27.13	SCZ database
(ED/outpatient)	(0.032)			
Disabarrand	0.822	Dete	α=0.29	CC7 database
Discharged	(0.164)	Beta	β=0.06	SCZ database
Referred to ED	0.152	Beta	α=5.14	SCZ database
Referred to ED	(0.030)	Бета	β=28.75	SCZ database
ED without referral	0.016		α=6.13	Dutton F et al., 2022, ⁵
	(0.003)	Beta	β=374.78	Montgomerry-Taylor et
(inappropriate)	(0.003)			al., 2015 ⁷
Standard Primary Care				
Did not attend	0.220	Beta	α=4.65	SCZ database
Dia not attend	(0.044)	Deta	β=16.50	SCZ database
			α=5.39	SCZ database,
Referrals from GP	0.118	Beta	β=40.45	Montgomerry-Taylor et
Referrals from Gr	(0.024)	Deta		al., 2015, ⁷ Action for
				children, 2020 ¹²
Referred to secondary			α=2.05	Assuming 10% more if
care services	0.579	Beta	β=1.49	without the ICS
	(0.116)	Dela		(Montgomerry-Taylor et
(ED/outpatient)				al., 2015) ⁷
Discharged	0.822	Beta	α=0.29	Assuming the same as
Discrial 8eu	(0.164)	Deta	β=0.06	SCZ
Referred to ED	0.156	Beta	α=5.12	Assuming 3% more if
Neierred to LD	(0.031)	Deta	β=27.58	without ICS

Parai	meter		Value Used (SE)	Distribution	Distribution Parameter	Source
						(Montgomerry-Taylor et
						al., 2015) ⁷
ED	without	referral	0.021	Beta	α=6.10	Dutton F et al., 2022 ⁵
(inap	propriate)		(0.004)	Бета	β=289.42	Dutton F et al., 2022

SCZ: Sparkbrook Children's Zone; ED: Emergency Department, SE: Standard Errors

Costs and resource use

The unit cost per SCZ consultation was derived from the SCZ business case. Other relevant unit costs were identified from established national sources, including the NHS Reference Costs¹⁷ and the Personal Social Services Research Unit (PSSRU) costs¹⁸. For parameters where costs were not available from these sources, unit costs were extracted from other studies.

To calculate total costs, relevant unit costs were multiplied by the corresponding resource use data. For Early Help and Children Social Care services, the total cost was calculated by weighting the average cost based on the proportion of patients receiving each stage of these services. All costs are reported in 2022-23 Great British pounds (GBP) values. Where necessary, costs were inflated using the Hospital and Community Health Services Pay and Prices Index¹⁸. Model costs are presented in Table 2, and the proportion of children in Early Help and Children Social Care services is presented in Table 3.

Benefit or effectiveness measure

The effectiveness measure was the proportion of patients attending an emergency department. In the analysis, an effectiveness value of "0" was assigned to pathways where patients did not attend the

emergency department and a value of "1" was assigned to pathways where patients did attend an emergency department.

Table 2. Model input costs

Resource use	Unit Cost (£)	Detail	Distribution	Distribution Parameter	Source
SCZ consultation	62	per child (25 min consultation)	Gamma	α=6.25 β=9.92	SCZ database/business case
GP consultation	45.48	per 10 mins consultation	Fixed		PSSRU 2022 ¹⁸
Outpatient Paediatric service	300.6	per consultation (weighted average outpatient care consultant led paediatric service code 420)	Fixed		National Cost Collection for the NHS 2021/2022 ¹⁷
ED Attendance (non-admission)	190.24	per consultation (weighted average the Emergency care code VB07Z - VB11Z (non- admission))	Fixed		National Cost Collection for the NHS 2021/2022 ¹⁷

Daga	Unit Cost	Datail	Distribution	Distribution	Cauras
Resource use	(£)	Detail	Distribution	Parameter	Source
ED Attendance (admission)	279.46	per admission (weighted average the Emergency care code VB07Z -	Fixed		National Cost Collection for the NHS 2021/2022 ¹⁷
		VB11Z (non- admission))			
Did Not Attend	32.49	per DNA	Gamma	α=6.25 β=5.20	NHSE, 2019, ¹⁹ Parsons et al., 2023 ²⁰
Referrals to Early Help worker within the SCZ (light touch/signposting)	62	Per child	Gamma	α=6.25 β=9.92	Covered in SCZ consultation
Early Help - Intensive support (Family connect form)	84.35	Per child	Gamma	α=6.25 β=13.50	Stanford and Lennon, 2019 ¹⁵
Decision of no further action without assessment (after referral to Children Social Care Service)	274.98	Per child	Gamma	α=6.25 β=44	Holmes et al., 2010 ²¹
Children not in need	400.64	Per child, assuming	Gamma	α=6.25 β=64.10	Holmes et al., 2010 ²¹

Resource use	Unit Cost (£)	Detail	Distribution	Distribution Parameter	Source
		just need			
		assessment			
Children in Need				α=6.25	Stanford and
(post Children	755.95	Per child	Gamma	β=120.95	Lennon, 2019 ¹⁵
Social Care Service	755.95	monthly			Leilion, 2019
referral)					
Child in need plan				α=6.25	Stanford and
(post Children	1,312.08	Per child	Gamma	β=209.93	Lennon, 2019 ¹⁵
Social Care Service	_,	monthly	G aa		
referral)					
Children in Child				α=6.25	
Protection Plan		Per child		β=139.45	Stanford and
(post Children	871.59		Gamma		Lennon, 2019 ¹⁵
Social Care Service		monthly			
referral)					
Children in				α=6.25	_
care/looked after		Per child		β=779.75	Stanford and
(post Children	4873.42		Gamma		Lennon, 2019 ¹⁵
Social Care Service		monthly			
referral)					

SCZ: the Sparkbrook Children's Zone; ED: Emergency Department; DNA: Did Not Attend

Table 3. Proportion of children in Early Help and Children Social Care Services

Parameter	Value Used	Distribution	Distribution Parameters	Source
Early Help within SCZ				
Referrals to Early Help	0.64	Dirichlet	α ₁ =237;	SCZ database
worker within the SCZ	0.04		$\alpha_1 + \alpha_2 + \alpha_3 = 370$	302 database

(light				
touch/signposting)				
Intensive support	0.34	Dirichlet	α ₂ =124;	SCZ database
(Family connect form)	0.54	Diricillet	$\alpha_1 + \alpha_2 + \alpha_3 = 370$	SCZ database
Referrals to children	0.02	Dirichlet	α ₃ =9;	SCZ database
social care	0.02	Diricinet	$\alpha_1+\alpha_2+\alpha_3=370$	362 database
Children Social Care				
Decision of no further		Dirichlet	α_1 =49421;	
action without		Diriotilee	$\alpha_1+\alpha_2+\alpha_3+$	
assessment (post	0.07		α_4 + α_5 +	Hoddinott et al., 2023 ²²
Children Social Care			$\alpha_6 = 639901$	
Service referral)				
Children not in need			α ₂ =187390;	
(require light			α_1 + α_2 + α_3 +	
touch/general		Dirichlet	α_4 + α_5 +	
guidance/minimal	0.27	Diricillet	α_6 =639901	Hoddinott et al., 2023 ²²
intervention) (post				
Children Social Care				
Service referral)				
Children in Need (post			α ₃ =164140;	
Children Social Care	0.24	Dirichlet	α_1 + α_2 + α_3 +	Hoddinott et al., 2023 ²²
Service referral)	0.24		α_4 + α_5 +	110001110tt Ct di., 2023
Service referrally			$\alpha_6 = 639901$	
Child in need plan			α ₄ =106000;	
(post Children Social	0.15	Dirichlet	$\alpha_1 + \alpha_2 + \alpha_3 +$	Children's Services Statistics
Care Service referral)	0.13		α_4 + α_5 +	team, 2024 ²³
care service referrally			$\alpha_6 = 639901$	
Children in Child			α ₅ =50780;	
Protection Plan (post	0.07	Dirichlet	α_1 + α_2 + α_3 +	Children's Services Statistics
Children Social Care	0.07		α_4 + α_5 +	team, 2024 ²³
Service referral)			α_6 =639901	
Children in	0.12	Dirichlet	α ₆ =82170;	Hoddinatt at al. 2022?
care/looked after	0.12			Hoddinott et al., 2023 ²²

01

ANALYSIS

Model-based analysis

A roll-back method was employed to estimate the anticipated costs and outcomes for both the SCZ

and standard primary care arms. The results were presented as the difference in costs and the disparity

in the proportion of patients attending the emergency department. A 30-day timeframe was used for

the evaluation. This short-term period aligns with the pilot nature of the SCZ program and the typical

monthly budget cycles of the program. Additionally, children's health can change rapidly, making a

shorter time frame suitable for observing the immediate effects of the intervention.

SENSITIVITY ANALYSIS

Probabilistic sensitivity analysis

Probabilistic sensitivity analysis (PSA) was conducted to assess uncertainty of the model inputs and

base case estimates. PSA presents uncertainty by simultaneously varying multiple input parameters,

repeatedly drawing random values to re-estimate differences in costs and outcomes based on

parameter distributions. 16 Parameters ranging between 0 and 1, such as model probabilities, were

assigned a Beta distribution. Meanwhile, parameters expected to have positively skewed values, like

cost data, were linked to a Gamma distribution. We also utilised the Dirichlet distribution to model the

proportions of patients across different stages of children's social care. This distribution is well-suited

for representing the distribution of multiple probabilities that sum up to one, such as the proportions

of patients receiving various levels of social care. Given the limited data availability, we intentionally

assigned a wide distribution around the estimates to identify the extent to which changes in these

values gave a different cost-effectiveness decision. We arbitrarily used a wide coefficient of variation

of 0.4 for parameters where standard error was not available. A higher coefficient of variation indicates

greater dispersion or variation around the mean value, implying greater uncertainty.²⁴ The PSA was run for 1,000 Monte Carlo simulations and result was visualised on a cost-effectiveness plane. 25,26

Deterministic sensitivity analysis

Deterministic Sensitivity Analysis (DSA) was performed to evaluate the sensitivity of the base case results to changes in specific input parameters. This involved altering the value of certain parameters at a time while keeping the rest constant. A parameter was deemed sensitive if its value change affected the base case cost-effectiveness results. We considered the lower and upper bounds for the probabilities and costs and for this study adjusted them arbitrarily by 20% to 80%. We also used values available from other studies. We additionally explored scenarios involving the variation of the proportion of Early Help (early years support) cases within the SCZ:

- a. Scenario A: In this scenario, 30% of the original light-touch cases in SCZ would require intensive support
- b. Scenario B: In this scenario, 30% of the original light-touch cases in SCZ would require intensive support, and 10% of those receiving intensive support would be referred to children's social care/trust.

Detailed information regarding the values utilised in the DSA is presented in Table 4.

Table 4. Parameter values used in deterministic sensitivity analysis

Variable	Base case	Lower	Upper	Source
	value	bound	bound	
MODEL PROBABILITIES				
Strategy: Sparkbrook's Children Zon	ie			
Did not attend	0.146	0.117	0.176	-20%, +20%
Referrals from SCZ	0.039	0.031	0.047	-20%, +20%
Referred to secondary care services	0.158	0.127	0.190	-20%, +20%
from SCZ (ED/outpatient)	2.200		21_00	==, -==

Variable	Base case	Lower	Upper	6
Variable	value	bound	bound	Source
Discharged	0.822	0.657	0.986	-20%, +20%
Referred to ED	0.152	0.122	0.182	-20%, +20%
ED without referral (inappropriate)	0.016	0.013	0.019	-20%, +20%
Strategy: Standard Primary Care				
Did not attend	0.220	0.176	0.264	-20%, +20%
Referrals from GP	0.118	0.094	0.141	-20%, +20%
Referred to secondary care services (ED/outpatient)	0.579	0.463	0.695	-20%, +20%
Discharged	0.822	0.657	0.986	-20%, +20%
Referred to ED	0.156	0.125	0.188	-20%, +20%
ED without referral (inappropriate)	0.021	0.017	0.025	-20%, +20%
COST INPUT				
SCZ Consultation	£62	£49.60	£74.40	-20%, +20%
Referrals to Early Help worker				
within the SCZ (light	£62	£49.60	£74.40	-20%, +20%
touch/signposting)				
Intensive support (Family connect	£84.35	£67.48	£101.22	-20%, +20%
form)	104.33	107.48	1101.22	-2070, +2070
Referrals to children social care	£1,146.94	£917.55	£1,376.32	-20%, +20%
Decision of no further action				
without assessment (post Children	£274.98	£219.98	£329.98	-20%, +20%
Social Care Service referral)				
Children not in need (require light				
touch/general guidance/minimal	£400.64	£320.51	£480.77	-20%, +20%
intervention) (post Children Social	1400.04	1320.31	1480.77	-2070, 12070
Care Service referral)				
Children in Need (post Children	£755.95	£604.76	£907.15	-20%, +20%
Social Care Service referral)	L/JJ.3J	1004.70	1907.13	-2U/0, T2U/0
Child in need plan (post Children	£1,312.08	£1,049.66	£1,574.50	-20%, +20%
Social Care Service referral)	11,312.00	L1,043.00	L1,3/4.3U	-20/0, F20/0

	Base case	Lower	Upper			
Variable	value	bound	bound	Source		
Children in Child Protection Plan						
(post Children Social Care Service	£871.59	£697.27	£1,045.91	-20%, +20%		
referral)						
Children in care/looked after (post				Lower bound -20%;		
Children Social Care Service	£4873.42	£3,898.74	£17,994.1	Upper bound from		
	140/3.42	13,030.74	9	Stanford and		
referral)				Lennon, 2019 ¹⁵		
PROPORTION OF EARLY HELP CASES WITHIN SCZ						
Scenario a. +30% light-touch cases v	vill need inte	nsive suppor	rt			
Light touch/signposting cases	0.641	0.448		-30%		
Intensive support (Family connect	0.335	0.527		+30% light-touch		
form)	0.555	0	027	cases		
Referrals to children social care	0.024	0.0	024	No change		
Scenario b. 30% light-touch cases w	ill need inten	sive support	and 10% fro	m the intensive		
support will be referred to children	social care/t	rust				
Light touch/signposting cases	0.641	0.4	148	-30%		
				+30% from light-		
Intensive support (Family connect	0.335	0.4	175	touch cases and -		
form)	0.555	0.475		10% to children		
				social care		
Referrals to children social care	0.024	0.0	777	+10% intensive		
vererials to chilufell social cafe	0.024	0.077		support cases		

SCZ: the Sparkbrook Children's Zone; ED: Emergency Department; EH: Early Help; GP: General Practitioner

RESULTS

Model-based analysis results

Base-case results

The base case result is presented in Table 5. The average cost of SCZ per patient was £66.22 which was cheaper compared to £110.36 for standard primary care. Additionally, the SCZ arm had a lower proportion of patients attending the emergency department (0.017 versus 0.029). Consequently, SCZ led to a cost-saving of £44.08 per patient and a reduced proportion of emergency department visits by 0.012, indicating that it dominated standard primary care in terms of being both cheaper and more effective.

Table 5. Base case results

Strategy	Mean Costs	Mean effect (Proportion attending ED)	Mean Incremental Costs	Mean Incremental Effect	ICER
SCZ	£66.28	0.017			
Standard			-£44.08	0.012	SCZ
Standard	£110.36	0.029	144.00	0.012	dominates
primary care					

SCZ: the Sparkbrook Children's Zone; ED: Emergency Department; ICER: Incremental Cost-Effectiveness Ratio

Probabilistic sensitivity analysis (PSA)

The PSA results are visualized in the cost-effectiveness plane (see Figure 2), plotting the difference in costs against the difference in the proportion of emergency department visits. The results indicate that most points fall within the South-East quadrant of the cost-effectiveness plane, indicating a 76.3% probability that the SCZ could be cost-effective, as it is associated with lower costs and fewer emergency department visits compared to standard primary care.




Figure 2. The Cost-Effectiveness Plane

Deterministic sensitivity analysis (DSA)

The results of the DSA (see appendix Table 1) consistently indicate that the SCZ dominates standard primary care, being both cheaper and resulting in a lower proportion of emergency department visits. The cost savings range from £2.99 to £102.81, while the difference in the proportion of emergency department visits ranges from 0.008 to 0.016.

DISCUSSION

This analysis was model based with limited data availability, which required us to seek expert opinions and make assumptions that were confirmed and agreed upon before the analysis began. While the preliminary results are indicative, they offer a promising indication that the Sparkbrook Children's Zone (SCZ) is likely to outperform standard primary care. The preliminary findings show that, on average, the total costs associated with the SCZ were lower compared to those of standard primary care, while also demonstrating a lower proportion of emergency department visits. The costsaving of £44.08 per patient achieved by the SCZ primarily stems from reduced referrals to children's social care and secondary medical care services compared to standard primary care. If we extrapolate this cost-saving to cover the entire 14,000 CYP eligible for SCZ services, it could potentially result in savings of £617,120 for the NHS. This could indicate that despite the slightly higher cost of integrated SCZ consultations (priced at £62 per SCZ consultation compared to £45.48 per GP consultation, which amounts to £16.5 more than standard GP consultations), the potential reduction in total costs resulting from fewer referrals to children's social care and secondary medical services, including the emergency department, could outweigh this additional expense. Our probabilistic sensitivity analysis supports these findings, indicating that under increased uncertainty surrounding input values, the SCZ 76.3% probability of being the preferable option. Furthermore, when deliberately inflating the cost of SCZ consultations by 50% and 80%, the results still indicate cost savings of £18.4 and £2.99 respectively, in comparison to the standard primary care. These findings underscore the potential advantages of integrating health and social care services within a community setting.

Building on these preliminary findings, Early Help emerges as a pivotal component of the SCZ's integrated approach. About 32% of children and young people attending SCZ face-to-face clinics are referred to this service, addressing the complex needs of families, particularly in areas such as diet

and nutrition, special educational needs and disabilities, and behavioural issues. Notably, 39% of these cases benefit from coordinated support via the Family Connect Form—a comprehensive tool for capturing detailed information on a family's circumstances and needs. This enables professionals from various disciplines to collaborate more effectively, allowing for the early identification of potential risks and the organization of multi-disciplinary team support around the family to address these issues before they escalate. Evidence indicates that investment in early help interventions reduces the number of children reaching the threshold of becoming 'in need and in care' under Section 17 of the Children's Act 1989, therefore saving local authorities money by preventing costly late intervention.12

The wider potential benefits of the integrated clinic offer

There are wider potential benefits of the SCZ that are difficult to quantify or assign a monetary value due to the current data constraints and the long-term timeframe required to observe them. The integrated approach of the SCZ attempts to tackle health inequalities by ensuring equitable access to health and social care services. It specifically targets factors such as social deprivation and limited access to services, which are known contributors to health disparities.^{3,27} Operating within an area characterized by high levels of diversity and deprivation, the SCZ aims to ensure that marginalised populations receive the necessary care and support, aligning with the CORE20PLUS5 approach advocated by NHS England to address healthcare inequalities. 11

The cornerstone of the SCZ's strategy is its emphasis on early years services. This support is designed to identify and address issues at an early stage and so preventing them from escalating into more complex problems that necessitate intensive social care interventions. This proactive stance stands in contrast to the reactive nature of traditional social care services, which typically intervene only after

issues have deteriorated. 12 Our modelling found a disparity in costs between Early Help support and the expenditures associated with children requiring more intensive social care intervention. While the Early Help services incur an average cost of approximately £900 per child annually, the financial burden increases considerably for children in need and in care, ranging from £8,300 to £192,000 per child annually. 15 By intervening early, the SCZ effectively mitigates the escalation of problems, thereby reducing the demand for more resource-intensive and costly social care interventions.

Meanwhile, the health promotion initiatives within the SCZ cover various aspects of CYP's health, including immunisation advice, opportunistic immunisations, distribution of Healthy Start vitamins, BMI measurement and referrals, dental health promotion, and smoking cessation advice. These activities collectively contribute to improved health outcomes and have the potential to generate significant cost savings. For instance, childhood immunisation effectively reduces the healthcare costs by preventing diseases like measles, diarrhoea and pneumonia, ^{28,29} which in turn reduces hospitalisation, medical expenses, and prevents productivity losses for caregivers. Similarly, promoting health eating and exercise addresses childhood obesity, reducing future healthcare costs associated with treating diabetes, cardiovascular, and other related disease and potentially saving the NHS over £37 billion and society over £202 billion by 2030. 30,31 Additionally, promoting dental health could reduce the incidence of dental caries, which can lead to gum disease and tooth loss and the high cost of ignoring preventive care (teeth extractions are the commonest reason a child undergoes a general anaesthetic in the UK).³²

Strengths of the Study

To the best of our knowledge, this is the first model-based economic evaluation conducted to analyse the cost-effectiveness of a clinic integrating health and early years support for CYP. We

conducted a range of sensitivity analyses to test the robustness of our findings and to explore the impact of different assumptions and inputs on the cost-effectiveness results. Evaluating integrated services is challenging due to the complexity of the program itself and the limited availability of data in a pilot program to capture its true potential impacts and benefits.

Limitations

The study's main limitation was the lack of robust available data, leading us to rely heavily on literature, expert opinion, and assumptions, especially for comparator data input. However, all assumptions were established and approved in advance to ensure they would not be swayed by the results of the analysis. The impact of these parameters on the model results was also tested through probabilistic and deterministic sensitivity analyses, which showed a positive indication that the SCZ could be a cost-effective intervention. Moreover, we also did not incorporate the potential long-term benefits of the SCZ, which could include additional cost-saving opportunities. Consequently, our analysis may underestimate the true cost-saving potential of the SCZ.

Future studies

The model-based analysis presented here relied on limited primary data and a number of assumptions and we deliberately attempted to undermine the benefits of the SCZ. Despite this, the preliminary results show potential for the SCZ to be cost-saving intervention. Alongside the potential cost savings, the study has also helped identify data and knowledge gaps that will need to be filled to ensure scale-up of integrated care both improves CYP wellbeing and is cost effective.

Future studies need more robust data to confirm these findings. They should also focus on identifying key indicators to measure impact and cost effectiveness of integrated care and evaluate

these over the longer term. Tracking patient outcomes and healthcare costs over extended periods through longitudinal studies will provide more solid evidence of sustained benefits, as demonstrated recently on the health benefits of the UK's Sure Start early years programme.³³ Research should also examine the specific components of integrated care that contribute most significantly to cost savings and improved outcomes. Understanding which aspects of the program are most effective can inform the design of future services and help optimise resource allocation.

CONCLUSION

The Sparkbrook Children's Zone demonstrates a potentially cost saving approach to delivering integrated health and social care within a community setting with high levels of deprivation. The inclusion of Early Help services enhances the cost-effectiveness and overall impact of the SCZ compared to standard primary care. This approach holds the potential to not only save costs but also to enhance the health and social well-being of children and their families. Policymakers and healthcare providers should explore adopting and expanding similar integrated care models to improve the efficiency and equity of healthcare delivery. However, further robust data and evaluation are essential to confirm these findings, ensuring the scalability and sustainability of such programs.

perpetuity.
It is made available under a CC-BY-NC-ND 4.0 International license .

References

- 1. Royal College of Paediatrics and Child Health. *Paediatrics 2040: Our Vision for the Future of Paediatrics in the UK.* (2021).
- 2. British Red Cross. *Nowhere Else to Turn: Exploring High Intensity Use of Accident and Emergency Services*. file:///C:/Users/aafla/Downloads/HIU-Summary-Report-Final.pdf (2021).
- 3. Kossarova Lucia, Cheung Ronny, Hargreaves Dougal & Keeble Eilís. *Admissions of Inequality: Emergency Hospital Use for Children and Young People*. https://www.nuffieldtrust.org.uk/sites/default/files/2017-12/nt-admissions-of-inequality-web.pdf (2017).
- Ruzangi, J. et al. Trends in healthcare use in children aged less than 15 years: a population-based cohort study in England from 2007 to 2017. BMJ Open 10, e033761 (2020).
- 5. Dutton, F., Wolhuter, C., Kaur, S., Rao, S. & Bird, C. 1066 A proposed pilot to integrate health and social care to tackle child inequality: the Sparkbrook Children's Zone, Birmingham. in *British Association of Child and Adolescent Public Health* A56–A56 (BMJ Publishing Group Ltd and Royal College of Paediatrics and Child Health, 2022). doi:10.1136/archdischild-2022-rcpch.91.
- 6. NHS Providers. *A GUIDE TO THE HEALTH AND CARE ACT 2022 for Trusts and Foundation Trusts*. https://nhsproviders.org/media/693819/nhs-providers-a-guide-to-the-health-and-care-act-2022.pdf (2022).
- 7. Montgomery-Taylor, S., Watson, M. & Klaber, R. Child Health General Practice Hubs: a service evaluation. *Arch Dis Child* **101**, 333–7 (2016).
- 8. Day, A. & Patel, S. Connecting Care Children's Hubs Project and Lessons Learnt Report . (2021).
- 9. Soley-Bori, M. *et al.* Cost-effectiveness of the children and young People's health partnership (CYPHP) model of integrated care versus enhanced usual care: analysis of a pragmatic cluster randomised controlled trial in South London. *The Lancet regional health. Europe* **42**, 100917 (2024).
- 10. Birmingham City Council. *Deprivation in Birmingham: Analysis of the 2019 Indices of Deprivation*.
 - $https://www.bing.com/ck/a?!\&\&p=316b2460a924e63aJmltdHM9MTcyMjY0MzlwMCZ\\ pZ3VpZD0wYjE0OTNkYy00OTZhLTYyNTktMzA5ZC04N2MzNDhkNzYzZWImaW5zaWQ9N\\ TlyNA\&ptn=3\&ver=2\&hsh=3\&fclid=0b1493dc-496a-6259-309d-$
 - 87c348d763eb&psq=COUNCIL+%2c+B.+C.+2019.+Deprivation+in+Birmingham%3a+An alysis+of+the+2019+Indices+of+Deprivation.+R+%2c+C.+S.+C.+G.+F.+C.+G.+2019.+The +Health+Effects+of+Sure+Start.&u=a1aHR0cHM6Ly93d3cuYmlybWluZ2hhbS5nb3Yud WsvZG93bmxvYWQvZG93bmxvYWRzL2lkLzl1MzMvaW5kZXhfb2ZfZGVwcml2YXRpb25f MjAxOS5wZGY&ntb=1 (2019).

perpetuity.
It is made available under a CC-BY-NC-ND 4.0 International license .

- 11. NHS England. Core20PLUS5 An approach to reducing health inequalities for children and young people. https://www.england.nhs.uk/about/equality/equality-hub/national-healthcare-inequalities-improvement-programme/core20plus5/core20plus5-cyp/ (2023).
- 12. Actions for Children. Too Little, Too Late: Early Help and Early Intervention Spending in England.
 https://media.actionforchildren.org.uk/documents/Too_Little_Too_Late_Report_Final.
 pdf (2022).
- 13. Bird Chris, Harper Lorraine, Muslim Syed, Yates Derick & Litchfield Ian. Exploring the impact of integrated health and social care services on child health and wellbeing in underserved populations: a systematic review. *medRxiv* (2024).
- 14. Kuntz, K. et al. Decision and Simulation Modeling in Systematic Reviews. (Agency for Healthcare Research and Quality (US), Rockville (MD), 2013).
- Stanford, M. & Lennon, M. Estimating Children's Services Spending on Vulnerable Children.
 https://assets.childrenscommissioner.gov.uk/wpuploads/2019/07/cco-vulnerability-2019-spend-report.pdf (2019).
- 16. Briggs, A., Claxton, K. & Sculpher, M. *Decision Modelling For Health Economic Evaluation*. (Oxford University PressOxford, 2006). doi:10.1093/oso/9780198526629.001.0001.
- 17. NHS England. 2021/22 National Cost Collection Data Publication. https://www.england.nhs.uk/publication/2021-22-national-cost-collection-data-publication/ (2022).
- 18. Jones, K. et al. Unit Costs of Health and Social Care 2022. (2022).
- 19. NHS England. Missed GP appointments costing NHS millions. https://www.england.nhs.uk/2019/01/missed-gp-appointments-costing-nhs-millions/(2019).
- 20. Parsons, J., Abel, G., Mounce, L. T. & Atherton, H. The changing face of missed appointments. *Br J Gen Pract* **73**, 134–135 (2023).
- 21. Holmes, L., McDermid, S., Soper, J., Sempik, J. & Ward, H. *Extension of the Cost Calculator to Include Cost Calculations for All Children in Need*. (2010).
- 22. Hoddinott, S., Davies, N., Fright, M., Nye, P. & Richards, G. *Performance Tracker 2023 Public Services as the UK Approaches: A General Election*. https://www.instituteforgovernment.org.uk/sites/default/files/2024-01/performance-tracker-2023.pdf (2023).
- 23. Children's Services Statistics Team. Children in need. https://explore-education-statistics.service.gov.uk/find-statistics/characteristics-of-children-in-need/2023 (2023).

perpetuity.

It is made available under a CC-BY-NC-ND 4.0 International license.

- 24. UCLA Statistical Consulting Group. What Is the Coefficient of Variation? . https://stats.oarc.ucla.edu/other/mult-pkg/faq/general/faq-what-is-the-coefficient-of-variation/.
- 25. Black, W. C. The CE plane: a graphic representation of cost-effectiveness. *Med Decis Making* **10**, 212–4 (1990).
- 26. Cohen, D. J. & Reynolds, M. R. Interpreting the results of cost-effectiveness studies. *J Am Coll Cardiol* **52**, 2119–26 (2008).
- 27. NICE. NICE and health inequalities. https://www.nice.org.uk/about/what-we-do/nice-and-health-inequalities#approaches-to-addressing-health-inequalities (2024).
- 28. Ozawa, S. *et al.* Return On Investment From Childhood Immunization In Low- And Middle-Income Countries, 2011-20. *Health Aff (Millwood)* **35**, 199–207 (2016).
- 29. Portnoy, A., Resch, S. C., Suharlim, C., Brenzel, L. & Menzies, N. A. What We Do Not Know About the Costs of Immunization Programs in Low- and Middle-Income Countries. *Value Health* **24**, 67–69 (2021).
- 30. Cercato, C. & Fonseca, F. A. Cardiovascular risk and obesity. *Diabetol Metab Syndr* **11**, 74 (2019).
- 31. Hochlaf, D. & Thomas, C. *The Whole Society Approach: Making a Giant Leap on Childhood Health.* https://ippr-org.files.svdcdn.com/production/Downloads/a-whole-society-approach-aug-2020.pdf (2020).
- 32. West, N. *Relationship between Caries and Gum Disease*. https://www.bsperio.org.uk/assets/downloads/Relationship_between_caries_and_gum_disease.pdf (2016).
- 33. Cattan, S., Conti, G., Farquharson, C. & Ginja, R. *The Health Effects of Sure Start*. https://ifs.org.uk/sites/default/files/output_url_files/R155-The-health-effects-of-Sure-Start.pdf (2019).

perpetuity.
It is made available under a CC-BY-NC-ND 4.0 International license .

Contributors

M was the lead author of the manuscript and was supervised in this role by TR and MM. This manuscript was written in collaboration with both TR and MM. All other authors contributed to editing and revising the manuscript.

Funding

Birmingham Women's and Children's NHS Foundation Trust

Competing Interests

All authors have completed the ICMJE uniform disclosure form at www.icmje.org/disclosure-of-interest/. CB declare receiving funding from the Birmingham Women's and Children's NHS Foundation Trust. IL and LH declare receiving funding by Birmingham Women's and Children's NHS Foundation Trust for this study, paid to their institution. All other authors declare no financial relationships with any organisations that might have an interest in the submitted work in the previous three years and no other relationships or activities that could appear to have influenced the submitted work.

Patient and public involvement

Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this economic evaluation research.

Patient consent for publication

Not required

Ethical approval

Ethical approval was granted by University of Birmingham Science Technology Engineering and Mathematics Ethics Committee

Provenance and peer review

Not commissioned; externally peer reviewed

Data availability statement

All reasonable data requests should be submitted to the lead author (m.melyda@bham.ac.uk)