medRxiv preprint doi: https://doi.org/10.1101/2024.09.27.24314273; this version posted September 30, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in

perpetuity. It is made available under a CC-BY 4.0 International license .

Atypical gut microbial ecosystem from athletes with very high exercise 1

capacity improves insulin sensitivity and muscle glycogen store in mice 2

Authors 3

- David Martin^{1,2}, Mathis Bonneau¹, Luz Orfila¹, Mathieu Horeau¹, Mathilde Hazon³, Romain Demay¹, 4
- Emmanuelle Lecommandeur⁴, Rufin Boumpoutou^{1,5}, Arthur Guillotel^{1,6}, Pierrick Guillemot⁷, Mickael 5
- Croyal^{8,9}, Pierre Cressard³, Chrystèle Cressard³, Anne Cuzol¹⁰, Valérie Monbet^{2#} and Frédéric Derbré^{1#}. 6

Affiliations 7

- 8 ¹ Univ Rennes, M2S - UR 1274, F-35000 Rennes, France
- 9 ² Univ Rennes, CNRS, IRMAR - UMR 6625, F-35000 Rennes, France
- 10 ³ Nahibu, Rennes, France
- 11 ⁴ ELS Consulting, Rennes, France
- 12 ⁵ Rennes Ortho Sport, Polyclinique Saint Laurent, Rennes, France
- 13 ⁶ Stade Rennais Football Club, Rennes, France
- 14 ⁷ CHU Pontchaillou, Rennes, France
- 15 ⁸ Institut du thorax, Nantes Université, CNRS, INSERM, Nantes, France
- 16 ⁹ UMS 016, UMS 3556, Nantes Université, Inserm, CNRS, Nantes, France
- ¹⁰ IUT Vannes, University of South Brittany, Vannes, France 17
- 18 [#]: The last co-authors equally contributed to this work
- 19 **Co-corresponding authors:**
- 20 Frederic Derbré, PhD, Assistant professor
- 21 Laboratory "Movement, Sport and Health Sciences" - EA7470
- 22 University of Rennes 2 / ENS Rennes, 35170 Bruz, France
- 23 Mail: Frederic.derbre@univ-rennes2.fr
- 24
- 25 Valerie Monbet, PhD, Full professor
- 26 **IRMAR - UMR CNRS 6625**
- 27 University of Rennes, 35000 Rennes, France
- 28 Mail : valerie.monbet@univ-rennes.fr

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

29 Abstract

- 30 Background. The gut bacterial ecosystem plays a key role in the host's energy metabolism, and
- potentially in the host's exercise capacity, recognized as a powerful predictor of health status and risk
 of mortality.
- Objective. To deepen our understanding of the gut bacterial ecosystem relationship with host's
 exercise capacity and energy metabolism, we characterized the gut microbiota in a cohort of healthy
- humans with heterogeneous exercise capacities, and next determined the impact of fecal microbiota
 transplantation (FMT) from donors of this cohort on energy metabolism and exercise capacity of
- 37 recipient mice.
- 38 Design. 50 male normo-weight participants (from inactive to elite endurance athletes) performed food
- 39 frequency questionnaire (FFQ) and exercise tests to determine exercise capacity parameters (VO_{2max},
- 40 fat oxidation, exercise energy expenditure). Metagenomic shotgun and metabolomic analyses were
- 41 performed to characterize gut microbiota ecosystem and short-chain fatty acids (SCFAs) on human and
- 42 mice fecal samples. Mice performed running exercise capacity tests and metabolic parameters were
- 43 determined in skeletal muscle and plasma samples.
- 44 Results. While our data support that the bacterial ecosystem appears to be modestly altered between
- 45 individuals with low to high exercise capacities, we report that gut bacterial α -diversity, density, and
- 46 functional richness are significantly reduced in athletes with very high exercise capacity. By using FMT,
- 47 we report that the engraftment of these atypical gut microbiota improves insulin sensitivity and muscle
- 48 glycogen stores into transfected mice.
- 49 **Conclusion.** Our data highlight promising therapeutic perspectives in fecal transplantation from human
- 50 donors selected based on exercise capacity parameters.
- 51 **KEYWORDS:** enterotype, propionate, training, nutrients, *Prevotella*

52 What is already known on this topic:

- 53 Gut microbiota ecosystem directly affects exercise capacity and muscle energy metabolism.
- 54 Athletes with high exercise capacity exhibit greater gut microbiota diversity and an over-
- 55 representation of some bacterial species compared to inactive individuals, but current available
- 56 data present major bias including the lack of consideration of dietary habits and body composition.

57 What this study adds:

- Exercise capacity is associated with atypical gut microbiota communities, independently of food
 habits and body composition.
- 60 Atypical gut microbial ecosystem from athletes with very high exercise capacity are related to high
- 61 fecal propionate content, but negatively associated with gut microbiota α -diversity, bacterial 62 density and gut microbiota functional abundance.
- 63 Depending on the donor's exercise capacity, gut bacterial ecosystems affect insulin sensitivity, but
- 64 not exercise capacity, of transfected mice.

65 How this study might affect research, practice or policy:

These data open promising research perspectives: 1) to improve the management of the gut
 microbiota ecosystem of elite athletes and patients performing adapted physical activity for
 therapeutic purposes, and 2) to personalize FMT in patients treated for non-communicable
 diseases by including exercise capacity parameters in the clinical criteria for donor selection.

70

71 INTRODUCTION

72 Exercise capacity is now considered as a powerful predictor of health status and risk of mortality [1,2]. 73 It refers to the maximum amount of physical exertion that an individual can sustain over a prolonged 74 period of time [3]. Exercise capacity can be assessed during exercise testing by measuring maximal 75 oxygen consumption ($\dot{V}O2_{max}$) and fat oxidation (FO) through indirect calorimetry. $\dot{V}O2_{max}$ characterizes 76 the maximum energy that an organism is able to produce during incremental exercise, while the 77 percentage of FO during fasted submaximal exercise provides insight into the whole-body metabolic 78 profile [4,5]. Specifically, during light to moderate submaximal exercise, a healthy organism 79 preferentially degrades fatty acids for its energy supply, derived from both adipose tissue and 80 intramuscular triglyceride breakdown [6]. When the intensity is high to maximal, the organism 81 switches to preferentially use carbohydrates (CHO) provided by muscle glycogen and blood glucose 82 derived from both liver glycogenolysis and gluconeogenesis [6]. However, unlike the almost unlimited 83 fatty acid stores, muscle glycogen stores are limited, and their depletion leads to a reduction in exercise 84 intensity or duration [7]. Increasing the proportion of energy substrates other than CHO during 85 exercise, particularly fatty acids, is thus an essential strategy to spare muscle glycogen improving 86 exercise capacity [8].

87 The gut bacterial ecosystem plays a key role in the host's energy metabolism [9–11], and potentially in 88 the host's exercise capacity. In germ-free or antibiotics-treated (ATB) mice, the absence of gut 89 microbiota directly reduces exercise capacity [12-14], notably due to concomitant reduction in 90 intestinal absorption of glucose and muscle glycogen stores [13,15]. Moreover, the absence or 91 reduction of short-chain fatty acids (SCFAs) production in an organism lacking gut microbiota or 92 exhibiting dysbiotic gut microbiota displays the crucial role of SCFAs in these alterations [16,17]. In 93 humans, cross-sectional studies reveal greater diversity and an over-representation of some bacterial 94 populations in the gut microbiota of elite athletes compared to less active individuals [18-21]. 95 However, all these clinical studies show some limitations including the lack of consideration of dietary 96 habits and body mass index which are known to modulate the gut microbiota composition and function 97 [22].

98 Herein, we report our characterization of the relationship between exercise capacity and the gut 99 bacterial ecosystem in humans, ranging from inactive to elite endurance athletes, independently from 100 diet habits or body composition (EXOMIC cohort). Our data highlight that exercise capacity impacts 101 gut bacterial ecosystem and fecal SCFA levels, independently from dietary habits. Surprisingly, we 102 observe that humans with very high exercise capacity exhibit reduced gut microbiota diversity, density 103 and functional pathways abundance. Using fecal microbiota transplantation (FMT) from humans of our

- 104 cohort to ATB-pretreated mice, we demonstrate that the gut microbiota associated to exercise 105
- capacity of donors affects insulin sensitivity and muscle glycogen store in transfected mice, highlighting
- 106 the essential role of the gut microbiota associated to donor's exercise capacity in the changes of
- 107 metabolic profile of the FMT recipients.

108 MATERIELS AND METHODS

109 Human clinical study

- 110 The EXOMIC pilot clinical study took place at the laboratory "Movement, Sport and health Sciences"
- 111 (M2S, Rennes, France). The study was approved by the national Comité de protection des personnes
- 112 Ouest IV Nantes (ID-RCB: 2021-A02496-35) and registered on ClinicalTrials.gov under NCT05220657.

113 **Mouse experiments**

- 114 C57BL/6J male mice (8-week-old; Janvier Labs, France) were randomly divided into the following two
- 115 experimental groups: control group (CTL, n = 12) versus mice transfected (n = 24) by heterogeneous
- 116 human donors (n=8). Except for running tests, mice were housed in individual standard cages without
- wheels, and thus were not subjected to daily exercise. 117

118 Medical inclusion visits, fecal samples collection and exercise testing

- 119 Each participant completed the food frequency questionnaire to estimate daily food and nutrient
- 120 intakes (FFQ) [23]. Daily energy expenditure was also calculated on a weekly basis (MET-minutes per
- 121 week) using the Global Physical Activity Questionnaire (GPAQ) [24]. Percentage body fat (%FM) was
- 122 measured by skinfold measurement using Harpenden® forceps based on the 4-skinfolds method. One
- 123 fecal sample was collected for each participant during the 15 days following the medical inclusion and
- 124 before the last visit to the laboratory. The participants performed two exercise tests on ergocycle to
- 125 respectively determine (1) maximal oxygen consumption ($\dot{V}O2_{max}$) and (2) carbohydrate (CHO) and fat
- 126 oxidation (FO) using an indirect calorimetry system (Ultima CardiO2, Medgraphics, United Kingdom).

127 Fecal Material Transfection (FMT) from humans to mice

- 128 After 10 days of treatment with an oral cocktail of broad-spectrum antibiotics [13], mice selected for 129 transfection (n=24) received 100 µg of fecal material by oral gavage every morning during 3 days, then 130 once per week for the next two weeks. Fecal samples from FMT and CTL mice were individually 131 collected before and after the antibiotic treatment, and 5 weeks post-FMT. Gastrocnemius muscles and
- 132 abdominal fat mass were dissected, weighed and then either frozen in liquid nitrogen.
- 133 Running exercise capacity in mice
- 134 All mice performed a submaximal running test on a treadmill (Ugo Basile, Gemonio, Italy) before and
- 135 after antibiotic treatment, and then each week during the next 2 weeks. Running exercise capacity was
- 136 determined by the time until exhaustion from a test adapted from Okamoto et al. [14].
- 137 **Metabolic** parameters

- 138 Glycemia was measured using Automated Beckman Coulter (Beckman Coulter, Brea, CA). Serum insulin
- 139 concentrations were measured by enzyme-linked immunosorbent assay (ELISA) according to
- manufacturer's instructions (Millipore, St Louis, MO, USA). The acid-hydrolysis method used for muscle 140
- 141 glycogen quantification was adapted from the protocols described by Adamo and Graham [25].

142 **Metagenomics data**

- 143 Fecal metagenomic shotgun sequences from both human and mice experiments underwent a pre-144 processing pipeline, where sequences were quality filtered using KneadData2 with default parameters 145 [26]. MetaPhIAn3 was used for quantitative profiling the taxonomic composition of the microbial 146 communities of all metagenomic samples [26]. HUMANn3 was used to profile pathway and gene family
- 147 abundances [26].

148 Metabolomic data

- 149 Measurement of short-chain fatty acids (SCFAs), amino acids and bile acids (BAs) concentrations were 150 determined in fecal samples by liquid chromatography-tandem mass spectrometry (LC-MS/MS) by the
- 151 CORSAIRE platform (Biogenouest, Nantes, France).

152 **Statistical analysis**

- 153 Data are presented as means ± SD. Statistical analysis between groups was analyzed according to the
- 154 data distribution and data type. The threshold for statistical significance was assumed to be p<0.05.
- 155 Additional experimental methods and information are provided in online supplemental materials files.

156 RESULTS

157 Exercise capacity is associated with atypical gut microbiota communities, independently of food and 158 nutrient intakes.

159 We first examined whether exercise capacity can affect gut microbiota compositions in healthy 160 humans. To this purpose, we recruited 50 young males to create a physiological and biological 161 database ranging from very low to very high exercise capacity populations, including non-athletes 162 (NoA, n=21), elite soccer players (ESP, n=15) and elite cyclists (EC, n=14). To best limit the differences 163 in food habits between experimental groups, we recruited the NoA participants by matching their food 164 intakes collected by Short Form Food Frequency Questionnaire to correspond qualitatively to the EC's 165 food intakes (Figure 1A, Supplemental Material). All participants recruited were normo-weight (Table 166 S1), completed a long form of FFQ, and performed two sessions of exercise testing (Figure 1A). 167 Participants included in this cohort presented $\dot{V}O2_{max}$ levels ranging from 35 to 85 ml/min/kg (Figure 168 1B), with a proportion of energy expenditure from fat oxidation during submaximal exercise, ranging from 0 to 85% (Figure 1C), characterizing a highly heterogeneous cohort in terms of energy expenditure 169 170 and energy substrates used during exercise.

171 All participants provided their fecal samples to characterize the gut bacterial ecosystem using 172 metagenomics sequencing [26]. We first observe that the two coordinates of the Principal Coordinates 173 Analysis (PCoA) of the pairwise dissimilarity matrix of samples (i.e. Bray Curtis dissimilarity metric) 174 account for 40% of the variability in gut microbiota composition within EXOMIC cohort (Figure 1D). 175 Specifically, individuals with negative coordinates on PCo1 are predominantly elite cyclists and exhibit 176 a gut microbiota composition characterized by a high relative abundance of species belonging to the 177 Prevotella genus, mainly represented by the high expression of Prevotella copri (Figure 1D). 178 Participants with positive coordinates on PCo 1 are mostly non-athletes and present a gut microbiota 179 composition characterized by a high relative abundance of species belonging to the Bacteroides genus, 180 associated with high expression of Bacteroides uniformis, Bacteroides vulgatus and Feacalibacterium 181 prausnitzii (Figure 1D). These observations are consistent with previous findings [21,27,28], and 182 support the concept that humans with high exercise capacity have an atypical gut microbiota 183 composition compared to their counterparts with low exercise capacity.

184 On the Figures 1E and 1F, each line symbolizes variables characterizing a participant included in the 185 EXOMIC cohort. The Hierarchical Ascendant Clustering (HAC) of gut microbiota compositions reveals 186 two distinct clusters related to exercise capacity parameters (Figure 1E), close to the Bacteroides and 187 Prevotella enterotypes concept [29]. The Figure 1F represents the environmental variables associated 188 to each gut microbiota composition, ordered by the HAC. We first estimated the 95% of the confidence 189 intervals for the r correlation coefficients between environmental variables (i.e. macronutrients, 190 physiological variables, body composition, SCFAs and diversity indexes) and coordinates that 191 summarize the variation of gut microbiota compositions within the EXOMIC cohort. Our results reveal 192 that relative macronutrient intakes, including carbohydrates (CHO), fibers, proteins, and lipids, exhibit 193 weak and non-significant association with the coordinates of the gut microbiota composition's PCoA 194 (Figures 1G and S1B). Employing non-parametric redundancy analysis (Canonical Analysis of Principal 195 Coordinates, CAP, Supplemental Material), we measured how strongly environmental variables (i.e. 196 diet, exercise capacity parameters, SCFAs) are linked to the structure and composition of the bacterial 197 ecosystem, computing the explained variance and its associated p value. CAP confirms no discernible 198 association between food or nutrient intakes and gut microbiota compositions (p>0.05, Figures S1C-199 F). Conversely, exercise capacity variables are significantly linked with 12% of the gut microbiota 200 compositions (p=0.01, Figure S1G). Notably, $\dot{V}O2_{max}$, EEE and FO display positive and significant 201 associations with the first Coordinate of the gut microbiota composition's PCoA (p<0.05, Figure 1G). 202 Together, these findings support that exercise capacity directly affects gut microbiota composition, 203 independently of food or nutrient intakes.

Atypical gut microbial ecosystem from athletes with very high exercise capacity are also related to high fecal propionate content.

206 We report in our cohort a significant association of 15% between fecal SFCAs content and gut 207 microbiota compositions (p=0.01, Figure S1H). Notably, fecal valerate and propionate content exhibit 208 the strongest associations (p<0.05, Figure 1G) with the first Coordinate of the gut microbiota 209 composition's PCoA, mainly driven by the Prevotella copri expression (Figure 1D). Species belonging to 210 the Prevotella genus are generally related to a non-western diet [30,31], where starch fermentation 211 leads to an increase in fecal SCFA content [32]. Here, we observe that the proportion of starch intakes 212 are not associated with fecal valerate and propionate content within the EXOMIC cohort (Figure S2A), 213 suggesting that the host's exercise capacity could impact fecal SCFA production independently of fiber 214 intakes. Interestingly, Phascolarctobacterium succinatutens, well-identified to transform succinate 215 into propionate [33], is the second main bacteria associated with the first Coordinate of the gut 216 microbiota composition's PCoA (Figure 1D). While Prevotella copri appears to be a succinate producer 217 rather than propionate producer [34], it seems consistent that fecal propionate content and those 218 species are associated with the first Coordinate of the gut microbiota composition's PCoA. These 219 associations between 1) high exercise capacity parameters, 2) gut microbiota composition dominated 220 by Prevotella copri and Phascolarctobacterium succinatutens and 3) high fecal propionate content are 221 finally consistent with previous studies supporting that propionate supplementation improves exercise 222 capacity in mice [19], as well as resting fat oxidation and energy expenditure in fasted humans [35].

High exercise capacity is negatively associated with gut microbiota α -diversity, bacterial density and gut microbiota functional abundance.

225 Over the last decade, clinical studies conducted in humans with low to moderate exercise capacity 226 have shown that $\dot{V}O2_{max}$ or training volume was positively correlated with gut microbiota α -diversity 227 [28,36,37]. Here, our data support that gut microbiota α -diversity would ultimately follow an inverted-228 U shape relationship with $\dot{V}O2_{max}$ (Figure 2A). For hosts with $\dot{V}O2_{max}$ between 35 and 60 ml/min/kg, the 229 association is positive and consistent with previous studies [36]. However, for hosts with $\dot{V}O2_{max}$ greater 230 than 60 ml/min/kg, the association becomes negative (Figure 2A). Fecal bacterial density is also 231 significantly reduced in hosts with a high capacity to consume energy during maximal exercise (Figure 232 2B). Interestingly, we also report a negative relationship between fat oxidation and gut microbiota α -233 diversity (p<0.05, Figure 2D), hosts with the lowest fecal bacterial density exhibiting the significant 234 highest fat oxidation during submaximal exercise (p<0.05, Figure 2E).

235 To refine our analysis, we explored gut microbiota functionality at different levels according to the 236 MetaCyc database. We observe that the ability of hosts to consume energy during maximal and 237 submaximal exercises are inversely proportional to the richness of gut microbiota functionality 238 (p<0.05, Figure 2C and S3C, respectively). Specifically, several classes of metabolic pathways, including 239 fermentation and carbohydrates degradation, show a negative association with the hosts' exercise 240 capacity parameters, especially $\dot{V}O2_{max}$ (Figure 2G). We then calculated the ratio between gut 241 microbiota capacity to produce energy from available nutrients (i.e. abundance of pathways involved 242 in the generation of precursor metabolites and energy), and its capacity to degrade substrates to serve 243 as sources of nutrients to their own growth (i.e. Degradation/Utilization/Assimilation pathways 244 abundance). Interestingly, this ratio is positively associated with the host's ability to consume energy 245 during maximal exercise (i.e. $\dot{V}O2_{max}$, r = 0.36, p=0.01, Figure 2H). Notably, the ratio of glycolysis and 246 fermentation pathways (i.e. pyruvate producers and fermenters) on carbohydrate degradation 247 pathways (i.e. degradation, storage and utilization of CHO essential for the bacterial growth) is also 248 positively associated with maximal energy consumption during exercise (r = 0.34, p=0.02, Figure 2H). 249 Together, these findings support that hosts with high exercise capacities exhibit a reduction in gut 250 functionality richness, but this reduction seems to be associated with a remodeling of the gut bacterial 251 ecosystem and its metabolic pathways, favoring bacterial energy production from available nutrients 252 over the production of energy from complex molecules (e.g. starch and fibers).

253 Fecal microbiota transplantation (FMT) from high to very high exercise capacity human donors elicits 254 distinct gut microbiota composition in transfected mice

255 To evaluate whether the atypical gut bacterial ecosystem observed in humans with very high exercise 256 capacity could directly affect host's energy metabolism and exercise capacity, we transfected gut 257 microbiota from high to very high exercise capacity humans into ATB-pretreated mice (Figure 3A). We 258 selected 8 physically active and non-sedentary donors exhibiting distinct exercise capacities (i.e. from 259 high to very high $\dot{V}O2_{max}$, from low to very high FO) (Table S2). The fecal bacterial density, species 260 engraftment rate, Bray-Curtis similarity between human donors and transfected mice post-FMT, as 261 well as the donors gut microbiota relative abundance fractions reported post-FMT, are in accordance 262 with current references to consider successful FMT engraftment in humans [38–40], thus supporting 263 the effectiveness of FMT from human donors to transfected mice in our experiments (Figures S5A-E). 264 The dissimilarity between mice reseeded with their own litter and transfected mice confirms such 265 conclusions (Figures S5G-F). We next investigated the link between the parameters of donor exercise 266 capacity with the gut microbiota compositions expressed in the transfected mice by using Canonical 267 Analysis of PCo (CAP, Supplemental Material). We show that $\dot{V}O2_{max}$, FO and EEE are associated with 268 17% of the variability of the gut microbiota compositions of transfected mice (p<0.01). The second axis 269 of the canonical analysis (CAP2) refers to the mice gut microbiota compositions explained by fat 270 oxidation (FO) and exercise energy expenditure (EEE) of the human donors (Figure 3B). Here, we 271 specifically identified *Akkermensia municiphila, Parabacteroides merdae* and *Bacteroides vulgatus* 272 among the bacteria of transfected mice well-associated with exercise capacity parameters (Figure 3B). 273 Together, these findings highlight that distinct gut microbiota compositions are transferred from 274 humans to transfected mice depending on the exercise capacity parameters of the donors.

The gut bacterial ecosystem from human donors affects insulin sensitivity and muscle glycogen stores in transfected mice, depending on the donor's exercise capacity.

Five weeks post-FMT, we assessed the running exercise capacity, HOMA-IR score and skeletal muscle glycogen content to characterize the exercise capacity and insulin sensitivity of each transfected mice (Table S3). We first observe that HOMA-IR score is positively correlated with relative abdominal fat mass, and negatively with skeletal muscle glycogen content (Figure 3C and 3D), indicating a heterogeneous continuum of metabolic profiles in the transfected mice.

282 On the Figures 3E and 3F, each line symbolizes variables characterizing a transfected mouse. The mice's 283 gut microbiota compositions are ordered depending on their positions in the second axis of the CAP 284 (i.e. linked to the FO and the EEE of the donors). The Figure 3F represents the functional and metabolic 285 variables (i.e. running exercise capacity, insulin sensitivity and body composition) associated to each 286 gut microbiota composition. We next determined the 95%-Cl of the r correlation between the second 287 coordinate of the gut microbiota composition's CAP and the physiological variables of the transfected 288 mice. Here, we show that muscle glycogen content is inversely correlated with the second coordinate 289 of the mice gut microbiota composition's CAP (r = -0.49, p < 0.05, Figure 3G), indicating that mice with 290 gut microbiota compositions with negative coordinates in CAP2 exhibit a higher muscle glycogen level 291 compared to those with positive coordinates. Since muscle glycogen content is directly associated with 292 host insulin sensitivity [41], we logically observe that the HOMA-IR score is positively correlated with 293 CAP2 (r = 0.41, p<0.05, Figure 3G), meaning that mice with gut microbiota compositions with negative 294 coordinates in CAP2 exhibit better whole-body insulin sensitivity compared to those with positive 295 coordinates. As previously described, Akkermensia municiphila, Parabacteroides merdae and 296 Bacteroides vulgatus are the main bacteria characterizing the negative coordinate of the second axis 297 of the CAP (Figure 3B). Our data are in line with previous studies where these bacteria were identified 298 to exert metabolic benefits, especially reducing insulin resistance and fat mass gain in obese 299 populations [42–44] and preventing atherosclerotic lesions in rodents [44,45]. In parallel to the results, 300 we report positive correlations between HOMA-IR score and gut microbiota α -diversity in transfected

301 mice (r = 0.41, p=0.046, Figure 3H), whereas muscle glycogen content is negatively correlated with gut 302 microbiota α -diversity (r = -0.36, p=0.08, Figure 3I). These results suggest that the metabolic benefits 303 of gut microbiota transplantation from human donors with very high exercise capacity do not 304 necessarily depend on the gut microbiota α -diversity.

305 We finally explored the gut metabolic functionality of transfected mice and the relationship between 306 functional pathways, exercise capacity, insulin sensitivity and muscle glycogen content. We report a 307 positive correlation between pyruvate fermentation into acetate and lactate with running exercise capacity, while these pathways are negatively associated with HOMA-IR score (p<0.05, Figure 3J). 308 309 Similarly, we observe a positive correlation between pathways of pyruvate synthesis (Glycolysis IV, 310 Glycolysis from glucose) and muscle glycogen content (p<0.05, Figure 3J). Our results are consistent 311 with previous studies reporting that high levels of the functional pathways involved in glycolysis (i.e. 312 pyruvate synthesis) in the gut increase the availability of pyruvate [46], a key substrate for functional pathways of fermentation synthetizing SCFAs from pyruvate degradation [47,48]. In agreement with 313 314 previous studies [49], all these findings highlight that transfection of gut microbiota associated with 315 specific donor characteristics, here exercise capacity parameters, can significantly remodel gut 316 microbiota composition and exert metabolic effects in transfected recipients.

317 **DISCUSSION**

318 During the last decade, substantial observational and interventional studies have been conducted to 319 characterize the impact of physical activity on the gut microbiota ecosystem in humans, providing 320 contradictory results concerning the effects on gut microbiota composition's α -diversity in men and 321 women [36,50–52]. The lack of consideration for the impact of diet when evaluating the effects of 322 exercise capacities on gut microbiota composition is probably the main factor that can explain such 323 discrepancies [28,36,51]. In our prospective clinical study, we rigorously recruited participants with 324 low exercise capacity by matching their food intakes to match qualitatively those of participants with 325 very high exercise capacity. Our data highlight that, for populations that do not present extreme 326 exercise capacities (very low or very high), the effects of host's exercise capacity on the gut bacterial 327 ecosystem appear to be very modest compared to those of food habits.

328 On the contrary, our results support that the gut bacterial ecosystem is remodeled in individuals with 329 a very high ability to consume energy and preferentially use fat during submaximal exercise. In these 330 athletes, we report a clear reduction of gut bacterial diversity (i.e. α -diversity, bacterial density and 331 richness of functionality) with a gut bacterial ecosystem dominated by *Prevotella copri*. This bacterial 332 ecosystem is associated to high fecal propionate levels, in accordance with the role of this SCFA on

333 exercise capacity and fat oxidation [19,35]. We also report that the ratio between functional pathways 334 characterizing the ability to produce energy from available nutrients, and its capacity to degrade 335 substrates to serve as sources of nutrients is higher in humans with a very high ability to consume 336 energy. Collectively, these findings suggest that the relationship between host energy metabolism and 337 the gut bacterial ecosystem may be influenced by the availability of energy resources, which decreases 338 when the host's physical activity levels and exercise capacity are high. In this context, the gut bacterial 339 ecosystem seems to specialize to cope with this situation, with some collateral implications, including 340 a reduction in bacterial diversity and density.

341 The FMT have been recognized as a therapeutic tool to treat non-communicable diseases including 342 obesity, diabetes, neuropsychiatric disorders or inflammatory bowel diseases [53]. One of the current 343 issues is to determine which clinical or metabolic characteristics the donor must possess to observe 344 clinical and/or metabolic effects in the recipient. Numerous studies have already assessed the impact of FMT from donors with distinct metabolic profiles (i.e. mice, brown bears or humans), reporting 345 346 benefits on transfected mice concerning glucose tolerance, insulin sensitivity or lipid metabolism 347 [49,54,55]. While exercise capacity is a clinical hallmark of health status [1,2], it is not yet considered 348 a donor's clinical parameter that could induce metabolic benefits in recipients. However, few recent 349 studies have highlighted that FMT from exercise-trained donor mice improves the metabolic profile of 350 the transfected mice compared to those from sedentary donor mice [56,57]. In the present study, we 351 intend for the first time to evaluate the consequences of FMT from human donors with high to very 352 high exercise capacity on both metabolic parameters and exercise capacity in recipient mice. When we 353 transfected ATB-pretreated mice with fecal samples from our human cohort, we observe in the 354 transfected mice that donors with very high exercise capacity promote an overexpression of 355 Akkermensia muciniphila, Parabacteroides merdae and Bacteroides vulgatus, recognized to induce 356 metabolic benefits for the host [42,43,45]. Interestingly, these bacterial ecosystems are associated 357 with improvements in whole-body insulin sensitivity and in muscle glycogen stores, supporting 358 metabolic benefits associated to the transfer of gut bacterial ecosystem from donors with very high 359 exercise capacity. Finally, mice exhibiting greater insulin sensitivity and higher muscle glycogen stores 360 display a reduction of α -diversity and a higher expression of SCFA producing-pathways, further 361 supporting the hypothesis of a specialization of the gut bacterial ecosystem in favor of metabolite 362 generation, but detrimental for bacterial growth and diversity.

363 In a therapeutic perspective, we selected human donors with high to very high $\dot{V}O2_{max}$ and normo-364 weight, constituting typical lean, healthy, very physically active donors. Despite gut microbiota 365 remodeling and metabolic adaptations, the running exercise capacity of transfected mice remained

unaffected. Importantly, these mice did not perform regular physical exercise during the 5 weeks of the

It is made available under a CC-BY 4.0 International license .

- 367 FMT protocol, and the gut bacterial ecosystem they received from human donors was consequently
- 368 not subjected to the high energy demands related to physical exercise. These results suggest that gut
- 369 microbiota remodeling from donors with high exercise capacity alone is not sufficient to improve the
- 370 exercise capacity of recipients. Overall, all these findings open promising research perspectives: 1) to
- 371 improve the management of the gut microbiota ecosystem of elite athletes and patients performing
- 372 adapted physical activity for therapeutic purposes, and 2) to personalize FMT in patients treated for
- 373 non-communicable diseases by including exercise capacity in the clinical criteria for donor selection.

374 DATA AND CODE AVAILIBILITY

366

Raw metagenomics sequencing data have been deposited into the NCBI under the accession numberPRJNA1115089. In addition, the scripts and others data (metabolomics, food intakes and physiological

- 377 data) used for the computational analyses described in this study are available at GitHub
- 378 <u>https://github.com/DJrMartin/Gut-Microbiota-and-Physical-Activity.git</u>. Any additional information
- 379 required to reanalyze the data reported in this paper is available from the lead contact upon request.

380 ACKNOWLEDGEMENTS

We thank all the participants and coaches from Stade Rennais FC and Sojasun Espoir for their committed efforts in participating in the present study. F.D. acknowledges fundings from the Brittany Council (N°21004090), the Société Française de Nutrition (SFN) and Nahibu. V.B. and F.D. acknowledge fundings from the University of Rennes. D.M, F.D and V.B acknowledge fundings from French National Research Agency within the framework of the PIA France 2030 programme for EUR DIGISPORT (ANR-18-EURE-0022) and CONTINUUM (ANR-21-ESRE-0030) projects.

387 AUTHORS CONTRIBUTION

- 388 Conceptualization and methodology, D.M., M.B., A.C., V.B. and F.D.; data curation, D.M., M.B., M.H.,
- 389 L.O., R.D., M.H., E.L., R.B., A.G., P.G., M.C., F.D.; formal analysis and investigation, D.M., M.B., M.H.,
- A.C., V.B. and F.D.; interpretation of data, D.M., M.B., V.B. and F.D.; writing original draft, D.M.; writ-
- ing review & editing, V.B. and F.D.; funding acquisition, P.C., C.C., V.B. and F.D.; project administra-
- tion, V.B. and F.D.; supervision, V.B. and F.D. All authors approved the final version of the manuscript.

393 DECLARATION OF INTERESTS

394 M.H., P.C. and C.C. are employees and skateholders of Nahibu.

FIGURE TITLES AND LEGENDS 395



396 Figure 1. Exercise capacity is associated with gut microbiota communities, independently of food 397 and nutrient intakes. (A) Experimental design (*non-athlete participants were selected to have their 398 dietary food intake match that of elite athletes). (B) E.E.E (kcal/min) with respect to $\dot{V}O2_{max}$ 399 (ml/min/kg). (C) Fat mass with respect to Fat Oxidation (% of the total energy substrate oxidation). (D) 400 Principal Coordinates Analysis (PCoA) from Bray Curtis dissimilarity of humans' gut microbiota 401 compositions. Subjects are colored by their ratio of *Bacteroides/Prevotella* within their metagenome. 402 The bar plots represent the strongest Spearman's r correlations (in absolute value) between species 403 and PCoA coordinates. (E) Hierarchical Ascendant Clustering computed from the Bray Curtis pairwise 404 dissimilarity matrix (left panel) with the gut microbiota composition of the participants (1 line = 1 405 participant; right panel) (F) Heatmap of relative macronutrient intakes, exercise capacity parameters, 406 body composition, fecal SCFA content and gut microbiota composition's diversities (1 line = 1 407 participant). The points' width depends on the intensity of the variable (i.e. normalization by the intra-408 variable maximum is applied). (G) 95% confidence interval of the r correlation between each variable

- 409 with PCo1 (in dark colors) and PCo2 (in light colors) (see panel **D**). (**H**) r correlation between exercise
- 410 energy parameters, gut microbiota diversity and bacterial density with fecal SCFA content. Statistical
- 411 significance of the correlation: * p<0.05; ** p<0.01; *** p<0.001.

medRxiv preprint doi: https://doi.org/10.1101/2024.09.27.24314273; this version posted September 30, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license .



412 Figure 2. Metagenomic and metabolomic data support a selective pressure between gut microbiota 413 and host energy metabolism. Local regression between $\dot{V}O2_{max}$ and gut microbiota α -diversity (A), 414 bacterial density (B) and gut microbiota functionality abundance (C). Local regression between fat 415 oxidation (FO %) and gut microbiota α -diversity (D), bacterial density (E) and gut microbiota 416 functionality abundance (F). The local green, blue and red regressions correspond respectively to the

- 417 α -diversity, bacterial density and functional abundance of the bacterial ecosystem within the EXOMIC
- 418 cohort. The boxplots represent the repartition of VO_{2max} (B) and FO (%) (E) within each quantile of
- 419 bacterial density. Quantiles are materialized by dashed lines on the local regression plots. (G)
- 420 Correlation between gut microbiota functional families and fecal SFCAs, and exercise capacity
- 421 parameters. Local regression between the ratio of Precursors of Energy metabolites on
- Degradation/Utilization/Assimilation pathways (gut microbiota functional superfamily) and $\dot{V}O2_{max}$ (H) 422
- 423 or FO (I). Local regression between the ratio Glycolysis and Fermentation on CHO degradation
- pathways (gut microbiota functional superfamily) and $\dot{V}O2_{max}$ (J) or FO (K). For local regressions, the 424
- 425 dark line represents the center of the local regression with the 95%-CI in dashed line.



426 Figure 3. The gut bacterial ecosystem from human donors affects insulin sensitivity and muscle 427 glycogen stores in transfected mice, depending on the donor's exercise capacity. (A) Experimental 428 design of FMT from human donors with heterogeneous exercise capacity into antibiotic-pretreated 429 mice. (B) Canonical Analysis of Principal Coordinates (CAP) between the Principal Coordinates of the 430 pairwise dissimilarity matrix (Bray Curtis) of mice's gut microbiota compositions and the donors' 431 exercise capacity parameters (blue arrows). The points represent the most associated species of the 432 second axis of the CAP. (C) Muscle glycogen content according to HOMA-IR. (D) Abdominal fat mass 433 according to HOMA-IR. (E) Ranked mice from the CAP2 with the corresponding gut microbiota 434 composition (1 line = 1 mice). (F) Heatmap of metabolic profile, body composition, food intake and gut 435 microbiota composition's diversities (1 line = 1 mice), the width of each point is normalized by the 436 intra-variable maximum. (G) 95% confidence interval of the r correlation between each variable either

- 437 with CAP1 (in dark colors) or with CAP2 (in light colors) determined in panel B. Local regression
- 438 between gut microbiota α -diversity and HOMA-IR score (H) or muscle glycogen content in mice. (J)
- 439 Correlations between gut microbiota functional families and metabolic parameters, exercise capacity
- 440 and body composition in mice. Statistical significance of the correlation: * p<0.05; ** p<0.01; ***
- 441 p<0.001.

442 **REFERENCES**

- 443 1 Myers J, Prakash M, Froelicher V, *et al.* Exercise Capacity and Mortality among Men
 444 Referred for Exercise Testing. *N Engl J Med.* 2002;346:793–801. doi:
 445 10.1056/NEJMoa011858
- 446 2 Mandsager K, Harb S, Cremer P, *et al.* Association of Cardiorespiratory Fitness With Long447 term Mortality Among Adults Undergoing Exercise Treadmill Testing. *JAMA Netw Open*.
 448 2018;1:e183605. doi: 10.1001/jamanetworkopen.2018.3605
- Goldstein RE. Exercise Capacity. In: Walker HK, Hall WD, Hurst JW, eds. *Clinical Methods: The History, Physical, and Laboratory Examinations*. Boston: Butterworths 1990.
- 4 Amaro-Gahete FJ, Sanchez-Delgado G, Jurado-Fasoli L, *et al.* Assessment of maximal fat
 oxidation during exercise: A systematic review. *Scand J Med Sci Sports*. 2019;29:910–21.
 doi: 10.1111/sms.13424
- 454 5 Clausen JSR, Marott JL, Holtermann A, *et al.* Midlife Cardiorespiratory Fitness and the
 455 Long-Term Risk of Mortality: 46 Years of Follow-Up. *J Am Coll Cardiol.* 2018;72:987–95.
 456 doi: 10.1016/j.jacc.2018.06.045
- 457 6 van Loon LJ, Greenhaff PL, Constantin-Teodosiu D, *et al.* The effects of increasing exercise
 458 intensity on muscle fuel utilisation in humans. *J Physiol.* 2001;536:295–304. doi:
 459 10.1111/j.1469-7793.2001.00295.x
- 460 7 Ørtenblad N, Westerblad H, Nielsen J. Muscle glycogen stores and fatigue. *J Physiol*.
 461 2013;591:4405–13. doi: 10.1113/jphysiol.2013.251629
- 462 8 Hargreaves M, Spriet LL. Skeletal muscle energy metabolism during exercise. *Nat Metab.*463 2020;2:817–28. doi: 10.1038/s42255-020-0251-4
- 464 9 Liu J, Tan Y, Cheng H, *et al.* Functions of Gut Microbiota Metabolites, Current Status and
 465 Future Perspectives. *Aging Dis.* 2022;13:1106–26. doi: 10.14336/AD.2022.0104
- Takeuchi T, Kubota T, Nakanishi Y, *et al.* Gut microbial carbohydrate metabolism
 contributes to insulin resistance. *Nature*. 2023;621:389–95. doi: 10.1038/s41586-02306466-x
- 469 11 Zarrinpar A, Chaix A, Xu ZZ, *et al.* Antibiotic-induced microbiome depletion alters
 470 metabolic homeostasis by affecting gut signaling and colonic metabolism. *Nat Commun.*471 2018;9:2872. doi: 10.1038/s41467-018-05336-9
- 472 12 Dohnalová L, Lundgren P, Carty JRE, *et al.* A microbiome-dependent gut-brain pathway
 473 regulates motivation for exercise. *Nature*. 2022;612:739–47. doi: 10.1038/s41586-022474 05525-z
- 13 Nay K, Jollet M, Goustard B, *et al.* Gut bacteria are critical for optimal muscle function: a
 potential link with glucose homeostasis. *Am J Physiol Endocrinol Metab.* 2019;317:E158–
 71. doi: 10.1152/ajpendo.00521.2018

478 14 Okamoto T, Morino K, Ugi S, *et al.* Microbiome potentiates endurance exercise through
479 intestinal acetate production. *Am J Physiol Endocrinol Metab.* 2019;316:E956–66. doi:
480 10.1152/ajpendo.00510.2018

- Kim HJ, Kim YJ, Kim YJ, *et al.* Microbiota influences host exercise capacity via modulation
 of skeletal muscle glucose metabolism in mice. *Exp Mol Med.* 2023;55:1820–30. doi:
 10.1038/s12276-023-01063-4
- 484 16 Hawley JA. Microbiota and muscle highway two way traffic. *Nat Rev Endocrinol*.
 485 2020;16:71–2. doi: 10.1038/s41574-019-0291-6
- 486 17 den Besten G, van Eunen K, Groen AK, *et al.* The role of short-chain fatty acids in the
 487 interplay between diet, gut microbiota, and host energy metabolism. *J Lipid Res.*488 2013;54:2325–40. doi: 10.1194/jlr.R036012
- Petersen LM, Bautista EJ, Nguyen H, *et al.* Community characteristics of the gut
 microbiomes of competitive cyclists. *Microbiome*. 2017;5:98. doi: 10.1186/s40168-0170320-4
- Scheiman J, Luber JM, Chavkin TA, *et al.* Meta-omics analysis of elite athletes identifies a
 performance-enhancing microbe that functions via lactate metabolism. *Nat Med.*2019;25:1104–9. doi: 10.1038/s41591-019-0485-4
- 495 20 Allen JM, Mailing LJ, Niemiro GM, *et al.* Exercise Alters Gut Microbiota Composition and
 496 Function in Lean and Obese Humans. *Med Sci Sports Exerc.* 2018;50:747–57. doi:
 497 10.1249/MSS.0000000001495
- 498 21 Barton W, Penney NC, Cronin O, *et al.* The microbiome of professional athletes differs
 499 from that of more sedentary subjects in composition and particularly at the functional
 500 metabolic level. *Gut.* 2018;67:625–33. doi: 10.1136/gutjnl-2016-313627
- 501 22 Greenhalgh K, Meyer KM, Aagaard KM, *et al.* The human gut microbiome in health:
 502 establishment and resilience of microbiota over a lifetime. *Environ Microbiol*.
 503 2016;18:2103–16. doi: 10.1111/1462-2920.13318
- Mulligan AA, Luben RN, Bhaniani A, *et al.* A new tool for converting food frequency
 questionnaire data into nutrient and food group values: FETA research methods and
 availability. *BMJ Open.* 2014;4:e004503. doi: 10.1136/bmjopen-2013-004503
- 507 24 Bull FC, Maslin TS, Armstrong T. Global physical activity questionnaire (GPAQ): nine
 508 country reliability and validity study. *J Phys Act Health*. 2009;6:790–804. doi:
 509 10.1123/jpah.6.6.790
- 510 25 Adamo KB, Graham TE. Comparison of traditional measurements with macroglycogen
 511 and proglycogen analysis of muscle glycogen. *J Appl Physiol Bethesda Md 1985*.
 512 1998;84:908–13. doi: 10.1152/jappl.1998.84.3.908

513 26 Beghini F, McIver LJ, Blanco-Míguez A, *et al.* Integrating taxonomic, functional, and strain514 level profiling of diverse microbial communities with bioBakery 3. *eLife*. 2021;10:e65088.
515 doi: 10.7554/eLife.65088

- 516 27 Clarke SF, Murphy EF, O'Sullivan O, *et al.* Exercise and associated dietary extremes impact 517 on gut microbial diversity. *Gut.* 2014;63:1913–20. doi: 10.1136/gutjnl-2013-306541
- 518 28 Baldanzi G, Sayols-Baixeras S, Ekblom-Bak E, *et al.* Accelerometer-based physical activity
 519 is associated with the gut microbiota in 8416 individuals in SCAPIS. *eBioMedicine*.
 520 2024;100. doi: 10.1016/j.ebiom.2024.104989
- S21 29 Arumugam M, Raes J, Pelletier E, *et al.* Enterotypes of the human gut microbiome.
 S22 Nature. 2011;473:174–80. doi: 10.1038/nature09944
- 30 Blanco-Míguez A, Gálvez EJC, Pasolli E, *et al.* Extension of the Segatella copri complex to
 13 species with distinct large extrachromosomal elements and associations with host
 conditions. *Cell Host Microbe*. 2023;31:1804-1819.e9. doi: 10.1016/j.chom.2023.09.013
- S1 Christensen L, Vuholm S, Roager HM, *et al. Prevotella* Abundance Predicts Weight Loss
 Success in Healthy, Overweight Adults Consuming a Whole-Grain Diet Ad Libitum: A Post
 Hoc Analysis of a 6-Wk Randomized Controlled Trial. *J Nutr.* 2019;149:2174–81. doi:
 10.1093/jn/nxz198
- den Besten G, van Eunen K, Groen AK, *et al.* The role of short-chain fatty acids in the
 interplay between diet, gut microbiota, and host energy metabolism. *J Lipid Res.*2013;54:2325–40. doi: 10.1194/jlr.R036012
- Watanabe Y, Nagai F, Morotomi M. Characterization of Phascolarctobacterium
 succinatutens sp. nov., an asaccharolytic, succinate-utilizing bacterium isolated from
 human feces. *Appl Environ Microbiol*. 2012;78:511–8. doi: 10.1128/AEM.06035-11
- 536 34 De Vadder F, Kovatcheva-Datchary P, Zitoun C, *et al.* Microbiota-Produced Succinate
 537 Improves Glucose Homeostasis via Intestinal Gluconeogenesis. *Cell Metab.* 2016;24:151–
 538 7. doi: 10.1016/j.cmet.2016.06.013
- 539 35 Chambers ES, Byrne CS, Aspey K, *et al.* Acute oral sodium propionate supplementation
 540 raises resting energy expenditure and lipid oxidation in fasted humans. *Diabetes Obes*541 *Metab.* 2018;20:1034–9. doi: 10.1111/dom.13159
- 542 36 Estaki M, Pither J, Baumeister P, *et al.* Cardiorespiratory fitness as a predictor of intestinal
 543 microbial diversity and distinct metagenomic functions. *Microbiome*. 2016;4:42. doi:
 544 10.1186/s40168-016-0189-7
- 545 37 Dupuit M, Rance M, Morel C, *et al.* Effect of concurrent training on body composition and
 546 gut microbiota in postmenopausal women with overweight or obesity. *Med Sci Sports*547 *Exerc.* 2022;54:517–29.

548 549 550	38	Ianiro G, Punčochář M, Karcher N, <i>et al.</i> Variability of strain engraftment and predictability of microbiome composition after fecal microbiota transplantation across different diseases. <i>Nat Med</i> . 2022;28:1913–23. doi: 10.1038/s41591-022-01964-3
551 552 553	39	Podlesny D, Durdevic M, Paramsothy S, <i>et al.</i> Identification of clinical and ecological determinants of strain engraftment after fecal microbiota transplantation using metagenomics. <i>Cell Rep Med</i> . 2022;3:100711. doi: 10.1016/j.xcrm.2022.100711
554 555 556	40	Porcari S, Benech N, Valles-Colomer M, <i>et al.</i> Key determinants of success in fecal microbiota transplantation: From microbiome to clinic. <i>Cell Host Microbe</i> . 2023;31:712–33. doi: 10.1016/j.chom.2023.03.020
557 558	41	Petersen MC, Shulman GI. Mechanisms of Insulin Action and Insulin Resistance. <i>Physiol Rev</i> . 2018;98:2133–223. doi: 10.1152/physrev.00063.2017
559 560 561 562	42	Dao MC, Everard A, Aron-Wisnewsky J, <i>et al.</i> Akkermansia muciniphila and improved metabolic health during a dietary intervention in obesity: relationship with gut microbiome richness and ecology. <i>Gut.</i> 2016;65:426–36. doi: 10.1136/gutjnl-2014-308778
563 564 565	43	Everard A, Belzer C, Geurts L, <i>et al.</i> Cross-talk between Akkermansia muciniphila and intestinal epithelium controls diet-induced obesity. <i>Proc Natl Acad Sci</i> . 2013;110:9066–71. doi: 10.1073/pnas.1219451110
566 567 568	44	Xu M, Lan R, Qiao L, <i>et al.</i> Bacteroides vulgatus Ameliorates Lipid Metabolic Disorders and Modulates Gut Microbial Composition in Hyperlipidemic Rats. <i>Microbiol Spectr</i> . 2023;11:e0251722. doi: 10.1128/spectrum.02517-22
569 570 571	45	Qiao S, Liu C, Sun L, <i>et al</i> . Gut Parabacteroides merdae protects against cardiovascular damage by enhancing branched-chain amino acid catabolism. <i>Nat Metab</i> . 2022;4:1271–86. doi: 10.1038/s42255-022-00649-y
572 573	46	Macfarlane S, Macfarlane GT. Regulation of short-chain fatty acid production. <i>Proc Nutr Soc</i> . 2003;62:67–72. doi: 10.1079/PNS2002207
574 575 576	47	Liu Y, Wang Y, Ni Y, <i>et al.</i> Gut Microbiome Fermentation Determines the Efficacy of Exercise for Diabetes Prevention. <i>Cell Metab.</i> 2020;31:77-91.e5. doi: 10.1016/j.cmet.2019.11.001
577 578	48	Louis P, Flint HJ. Formation of propionate and butyrate by the human colonic microbiota. <i>Environ Microbiol</i> . 2017;19:29–41. doi: 10.1111/1462-2920.13589
579 580 581	49	de Groot P, Scheithauer T, Bakker GJ, <i>et al.</i> Donor metabolic characteristics drive effects of faecal microbiota transplantation on recipient insulin sensitivity, energy expenditure and intestinal transit time. <i>Gut</i> . 2020;69:502–12. doi: 10.1136/gutjnl-2019-318320
582 583 584	50	Barton W, Penney NC, Cronin O, <i>et al.</i> The microbiome of professional athletes differs from that of more sedentary subjects in composition and particularly at the functional metabolic level. <i>Gut.</i> 2018;67:625–33. doi: 10.1136/gutjnl-2016-313627

- 585 51 Clarke SF, Murphy EF, O'Sullivan O, *et al.* Exercise and associated dietary extremes impact 586 on gut microbial diversity. *Gut.* 2014;63:1913–20. doi: 10.1136/gutjnl-2013-306541
- 52 Shah S, Mu C, Moossavi S, *et al.* Physical activity-induced alterations of the gut
 microbiota are BMI dependent. *FASEB J Off Publ Fed Am Soc Exp Biol.* 2023;37:e22882.
 doi: 10.1096/fj.202201571R
- 53 Hanssen NMJ, de Vos WM, Nieuwdorp M. Fecal microbiota transplantation in human
 metabolic diseases: From a murky past to a bright future? *Cell Metab*. 2021;33:1098–
 110. doi: 10.1016/j.cmet.2021.05.005
- 54 Kootte RS, Levin E, Salojärvi J, *et al.* Improvement of Insulin Sensitivity after Lean Donor
 Feces in Metabolic Syndrome Is Driven by Baseline Intestinal Microbiota Composition. *Cell Metab.* 2017;26:611-619.e6. doi: 10.1016/j.cmet.2017.09.008
- 55 Sommer F, Ståhlman M, Ilkayeva O, *et al.* The Gut Microbiota Modulates Energy
 Metabolism in the Hibernating Brown Bear Ursus arctos. *Cell Rep.* 2016;14:1655–61. doi:
 10.1016/j.celrep.2016.01.026
- 56 Aoi W, Inoue R, Mizushima K, *et al.* Exercise-acclimated microbiota improves skeletal
 muscle metabolism via circulating bile acid deconjugation. *iScience*. 2023;26:106251. doi:
 10.1016/j.isci.2023.106251
- Lai Z-L, Tseng C-H, Ho HJ, *et al.* Fecal microbiota transplantation confers beneficial
 metabolic effects of diet and exercise on diet-induced obese mice. *Sci Rep.* 2018;8:15625.
 doi: 10.1038/s41598-018-33893-y

605