#### Prevalence and Trends in Cardiovascular Risk Factors Among Middle Aged Persons from Five Race and Ethnic Groups in the United States: A Longitudinal Analysis of Two Cohort **Studies**

- Havisha Pedamallu\*, BA<sup>1</sup>, Zeynab Aghabazaz\*, PhD<sup>1</sup>, Nicola Lancki, MPH<sup>1</sup>, Luis A. Rodriguez, PhD, MPH, RD<sup>2,3,4</sup>, Juned Siddique, DrPH<sup>1</sup>, Meena Moorthy, BA<sup>1</sup>, Nilay S. Shah,
- MD, MPH<sup>1</sup>, Norrina B. Allen, PhD<sup>1</sup>, Alka M. Kanaya, MD<sup>2</sup>, Namratha R. Kandula, MD, MPH<sup>1</sup>

- <sup>1</sup> Northwestern University Feinberg School of Medicine, Chicago, Illinois
- <sup>2</sup> University of San Francisco in California, San Francisco, California
- <sup>3</sup> Kaiser Permanente Northern California Division of Research, Oakland, California
- <sup>4</sup> Kaiser Permanente Bernard J. Tyson School of Medicine, Oakland, California
- \*These authors contributed equally to this work
- Word Count: 3,263

18	<b>Corresponding Author:</b>	Namratha R. Kandula, MD, MPH
19		675 N St Clair St Ste 18-200,
20		Chicago, IL 60611
21		Phone: 312-695-8630
22		Email: n-kandula@northwestern.edu
23		

29	Key Points (90/100 words)
30	
31	Question: Do South Asian adults have a higher burden of cardiovascular risk factors at age 45
32	years compared adults from other race and ethnic groups?
33	
34	Findings: In this study of 2754 adults from two cohort studies, the prevalence of prediabetes and
35	diabetes at age 45 years was higher among South Asians than in Black, Chinese, Hispanic and
36	White adults; hypertension prevalence was higher among South Asians than all groups except
37	Black adults.
38	
39	Meaning: South Asian adults have a higher prevalence of several clinical cardiovascular risk
40	factors at a younger age.
41	
42	

43	Abstract (348/350 words)
44	
45	Importance: It is well established that people of South Asian background have a high burden of
46	atherosclerotic cardiovascular disease (ASCVD). However, few studies have comprehensively
47	examined if South Asian adults in the United States (US) develop cardiovascular risk factors at
48	younger ages than adults from other race and ethnic groups.
49	
50	Objective: To compare the prevalence and change in ASCVD risk factors across age strata by
51	race and ethnic group.
52	
53	Design: We combined data from the Mediators of Atherosclerosis in South Asians Living in
54	America (MASALA) and the Multi-Ethnic Study of Atherosclerosis (MESA) cohort studies.
55	Longitudinal data from all eligible participants at all available exam visits were used to estimate
56	the prevalence of risk factors at ages 45 and 55 years for each race and ethnic group.
57	
58	Setting: Multicenter longitudinal cohort study in 7 field centers across the U.S.
59	
60	Participants: The baseline study sample included individuals free of clinical ASCVD; 554 South
61	Asians, 796 White, 588 Black, 517 Hispanic/Latino, and 245 Chinese adults aged 45-55 years
62	were included.
63	
64	Exposures: Self-identified race (Black, Chinese, South Asian, or White) or ethnic group
65	(Hispanic/Latino).

66

67	Main Outcome(s) and Measure(s): Prevalence of clinical (prediabetes & diabetes,
68	hypertension, dyslipidemia, BMI) and behavioral (diet quality, alcohol use, exercise) ASCVD
69	risk factors at age 45 and age 55.
70	
71	Results: At age 45, South Asian men and women had the highest prevalence of pre-diabetes and
72	diabetes and higher prevalence of hypertension compared to White, Chinese, and
73	Hispanic/Latino men and women. South Asian men had a higher prevalence of dyslipidemia than
74	White, Chinese, and Black men, and South Asian women had a higher prevalence than Chinese
75	and Black women. All groups had worse diet quality than South Asian men and women at age
76	45, and most also had higher rates of alcohol use.
77	
78	Conclusions and Relevance: We observed significant differences in the prevalence of risk
78 79	<b>Conclusions and Relevance:</b> We observed significant differences in the prevalence of risk factors for South Asian adults compared to adults from other U.S. race and ethnic groups at age
78 79 80	Conclusions and Relevance: We observed significant differences in the prevalence of risk factors for South Asian adults compared to adults from other U.S. race and ethnic groups at age 45 years. Understanding trends and disparities in cardiovascular risk and protective factors
78 79 80 81	Conclusions and Relevance: We observed significant differences in the prevalence of riskfactors for South Asian adults compared to adults from other U.S. race and ethnic groups at age45 years. Understanding trends and disparities in cardiovascular risk and protective factorsacross the life course can help equitably improve prevention and treatment strategies for US
78 79 80 81 82	Conclusions and Relevance: We observed significant differences in the prevalence of risk factors for South Asian adults compared to adults from other U.S. race and ethnic groups at age 45 years. Understanding trends and disparities in cardiovascular risk and protective factors across the life course can help equitably improve prevention and treatment strategies for US populations.
78 79 80 81 82 83	Conclusions and Relevance: We observed significant differences in the prevalence of risk factors for South Asian adults compared to adults from other U.S. race and ethnic groups at age 45 years. Understanding trends and disparities in cardiovascular risk and protective factors across the life course can help equitably improve prevention and treatment strategies for US populations.
78 79 80 81 82 83 83	Conclusions and Relevance: We observed significant differences in the prevalence of risk factors for South Asian adults compared to adults from other U.S. race and ethnic groups at age 45 years. Understanding trends and disparities in cardiovascular risk and protective factors across the life course can help equitably improve prevention and treatment strategies for US populations.
78 79 80 81 82 83 83 84 85	Conclusions and Relevance: We observed significant differences in the prevalence of risk factors for South Asian adults compared to adults from other U.S. race and ethnic groups at age 45 years. Understanding trends and disparities in cardiovascular risk and protective factors across the life course can help equitably improve prevention and treatment strategies for US populations.
78 79 80 81 82 83 83 84 85 86	Conclusions and Relevance: We observed significant differences in the prevalence of risk factors for South Asian adults compared to adults from other U.S. race and ethnic groups at age 45 years. Understanding trends and disparities in cardiovascular risk and protective factors across the life course can help equitably improve prevention and treatment strategies for US populations.
78 79 80 81 82 83 83 84 85 86 86	Conclusions and Relevance: We observed significant differences in the prevalence of risk factors for South Asian adults compared to adults from other U.S. race and ethnic groups at age 45 years. Understanding trends and disparities in cardiovascular risk and protective factors across the life course can help equitably improve prevention and treatment strategies for US populations.

# 89 INTRODUCTION

90	South Asian individuals (people who trace their ancestry from Bangladesh, Bhutan, India,
91	Maldives, Nepal, Pakistan, and Sri Lanka) have high rates of atherosclerotic cardiovascular
92	disease (ASCVD), with estimates approximating rates 2.5 times higher compared to non-
93	Hispanic White and East Asian populations <sup>1</sup> . On average, South Asians develop coronary artery
94	disease (CAD) up to a decade earlier than the average age at which CAD develops in most
95	people in the United States (US) <sup>2</sup> . A quarter of heart attacks occur under age 40 for South Asians
96	and 50% occur under age 50. In clinical guidelines, South Asian ancestry is considered a "risk
97	enhancing factor" <sup>3</sup> .
98	While South Asian in the US have an overall higher prevalence of ASCVD compared to
99	non-Hispanic White US populations that manifests at younger ages, there is limited research to
100	know if the prevalence of ASCVD risk factors at younger ages is higher among South Asian
101	populations compared to other race and ethnic groups. Therefore, the objective of this study was
102	to combine longitudinal data from the Mediators of Atherosclerosis in South Asians Living in
103	America (MASALA) study and the Multiethnic Study of Atherosclerosis (MESA) to estimate the
104	prevalence of cardiovascular risk factors during middle age, and to test whether the prevalence of
105	these risk factors and their change over time differ between South Asians compared to Black,
106	Chinese, Hispanic/Latino and White adults. If South Asians develop ASCVD risk factors at
107	younger ages than other race and ethnic groups, this finding could lead to earlier screening,
108	prevention, and treatment for this at-risk group.
109	
110	

111

## 112 METHODS

113	Study Population. This study was conducted using longitudinal data from all available
114	exams of the Mediators of Atherosclerosis in South Asians Living in America (MASALA) and
115	the Multi-Ethnic Study of Atherosclerosis (MESA) studies, which are both multicenter
116	prospective cohort studies examining the development and determinants of clinical ASCVD risk
117	factors, subclinical ASCVD, and ASCVD outcomes.
118	The study design for MASALA has been published elsewhere <sup>4</sup> . Briefly, MASALA
119	included 1164 South Asian (defined as those with Indian, Pakistani, Bangladeshi, Nepali, and Sri
120	Lankan ancestry) participants who were CVD free between 40 to 84 years of age recruited
121	through two field centers in San Francisco, CA and Chicago, IL; Data were collected at baseline
122	(Exam 1) for 906 participants in 2010-2013 with follow up Exam 2 data collection in 2016-2018
123	for 748 participants. An additional 258 participants were recruited in 2017-2018 and their
124	baseline (Exam 1A) data are included.
125	The study design for MESA has also been published <sup>5</sup> . Briefly, MESA included 6,814
126	White, Black, Hispanic/Latino, and Chinese men and women free from clinical CVD between
127	45-84 years of age. Baseline exam 1 data were collected between 2000-2002 with repeat
128	examinations every 2 years. There have been 6 exams, the most recent being 2016-2018.
129	Participants were recruited through 6 field centers: Forsyth County, NC; Northern Manhattan and
130	the Bronx, NY; Baltimore City and Baltimore County, MD; St. Paul, MN; Chicago and the
131	village of Maywood, IL; and Los Angeles County, CA.
132	Study Sample and Exclusion Criteria. The analysis was restricted to participants
133	between ages 45 to 55 at the baseline exam to minimize bias related to the cohorts' inclusion
134	criteria of being free from ASCVD at baseline. It is likely that the older individuals enrolled in

135 MASALA and MESA may not be representative of the general older population since they were 136 required to be free of clinical symptoms of ASCVD and ASCVD is more common at older ages. 137 Therefore, we included 554 South Asians from the MASALA study and 796 White, 588 Black, 138 517 Hispanic/Latino, and 245 Chinese adults from the MESA study. Harmonized data from MASALA Exams 1/1A and 2 and MESA Exams 1 through 6 were combined. On average, our 139 140 data set consisted of 3.3 exams per MESA participant and 1.4 exams per MASALA participant. 141 Primary Exposure. The primary exposure was self-reported race (Black, Chinese, South 142 Asian, or White) or ethnic group (Hispanic/Latino). **Primary Outcome.** The primary outcomes were clinical and behavioral ASCVD risk 143 144 factors. These risk factors were chosen based on prior research establishing important risk factors in the development of ASCVD and the American Heart Association's Life's Essential 8<sup>6</sup>. Risk 145 146 factor definitions were previously harmonized between the two data sets using a standardized set

of definitions<sup>7-9</sup>. All risk factors were available at all exams, except diet and exercise at MESA 147

148 exam 4. Diet was assessed for 87% of MESA participants at Exam 1, 72% of MESA participants

149 at exam 5, 99% of MASALA participants at Exam 1, and 71% of MASALA participants at Exam

150 2.

151 Clinical risk factors include hypertension, pre-diabetes or diabetes status, dyslipidemia, 152 and body mass index (BMI). Hypertension was defined as having a systolic blood pressure 153  $\geq$ 140 mmHg, diastolic blood pressure  $\geq$ 90 mmHg, or use of any antihypertensive medication according to guidelines in use at the time of the baseline exam<sup>10</sup>. Prediabetes was defined as 154 155 having a fasting plasma glucose level between 100 and 125 mg/dL. Diabetes status was defined by self-reported use of glucose-lowering medications or fasting plasma glucose  $\geq 126$  mg/dL<sup>11</sup>. 156 157 Dyslipidemia was determined using standard lipid level cut points (total cholesterol  $\geq 200$ 

158	mg/dL, low-density lipoprotein $\geq 160$ mg/dL, triglycerides $\geq 150$ mg/dL, or high-density
159	lipoprotein < 40 mg/dL) using blood samples or self-report of currently taking lipid lowering
160	therapy- statin/fibrate/niacin <sup>12</sup> . BMI was calculated using weight and height measurements at
161	each exam visit and was treated as a continuous variable in the analysis. BMI at Exam 1 was
162	reported as a categorical variable using race and ethnic group-specific cut-points (Chinese and
163	South Asian Individuals: underweight BMI < 18.5, normal BMI 18.5 to 22.9, overweight 23 to
164	24.9, and obese BMI $\geq$ 25; All others: underweight BMI < 18.5, normal BMI 18.5 to 24.9,
165	overweight 25 to 29.9, and obese $BMI \ge 30$ ) <sup>13, 14</sup> .
166	Behavioral risk factors included alcohol use, physical activity level, and diet quality.
167	Alcohol use was defined as currently drinking 1 or more drinks per week. Participants reported
168	exercise using the Typical Week's Physical Activity Questionnaire which was quantified as MET
169	minutes/week <sup>15</sup> . Diet quality was determined by calculating the Alternative Health Eating Index-
170	2010 (AHEI) score <sup>9</sup> . This is a dietary quality index that predicts the risk of chronic disease based
171	on nutrient and food intake, with a range between 0 (worst) to 110 (best) points <sup>9</sup> .
172	Statistical Analysis. Baseline characteristics (age, smoking status, education, insurance
173	status, BMI) were calculated for each race and ethnic group. Comparisons across groups were
174	tested using one-way ANOVAs for continuous variables and chi-squared tests of independence
175	for categorical variables.
176	Our estimands of interest are the risk factor prevalences (for binary outcomes) or means
177	(for continuous outcomes) at age 45 and their change over 10 years for each race and ethnic
178	group and their differences between race and ethnic groups.
179	Using data from all eligible MESA and MASALA participants, we modeled clinical and
180	behavioral risk factors as a function of age using mixed-effects linear regression models for

continuous outcomes (BMI, physical activity, diet quality) and mixed-effects logistic regression
models for binary outcomes (all other outcomes). Smoking status was not modeled as a
longitudinal risk factor because it tended not to vary over time and cannot be modeled at age 45
and 55. All models controlled for baseline education and baseline insurance status and were fit
separately by risk factor and sex.
To estimate risk factor prevalences at age 45 and their 10-year change, we used the

method of marginal predicted means<sup>16</sup> - the average value of the outcome averaging over all
covariates—to estimate the prevalence of each risk factor at ages 45 and 55 for each race and
ethnic group. These marginal predicted means were obtained from our mixed-effects models that
use data from all eligible MESA and MASALA participants.

191 To compare risk factor prevalence between South Asians and the other race and ethnic 192 groups, we first performed—separately by risk factor—an overall test at age 45 of whether the 193 South Asian prevalence was significantly different from any of the other race and ethnic groups. 194 If that test was significant, we subsequently performed the four pairwise tests comparing South 195 Asians to each of the other race and ethnic groups.

196 To test if the change in risk factors between ages 45 and 55 differed for South Asians 197 compared to the other groups, we calculated the difference (slope) between marginal means at 198 ages 45 and 55 (i.e 55 minus 45) for all five race and ethnic groups. Then, we again performed 199 an overall test of whether the South Asian slope was significantly different from the slopes of 200 any of the other race and ethnic groups. If this test was significant, we then performed the four 201 pairwise tests comparing the South Asian slope to the slope of each race and ethnic group. More 202 information on estimation of risk factor prevalences, their change over time, and their 203 comparison across race and ethnic groups in included in the Appendix. Since this is an

204 exploratory analysis of observational data, we did not correct for multiple comparisons. All

analyses and post-estimation processing were performed using STATA version  $18.0^{17}$ .

206 RESULTS

207 **Baseline Characteristics.** The average age of South Asian participants was 48.5 years at 208 the baseline exam which was similar when compared to the other race and ethnic groups (Table 209 1). South Asian participants overall had higher education levels compared to the other race and 210 ethnic groups. A majority (60.8%) of South Asians had more than a college level education 211 compared to 32.9% of White participants, the next highest educated group. South Asians also 212 had the highest never smoking prevalence (85.2%) at baseline with Chinese adults having a 213 slightly lower prevalence (76.7%). Hispanic/Latino (53.0%), Black (48.0%), and White (45.5%) 214 had much lower never smoking prevalences in comparison. A higher proportion of South Asian 215 (86.8%), White (91.7%), and Black (81.0%) participants had private insurance compared to 216 Chinese (68.2%) and Hispanic/Latino (67.3%) participants. When using race and ethnic group-217 specific BMI cut points, 20.9% of South Asian, 34.6% of White, 35.5% of Chinese, 15.8% of 218 Black, and 16.1% of Latino participants had an underweight or normal BMI, 219 Risk Factor Prevalence at Age 45 among Men. South Asian men had the highest

prevalence of pre-diabetes and diabetes (41.8%; Table 2) when compared to men of all other groups (White: 7.2%, Chinese: 18.9%, Black: 16.0%, Hispanic/Latino: 14.7%; p < 0.001). They had a significantly greater prevalence of hypertension compared to Chinese and Hispanic/Latino men and a significantly greater prevalence of dyslipidemia compared to Black men. They also had a significantly lower BMI than White, Black, and Hispanic/Latino men. South Asian men had significantly higher diet quality scores compared to men of all other race and ethnic groups

(Table 4). They also had significantly lower alcohol use than Chinese men and higher level ofexercise than White and Chinese men.

228 Trends Across Age Among Men. There were no significantly different changes in 229 prevalence of clinical risk factors between age 45 and 55 for South Asian men versus men from 230 the other groups, with the exception of White men having a smaller increase in the prevalence of 231 hypertension between age 45 and 55 (21.9% for South Asian men, 11.6% for White men, p =232 0.05). Among the behavioral risk factors, White men also had a larger improvement in AHEI diet 233 score (7.1) than South Asian men (3.4, p = 0.006) over time. Chinese men had a larger increase in exercise (1605 MET min/wk) than South Asian men (355 MET min/wk, p = 0.011). 234 235 Risk Factor Prevalence at Age 45 among Women. Similar to South Asian men, South 236 Asian women had the highest prevalence of pre-diabetes and diabetes of women of all race and 237 ethnic groups (Table 3). South Asian women also had the highest hypertension prevalence, and a 238 significantly higher prevalence of dyslipidemia compared to Chinese and Black women. BMI 239 among South Asian women was significantly higher than BMI in Chinese women and 240 significantly lower than Black and Hispanic/Latino women. South Asian women had the highest 241 AHEI score (68.5) of all groups (White: 51.5, Chinese: 58.6, Black: 47.7, Hispanic/Latino: 50.3), 242 the lowest alcohol use, and the most exercise.

243Trends Across Age Among Women. South Asian women had a significantly smaller244increase in HTN prevalence (4.7%) from age 45 to 55 compared to Chinese (17.2%, p < 0.01),245Black (31.0%, p < 0.001), and Latina (14.9%, p < 0.01) women. South Asian women had larger246increases in BMI (2.0) compared to White (1.0, p = 0.018) and Chinese (-0.2, p < 0.001) women.247White women had a larger increase in AHEI diet score (7.0) compared to South Asian women

(3.4, p = 0.006). In addition, Black women had a larger increase in exercise (962 MET min/wk)
than South Asian women (203 MET min/wk).

250 DISCUSSION

251 **Principal Findings.** In this longitudinal multi-race and ethnic population-based cohort 252 study of 2,700 adults ages 45-55 at baseline in the U.S., we observed significant differences in 253 the prevalence of risk factors for South Asians compared to other U.S. race and ethnic groups at 254 age 45 years. At age 45, South Asian adults had a higher prevalence of prediabetes, diabetes, and 255 hypertension than White adults. Namely, the prevalence of pre-diabetes and diabetes among 256 South Asian men and women is almost twice the prevalence of Black and White individuals. 257 Despite observing a higher prevalence of clinical risk factors among South Asian adults at 45 258 years, we also found that South Asian adults had the highest (i.e. best) diet quality of all race and 259 ethnic groups, as well as lower alcohol use and higher exercise levels than most other groups. 260 Interestingly, these findings differ from past research showing that South Asians are less likely to meet physical activity recommendations and eat less healthful diets.<sup>18</sup> However, prior studies 261 262 included participants across the adult lifespan, , and the present analysis only included South 263 Asian adults during middle-age who we found to have more cardioprotective behaviors.

Secondary Findings. Overall, it appears that as age increases, the prevalence of clinical risk factors increases significantly among men and women. However, as age increased, we also observed a favorable increase in diet quality among all groups and exercise in all groups except for South Asian women. These findings indicate that the increase in ASCVD risk factors among South Asians as they age is in alignment with other groups, but that South Asians may develop some risk factors earlier than other groups. In addition, while health behaviors improved with

age among White, Black, Hispanic/Latino, and Chinese adults, the same level of improvementwas not seen among South Asian adults.

272 There are multiple factors that could be contributing to this trend. For example, there may 273 be social and cultural underpinnings to differences in health behaviors as individuals age. The 274 majority of middle- and older-age South Asians in the US are immigrants, and immigration and acculturation have been shown to impact diet and physical activity <sup>19-21</sup>. Language and cultural 275 276 barriers may also contribute to reduced health literacy among South Asians living in the U.S. and 277 preventive health messaging reaching other race and ethnic groups may not be reaching South Asians effectively<sup>22</sup>. In addition, clinical ASCVD risk factors are chronic health conditions 278 279 which are associated with disability and functional decline. The increased burden of ASCVD risk 280 factors among South Asians in middle-age compared to other groups may be impacting their 281 ability to engage in positive health behaviors like increased physical activity.

282 Previous studies have shown that the number of positive health behaviors at middle-age is associated with less frailty at older ages<sup>23</sup>. In addition, increasing the number of positive health 283 behaviors a person engages in during middle-age is associated with a lower risk of frailty<sup>23</sup>. 284 285 Given this finding, strategies to enhance healthy behaviors among aging South Asian adults may 286 be needed to reduce disparities. Future studies should replicate this study design in younger 287 populations to determine the prevalence of clinical and behavioral cardiovascular risk factors 288 prior to middle-age and how behavioral and clinical risk factors may change over time to 289 influence ASCVD outcomes. It is possible that health behaviors at younger ages are more 290 predictive of health status at older ages compared to behaviors at middle-age.

Implications. This is the first study, to our knowledge, to show that the increased burden
of ASCVD in South Asians may be attributed to a higher prevalence of clinical risk factors at

younger ages. The results of this study indicate that screening and prevention of ASCVD risk
must begin early for people of South Asian background given the increased burden of risk factors
at younger ages. Despite having the highest diet quality of all the groups included in this study
and high levels of exercise at younger ages, South Asians still had a high prevalence of ASCVD
risk factors at young ages, especially diabetes and hypertension. Determining the age at which
risk factors screening, prevention strategies, and treatment can optimize health outcomes for
South Asian adults is fundamental to addressing their increased ASCVD risk.

300 **Limitations.** In this study, participants had to be free of cardiovascular disease when they 301 were enrolled in the study, so this data is potentially excluding individuals who may have early 302 ASCVD and could add to the burden of risk factors. Also, whereas MASALA and MESA offer 303 longitudinal information regarding patient populations that have been traditionally 304 underrepresented in research, the participants in these studies are likely not fully representative 305 of all people of that race or ethnic group living in the United States. For example, MASALA 306 only recruited from two study sites, all of which are large urban areas, excluding many 307 individuals who may add more diversity to the data collected. In addition, longitudinal data 308 heavily relies on patient follow up. Participants with higher educational attainment, have higher socioeconomic status, and have higher health motivation will be more likely to follow up<sup>24</sup>. It is 309 310 also likely that differences between participants in MESA and MASALA are confounded by time 311 period-based differences, as there is a decade difference between the baseline exams of both studies. 312

313 Conclusions. This study suggests that South Asians have a higher prevalence of several
314 clinical risk factors at age 45, while having more favorable cardiovascular health behaviors,
315 compared to other race and ethnic groups. Given the relatively high observed burden of clinical

316	risk factors in South Asians, future studies should examine the development of risk factors
317	among younger South Asians to better characterize the development and burden of risk earlier in
318	the life course. Given that heterogeneity in risk factors already existed at age 45, future studies
319	should aim to examine the entire life course and identify critical time periods where the burden
320	of risk factors stratifies.
321	Further research is needed to identify mechanisms of the higher observed burden of
322	clinical risk factors among U.S. South Asians at younger ages despite more favorable
323	cardiovascular health behaviors. Understanding trends and disparities in risk and protective
324	factors across the life course can help equitably improve prevention and treatment strategies for
325	US populations.
326	ACKNOWLEDGEMENTS
327	The authors thank the other investigators, the staff, and the participants of the MESA and
328	MASALA studies for their valuable contributions.
329	Funding Sources: The MESA study was supported by contracts N01-HC-95159, N01-HC-
330	95160, N01-HC-95161, N01-HC-95162, N01-HC-95163, N01-HC-95164, N01-HC-95165, N01-
331	HC-95166, N01-HC-95167, N01-HC-95168 and N01-HC-95169 from the National Heart, Lung,
332	and Blood Institute and by grants UL1-TR-000040, UL1 TR 001079 and UL1-RR-025005 from
333	NCRR. The MASALA study was supported by Grants R01HL093009 and R01HL120725 from
334	the National Heart, Lung, and Blood Institute, the National Center for Research Resources, and
335	the National Center for Advancing Translational Sciences, the National Institutes of Health
336	(NIH); and through the UCSF-CTSI Grants UL1RR024131 and UL1TR001872.
337	Human Subjects Statement: The Multi-Ethnic Study of Atherosclerosis (MESA) was approved
338	by the institutional review boards at each of the study sites.

Table 1: Baseline	Characteristics at E	xam 1, MASALA	A and MESA studies.	Values are N (%)	unless otherwise
reported.					

	South Asian	White	Chinese	Black	Hispanic/Latino	P-value
	(n =554)	(n = 796)	(n = 245)	(n = 588)	(n = 517)	
Age, Mean (SD)	48.5 (4.4)	50.4 (3.1)	50.1 (3.2)	50.3 (3.0)	49.9 (3.3)	< 0.001
Sex						0.665
Female	287 (51.8)	426 (53.5)	125 (51.0)	325 (55.3)	267 (51.6)	
Male	267 (48.2)	370 (46.5)	120 (49.0)	263 (44.7)	250 (48.4)	
Education						< 0.001
< HS	15 (2.7)	11 (1.4)	40 (16.3)	34 (5.8)	170 (32.9)	
HS	22 (4.0)	75 (9.4)	35 (14.3)	90 (15.3)	101 (19.5)	
< College	30 (5.4)	239 (30.0)	49 (20.0)	235 (34.0)	178 (34.4)	
College	150 (27.1)	208 (26.1)	65 (26.5)	133 (22.6)	40 (7.7)	
> College	337 (60.8)	262 (32.9)	56 (22.9)	93 (15.8)	28 (5.4)	
Missing	0	1 (0.1)	0	3 (0.5)	0	
Insurance						< 0.001
None	48 (8.7)	30 (3.8)	60 (24.5)	70 (11.9)	123 (23.8)	
Public	25 (4.5)	35 (4.4)	18 (7.4)	39 (6.6)	46 (8.9)	
Private	481 (86.8)	730 (91.7)	167 (68.2)	476 (81.0)	348 (67.3)	
Missing	0	1 (0.1)	0	3 (0.51)	0	
BMI						< 0.001
Under/Normal						
weight	116 (20.9)	275 (34.6)	87 (35.5)	93 (15.8)	83 (16.1)	
Overweight	110 (19.9)	276 (34.7)	65 (26.5)	197 (33.5)	226 (43.7)	
Obese	328 (59.2)	245 (30.8)	93 (38.0)	298 (50.7)	208 (40.2)	
Smoking						< 0.001
Never	472 (85.2)	362 (45.48)	188 (76.73)	282 (47.96)	274 (53.00)	
Former	63 (11.4)	297 (37.31)	39 (15.92)	165 (28.06)	133 (25.73)	
Current	19 (3.43)	136 (17.09)	18 (7.35)	138 (23.47)	110 (21.28)	
Missing	0	1 (0.13)	0	3 (0.51)	0	

	Age	Group vs SA at age	P-value*	Age	Change from age	Difference in	P-value**
	45	45 (CI)		55	45 to 55 (CI)	Change (CI)	
Pre-diabetes &							
Diabetes(%)							0.374
South Asian	41.8	(Ref)		69.4	27.6 (18.9,36.4)	(Ref)	
White	7.2	-34.6 (-43.3,-25.9)	< 0.001	27.8	20.6 (17.3,24.0)	-7.0 (-16.4,2.4)	
Chinese	18.9	-22.9 (-34.8,-11.0)	< 0.001	35.6	16.7 (8.2,25.3)	-10.9 (-23.4,1.5)	
Black	16.0	-25.7 (-36.3,-15.2)	< 0.001	33.3	17.2 (10.8,23.7)	-10.3 (-21.4,0.7)	
Hispanic/Latino	14.7	-27.0 (-37.6,-16.4)	< 0.001	36.5	21.7 (15.5,28.0)	-5.9 (-16.7,5.0)	
Hypertension(%)							
South Asian	25.2	(Ref)		47.2	21.9 (12.7,31.1)	(Ref)	
White	18.7	-6.5 (-13.0,-0.1)	0.047	30.3	11.6 (6.3,16.9)	-10.3 (-20.6,0.0)	0.050
Chinese	6.7	-18.5 (-25.2,-11.9)	< 0.001	26.2	19.4 (14.1,24.8)	-2.5 (-13.1,8.2)	0.649
Black	27.7	2.5 (-3.3,8.3)	0.401	52.4	24.6 (18.0,31.2)	2.7 (-8.6,14.0)	0.642
Hispanic/Latino	10.9	-14.3 ( -21.5,-7.2)	< 0.001	28.9	18.0 (12.4,23.5)	-3.9 (-14.5,6.6)	0.464
Dyslipidemia(%)							0.307
South Asian	78.5	(Ref)		83.0	4.4 (-2.8,11.8)	(Ref)	
White	73.7	-4.7 (-12.2,2.7)	0.213	76.3	2.5 (-2.7,7.7)	-1.9 (-10.9,7.0)	
Chinese	67.3	-11.2 (-22.4,0.1)	0.051	71.1	3.8 (-5.8,13.4)	-0.6 (-12.8,11.3)	
Black	59.4	-19.0 (-2.9,-8.5)	< 0.001	69.9	10.4 (1.6,19.2)	5.9 (-5.5,17.3)	
Hispanic/Latino	79.2	0.7 (-7.9,9.3)	0.869	77.8	-1.4 (-7.8,5.0)	-5.9 (-15.6,3.8)	
BMI (kg/m <sup>2</sup> )							0.213
South Asian	26.4	(Ref)		27.0	0.7 (0.1,1.2)	(Ref)	
White	27.7	1.3 (0.6,2.1)	< 0.001	28.6	0.9 (0.7,1.2)	0.3 (-0.3,0.9)	
Chinese	24.5	-1.9 (-2.9,-0.8)	< 0.001	25.1	0.5 (0.1,1.0)	-0.1 (-0.8,0.6)	
Black	28.3	1.9 (1.1,2.8)	< 0.001	29.5	1.2 (0.8,1.5)	0.5 (-0.1,1.1)	
Hispanic/Latino	28.4	2.1 (1.1,3.0)	< 0.001	29.2	0.8 (0.5,1.1)	0.3 (-0.5,0.8)	

**Table 2:** Clinical Risk Factors Stratified by Age and Race and Ethnic Group in Men, MASALA vs. MESA Groups

\* P-value < 0.05 from a pairwise test at age 45 vs. South Asians; \*\* P-value from a test of change from age 45 to 55 vs. South Asian. When the multivariate test was not significant, pairwise tests were not performed.

Age	Group vs SA at age	P-value*	Age	Change from age	Difference in	P-value**
45	45 (CI)		55	45 to 55 (CI)	Change (CI)	
27.2	(Ref)		45.4	18.2 (10.1,26.3)	(Ref)	0.150
8.1	-19.1 (-24.2,-13.9)	< 0.001	19.8	11.7 (8.1,15.2)	-6.6 (-15.5,2.4)	0.444
5.8	-21.4 (-26.9,-15.8)	< 0.001	27.6	21.8 (17.5,26.1)	3.5 (-5.6,12.7)	0.054
17.4	-9.7 (-16.3,-3.2)	0.003	26.4	9.0 (0.039,14.0)	-9.3 (-18.7,0.1)	0.400
8.1	-19.1 (-24.8,-13.3)	< 0.001	30.2	22.1 (17.8,26.4)	3.9 (-5.2,13.0)	
24.6	(Ref)		29.3	4.7 (0.1,9.2)	(Ref)	
18.6	-6.0 (-11.8,-0.3)	0.040	28.5	9.8 (5.5,14.3)	52 (-1.0,11.3)	0.095
7.5	-17.1 (-24.6,-9.6)	< 0.001	24.7	17.2 (10.9,23.5)	12.4 (4.7,20.3)	0.002
30.1	5.5 (0.1,10.9)	0.047	61.2	31.0 (26.1,36.0)	26.4 (19.5,33.2)	< 0.001
15.1	-9.5 (-16.7,-2.3)	0.010	30.0	14.9 (9.1,20.8)	10.2 (3.0,17.4)	0.005
						0.147
54.3	(Ref)		74.7	20.4 (11.0,29.8)	(Ref)	
55.5	1.2 (-9.6,12.0)	0.827	71.4	15.9 (8.9,22.9)	-4.5 (-16.2,7.1)	
39.4	-14.9 (-29.4,-0.3)	0.046	68.6	29.2 (17.0,41.4)	8.8 (-6.6,24.2)	
39.2	-15.0 (-26.3,-3.7)	0.009	62.1	22.8 (15.0,30.7)	2.4 (-9.8,14.6)	
54.9	0.6 (-11.6,12.9)	0.917	67.4	12.5 (4.2,20.8)	-7.9 (-20.5,4.6)	
26.4	(Ref)		28.3	2.0 (1.2,2.7)	(Ref)	
27.3	0.9 (-0.1,2.0)	0.086	28.3	1.0 (0.7,1.3)	-1.0 (-1.9,-0.2)	0.018
23.9	-2.4 (-3.9,-1.0)	0.001	23.7	-0.2 (-0.8,0.4)	-2.2 (-3.2,-1.3)	< 0.001
31.3	5.0 (3.8,6.1)	< 0.001	32.7	1.4 (1.0,1.8)	-0.6 (-1.5,0.2)	0.149
29.6	3.3 (2.0,4.6)	< 0.001	30.8	1.2 (0.8,1.6)	-0.8 (-1.7,0.0)	0.062
	Age         45         27.2         8.1         5.8         17.4         8.1         24.6         18.6         7.5         30.1         15.1         54.3         55.5         39.4         39.2         54.9         26.4         27.3         31.3         29.6	AgeGroup vs $3A$ at age4545 (CI)27.2(Ref)8.1-19.1 (-24.2,-13.9)5.8-21.4 (-26.9,-15.8)17.4-9.7 (-16.3,-3.2)8.1-19.1 (-24.8,-13.3)24.6(Ref)18.6-6.0 (-11.8,-0.3)7.5-17.1 (-24.6,-9.6)30.15.5 (0.1,10.9)15.1-9.5 (-16.7,-2.3)54.3(Ref)55.51.2 (-9.6,12.0)39.4-14.9 (-29.4,-0.3)39.2-15.0 (-26.3,-3.7)54.90.6 (-11.6,12.9)26.4(Ref)27.30.9 (-0.1,2.0)23.9-2.4 (-3.9,-1.0)31.35.0 (3.8,6.1)29.63.3 (2.0,4.6)	AgeGloup vs SA at ageF-value $45$ $45$ (CI) $27.2$ (Ref) $8.1$ $-19.1$ (-24.2,-13.9) $<0.001$ $5.8$ $-21.4$ (-26.9,-15.8) $<0.001$ $17.4$ $-9.7$ (-16.3,-3.2) $0.003$ $8.1$ $-19.1$ (-24.8,-13.3) $<0.001$ $24.6$ (Ref) $18.6$ $-6.0$ (-11.8,-0.3) $0.040$ $7.5$ $-17.1$ (-24.6,-9.6) $<0.001$ $30.1$ $5.5$ (0.1,10.9) $0.047$ $15.1$ $-9.5$ (-16.7,-2.3) $0.010$ $54.3$ (Ref) $55.5$ $1.2$ (-9.6,12.0) $0.827$ $39.4$ $-14.9$ (-29.4,-0.3) $0.046$ $39.2$ $-15.0$ (-26.3,-3.7) $0.009$ $54.9$ $0.6$ (-11.6,12.9) $0.917$ $26.4$ (Ref) $27.3$ $0.9$ (-0.1,2.0) $0.086$ $23.9$ $-2.4$ (-3.9,-1.0) $0.001$ $31.3$ $5.0$ (3.8,6.1) $<0.001$ $29.6$ $3.3$ (2.0,4.6) $<0.001$	AgeGloup vs SA at ageF-valueAge $45$ $45$ (CI) $55$ $27.2$ (Ref) $45.4$ $8.1$ $-19.1$ (-24.2,-13.9) $<0.001$ $5.8$ $-21.4$ (-26.9,-15.8) $<0.001$ $27.4$ $-9.7$ (-16.3,-3.2) $0.003$ $26.4$ $8.1$ $-19.1$ (-24.8,-13.3) $24.6$ (Ref) $29.3$ $18.6$ $-6.0$ (-11.8,-0.3) $0.040$ $28.5$ $7.5$ $-17.1$ (-24.6,-9.6) $7.5$ $-17.1$ (-24.6,-9.6) $<0.001$ $24.7$ $30.1$ $5.5$ ( $0.1,10.9$ ) $0.047$ $61.2$ $15.1$ $-9.5$ (-16.7,-2.3) $0.010$ $30.0$ $30.0$ $54.3$ (Ref) $74.7$ $55.5$ $1.2$ (-9.6,12.0) $0.827$ $71.4$ $39.4$ $-14.9$ (-29.4,-0.3) $9.4$ $-14.9$ (-29.4,-0.3) $0.046$ $68.6$ $39.2$ $-15.0$ (-26.3,- $3.7$ ) $9.06$ (-11.6,12.9) $0.917$ $57.3$ $0.9$ (- $0.1,2.0$ ) $0.086$ $28.3$ $23.9$ $-2.4$ ( $-3.9,-1.0$ ) $0.001$ $23.7$ $31.3$ $5.0$ ( $3.8,6.1$ ) $<0.001$ $32.7$ $29.6$ $3.3$ ( $2.0,4.6$ ) $<0.001$ $32.7$	AgeChoup vs 3A at ageF-valueAgeChange from age $45$ $45$ (CI) $55$ $45$ to $55$ (CI) $27.2$ (Ref) $45.4$ $18.2$ (10.1,26.3) $8.1$ $-19.1$ (-24.2,-13.9) $<0.001$ $19.8$ $11.7$ (8.1,15.2) $5.8$ $-21.4$ (-26.9,-15.8) $<0.001$ $27.6$ $21.8$ (17.5,26.1) $17.4$ $-9.7$ (-16.3,-3.2) $0.003$ $26.4$ $9.0$ (0.039,14.0) $8.1$ $-19.1$ (-24.8,-13.3) $<0.001$ $30.2$ $22.1$ (17.8,26.4) $24.6$ (Ref) $29.3$ $4.7$ (0.1,9.2) $18.6$ $-6.0$ (-11.8,-0.3) $0.040$ $28.5$ $9.8$ (5.5,14.3) $7.5$ $-17.1$ (-24.6,-9.6) $<0.001$ $24.7$ $17.2$ (10.9,23.5) $30.1$ $5.5$ (0.1,10.9) $0.047$ $61.2$ $31.0$ (26.1,36.0) $15.1$ $-9.5$ (-16.7,-2.3) $0.010$ $30.0$ $14.9$ (9.1,20.8) $54.3$ (Ref) $74.7$ $20.4$ (11.0,29.8) $55.5$ $1.2$ (-9.6,12.0) $0.827$ $71.4$ $15.9$ (8.9,22.9) $39.4$ $-14.9$ (-29.4,-0.3) $0.046$ $68.6$ $29.2$ (17.0,41.4) $39.2$ $-15.0$ (-26.3,-3.7) $0.009$ $62.1$ $22.8$ (15.0,30.7) $54.9$ $0.6$ (-11.6,12.9) $0.917$ $67.4$ $12.5$ (4.2,20.8) $26.4$ (Ref) $28.3$ $2.0$ (1.2,2.7) $27.3$ $0.9$ (-0.1,2.0) $0.086$ $28.3$ $1.0$ (0.7,1.3) $23.9$ $-2.4$ (-3.9,-1.0) $0.001$ $23.7$ $-0.2$ (-0.8,0.4) $31.3$ <	AgeGroup vs SA at ageF-valueAgeChange (Change Holn ageDifference in Change (CI)27.2(Ref)45.418.2 (10.1,26.3)(Ref)8.1-19.1 (-24.2,-13.9)<0.001

# **Table 3:** Clinical Risk Factors Stratified by Age and Race and Ethnic Group in Women, MASALA vs. MESA Groups

\* P-value < 0.05 from a pairwise test at age 45 vs. South Asians; \*\* P-value from a test of change from age 45 to 55 vs. South Asian. When the multivariate test was not significant, pairwise tests were not performed.

	Age	Group vs SA at age	P-value*	Age	Change from age	Difference in	P-value**		
	45	45 (CI)		55	45 to 55 (CI)	Change (CI)			
AHEI Diet Score									
South Asian	66.1	(Ref)		69.5	3.4 (1.5,5.2)	(Ref)			
White	48.3	-17.8 (-20.1,-15.6)	< 0.001	55.4	7.1 (5.1,9.1)	3.8 (1.1,6.4)	0.006		
Chinese	55.8	-10.3 (-13.8,-6.8)	< 0.001	60.5	4.7 (1.0,8.3)	1.3 (-2.7,5.4)	0.523		
Black	49.8	-16.3 (-19.2,-13.5)	< 0.001	53.6	3.8 (0.9,6.7)	0.5 (-2.9,3.9)	0.792		
Hispanic/Latino	50.2	-16.0 (-18.6,-13.3)	< 0.001	55.7	5.5 (3.2,7.9)	2.2 (-0.8,5.2)	0.150		
Alcohol Use (%)							0.728		
South Asian	39.3	(Ref)		41.1	1.8 (-7.0,10.6)	(Ref)			
White	64.2	24.9 (15.7,34.2)	< 0.001	62.0	-2.2 (-9.4,5.0)	-4.0 (-15.4,7.3)			
Chinese	34.5	-4.8 (-19.3,9.8)	0.521	29.4	-5.2 (-18.9,8.5)	-7.0 (-23.3,9.3)			
Black	50.5	11.2 (-1.9, 24.3)	0.094	42.4	-8.1 (-20.5,4.3)	-9.9 (-25.2,5.3)			
Hispanic/Latino	55.6	0.163 (4.3,28.2)	0.008	50.7	-4.8 (-15.4,5.6)	-6.7 (-20.4,7.0)			
Exercise (MET									
min/wk)									
South Asian	1082	(Ref)		1437	355 (-194,905)	(Ref)			
White	878	-204 (-801,392)	0.502	1873	995 (525,1464)	639 (-83,1362)	0.083		
Chinese	105	-977 (-1822,-132)	0.023	1711	1605 (818,2393)	1250 (289,2211)	0.011		
Black	1604	522 (-172,1216)	0.140	2686	1082 (478,1685)	726 (-89,1542)	0.081		
Hispanic/Latino	1604	521 (-154,1197)	0.130	2277	674 (126,1222)	319 (-457,1095)	0.421		
* P-value < 0.05 from a pairwise test at age 45 vs. South Asians; ** P-value from a test of change from age 45 to 55 vs. South									
A * 33.71 (1	1				· · · 1				

Table 4: Behavioral Risk Factors Stratified by Age and Race and Ethnic Group in Men, MASALA vs. MESA Groups

Asian. When the multivariate test was not significant, pairwise tests were not performed.

	Age	Group vs SA at age	P-value*	Age	Change from age	Difference in	P-value**
	45	45 (CI)	1 varae	55	45 to 55 (CI)	Change (CI)	1 (1140
AHEI Diet Score							
South Asian	68.5	(Ref)		71.8	3.35 (1.5,5.2)	(Ref)	
White	51.5	-17 (-19.2,-14.7)	< 0.001	58.5	7.02 (5.2,8.9)	3.67 (1.0,6.3)	0.006
Chinese	58.6	-9.8 (-13.1,-6.6)	< 0.001	64.6	5.94 (2.7,9.2)	2.59 (-1.2,6.3)	0.176
Black	47.7	-21 (-23.3,-18.2)	< 0.001	53.5	5.78 (3.5,8.1)	2.43 (-0.5,5.4)	0.106
Hispanic/Latino	50.3	-18.2 (-20.8,-15.6)	< 0.001	56.6	6.32 (4.1,8.5)	2.97 (0.1,5.9)	0.045
Alcohol Use (%)							0.051
South Asian	18.6	(Ref)		17.4	-1.2 (-7.0,4.6)	(Ref)	
White	42.8	24.1 (16.0,32.4)	< 0.001	46.7	3.8 (-3.8,11.5)	5.0 (-4.5,14.6)	
Chinese	25.8	0.072 (-4.4,18.7)	0.226	14.8	-11.0 (-23.4,1.4)	-9.8 (-23.5,3.9)	
Black	30.7	0.120 (4.1,20.0)	0.003	21.4	-9.2 (-16.4, -2.1)	-8.1 (-17.3,1.1)	
Hispanic/Latino	32.4	0.137 (4.8,22.7)	0.002	23.8	-8.6 (-16.5, -0.7)	-7.4 (-17.2,2.3)	
Exercise (MET							
min/wk)							
South Asian	1226	(Ref)		1429	203 (-263, 670)	(Ref)	
White	989	-237 (-718, 243)	0.334	1597	608 (242, 974)	405 (-188, 998)	0.181
Chinese	407	-819 (-1503, -136)	0.019	1267	861 (204, 1518)	657 (-148, 1463)	0.110
Black	1063	-163 (-689, 362)	0.542	2025	962 (525, 1400)	759 (119, 1399)	0.020
Hispanic/Latino	950	-276 (-815, 264)	0.316	1609	659 (219, 1098)	455 (-185, 1096)	0.164
* P-value < 0.05 fro	m a pair	wise test at age 45 vs. S	South Asians	: ** P-val	ue from a test of chan	ge from age 45 to 55	vs. South

# **Table 5:** Behavioral Risk Factors Stratified by Age and Race and Ethnic Group in Women, MASALA vs. MESA groups

\* P-value < 0.05 from a pairwise test at age 45 vs. South Asians; \*\* P-value from a test of change from age 45 to 55 vs. South Asian. When the multivariate test was not significant, pairwise tests were not performed.

#### **APPENDIX: STATISTICAL METHODS**

#### **A CONTINUOUS RISK FACTORS**

#### A.1 MODEL

Let  $y_{ij}$  be the risk factor for participant i (i = 1, ..., n) measured at exam j ( $j = 1, ..., J_i$ ). For continuous risk factors (BMI, diet score, physical activity) we fit the following linear mixed-effects model:

$$y_{ij} = \beta_{0} + \beta_{1}age_{ij} + \beta_{2}age_{ij}^{2} + \beta_{3}edu_{i} + \beta_{4}ins_{i} + \beta_{5}white_{i} + \beta_{6}black_{i} + \beta_{7}chinese_{i} + \beta_{8}hisp_{i} + \beta_{9}(age_{ij} * white_{i}) + \beta_{10}(age_{ij} * black_{i}) + \beta_{11}(age_{ij} * chinese_{i}) + \beta_{12}(age_{ij} * hisp_{i}) + \beta_{13}(age_{ij}^{2} * white_{i}) + \beta_{14}(age_{ij}^{2} * black_{i}) + \beta_{15}(age_{ij}^{2} * chinese_{i}) + \beta_{16}(age_{ij}^{2} * hisp_{i}) + b_{0i} + \varepsilon_{ij}$$
(1)

where  $age_{ij}$  is the age of participant *i* at exam *j*,  $edu_i$  is a vector of indicator variables for the education categories in Table 1,  $ins_i$  is a vector of indicator variables for the insurance categories in Table 1, and  $white_i$ ,  $black_i$ ,  $chinese_i$ , and  $hisp_i$  are indicator variables for race (the reference race is South Asians). The model also includes age by race interaction terms as well as age squared by race interaction terms. The parameters  $b_{0i}$  and  $\varepsilon_{ij}$  are independent random subject-specific intercept and error terms each normally distributed with variance  $\sigma_b^2$  and,  $\sigma_{\varepsilon}^2$  respectively.

#### **A.2 PREDICTIVE MARIGINAL MEANS**

To calculate the risk factor means by age and race controlling for the effects of education and insurance status, we use the method of predictive marginal means [1]. For example, to obtain the mean risk factor level at age 45 for whites, we set  $age_{ij} = 45$ ,  $age_{ij}^2 = 45^2$  and  $white_i = 1$  and perform the following calculation, using the estimated coefficients ( $\beta$ 's) from the model in (1) and the empirical Bayes estimates of the random effects ( $\hat{b}_{0i}$ ). That is,

$$\hat{\mu}_{W}^{45} = \frac{1}{n} \sum_{i=1}^{n} \hat{\beta}_{0} + \hat{\beta}_{1} 45 + \hat{\beta}_{2} (45^{2}) + \hat{\beta}_{3} e du_{i} + \hat{\beta}_{4} i n s_{i} + \hat{\beta}_{5} + \hat{\beta}_{9} 45 + \hat{\beta}_{13} (45^{2}) + \hat{b}_{0i}.$$
(2)

Similar calculations can be performed to obtain predicted means for the other race/ethnic groups at age 45 and age 55.

#### **B BINARY RISK FACTORS**

#### **B.1 MODEL**

The model for binary risk factors (hypertension, dyslipidemia, diabetes, alcohol use) is similar to that in (1) except we now use a mixed-effects logistic regression model of the probability  $(p_{ij})$  that participant *i* has the risk factor at exam *j*. That is,

$$log\left(\frac{p_{ij}}{1-p_{ij}}\right) = \beta_{0} + \beta_{1}age_{ij} + \beta_{2}age_{ij}^{2} + \beta_{3}edu_{i} + \beta_{4}ins_{i} + \beta_{5}White_{i} + \beta_{6}Black_{i} + \beta_{7}Chinese_{i} + \beta_{8}Hsp_{i} + \beta_{9}(age_{ij} * White_{i}) + \beta_{10}(age_{ij} * Black_{i}) + \beta_{11}(age_{ij} * Chinese_{i}) + \beta_{12}(age_{ij} * Hisp_{i}) + \beta_{13}(age_{ij}^{2} * White_{i}) + \beta_{14}(age_{ij}^{2} * Black_{i}) + \beta_{15}(age_{ij}^{2} * Chinese_{i}) + \beta_{16}(age_{ij}^{2} * Hisp_{i}) + b_{0i}$$
(3)

where as in (1) the parameter  $b_{0i}$  is a random intercept term for participant *i*.

#### **B.2 PREDICTIVE MARIGINAL MEANS (PROBABILITIES)**

Predictive marginal means for the model in (3) can be calculated similar to that in (2). Again, using the example of a White participant at age 45 we calculate the following quantity:

$$\hat{\eta}_{W}^{45} = \frac{1}{n} \sum_{i=1}^{n} \hat{\beta}_{0} + \hat{\beta}_{1} 45 + \hat{\beta}_{2} (45^{2}) + \hat{\beta}_{3} e du_{i} + \hat{\beta}_{4} ins_{i} + \hat{\beta}_{5} + \hat{\beta}_{9} 45 + \hat{\beta}_{13} (45^{2}) + \hat{b}_{0i}.$$

which is the log odds that a white participant has the risk factor at age 45. To put it on the probability scale, we apply the inverse logit transformation:

$$\hat{\mu}_W^{45} = \frac{e^{\hat{\eta}_W^{45}}}{1 + e^{\hat{\eta}_W^{45}}}$$

### **C CONTRASTS**

#### C.1 CONTRASTS AT AGE 45

The first set of contrasts compare predicted marginal mean outcomes at age 45 of White (W), Chinese (C), Black (B), and Hispanics/Latino (H) versus South Asians (SA). We first performed an overall (Wald) test of whether any of these contrasts were significant. That is

$$H_{0}: (\mu_{W}^{45} - \mu_{SA}^{45}) = (\mu_{C}^{45} - \mu_{SA}^{45}) = (\mu_{B}^{45} - \mu_{SA}^{45}) = (\mu_{H}^{45} - \mu_{SA}^{45}) = 0$$
$$H_{1}: (\mu_{W}^{45} - \mu_{SA}^{45})or (\mu_{C}^{45} - \mu_{SA}^{45})or (\mu_{B}^{45} - \mu_{SA}^{45})or (\mu_{H}^{45} - \mu_{SA}^{45}) \neq 0$$
(4)

Only if the overall test in (4) is rejected did we perform the four separate comparisons.

#### **C.1 SLOPE CONTRASTS**

The second set of contrasts compares the change in risk factor levels from age 45 to 55 in South Asians versus the four other race/ethnic groups. Let  $\Delta_R$  be the change in predicted risk factor means from age 45 to age 55 for race/ethnic group *R*. That is

$$\Delta_R = \mu_R^{55} - \mu_R^{45}$$

As with the baseline contrasts, we first performed an overall (Wald) test of whether any of the slopes in White, Chinese, Black, and Latino people were significantly different from the South Asian slope, that is

$$H_0: (\Delta_W - \Delta_{SA}) = (\Delta_C - \Delta_{SA}) = (\Delta_B - \Delta_{SA}) = (\Delta_H - SA) = 0$$
  
$$H_1: (\Delta_W - \Delta_{SA}) \text{ or } (\Delta_C - \Delta_{SA}) \text{ or } (\Delta_B - \Delta_{SA}) \text{ or } (\Delta_H - SA) \neq 0$$
(5)

Only if the overall test in (5) is rejected did we perform the four separate tests comparing the rate of change in the four MESA race/ethnic groups to South Asians.

## **Sources Cited**

1. Prasad DS, Kabir Z, Dash AK and Das BC. Childhood cardiovascular risk factors in South Asians: A cause of concern for adult cardiovascular disease epidemic. *Ann Pediatr Cardiol*. 2011;4:166-71.

2. Stanford South Asian Translational Heart Initiative.

3. Gupta K, Modi S and Ananthasubramaniam K. Toward Understanding Cardiovascular Risk Burden in South Asians: A Major Step Forward. *JACC Asia*. 2022;2:912-915.

4. Kanaya AM, Kandula N, Herrington D, Budoff MJ, Hulley S, Vittinghoff E and Liu K. Mediators of Atherosclerosis in South Asians Living in America (MASALA) study: objectives, methods, and cohort description. *Clin Cardiol*. 2013;36:713-720.

5. Bild DE, Bluemke DA, Burke GL, Detrano R, Diez Roux AV, Folsom AR, Greenland P, Jacob DR, Jr., Kronmal R, Liu K, Nelson JC, O'Leary D, Saad MF, Shea S, Szklo M and Tracy RP. Multi-Ethnic Study of Atherosclerosis: objectives and design. *Am J Epidemiol*. 2002;156:871-81.

6. Lloyd-Jones DM, Allen NB, Anderson CAM, Black T, Brewer LC, Foraker RE, Grandner MA, Lavretsky H, Perak AM, Sharma G, Rosamond W and American Heart A. Life's Essential 8: Updating and Enhancing the American Heart Association's Construct of Cardiovascular Health: A Presidential Advisory From the American Heart Association. *Circulation*. 2022;146:e18-e43.

7. Kanaya AM, Herrington D, Vittinghoff E, Ewing SK, Liu K, Blaha MJ, Dave SS, Qureshi F and Kandula NR. Understanding the high prevalence of diabetes in U.S. south Asians compared with four racial/ethnic groups: the MASALA and MESA studies. *Diabetes Care*. 2014;37:1621-8.

8. Gujral UP, Vittinghoff E, Mongraw-Chaffin M, Vaidya D, Kandula NR, Allison M, Carr J, Liu K, Narayan KMV and Kanaya AM. Cardiometabolic Abnormalities Among Normal-Weight Persons From Five Racial/Ethnic Groups in the United States: A Cross-sectional Analysis of Two Cohort Studies. *Ann Intern Med*. 2017;166:628-636.

9. Rodriguez LA, Jin Y, Talegawkar SA, Otto MCO, Kandula NR, Herrington DM and Kanaya AM. Differences in Diet Quality among Multiple US Racial/Ethnic Groups from the Mediators of Atherosclerosis in South Asians Living in America (MASALA) Study and the Multi-Ethnic Study of Atherosclerosis (MESA). *J Nutr*. 2020;150:1509-1515.

10. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, Jr., Jones DW, Materson BJ, Oparil S, Wright JT, Jr., Roccella EJ, Joint National Committee on Prevention DE, Treatment of High Blood Pressure. National Heart L, Blood I and National High Blood Pressure Education Program Coordinating C. Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Hypertension*. 2003;42:1206-52.

11. Genuth S, Alberti KG, Bennett P, Buse J, Defronzo R, Kahn R, Kitzmiller J, Knowler WC, Lebovitz H, Lernmark A, Nathan D, Palmer J, Rizza R, Saudek C, Shaw J, Steffes M, Stern M, Tuomilehto J, Zimmet P, Expert Committee on the D and Classification of Diabetes M. Follow-up report on the diagnosis of diabetes mellitus. *Diabetes Care*. 2003;26:3160-7.

12. Expert Panel on Detection E and Treatment of High Blood Cholesterol in A. Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, And Treatment of High Blood Cholesterol In Adults (Adult Treatment Panel III). *JAMA*. 2001;285:2486-97. 13. Consultation WHOE. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet*. 2004;363:157-63.

14. Defining Adult Overweight & Obesity.

15. Ainsworth BE, Irwin ML, Addy CL, Whitt MC and Stolarczyk LM. Moderate physical activity patterns of minority women: the Cross-Cultural Activity Participation Study. *J Womens Health Gend Based Med*. 1999;8:805-13.

16. Williams R. Using the margins command to estimate and interpret adjusted predictions and marginal effects. *The Stata Journal*. 2012;12:308–331.

Stata 18 Base Reference Manual [computer program]. College Station, TX: Stata Press;
 2023.

18. Nair MP, D. Why Do South Asians Have High Risk for CAD? *Global Heart* 2012;7:307-314.

19. Patel M, Phillips-Caesar E and Boutin-Foster C. Barriers to lifestyle behavioral change in migrant South Asian populations. *J Immigr Minor Health*. 2012;14:774-85.

20. Zhu X, Liu J, Sevoyan M and Pate RR. Acculturation and leisure-time physical activity among Asian American adults in the United States. *Ethn Health*. 2022;27:1900-1914.

21. Dave SS, Craft LL, Mehta P, Naval S, Kumar S and Kandula NR. Life stage influences on U.S. South Asian women's physical activity. *Am J Health Promot*. 2015;29:e100-8.

22. Maresova P, Javanmardi E, Barakovic S, Barakovic Husic J, Tomsone S, Krejcar O and Kuca K. Consequences of chronic diseases and other limitations associated with old age - a scoping review. *BMC Public Health*. 2019;19:1431.

23. Gil-Salcedo A, Dugravot A, Fayosse A, Dumurgier J, Bouillon K, Schnitzler A, Kivimaki M, Singh-Manoux A and Sabia S. Healthy behaviors at age 50 years and frailty at older ages in a 20-year follow-up of the UK Whitehall II cohort: A longitudinal study. *PLoS Med*. 2020;17:e1003147.

24. Guo A, Jin H, Mao J, Zhu W, Zhou Y, Ge X and Yu D. Impact of health literacy and social support on medication adherence in patients with hypertension: a cross-sectional community-based study. *BMC Cardiovasc Disord*. 2023;23:93.