

The Interplay of Perceived Mental Health and Suicidal Ideation among Aboriginal Peoples in Canada: A Secondary Analysis of Aboriginal Peoples Survey

Olasehinde Ben Adebayo

Affiliation: *College of Health Professions, Bellarmine University, Louisville, KY, USA*

Abstract

The disproportionately high suicide rates among Aboriginal populations in Canada, linked to historical trauma and systemic inequities, necessitate targeted mental health interventions. This study investigated the relationship between perceived mental health and suicidal ideation among the Aboriginal population in Canada, First Nations, Inuit, and Métis, using data from the 2012 Aboriginal Peoples Survey (APS) (n=4,725). The analysis employed chi-square tests and bivariate logistic regression to examine the association between self-perceived mental health, suicidal ideation, and socio-demographic factors. Results revealed significant associations between suicidal ideation, age group ($p < .001$), and sex ($p < .001$), with younger individuals (18-24 years) and females reporting a higher prevalence of suicidal thoughts. Also, bivariate logistic regression showed that being between the ages of 25 and 34 was a strong predictor of suicidal thoughts (OR = 1.954, $p = .000$), while being female (OR = .648, $p = .000$) and having a higher socioeconomic status (OR = .678, $p = .000$) were weaker predictors. Although Aboriginal identity was not significantly associated with suicidal ideation, self-perceived mental health was. These results made it clear that focused and targeted mental health interventions and culturally appropriate suicide prevention strategies are needed right away in Aboriginal communities in Canada. These strategies need to meet the needs of vulnerable groups, like younger people and women.

Keywords: Suicide Ideation, Mental Health, Aboriginal Peoples, Socioeconomic Status

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Introduction

Between 2011 and 2016 in Canada, suicide rates were noticeably higher among Aboriginal populations in comparison to non-Aboriginal populations (Kumar & Tjepkema, 2019). The disproportionately high rates of suicide among the Canadian Aboriginal population, comprising the First Nations, Inuit, and Métis populations, compared to the non-Aboriginal population necessitate a deeper understanding of the underlying factors contributing to this mental health crisis. Existing research has revealed a significant burden of mental health problems and suicides among Aboriginal Canadians (Kral, 2016; Nelson & Wilson, 2017) with First Nations and Inuit youth experiencing particularly alarming suicide rates (Nelson & Wilson, 2017; Orkin et al., 2013). These disparities are often attributed to a complex interplay of socio-economic, cultural, and historical factors, which include poverty, discrimination, and the enduring effects of colonisation and residential school experiences (Kral, 2016; Toombs et al., 2023; Wilk et al., 2017). The economic disadvantages faced by many Indigenous communities, coupled with historical trauma and systemic racism, can significantly impact mental health status and increase the likelihood of suicidal ideation (Kral, 2016; Pandeya et al., 2021; Smye et al., 2023), making it a crucial area for investigation.

This study examines the relationship between perceived mental health and suicidal ideation within these Aboriginal populations through a secondary data analysis. This study aims to determine the extent to which perceived poor mental health is associated with suicidal ideation among First Nations, Inuit, and Métis populations in Canada. By addressing the research question, "To what extent does perceived poor mental health correlate with suicidal thinking among First Nations, Inuit, and Métis populations in Canada?", this study seeks to identify a critical intervention point for targeted mental health programmes and suicide prevention efforts. This study, by elucidating the relationship between perceived mental

health and suicidal thoughts within these communities, can inform the development of culturally sensitive mental health interventions and suicide prevention strategies tailored to the specific needs of Aboriginal peoples in Canada.

Methodology

Participants

The secondary data for this research was derived from the 2012 Aboriginal Peoples Survey (APS), a comprehensive survey conducted by Statistics Canada (Government of Canada, 2023). The Aboriginal Peoples are defined as the First Nations, Inuit, and Métis people. The Aboriginal people surveyed are 18 years of age and older. The sample consisted of 55.2% females and 44.8% males. 39.3% of the participants identified as First Nations, 36.1% as Métis and 24.6% as Inuit (Government of Canada, 2023). The largest age group in the study was young adults aged 18 to 24 years (31.7%) (Government of Canada, 2023).

Procedure

This study used secondary data analysis methods to analyze existing APS data. The secondary data in APS is cost-effective, timesaving, and contains a wider range of information. The APS data was selected to ensure reliable and valid responses. The survey included standardised questions across various domains, including health, education, employment, and cultural participation. The following variables of interest were extracted from the APS dataset for this research: perceived mental health, suicidal thoughts, and socioeconomic status (SES).

The analysis was conducted using data obtained from Statistics Canada's official repository at <https://www150.statcan.gc.ca/n1/tbl/csv/41100011-eng.zip> (Government of Canada, 2023). Extensive data cleaning was performed using R-Studio 4.1.2 due to the initial disarray of the dataset. Subsequently, data analysis was carried out using SPSS version 22 (Appendix A). Descriptive statistics were used to examine perceived mental health, suicidal

thoughts, and SES across different Aboriginal identity groups. To investigate the association between sociodemographic variables and suicidal ideation and self-perceived mental health among Aboriginal people in Canada. A chi-square cross-tabulation analysis was performed to identify potential relationships between categorical variables due to its suitability and ability to identify relationships between categorical variables (Momeni et al., 2017). The researcher used the chi-square to test or identify the relationship between the categorical variables (suicide ideation and perceived mental health) and the socio-demographic variables (Aboriginal identity, sex, and age group).

Furthermore, bivariate logistic regression analysis was employed to examine the association between perceived mental health and suicidal ideation among Aboriginal people in Canada. This method was selected as logistic regression is well-suited for modelling binary outcomes (Starbuck, 2023). The presence or absence of suicidal ideation is a dichotomous variable. The researcher calculated and utilized odds ratios (OR) to understand the strength and direction of the relationship between the outcomes. I conducted all statistical analyses using SPSS software (Appendix A) to ensure robust and reliable results.

Data Preparation

The dataset underwent filtering, recoding, and transformation processes to maintain data integrity and relevance to the study's focus on Aboriginal identity, socioeconomic status (SES), and mental health outcomes, particularly suicidal thoughts. These steps were taken to convert the raw data into a suitable format for statistical analysis. The data required extensive cleaning due to its messy state. The dataset was filtered to include only relevant categories for Aboriginal identity (First Nations, Métis, and Inuit). Irrelevant categories were removed, such as adult age cohorts not pertinent to the study's socioeconomic and mental health indicators, specifically those below 18 years of age. The "Both sexes" category was also eliminated, focusing on males and females separately.

A crucial step in the data-cleaning process involved creating a variable indicating suicidal ideation. This was achieved by recording responses from the self-perceived mental health and suicidal thoughts Likert measures. The researcher coded 1 (yes) for participants who selected responses indicating suicidal ideation, and 0 (no) for others. This resulted in a dichotomous variable named "Suicidal ideation," which served as the dependent variable in subsequent analyses. Lastly, missing observations were removed to ensure data reliability for analysis.

Measures

Perceived Mental Health

The overall mental health of individuals was assessed using a self-report question in the APS. Responses ranged from "excellent" to "poor," providing a subjective measure of perceived mental well-being. For analysis, perceived mental health was dichotomized into "good" (combining excellent/very good/good) and "poor" (combining fair/poor).

Suicidal Thoughts

Suicidal thoughts were assessed by inquiring whether respondents had any thoughts of suicide within the past year. This dichotomous variable (yes/no) represents the presence or absence of suicidal thoughts and served as the outcome variable in the logistic regression analysis.

Socioeconomic Status (SES)

Socioeconomic status was determined based on reported income and education levels in the APS dataset. The education level was determined based on the highest grade completed. The analysis included SES as a covariate to control for its potential confounding effect on the relationship between perceived mental health and suicidal thoughts.

Sociodemographic Characteristics

Age, gender, and Aboriginal identity were included as socio-demographic variables in the survey. Age groups were consolidated to include those 18 years and older to assess changes

in mental health attitudes across different life stages. Both sexes were included to identify potential gender-specific differences in mental health outcomes. The Aboriginal identity variable comprised three categories: First Nations, Métis, and Inuit

Results

Table 1: Frequency distribution of the variables used in the study

Variables	Frequency	Percentage, %
Aboriginal identity		
First nation	1858	39.3
Inuit	1160	24.6
Metis	1707	36.1
Age group		
18-24 years	1496	31.7
25-34 years	1068	22.6
35-44 years	1072	22.7
45-54 years	1089	23.0
Sex		
Male	2117	44.8
Female	2608	55.2
Suicidal thoughts		
No	2872	60.8
Yes	1853	39.2

The initial analysis of the dataset presented in Table 1 shows the frequency distribution of the variables used in the study. The results indicate that 39.3% of the respondents identified as First Nation (n = 1858), 24.6% as Inuit (n = 1160), and 36.1% as Métis (n = 1707). Similarly, the result shows that among the age groups, 31.7% of the respondents (n = 1496) were aged

18 to 24 years, followed by individuals aged between 45 and 54 years, constituting 23.0% of the sample (n = 1089), individuals aged 35 to 44 years constituted 22.7% (n = 1072), and 22.6% (n = 1068) were those aged 25 to 34 years. The result further shows the sample was slightly skewed towards females (55.2%) while 44.8% (n = 2117) were males. The age group with the highest representation was 18-24 years (31.7%), indicating a focus on younger individuals. A significant portion of the sample (39.2%) reported having experienced suicidal thoughts.

Chi-Square Test of Association (Suicide Ideation)

Table 2: Cross-tabulations of socio-demographic variables with suicide ideation in past years among Aboriginal people in Canada

	Suicide ideation, %		p-value
	No	Yes	
Aboriginal identity			0.507
First nation	60.3	39.7	
Inuit	59.9	40.1	
Metis	61.9	38.1	
Age group			0.000
18-24 years	49.7	50.3	
25-34 years	65.3	34.7	
35-44 years	66.6	33.4	
45-54 years	65.9	34.1	
Sex			0.000
Male	66.0	34.0	
Female	56.5	43.5	

$p < .001$

Table 2 presents the cross-tabulation results of socio-demographic variables associated with suicide ideation among Aboriginal people in Canada. Of the respondents, 39.7% of the first nation individuals indicated they have experienced suicide ideation in the past 12 months, 40.1% of the Inuit individuals have experienced suicide thoughts, 38.1% of the Metis individuals have experienced suicide thoughts in the past 12 months, and 60.3%, 59.9% and 61.9% respectively, indicated they did not. There was no significant association between suicide ideation and aboriginal identity ($p = .507$). Individuals aged between 18 and 24 years had a higher prevalence of suicide ideation (50.3%) compared to other age groups. The result shows that there is a statistically significant association between suicide ideation and age group ($p = .000$). Females had a higher prevalence of suicide ideation than males, at 43.5% versus 34.0%. Among males, 66.0% reported no suicide ideation, while among females, 56.5% reported no suicide ideation. The result also shows a significant association between gender and suicide ideation ($p = .000$).

Table 3: Cross Tabulations of Socio-demographic variables with perceived mental health in past years among Aboriginal people in Canada (self-perceived mental health)

	Self-perceived mental health and suicidal thoughts, %							p-value
	Ever seriously	Excellent or very good	Fair or poor	Good	Never seriously	Seriously contemplated (past 12 months)	Seriously contemplated (not past 12 months)	
Aboriginal identity								0.032
First nation	39.7	39.7	41.9	39.8	37.7	40.9	37.2	

Inuit	27.2	23.6	19.4	22.2	26.5	22.4	29.2
Metis	33.1	36.8	38.7	37.9	35.8	36.6	33.6
Age group							0.000
18-24	34.0	25.9	40.7	26.6	25.1	63.4	38.4
years							
25-34	23.1	24.3	17.8	23.6	24.9	13.4	21.4
years							
35-44	21.8	25.1	19.8	24.4	25.0	13.8	18.2
years							
45-54	21.0	24.7	21.7	25.4	25.0	9.5	22.0
years							
Sex							0.000
Male	41.0	48.7	34.6	48.0	49.3	37.1	44.8
Female	59.0	51.3	65.4	52.0	50.7	62.9	55.2

$p < .001$

All the socio-demographic characteristics variables were statistically associated with self-perceived mental health ($p < .001$). Among the age groups, the results were highly significant ($p < .001$), with younger age groups (18-24 years) showing higher percentages of serious suicidal thoughts in the past 12 months compared to older age groups. This suggests that younger individuals experience significant changes such as puberty, life expectations, and social pressure, which can contribute to stress and anxiety Backes et al., 2024)

Furthermore, there was a significant association between sex, self-perceived mental health, and suicidal thoughts ($p < .001$). A higher percentage of males reported excellent or very good self-perceived mental health (48.7%), while females reported higher percentages of fair or poor self-perceived mental health and suicidal thoughts (65.4%). The researcher found

a significant association between Aboriginal identity and self-perceived mental health ($p = .032$). A significant association was found between age group and self-perceived mental health ($p < .001$). Younger individuals (18-24 years) reported higher rates of poor mental health and suicidal thoughts. A significant association was found between sex and self-perceived mental health ($p < .001$). Females reported higher rates of fair or poor mental health and suicidal thoughts compared to males.

Bivariate Logistic Regression

The result of the bivariate logistic regression model examining the relationship between perceived mental health and suicidal thoughts is presented in Table 4. The result shows that individuals with poor perceived mental health have a high likelihood of contemplating suicidal thoughts, although this was not statistically significant ($OR = 1.019$, $p = 0.248$). This suggests that the likelihood of contemplating suicidal thoughts was not meaningfully different across varying levels of perceived mental health.

Regarding Aboriginal identity, the odds ratios for Inuit ($OR = 1.069$, $p = 0.344$) and Métis ($OR = 1.020$, $p = 0.800$) were not statistically significant, meaning that Inuit and Métis individuals were not more likely to experience suicidal thoughts compared to First Nations individuals. The result suggests that the odds of experiencing suicidal thoughts were essentially the same across these Aboriginal identities.

Among the age groups, individuals between the ages of 25 and 34 ($OR = 1.954$, $p = 0.00$) are more likely to experience suicidal thoughts than others. This implies that an individual between the ages of 25 and 34 is associated with a 1.954 increase in the odds of being involved in suicidal ideations. Furthermore, the result shows that females have a lower odd of experiencing suicidal thoughts than males ($OR = 0.648$, $p = 0.000$). Lastly, socioeconomic status also showed a significant negative association with suicidal thoughts ($OR = 0.678$, $p = 0.000$), suggesting that higher socioeconomic status (reported income and education levels) is

associated with lower odds of experiencing suicidal thoughts. The model showed a net improvement in classification, with the percentage of correct classifications increasing from the null model to the fitted model. This indicates that the inclusion of perceived mental health, Aboriginal identity, age group, sex, and socioeconomic status as predictors improved the model's ability to correctly classify individuals with and without suicidal thoughts

The model fit statistics show that the model is a moderately good fit for the data (Nagelkerke's $R^2 = 0.058$ and Cox & Snell's $R^2 = 0.043$). The model chi-square test is statistically significant ($\chi^2 = 65.353$, $p < 0.001$), indicating that the model significantly predicts the outcome variable.

Table 4: Bivariate Logistic Regression model:

A two-variable logistic regression model was used to look at the link between how mentally healthy someone thinks they are and suicidal thoughts while considering sociodemographic factors.

Variables included	B	Wald	p-value	Odds ratio (OR)	95% CI for OR	
					Lower	Upper
Constant	-.609	36.629	.000	.544		
Perceived mental health and suicidal	.019	1.337	.248	1.019	.987	1.053
Aboriginal identity						
¹ First nation		.938	.626			
Inuit	.067	.895	.344	1.069	.931	1.227
Metis	.020	.064	.800	1.020	.872	1.194
Age group						

18-24 years		109.713	.000			
25-34 years	.670	64.896	.000	1.954	1.660	2.300
35-44 years	.032	.121	.728	1.032	.863	1.235
45-54 years	-.043	.215	.643	.958	.800	1.147
Sex						
Male						
Female	-.434	49.454	.000	.648	.574	.731
Socioeconomic status	-.389	29.894	.000	.678	.589	.779

$R^2 = 0.058$ (Nagelkerke), 0.043 (Cox & Snell). Model $\chi^2_8 = 65.353$, $p = 0.000$.

Reference categories : Aboriginal identity – First Nation ;

Age group – 18-24 years;

Sex: Male

4. Discussion

Findings from the study showed that there was no significant association between suicide ideation and Aboriginal identity. Furthermore, there was a statistically significant association between suicide ideation and age group, as well as a significant association between gender and suicide ideation. This aligns with the findings of Zhang et al., who reported a 60% increase in female suicide rates over male suicide rates in their study on gender differences in suicidal ideation and health risk behaviour (Zhang et al., 2019). However, this contradicts the findings of Lemstra et al. (2013), who found no significant influence of age or gender on the likelihood of suicide ideation. Their study revealed that suicidal ideation disproportionately affected girls, with 27.8% reporting such thoughts among youths in Canada, compared to 15.2% for boys, a finding that aligns with my findings.

The data suggests that age and sex are significant factors influencing suicide ideation and self-perceived mental health among Aboriginal people in Canada. Younger individuals and

females appear to be particularly vulnerable groups. While Aboriginal identity was not significantly associated with suicide ideation in Table 2, it was significantly associated with self-perceived mental health in Table 3, suggesting a more nuanced relationship. The findings further showed that individuals with poor perceived mental health have a high likelihood of contemplating suicidal thoughts, although this was not statistically significant. This agrees with the findings of Akram et al. (2020) which analysed the link between suicidal ideation and psychiatric symptoms like depression, mania, psychosis, and stress in UK university students. The result concluded that suicidal ideation was significantly associated with symptoms of depression, mania, psychosis, and stress (Akram et al., 2020).

Among Aboriginal identities, being an Inuit individual is associated with an increase in the odds of experiencing suicidal thoughts compared to non-Inuit individuals. Similarly, there are higher odds of Metis individuals experiencing suicidal thoughts than non-Metis individuals. The lack of a significant association between perceived mental health and suicidal thoughts indicates that interventions should not solely focus on improving perceived mental health but should also address broader sociodemographic factors such as occupational status, sexual orientation, educational level, immigration status, and family structure. This study has several limitations. The use of secondary data analysis restricts the analysis to the variables available within the APS, potentially omitting other important factors. Finally, stigma, fear of judgment, or memory errors may lead to underreporting or overreporting, compromising the accuracy of self-reported mental health data.

Ethical Considerations

Conflict of interest: None.

Ethical Statement: The submitted manuscript did not involve human or animal research.

This study utilised secondary data. No ethical approval was sought or required for this study.

It is a research review submitted in Partial fulfilment of the requirement for PhD Health

Professions Education, 2024SU Statistics in Health Professions Education II (HLTH-862-ON)/Summer 2024 at Bellarmine University Kentucky, United States of America.

Conclusions

This analysis reveals the complex interplay of perceived mental health, suicidal ideation, and socio-demographic factors among Aboriginal peoples in Canada. The findings from the secondary data used accentuate the heightened susceptibility of younger individuals, particularly those aged 25-34 years old and females, to suicidal thoughts. This emphasises the need for targeted mental health interventions and culturally sensitive suicide prevention strategies for the population. While Aboriginal identity alone may not directly predict suicidal ideation, its significant association with self-perceived mental health points to a subtle interplay that warrants further investigation.

The research utilised secondary data, meaning the researcher did not participate in the data collection process. This lack of control raises concerns regarding potential inconsistencies, errors, or biases within the data, which may affect the quality, reliability, and validity of the findings. Additionally, the reliance on self-reported data highlights the need for future research to employ prospective designs that incorporate a wider array of variables, including cultural determinants and historical trauma. This approach would facilitate a more comprehensive understanding of the multifaceted factors influencing mental well-being and suicidal ideation among Aboriginal communities in Canada. To improve understanding, future investigations should prioritize prospective research designs to elucidate causal relationships between the variables examined herein and those identified for future inquiry. Additionally, exploring the impact of cultural determinants, such as perceived stress, religious affiliation, and historical

trauma, on mental well-being and suicidal ideation within these populations is warranted.

By addressing these knowledge gaps, researchers and policymakers can collaboratively develop more effective, culturally sensitive interventions to alleviate the mental health crisis and reduce suicide rates within these populations.

References

- Akram, U., Ypsilanti, A., Gardani, M., Irvine, K., Allen, S., Akram, A., Drabble, J., Bickle, E., Kaye, L., Lipinski, D., Matuszyk, E., Sarlak, H., Steedman, E., & Lazuras, L. (2020). Prevalence and psychiatric correlates of suicidal ideation in UK university students. *Journal of Affective Disorders*, 272, 191–197.
<https://doi.org/10.1016/j.jad.2020.03.185>
- Backes, E. P., & Bonnie, R. J. (2019). *Adolescent development*. National Library of Medicine. <https://www.ncbi.nlm.nih.gov/books/NBK545476/>
- Government of Canada (2023). *Perceived mental health and suicidal thoughts by Aboriginal identity - Dataset* <https://open.canada.ca/data/en/dataset/200fbda4-ca32-4725-88ae-dee57e5d372c/resource/c982d4dd-5edf-487f-82e7-4cead4bf460f>
- Kral, M. J. (2016). Suicide and Suicide Prevention among Inuit in Canada. *The Canadian Journal of Psychiatry/Canadian Journal of Psychiatry*, 61(11), 688–695.
<https://doi.org/10.1177/0706743716661329>
- Kumar, M. B., & Tjepkema, M. (2019, June 28). *Suicide among First Nations people, Métis and Inuit (2011-2016): Findings from the 2011 Canadian Census Health and Environment Cohort (CanCHEC)*. Statistics Canada; Statistics Canada. Retrieved July 20, 2024, from <https://www150.statcan.gc.ca/n1/pub/99-011-x/99-011-x2019001-eng.htm>

Lemstra, M., Rogers, M., Moraros, J., & Grant, E. (2013). Risk indicators of suicide ideation among on-reserve First Nations youth. *Paediatrics & Child Health*, 18(1), 15–20.

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3680266/>

Momeni, A., Pincus, M., & Libien, J. (2017). Cross Tabulation and Categorical Data Analysis. In *Springer eBooks* (pp. 93–120). https://doi.org/10.1007/978-3-319-60543-2_5

Nelson, S. E., & Wilson, K. (2017). The mental health of Indigenous peoples in Canada: A critical review of research. *Social Science & Medicine*, 176, 93–112.
<https://doi.org/10.1016/j.socscimed.2017.01.021>

Orkin, A. M., Rajaram, N., & Schwandt, M. (2013). Aboriginal populations and youth suicide. *Canadian Medical Association Journal*, 185(15), 1347–1347.
<https://doi.org/10.1503/cmaj.113-2139>

Pandeya, N. A., Schluter, P. J., Spurling, G. K., Tyson, C., Hayman, N. E., & Askew, D. A. (2021). Factors Associated with Thoughts of Self-Harm or Suicide among Aboriginal and Torres Strait Islander People Presenting to Urban Primary Care: An Analysis of De-Identified Clinical Data. *International Journal of Environmental Research and Public Health*, 19(1), 153. <https://doi.org/10.3390/ijerph19010153>

Smye, V., Browne, A. J., Josewski, V., Keith, B., & Mussell, W. (2023). Social Suffering: Indigenous Peoples' Experiences of Accessing Mental Health and Substance Use Services. *International Journal of Environmental Research and Public Health*, 20(4), 3288. <https://doi.org/10.3390/ijerph20043288>

Starbuck, C. (2023). Logistic Regression. In *Springer eBooks* (pp. 223–238).
https://doi.org/10.1007/978-3-031-28674-2_12

Toombs, E., Lund, J. I., Mushquash, A. R., & Mushquash, C. J. (2023). Intergenerational residential school attendance and increased substance use among First Nation adults

living off-reserve: An analysis of the aboriginal peoples survey 2017. *Frontiers in Public Health*, 10(10). <https://doi.org/10.3389/fpubh.2022.1029139>

Wilk, P., Maltby, A., & Cooke, M. (2017). Residential schools and the effects on Indigenous health and well-being in Canada—a scoping review. *Public Health Reviews*, 38(1). <https://doi.org/10.1186/s40985-017-0055-6>

Zhang, Y.-Y., Lei, Y.-T., Song, Y., Lu, R.-R., Duan, J.-L., & Prochaska, J. J. (2019). Gender differences in suicidal ideation and health-risk behaviors among high school students in Beijing, China. *Journal of Global Health*, 9(1). <https://doi.org/10.7189/jogh.09.010604>

APPENDIX A

SPSS Output File

CROSSTABS

/TABLES=Self_perceived_mental_health_and_suicidal_thoughts BY Aboriginal_identity

Age_group Sex

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ CORR

/CELLS=COUNT ROW COLUMN

/COUNT ROUND CELL.

Crosstabs

Self_perceived_mental_health_and_suicidal_thoughts * Aboriginal_identity

Crosstab

			Aboriginal_identity			Total
			First nation	Inuit	Metis	
Self_perceived_mental_health_and_suicidal_thoughts	Ever seriously contemplated suicide	Count	251	172	209	632
		% within Self_perceived_mental_health_and_suicidal_thoughts	39.7%	27.2%	33.1%	100.0%
		% within Aboriginal_identity	13.5%	14.8%	12.2%	13.4%
	Excellent or very good self-perceived mental health	Count	384	228	356	968
		% within Self_perceived_mental_health_and_suicidal_thoughts	39.7%	23.6%	36.8%	100.0%
		% within Aboriginal_identity	20.7%	19.7%	20.9%	20.5%
	Fair or poor self-perceived mental health	Count	205	95	189	489
		% within Self_perceived_mental_health_and_suicidal_thoughts	41.9%	19.4%	38.7%	100.0%

	% within Aboriginal_identity	11.0%	8.2%	11.1%	10.3%
Good self-perceived mental health	Count	353	197	336	886
	% within Self_perceived_mental_health_and_suicidal_thoughts	39.8%	22.2%	37.9%	100.0%
	% within Aboriginal_identity	19.0%	17.0%	19.7%	18.8%
Never seriously contemplated suicide	Count	384	270	364	1018
	% within Self_perceived_mental_health_and_suicidal_thoughts	37.7%	26.5%	35.8%	100.0%
	% within Aboriginal_identity	20.7%	23.3%	21.3%	21.5%
Seriously contemplated suicide in the past 12 months	Count	95	52	85	232
	% within Self_perceived_mental_health_and_suicidal_thoughts	40.9%	22.4%	36.6%	100.0%
	% within Aboriginal_identity	5.1%	4.5%	5.0%	4.9%
Seriously contemplated suicide but not in the past 12 months	Count	186	146	168	500
	% within Self_perceived_mental_health_and_suicidal_thoughts	37.2%	29.2%	33.6%	100.0%
	% within Aboriginal_identity	10.0%	12.6%	9.8%	10.6%
Total	Count	1858	1160	1707	4725
	% within Self_perceived_mental_health_and_suicidal_thoughts	39.3%	24.6%	36.1%	100.0%
	% within Aboriginal_identity	100.0%	100.0%	100.0%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
--	-------	----	-----------------------

Pearson Chi-Square	22.519 ^a	12	.032
Likelihood Ratio	22.686	12	.031
Linear-by-Linear Association	.308	1	.579
N of Valid Cases	4725		

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 56.96.

Symmetric Measures

		Value	Asymp. Std. Error ^a	Approx. T ^b	Approx. Sig.
Interval by Interval	Pearson's R	.008	.014	.555	.579 ^c
Ordinal by Ordinal	Spearman Correlation	.009	.014	.619	.536 ^c
N of Valid Cases		4725			

- a. Not assuming the null hypothesis.
b. Using the asymptotic standard error assuming the null hypothesis.
c. Based on normal approximation.

Self_perceived_mental_health_and_suicidal_thoughts * Age_group

Crosstab

			Age_group		
			18 to 24 years	25 to 34 years	35 to 44 years
Self_perceived_mental_health_and_suicidal_thoughts	Ever seriously contemplated suicide	Count	215	146	138
		% within Self_perceived_mental_health_and_suicidal_thoughts	34.0%	23.1%	21.8%
		% within Age_group	14.4%	13.7%	12.9%
	Excellent or very good self-perceived mental health	Count	251	235	243
		% within Self_perceived_mental_health_and_suicidal_thoughts	25.9%	24.3%	25.1%
		% within Age_group	16.8%	22.0%	22.7%
	Fair or poor self-perceived mental health	Count	199	87	97
		% within Self_perceived_mental_health_and_suicidal_thoughts	40.7%	17.8%	19.8%

	% within Age_group	13.3%	8.1%	9.0%
Good self-perceived mental health	Count	236	209	216
	% within Self_perceived_mental_health_and_suicidal_thoughts	26.6%	23.6%	24.4%
	% within Age_group	15.8%	19.6%	20.1%
Never seriously contemplated suicide	Count	256	253	255
	% within Self_perceived_mental_health_and_suicidal_thoughts	25.1%	24.9%	25.0%
	% within Age_group	17.1%	23.7%	23.8%
Seriously contemplated suicide in the past 12 months	Count	147	31	32
	% within Self_perceived_mental_health_and_suicidal_thoughts	63.4%	13.4%	13.8%
	% within Age_group	9.8%	2.9%	3.0%
Seriously contemplated suicide but not in the past 12 months	Count	192	107	91
	% within Self_perceived_mental_health_and_suicidal_thoughts	38.4%	21.4%	18.2%
	% within Age_group	12.8%	10.0%	8.5%
Total	Count	1496	1068	1072
	% within Self_perceived_mental_health_and_suicidal_thoughts	31.7%	22.6%	22.7%
	% within Age_group	100.0%	100.0%	100.0%

Crosstab

			Age_group	
			45 to 54 years	Total
Self_perceived_mental_health_and_suicidal_thoughts	Ever seriously contemplated suicide	Count	133	632
		% within Self_perceived_mental_health_and_suicidal_thoughts	21.0%	100.0%

	% within Age_group	12.2%	13.4%
Excellent or very good self-perceived mental health	Count	239	968
	% within Self_perceived_mental_health_and_suicidal_thoughts	24.7%	100.0%
	% within Age_group	21.9%	20.5%
Fair or poor self-perceived mental health	Count	106	489
	% within Self_perceived_mental_health_and_suicidal_thoughts	21.7%	100.0%
	% within Age_group	9.7%	10.3%
Good self-perceived mental health	Count	225	886
	% within Self_perceived_mental_health_and_suicidal_thoughts	25.4%	100.0%
	% within Age_group	20.7%	18.8%
Never seriously contemplated suicide	Count	254	1018
	% within Self_perceived_mental_health_and_suicidal_thoughts	25.0%	100.0%
	% within Age_group	23.3%	21.5%
Seriously contemplated suicide in the past 12 months	Count	22	232
	% within Self_perceived_mental_health_and_suicidal_thoughts	9.5%	100.0%
	% within Age_group	2.0%	4.9%
Seriously contemplated suicide but not in the past 12 months	Count	110	500
	% within Self_perceived_mental_health_and_suicidal_thoughts	22.0%	100.0%
	% within Age_group	10.1%	10.6%
Total	Count	1089	4725
	% within Self_perceived_mental_health_and_suicidal_thoughts	23.0%	100.0%
	% within Age_group	100.0%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
--	-------	----	-----------------------

Pearson Chi-Square	189.001 ^a	18	.000
Likelihood Ratio	181.437	18	.000
Linear-by-Linear Association	7.400	1	.007
N of Valid Cases	4725		

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 52.44.

Symmetric Measures

		Value	Asymp. Std. Error ^a	Approx. T ^b	Approx. Sig.
Interval by Interval	Pearson's R	-.040	.015	-2.722	.007 ^c
Ordinal by Ordinal	Spearman Correlation	-.037	.015	-2.530	.011 ^c
N of Valid Cases		4725			

- a. Not assuming the null hypothesis.
b. Using the asymptotic standard error assuming the null hypothesis.
c. Based on normal approximation.

Self_perceived_mental_health_and_suicidal_thoughts * Sex

Crosstab

			Sex		Total
			Male	Female	
Self_perceived_mental_health_and_suicidal_thoughts	Ever seriously contemplated suicide	Count	259	373	632
		% within Self_perceived_mental_health_and_suicidal_thoughts	41.0%	59.0%	100.0%
		% within Sex	12.2%	14.3%	13.4%
	Excellent or very good self-perceived mental health	Count	471	497	968
		% within Self_perceived_mental_health_and_suicidal_thoughts	48.7%	51.3%	100.0%
		% within Sex	22.2%	19.1%	20.5%
	Fair or poor self-perceived mental health	Count	169	320	489
		% within Self_perceived_mental_health_and_suicidal_thoughts	34.6%	65.4%	100.0%
		% within Sex	8.0%	12.3%	10.3%
		Count	425	461	886

	Good self-perceived mental health	% within Self_perceived_mental_health_and_suicidal_thoughts	48.0%	52.0%	100.0%
		% within Sex	20.1%	17.7%	18.8%
		Count	502	516	1018
	Never seriously contemplated suicide	% within Self_perceived_mental_health_and_suicidal_thoughts	49.3%	50.7%	100.0%
		% within Sex	23.7%	19.8%	21.5%
		Count	86	146	232
	Seriously contemplated suicide in the past 12 months	% within Self_perceived_mental_health_and_suicidal_thoughts	37.1%	62.9%	100.0%
		% within Sex	4.1%	5.6%	4.9%
		Count	205	295	500
	Seriously contemplated suicide but not in the past 12 months	% within Self_perceived_mental_health_and_suicidal_thoughts	41.0%	59.0%	100.0%
		% within Sex	9.7%	11.3%	10.6%
		Count	2117	2608	4725
Total		% within Self_perceived_mental_health_and_suicidal_thoughts	44.8%	55.2%	100.0%
		% within Sex	100.0%	100.0%	100.0%
		Count			

Chi-Square Tests

	Value	Df	Asymp. Sig. (2-sided)
Pearson Chi-Square	50.788 ^a	6	.000
Likelihood Ratio	51.298	6	.000
Linear-by-Linear Association	.000	1	.988
N of Valid Cases	4725		

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 103.95.

Symmetric Measures

	Value	Asymp. Std. Error ^a	Approx. T ^b	Approx. Sig.
Interval by Interval Pearson's R	.000	.014	-.015	.988 ^c

Ordinal by Ordinal N of Valid Cases	Spearman Correlation	-.004 4725	.015	-.296	.767 ^c
---	-------------------------	---------------	------	-------	-------------------

- Not assuming the null hypothesis.
- Using the asymptotic standard error assuming the null hypothesis.
- Based on normal approximation.

CROSSTABS

```

/TABLES=Aboriginal_identity Age_group Sex BY Suicidal_Thoughts
/FORMAT=AVALUE TABLES
/STATISTICS=CHISQ CORR
/CELLS=COUNT ROW COLUMN
/COUNT ROUND CELL.

```

Crosstabs

Notes

Output Created		27-JUN-2024 20:19:25
Comments		
Input	Data	C:\Users\Ayeni Olabisi\Desktop\Research Methodology\Mr Ben Project\aps_cleaned.csv
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	4725
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
Syntax		CROSSTABS /TABLES=Aboriginal_identity Age_group Sex BY Suicidal_Thoughts /FORMAT=AVALUE TABLES /STATISTICS=CHISQ CORR /CELLS=COUNT ROW COLUMN /COUNT ROUND CELL.
Resources	Processor Time	00:00:00.02
	Elapsed Time	00:00:00.05
	Dimensions Requested	2

Cells Available

131029

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Aboriginal_identity * Suicidal_Thoughts	4725	100.0%	0	0.0%	4725	100.0%
Age_group * Suicidal_Thoughts	4725	100.0%	0	0.0%	4725	100.0%
Sex * Suicidal_Thoughts	4725	100.0%	0	0.0%	4725	100.0%

Aboriginal_identity * Suicidal_Thoughts

Crosstab

			Suicidal_Thoughts		Total
			No	Yes	
Aboriginal_identity	First nation	Count	1121	737	1858
		% within Aboriginal_identity	60.3%	39.7%	100.0%
		% within Suicidal_Thoughts	39.0%	39.8%	39.3%
	Inuit	Count	695	465	1160
		% within Aboriginal_identity	59.9%	40.1%	100.0%
		% within Suicidal_Thoughts	24.2%	25.1%	24.6%
	Metis	Count	1056	651	1707
		% within Aboriginal_identity	61.9%	38.1%	100.0%
		% within Suicidal_Thoughts	36.8%	35.1%	36.1%
Total	Count	2872	1853	4725	
	% within Aboriginal_identity	60.8%	39.2%	100.0%	
	% within Suicidal_Thoughts	100.0%	100.0%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.360 ^a	2	.507
Likelihood Ratio	1.362	2	.506
Linear-by-Linear Association	.845	1	.358
N of Valid Cases	4725		

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 454.92.

Symmetric Measures

		Value	Asymp. Std. Error ^a	Approx. T ^b	Approx. Sig.
Interval by Interval	Pearson's R	-.013	.015	-.919	.358 ^c
Ordinal by Ordinal	Spearman Correlation	-.013	.015	-.910	.363 ^c
N of Valid Cases		4725			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.

Age_group * Suicidal_Thoughts

Crosstab

			Suicidal_Thoughts		Total
			No	Yes	
Age_group	18 to 24 years	Count	743	753	1496
		% within Age_group	49.7%	50.3%	100.0%
		% within Suicidal_Thoughts	25.9%	40.6%	31.7%
	25 to 34 years	Count	697	371	1068
		% within Age_group	65.3%	34.7%	100.0%
		% within Suicidal_Thoughts	24.3%	20.0%	22.6%
	35 to 44 years	Count	714	358	1072
		% within Age_group	66.6%	33.4%	100.0%
		% within Suicidal_Thoughts	24.9%	19.3%	22.7%
	45 to 54 years	Count	718	371	1089
		% within Age_group	65.9%	34.1%	100.0%
		% within Suicidal_Thoughts	25.0%	20.0%	23.0%
Total	Count	2872	1853	4725	
	% within Age_group	60.8%	39.2%	100.0%	
	% within Suicidal_Thoughts	100.0%	100.0%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	113.907 ^a	3	.000

Likelihood Ratio	112.706	3	.000
Linear-by-Linear Association	77.687	1	.000
N of Valid Cases	4725		

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 418.84.

Symmetric Measures

		Value	Asymp. Std. Error ^a	Approx. T ^b	Approx. Sig.
Interval by Interval	Pearson's R	-.128	.015	-8.886	.000 ^c
Ordinal by Ordinal	Spearman Correlation	-.132	.015	-9.138	.000 ^c
N of Valid Cases		4725			

- a. Not assuming the null hypothesis.
b. Using the asymptotic standard error assuming the null hypothesis.
c. Based on normal approximation.

Sex * Suicidal_Thoughts

Crosstab

			Suicidal_Thoughts		Total
			No	Yes	
Sex	Male	Count	1398	719	2117
		% within Sex	66.0%	34.0%	100.0%
		% within Suicidal_Thoughts	48.7%	38.8%	44.8%
	Female	Count	1474	1134	2608
		% within Sex	56.5%	43.5%	100.0%
		% within Suicidal_Thoughts	51.3%	61.2%	55.2%
Total	Count	2872	1853	4725	
	% within Sex	60.8%	39.2%	100.0%	
	% within Suicidal_Thoughts	100.0%	100.0%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	44.412 ^a	1	.000		
Continuity Correction ^b	44.014	1	.000		
Likelihood Ratio	44.634	1	.000		
Fisher's Exact Test				.000	.000

Linear-by-Linear Association	44.403	1	.000		
N of Valid Cases	4725				

- a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 830.22.
b. Computed only for a 2x2 table

Symmetric Measures

		Value	Asymp. Std. Error ^a	Approx. T ^b	Approx. Sig.
Interval by Interval	Pearson's R	.097	.014	6.694	.000 ^c
Ordinal by Ordinal	Spearman Correlation	.097	.014	6.694	.000 ^c
N of Valid Cases		4725			

- a. Not assuming the null hypothesis.
b. Using the asymptotic standard error assuming the null hypothesis.
c. Based on normal approximation.

LOGISTIC REGRESSION VARIABLES Suicidal_Thoughts
/METHOD=ENTER Self_perceived_mental_health_and_suicidal_thoughts
Aboriginal_identity Age_group Sex SES
/CONTRAST (Aboriginal_identity)=Indicator
/PRINT=GOODFIT CI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

Logistic Regression

Case Processing Summary

Unweighted Cases ^a	N	Percent
Selected Cases		
Included in Analysis	4725	100.0
Missing Cases	0	.0
Total	4725	100.0
Unselected Cases	0	.0
Total	4725	100.0

- a. If weight is in effect, see classification table for the total number of cases.

Dependent Variable Encoding

Original Value	Internal Value
No	0
Yes	1

Categorical Variables Codings

		Frequency	Parameter coding	
			(1)	(2)
Aboriginal_identity	First nation	1858	1.000	.000
	Inuit	1160	.000	1.000
	Metis	1707	.000	.000

Block 0: Beginning Block

Classification Table^{a,b}

			Predicted		
			Suicidal_Thoughts		Percentage Correct
			No	Yes	
Step 0	Suicidal_Thoughts	No	2872	0	100.0
		Yes	1853	0	.0
	Overall Percentage				60.8

a. Constant is included in the model.

b. The cut value is .500

Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)
Step 0 Constant	-.438	.030	216.276	1	.000	.645

Variables not in the Equation

			Score	df	Sig.
Step 0	Variables	Self_perceived_mental_health_and_suicidal_thoughts	2.822	1	.093
		Aboriginal_identity	1.360	2	.507
		Aboriginal_identity(1)	.259	1	.611
		Aboriginal_identity(2)	.487	1	.485
		Age_group	77.704	1	.000
		Sex	44.412	1	.000
		SES	38.549	1	.000
Overall Statistics			161.746	6	.000

Block 1: Method = Enter

Omnibus Tests of Model Coefficients

		Chi-square	df	Sig.
Step 1	Step	174.777	6	.000
	Block	174.777	6	.000
	Model	174.777	6	.000

Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	6153.969 ^a	.036	.049

a. Estimation terminated at iteration number 5 because parameter estimates changed by less than .001.

Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	65.715	8	.000

Contingency Table for Hosmer and Lemeshow Test

		Suicidal_Thoughts = No		Suicidal_Thoughts = Yes		Total
		Observed	Expected	Observed	Expected	
Step 1	1	361	365.602	112	107.398	473
	2	320	334.089	153	138.911	473
	3	293	315.235	180	157.765	473
	4	369	302.442	104	170.558	473
	5	267	292.186	206	180.814	473
	6	281	279.155	192	193.845	473
	7	283	271.826	190	201.174	473
	8	250	257.546	223	215.454	473
	9	209	237.023	264	235.977	473
	10	239	216.896	229	251.104	468

Classification Table^a

			Predicted		
			Suicidal_Thoughts		Percentage Correct
			No	Yes	
Step 1	Suicidal_Thoughts	No	2553	319	88.9
		Yes	1470	383	20.7
	Overall Percentage				62.1

a. The cut value is .500

Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)
Step 1 ^a Self_perceived_mental_health_and_suicidal_thoughts	.022	.016	1.807	1	.179	1.022
Aboriginal_identity			.992	2	.609	
Aboriginal_identity(1)	.065	.070	.846	1	.358	1.067
Aboriginal_identity(2)	.007	.080	.007	1	.931	1.007
Age_group	-.231	.027	75.281	1	.000	.794
Sex	.425	.061	47.896	1	.000	1.530
SES	-.398	.071	31.159	1	.000	.672
Constant	-.712	.139	26.409	1	.000	.490

Variables in the Equation

		95% C.I. for EXP(B)	
		Lower	Upper
Step 1 ^a Self_perceived_mental_health_and_suicidal_thoughts		.990	1.056
Aboriginal_identity			
Aboriginal_identity(1)		.930	1.224
Aboriginal_identity(2)		.861	1.177
Age_group		.753	.836
Sex		1.356	1.725
SES		.584	.772
Constant			

a. Variable(s) entered on step 1: Self_perceived_mental_health_and_suicidal_thoughts, Aboriginal_identity, Age_group, Sex, SES.

LOGISTIC REGRESSION VARIABLES Suicidal_Thoughts

/METHOD=ENTER Self_perceived_mental_health_and_suicidal_thoughts

Aboriginal_identity Age_group Sex SES

/CONTRAST (Aboriginal_identity)=Indicator

/CONTRAST (Age_group)=Indicator

/CONTRAST (Sex)=Indicator

/PRINT=GOODFIT CI(95)

/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

Case Processing Summary

Unweighted Cases ^a	N	Percent
-------------------------------	---	---------

Selected Cases	Included in Analysis	4725	100.0
	Missing Cases	0	.0
	Total	4725	100.0
Unselected Cases		0	.0
Total		4725	100.0

a. If weight is in effect, see classification table for the total number of cases.

Dependent Variable Encoding

Original Value	Internal Value
No	0
Yes	1

Categorical Variables Codings

		Frequency	Parameter coding		
			(1)	(2)	(3)
Age_group	18 to 24 years	1496	1.000	.000	.000
	25 to 34 years	1068	.000	1.000	.000
	35 to 44 years	1072	.000	.000	1.000
	45 to 54 years	1089	.000	.000	.000
Aboriginal_identity	First nation	1858	1.000	.000	
	Inuit	1160	.000	1.000	
	Metis	1707	.000	.000	
Sex	Male	2117	1.000		
	Female	2608	.000		

Block 0: Beginning Block

Classification Table^{a,b}

Observed			Predicted		
			Suicidal_Thoughts		Percentage Correct
			No	Yes	
Step 0	Suicidal_Thoughts	No	2872	0	100.0
		Yes	1853	0	.0
Overall Percentage					60.8

a. Constant is included in the model.

b. The cut value is .500

Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)
Step 0 Constant	-.438	.030	216.276	1	.000	.645

Variables not in the Equation

	Score	df	Sig.
Step 0 Variables Self_perceived_mental_health_and_suicidal_thoughts	2.822	1	.093
Aboriginal_identity	1.360	2	.507
Aboriginal_identity(1)	.259	1	.611
Aboriginal_identity(2)	.487	1	.485
Age_group	113.907	3	.000
Age_group(1)	113.503	1	.000
Age_group(2)	11.614	1	.001
Age_group(3)	19.713	1	.000
Sex(1)	44.412	1	.000
SES	38.549	1	.000
Overall Statistics	196.893	8	.000

Block 1: Method = Enter

Omnibus Tests of Model Coefficients

	Chi-square	df	Sig.
Step 1 Step	208.362	8	.000
Block	208.362	8	.000
Model	208.362	8	.000

Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	6120.384 ^a	.043	.058

a. Estimation terminated at iteration number 5 because parameter estimates changed by less than .001.

Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	65.353	8	.000

Contingency Table for Hosmer and Lemeshow Test

		Suicidal_Thoughts = No		Suicidal_Thoughts = Yes		Total
		Observed	Expected	Observed	Expected	
Step 1	1	340	361.532	133	111.468	473

2	338	334.065	135	138.935	473
3	380	328.165	93	144.835	473
4	255	307.593	218	165.407	473
5	309	289.679	164	183.321	473
6	299	284.246	174	188.754	473
7	271	277.696	202	195.304	473
8	251	260.149	222	212.851	473
9	221	228.368	252	244.632	473
10	208	200.508	260	267.492	468

Classification Table^a

Observed			Predicted		
			Suicidal Thoughts		Percentage Correct
			No	Yes	
Step 1	Suicidal_Thought	No	2521	351	87.8
	s	Yes	1446	407	22.0
	Overall Percentage				62.0

a. The cut value is .500

Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)
Step 1 ^a Self_perceived_mental_health_and_suicidal_thoughts	.019	.016	1.337	1	.248	1.019
Aboriginal_identity			.938	2	.626	
Aboriginal_identity(1)	.067	.070	.895	1	.344	1.069
Aboriginal_identity(2)	.020	.080	.064	1	.800	1.020
Age_group			109.713	3	.000	
Age_group(1)	.670	.083	64.896	1	.000	1.954
Age_group(2)	.032	.092	.121	1	.728	1.032
Age_group(3)	-.043	.092	.215	1	.643	.958
Sex(1)	-.434	.062	49.454	1	.000	.648
SES	-.389	.071	29.894	1	.000	.678
Constant	-.609	.101	36.629	1	.000	.544

Variables in the Equation

		95% C.I.for EXP(B)	
		Lower	Upper
Step 1 ^a	Self_perceived_mental_health_and_suicidal_thoughts	.987	1.053
	Aboriginal_identity		
	Aboriginal_identity(1)	.931	1.227
	Aboriginal_identity(2)	.872	1.194
	Age_group		
	Age_group(1)	1.660	2.300
	Age_group(2)	.863	1.235
	Age_group(3)	.800	1.147
	Sex(1)	.574	.731
	SES	.589	.779
	Constant		

a. Variable(s) entered on step 1: Self_perceived_mental_health_and_suicidal_thoughts, Aboriginal_identity, Age_group, Sex, SES.

FREQUENCIES VARIABLES=Aboriginal_identity Age_group Sex
Self_perceived_mental_health_and_suicidal_thoughts Suicidal_Thoughts
/ORDER=ANALYSIS.

Frequencies

Statistics

		Aboriginal_i dentity	Age_group	Sex	Self_perceive d_mental_he alth_and_suic idal_thoughts	Suicidal_Tho ughts
N	Valid	4725	4725	4725	4725	4725
	Missing	0	0	0	0	0

Frequency Table

Aboriginal_identity

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	First nation	1858	39.3	39.3	39.3
	Inuit	1160	24.6	24.6	63.9
	Metis	1707	36.1	36.1	100.0
	Total	4725	100.0	100.0	

Age_group

		Frequency	Percent	Valid Percent	Cumulative Percent
--	--	-----------	---------	------------------	-----------------------

Valid	18 to 24 years	1496	31.7	31.7	31.7
	25 to 34 years	1068	22.6	22.6	54.3
	35 to 44 years	1072	22.7	22.7	77.0
	45 to 54 years	1089	23.0	23.0	100.0
	Total	4725	100.0	100.0	

Sex

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Male	2117	44.8	44.8	44.8
Female	2608	55.2	55.2	100.0
Total	4725	100.0	100.0	

Self_perceived_mental_health_and_suicidal_thoughts

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Ever seriously contemplated suicide	632	13.4	13.4	13.4
Excellent or very good self-perceived mental health	968	20.5	20.5	33.9
Fair or poor self-perceived mental health	489	10.3	10.3	44.2
Good self-perceived mental health	886	18.8	18.8	63.0
Never seriously contemplated suicide	1018	21.5	21.5	84.5
Seriously contemplated suicide in the past 12 months	232	4.9	4.9	89.4
Seriously contemplated suicide but not in the past 12 months	500	10.6	10.6	100.0
Total	4725	100.0	100.0	

Suicidal_Thoughts

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid No	2872	60.8	60.8	60.8

Yes	1853	39.2	39.2	100.0
Total	4725	100.0	100.0	