

Estimated dietary intakes of vitamin A5

Torsten Bohn (1), Marta Despotovic (2), Farhad Vahid (1), and Ralph Rühl (3)

- (1) Nutrition and Health Research Group, Department of Precision Health, Luxembourg Institute of Health, L-1445 Strassen, Luxembourg
- (2) Institute for Medical Research, Department of Nutrition and Metabolism, Belgrade, Serbia
- (3) CISCAREX UG, Berlin, Germany

Correspondence:

Dr. Ralph Rühl
CISCAREX UG
Transvaalstr. 27c
D-13351 Berlin
Email: ralphruehl@ciscarex.com

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Abstract:

A new vitamin concept, termed vitamin A5, was recently identified. Furthermore, dietary recommendations in the range of 1,1 (0,5 – 1,8) mg vitamin A5 / day were suggested by an international expert consortium. The ensuing question arises as to the current daily dietary intake amounts in the Western civilization. Addressing this answer included calculating the intake based on known amounts of vitamin A5 in frequently consumed food items of the human diet high in this vitamin as well as the known daily dietary intake amounts of those selected food components in Westernized countries. Regarding food items, amounts of vitamin A5 in the form of provitamin A5 (i.e. 9-*cis*-beta-carotene (9CBC)), the predominant form in the diet, were found to range from 0,1 to 39 µg 9CBC / g for individual fruits and vegetables, with highest concentrations in leafy vegetables. The average intake amounts of vitamin A5 in adults of the general population following a Western lifestyle in Europe, averaged 0,9 with a range of 0,5 (for Austria) – 1,3 (for Italy) mg 9CBC / day. Furthermore, based on our calculations, large parts, i.e. ~ 2/3rd of the population are low, even too low (<1.1 mg/day), in daily vitamin A5 intake. In addition to the importance of nudging the population towards a regrettably non-well accepted higher intake of fruits and vegetables, an additional fortification and supplementation of vitamin A5 could be considered, similar as to other micronutrients that are low a Westernized diet.

Key-words: Carotenoids, 9-*cis*-beta-carotene, dietary intake recommendations, fruit and vegetable consumption, intake stratification, supplementation.

Introduction:

A new vitamin concept was recently proposed for vitamin A5, as the first novel identified vitamin for 80 years [1]. The active form of this vitamin is formed out of its precursors, 9-*cis*-beta-carotene (9CBC), originating from plant derived food items [2], especially from fruits and vegetables, as well as out of 9-*cis*-13,14-dihydroretinol, originating from animal sources [3]. These two vitamin A5 derivatives are the nutritional relevant precursors of 9-*cis*-13,14-dihydroretinoic acid, the molecule relevant for the endogenous activation of the nuclear hormone receptor, the retinoid X receptors (RXRs) [2].

Historically, it was only after clearly observable vitamin deficiencies were identified in humans that the underlying individual and health-relevant mechanism of action was identified [1]. Later on, further detailed dietary recommendations were finally suggested by national and international authorities, such as the USDA in the US [1] or EFSA for Europe [4]. Traditionally, vitamin intake recommendations were based on studies – human observational or even intervention studies or animal ones – that prevented typical, i.e. vitamin related deficiency symptoms [1, 5]. Today, after 80 years since the discovery of the last vitamin [6], different strategies for the identification and establishment of potential novel vitamin concepts are more appropriate. Such a novel strategy is needed, mostly due to ethical restrictions for deprivation and supplemental dose-escalating studies of perceived important dietary constituents in humans [7] and regarding animal studies in sight of difficulties to transfer findings from animal models to humans, due to different metabolism and body weight [7, 8]. In addition, strategies performed around 100 years ago included randomly supplementing people with diseases in a non-ethical and non-standardized way and are not possibly anymore [9], and have been replaced by extremely work intensive and highly expensive standardized clinical procedures [10]. Targeted supplementation trials that exclude specific food components rich in a specific vitamin in a person's diets, such as performed earlier for vitamin B3 employing war prisoners [11], are for obvious reasons not ethically possible anymore.

Instead, today, a backwards strategy, starting i) from a unique mechanism of action involving the active vitamin form in blood / tissues and ii) relevant cellular concentrations related to normal physiological metabolism towards iii) considering nutritional precursors from food components and calculation of their daily intake amounts, also accounting for iv) aspects of their bioavailability, are a more feasible and logical strategy [12, 13].

In summary, establishing a new vitamin category today requires different strategies and measures than a century ago, especially using state-of-the-art laboratory experiments for establishing a novel food-dependent mechanism of action with crucial health benefits.

Regarding the large picture for vitamin A5, many of these puzzle pieces have been sorted out [1, 2], including its biological relevance as the ligand for the RXR nuclear receptor and the presence of precursor molecules in the diet. However, some crucial parts along the way of establishing intake recommendations can only be addressed following a logical step-by-step procedure connecting the various aspects of tissue concentrations, bioavailability and bioconversion aspects and occurrence in food items together [1].

In this article, we add another important piece to the vitamin A5 story, based on a calculated daily intake of this vitamin in Westernized countries. Based on these estimations, the nutritional situation of these countries with respect to vitamin A5 supply can be evaluated and we can further stratify which subgroups regarding age, gender, or country, may be potentially low in their daily intake levels of this novel vitamin.

Materials and methods:

Vitamin A5 / Provitamin A5 levels in food items

First, we summarized, based on published studies and databases [14, 15], the vitamin A5 levels in the form of provitamin A5 (9CBC) in food items, focusing on fruits and vegetables as provitamin A5-rich sources that are also frequently consumed. For this purpose, a review of published peer-reviewed literature was conducted to identify relevant data on vitamin A5 levels in food items [1].

In addition to the EFSA food consumption database [15], one additional study that received particular attention was the EPIC cohort [14], as fruit and vegetable intake was available at relevant sub-categories, i.e. leafy vegetables, cabbages, root vegetables, fruiting vegetables, onions/garlic as well as other vegetables, which was not the case for the EFSA food consumption database [15]. We combined the averaged data of 9CBC concentrations in vegetable and fruit subgroups from Table 1A to calculate a weighted average of 0.2 µg 9CBC / g fruits and 5.2 µg 9CBC / g vegetables (Table 1B).

Consumption of fruits and vegetables as well as their subgroups

In a second step, we summarized, based on European food consumption data [15], the average amounts of these food items (i.e. fruits and vegetables) consumed by different populations in individual European countries. For this purpose, due to the large amount of available data and due to the standardized approach by which the data was obtained, we have focused largely on EFSA's food consumption database summarizing food consumption patterns across member states (displayed in Table 2) [16] and the EPIC cohort based database displayed in Table 3 [14].

The EFSA food consumption database provided detailed data on food intake patterns, including frequency and quantity of fruit and vegetable consumption, across various demographic groups (age and gender based). When several populations or studies were available for a European country, the

arithmetic mean per each study was first calculated. Thereafter, the average consumption per country level was calculated, which was then termed “mean of means”.

In addition to food intake data, information was also collected on fruit and vegetable intake recommendations per country, in order to compare estimated intakes with the respective countries’ food based dietary guidelines.

Calculation of provitamin A5 intake

As a third step, Further we calculated the intake of 9CBC by means of the more detailed EPIC cohort data, by combining the average 9CBC contents from Table 1B with the food intake data from EPIC (Table 3) per country. Furthermore, we calculated, based on these estimations and the previous intake estimates the amount of provitamin A5 consumed per day based in each European nation (Figure 3 and 4).

Statistical Analysis

Descriptive statistics, including mean and standard deviation (SD), were computed to summarize the average daily intake levels of provitamin A5 across different population groups. Frequency distributions were also examined to identify patterns of vitamin A5 intake within each population.

Results

Concentration of vitamin A5 in the form of provitamin A5 in individual food items

A limited number of publications exists reporting concentrations of vitamin A5 / 9CBC, expressed as provitamin A5, i.e. 9CBC, in fruits and vegetables (Table 1). The average concentration of 9CBC in fresh fruits was found to be 0.2 µg 9CBC / g fruits and the estimated concentration in vegetables was 4.7 µg 9CBC / g vegetables, while a weighted average estimation, considering different intake amounts of individual vegetable subgroups, resulted in an average of 5.2 µg 9CBC / g vegetables, which was used for all further calculations.

When comparing different sub-groups of vegetables, the highest concentrations appeared to be present in leafy vegetables, followed by root vegetables and cabbages. It should be noted that fruit juices, for lack of data regarding 9CBC, were not included in this evaluation.

Data regarding vitamin A5 intake in the form of 9CDHROL and 9CDHROL-esters from animal based products were not reported, as there are currently no data available and concentrations have been reported to be very low [2].

Summary of current intake recommendations for fruits and vegetables

Daily intake of fruit and vegetables in Europe

A. Based on the EFSA food consumption database

Data from 24 countries, 23 based on EFSA's food consumption database, encompassing a total of 26.600 participants aged 18-64 years were included in this investigation. As shown in figure 1 and table 2, fruit intake ranged from 91 g / day (Sweden and Ireland) to 287 g/day (Luxembourg) with an

average of 147 g / day. Vegetable intake was reported lowest in Austria (90 g / day) and highest in Italy (236 g / day) and an average of 164 g / day. This translates into a highest combined intake of fruits and vegetables with 503 g / day in Luxembourg, lowest in Sweden with 218 g / day and an average of 312 g / day.

Most countries did not even reach the minimum WHO recommended 400 g fruits and vegetables / day: According to EFSA's food consumption database, out of the 23 countries listed in table 2, with only Italy and Luxembourg reaching the recommended 400 g of fruits and vegetables / day (i.e. 5 portions per day) target. Especially Southern European countries such as Spain, Greece, Portugal, Italy and Cyprus showed generally higher combined fruit and vegetable intakes compared to Northern European countries such as Ireland, Denmark, Sweden, or Finland.

B. The EPIC cohort

When comparing findings by EFSA to the EPIC cohort, in which data was generally also obtained by 24 h recalls (though not by multiple ones), similar tendencies were found for fruit and vegetable intakes with a higher average of 417 g / day in comparison to 312 g / day in the EFSA database. The average vegetable intake in the EPIC database was 175 g / day and comparable to the 164 g / day in the EFSA database. The average fruit intake was 242 g / day in the EPIC cohort while the EFSA values were much lower with 147 g / day.

Highest intake of fruits / vegetables per capita (Table 3) were found comparable to the EFSA database in the Southern European Nations like Spain (573 g / day), followed by Italy (569 g / day) and Greece (496 g / day), while lowest once were mainly determined also comparable in Northern European countries like Sweden (262 g / day) and the Netherlands (313 g / day). With regard to the recommended 400 g of fruit and vegetables per day, only 4 out of 8 countries (Table 3 and Figure 1), for which data were available, for both men and women met this intake recommendation.

Calculation of individual intake of vegetable subgroups

Regarding vegetable subgroups as determined based on the EPIC cohort study, a total of 22 g leafy vegetables (men and women combined, out of 175 g total vegetables) were consumed, based on data from 8 countries reporting both data of men and women (Table 3).

Regarding fruits, an average of 242 g / day were consumed. Southern countries such as Italy, Spain and Greece showed consistently higher intakes for vegetables, including leafy ones and fruits compared to Northern European countries such as Denmark, the Netherlands, Germany and Sweden. An exception was noted for cabbages, which did not follow this trend, as high concentrations were consumed in the UK, Denmark and Germany. When looking at subgroups in man or woman, no remarkable overall differences were apparent for the intake of total vegetables, leafy vegetables, and individual subgroups (Figure 2).

Estimating the daily intake of provitamin A5

Based on the EPIC cohort about half of the countries would fall below the earlier proposed 1.1 mg / day intake of provitamin A5 (Figure 2 and 3). Only the Southern European countries, due to their generally higher intake of fruits and vegetables (Tables 2-3 and Figure 2) were estimated to reach the 1.1 mg / day target, while Northern and Central European countries had lower intakes as 0.5 mg / day. Employing data from EFASs Food Consumption database, and also assuming an approx. content of 9CBC of 0.2 µg / g for fruits and 5.2 µg / g for vegetables (based on Table 1B) based on this cruder estimate, only few countries like Italy, Hungary, Luxembourg and Serbia reached the 1.1 mg / day 9CBC target while the majority of 19 countries did not reach the suggested 1,1 mg 9CBC intake / day. The European average was comparable with 0.9 mg 9CBC / day in the EPIC cohort and 0.9 mg 9CBC in the EFSA cohort (Figure 3).

218
219 Due to their high content of 9CBC combined with their relatively frequent consumption, leafy
220 vegetables were the vegetable subgroup most largely contributing to total provitamin A5 intake
221 (contributing to often >50% of total 9CBC intake), followed by root vegetables, cabbages and other
222 vegetables (Figure 2). Fruits contributed, due to their low content of 9CBC, very little to overall
223 intake, below 5% (Figure 2). Slight differences between men and women were noted, with women
224 showing slightly lower average intakes of 9CBC.

225
226 When looking at 9CBC intake by age in the EFSA cohort (Figure 3C), the differences between the age
227 groups appeared to be small, with the highest intakes in elderly adults aged 65-74 years. This is
228 somewhat confirmed by EFSA data [15], as fruit and especially vegetable consumption is highest also
229 for the elderly compared to adolescents and the elderly.

230
231 **Comparison of provitamin A5 intake in persons consuming low, medium and high amounts of fruits**
232 **and vegetables (Figure 4)**

233 Based on data from a study just performed in France [17], the majority (>50%) consumed less than
234 3.5 portions fruits and vegetables / day with an average of 2 portions, reach in consequence an
235 average intake of only 0.6 mg provitamin A5 / day. Furthermore, only 21% of the French population
236 consumed between 3,5 and 5 portions of fruits and vegetables per day, based on an average of 4,2
237 portions of fruits and vegetables / day and consequently an estimated provitamin A5 intake of 1,3
238 mg 9CBC / day [18]. Only 25 % of the French population consumed sufficient fruits and vegetables
239 with 5 and more portions / day and an estimated provitamin A5 intake of 2,1 mg 9CBC / day [18, 19].
240 In consequence, this means that ~64% (~2/3rd) of the French population consume too few fruits and
241 vegetables per day and in consequence also a too low daily intake of provitamin A5.

242

Discussion:

In this article we present, for the first time, an estimation of vitamin A5 intake in various European countries. This compound has been identified and categorized as a novel vitamin and / or an independent subcategory of vitamin A and an intake estimate across various European populations was required. Toward this end, we calculated vitamin A5 intakes independently from vitamin A(1) from the background diet. An average intake of vitamin A5 in the adult population ranging from 0.5 to 1.3 mg per day was found, with a tendency for higher intakes in Southern European countries. It was also estimated that approx. 2/3 of the population would consume less than the proposed recommended intake of 1.1 mg provitamin A5 / day.

Comparable to other vitamins, we make use of available data from cohort studies, in our case from the EPIC [14] and EFSA cohorts [15], allowing to calculate the daily intake specified for several European countries, sex and partly age groups, which are currently available in these databases. As a first step, based on available data, we calculated the average intake of vitamin A5 in food categories that are rich in vitamin A5. Here we focused on fruits and vegetables, where provitamin A5, i.e. 9-*cis*- β -carotene, is mainly present. The levels of 9-*cis*-13,14-dihydroretinol, as well as its esters in the human food matrix have thus far either not been determined or are deemed very low [3], suggesting that the main intake of vitamin A5 is via plant foods rich in provitamin A5 [20]. At first, an average provitamin A5 concentration was calculated, with 0.2 μ g provitamin A5 / day for fruits and 5.2 μ g provitamin A5 / day for vegetables, which were then employed for further calculations. This was required, as most databases do not contain further details on the consumption of individual fruits and vegetable subgroups.

The highest daily intake amounts of fruits and vegetables was determined in Southern European countries such as Greece, Spain and Italy, and lowest in Northern European countries including Sweden, the Netherlands and Denmark. Remarkable is that the large majority of Europeans was

consuming less than the suggested “5 A DAY”-intake recommendations of 5 portions of each 80 g fruits and vegetables by national and international food organizations [14, 15, 21]. However, this has been remarked in earlier consumption surveys in Europe [14].

Regarding the best sources of provitamin A5 per weight, the main source of provitamin A5 intake originates from the vegetable sub-group leafy vegetables, with 20.3 µg provitamin A5 / g food matrix, followed by cabbages (6.4 µg/g) and root vegetables (6.4 µg/g), with low concentrations in fruiting vegetables (1.0 µg/g) and onions / garlic (0.1 µg/g). The European country-specific daily intake amount of these leafy vegetables was highest in Southern European countries such as Greece, Spain and Italy and lowest in Northern / Central European countries including UK, Norway, Denmark and Sweden. This is in line with studies reporting higher fruit and vegetable intake in general in Southern European countries [14].

Based on the fruit and vegetable intake amounts from the EPIC and the EFSA database [14, 15], summarizing 65,544 individuals, we calculated for 27 European countries country specific daily vitamin A5 intake amounts with an average of 0.9 mg provitamin A5 / day, which is lower than the previously suggested recommendation of 1.1 mg provitamin A5 / day, calculated by two independent methods [1]. These estimated average intake amounts of provitamin A5 on average of European countries sound moderate / fair compared to the suggested intake amounts. While when stratifying for different countries, we observed that the average in the non-Southern European countries such as especially Austria (0.5 mg/day), Germany (0.5 mg/day), Czech Republic (0.6 mg/day), Sweden (0.7 mg/day EFSA data / 0.6 mg/day EPIC data) were clearly below the suggested intake amounts of 1.1 mg provitamin A5 / day. Based on the estimations of daily provitamin A5 intake from the 27 European countries, only 4 out of 24 countries for the EFSA database calculations and 3 out of 8 from the EPIC database calculations reach sufficient high daily provitamin A5 intake levels on average, while the large majority of the European countries 3 (based on the EPIC database) and 8 (based on

the EFSA database) being even below 0.8 mg provitamin A5 / day on average and thereby far below the suggested daily intake amounts of provitamin A5. This means that the large majority of European countries on average have a too low daily intake of daily vitamin A5. However, it also needs to be specified that both the intake recommendation as well as the calculated intake are based on average values for adults, and due to the limited amount of data available. Further studies should aim to determine detailed estimations such as allowing estimating an intake that would cover over 97% of the population's need, such as necessary to determine the population reference intake (PRI).

Further based on evaluations in France [17] it was calculated that 54% of the French population eat less than 3.5 daily portions of fruits and vegetables and a further considerable fraction (21 %) consume between 3.5 – 5 portions, which means that ca. 64% of the French population are below the suggested “5 A DAY” recommendations. This means as a consequence that ~2/3 of the French population, as well as transferable to many other European populations are below the suggested daily intake levels of vitamin A5 which were based on the “5 A DAY”-concept.

In summary, due to the low intake of fruits and vegetables [22], especially vegetables [14] and with special focus on low intake of leafy vegetables [14], there is an estimated low, even too low, daily dietary intake of vitamin A5 present in Europeans. This data is likely relevant for other rather Western-lifestyle based societies following western based dietary patterns, which includes large populations in many developing countries [1, 23]. We recommend, based on these calculated parameters that, in order to improve dietary patterns and assure also sufficient intake of vitamin A5 that a) novel dietary recommendations should focus mainly on a higher daily dietary intake of vegetables and especially leafy vegetables and b) consider additional food fortification with vitamin A5 and / or dietary supplements to meet the suggested daily dietary intake recommendations for vitamin A5 to avoid a potential vitamin A5 deficiency [20].

Whether a low daily intake of vitamin A5 is associated with mental problems, which are prevalent in our Western hemisphere [20] is currently merely predicted and requires further attention based on European-wide epidemiological databases with sufficient details for specific European countries regarding dietary intake and mental health issues. Not surprisingly, in Southern European countries like Greece, Italy and Croatia with a high average estimated vitamin A5 intakes, the mental problems appear to have a lower prevalence, while in Northern European countries like the Netherlands, Finland, Denmark, Sweden and Germany these mental problems are reported to have a high prevalence [24]. A direct correlation and causal connection must still be elaborated and evaluated in larger databases, together with deeper functional mechanistic evaluations. Furthermore, described and suggested long-term or medium-term vitamin A5-deficiencies such as poor mental health and high incidence of numerous neurological diseases should be evaluated with respect to nutritional or functional vitamin A5-signalling pathways [20]. While poor mental health is described to be based on low daily intake of fruits and vegetables, especially vegetables [25] and with a special focus on leafy vegetables [26]. A detailed clinical mechanistic step-by-step approach how fruits and vegetables, with a focus on vegetables and leafy vegetables rich in vitamin A5, transmit their health-promoting and –protecting activity, via the retinoid X receptor, was summarized in a recent article [1], which is based on dozens of mechanistic studies performed using laboratory models as a base.

In this study, a more detailed and data-based intake of vitamin A5 in several European countries was attempted and results were compared against a first recommendation for vitamin A5 intake. The calculated data were evaluated in an ideal case dietary scenario as well as in a reality-based scenario [1], showing that large parts of the European society may be too low in daily intake of vitamin A5.

Author contributions

RR was responsible for the idea and outline and writing of the article, FV did export and calculate food consumption data, TB was involved in writing the article and critically reviewing the concept.

347

348 **Conflict of interest**

349 RR is CEO and shareholder of CISCAREX UG.

350

351

352

Table 1: Concentration of 9CBC in µg 9CBC/g food matrix

- A -	9CBC dry weight in µg/g	9CBC fresh weight in µg/g
food item		
fruits		
apricot	4.4	0.6
banana	0.3	0.1
cherry	-	-
date	-	-
grape	-	-
guava	-	-
mango	-	-
nectarine	-	-
papaya	7.0	1.3
peach	-	-
pear	-	-
prune	-	-
peach	-	0.3
average of all fruits	0.9	0.2
vegetables		
lettuce #	41	2.0
spinach #		38.6
cabbage ##	-	-
kale ##		14.1
broccoli ##		5.0
carrot §	57.1	6.5
cucumber §§	-	-
eggplant §§	-	-
pumpkin (1) §§	2.7	0.2
tomato (1) §§	-	-
zucchini §§	-	-
pumkin (2) §§		2.5
tomato (2) §§		4.8
pepper (red) §§	6.0	0.6
pepper (red hot) §§	9.6	1.0
onion (green) §	-	-
parsley §§	111.2	11.6
dill §§	27.7	4.3
potato §§		-
sweet potato 1 §§	65.1	10.2
sweet corn §§	-	-
sweet potato 2 §§		1.5
average of all vegetables	20.1	4.7
- B -		
# - average for leafy vegetables		20.3
## - average for cabbages		6.4
§ - average for root vegetables		6.5
§§ - average for fruiting vegetables		1.0
\$ - average for onions / garlic		0.1
\$\$ - average for other vegetables		4.6
adjusted average of all vegetables		5.2 *

Some data were not measured indicated by a "blank", some data were under the quantification limit and indicated by a "-" which was set with 0.05 as half of the quantification limit for further calculation, * - this calculation is based on the listed distribution of different vegetable sub-groups given in the EPIC cohort study [14] with a calculated 13% leafy vegetables, 12% cabbages, 13% root vegetables, 42 % fruiting vegetables, 7% onion / garlic and 14% other vegetables. Numbers 1 [27] and 2 [28] indicate different levels of 9CBC based on different studies.

Table 2: Fruits (F) and vegetables (V) intake (g per day) from the EFSA database of different European countries[†].

Countries	Fruits	Vegetables	F + V	% V	Measurement method and Survey name	Sample size	Year
Italy	171 ± 143	236 ± 155	407	58	Italian national dietary survey on the adult population from 10 up to 74 years old	726	2018
Hungary	138 ± 139	235 ± 166	373	63	Hungarian national food consumption survey	529	2018
Serbia	137 ± 164	217 ± 165	354	62	Serbian Food Consumption Survey on adults	1150	2019
Luxembourg*	287 ± 268	216 ± 172	503	43	ORISCAV-LUX 2 [29]	1326	2017
Latvia	141 ± 170	203 ± 133	345	59	Latvian National Dietary Survey	1080	2012
Cyprus	111 ± 135	201 ± 145	312	65	National dietary survey of the adult population of Cyprus	272	2014
Greece	121 ± 164	192 ± 143	313	61	The EFSA-funded collection of dietary and related data in the general population aged 10-74 years in Greece	260	2014
Portugal	161 ± 140	182 ± 130	343	53	National Food, Nutrition, and Physical Activity Survey of the Portuguese general population	3102	2015
Croatia	136 ± 153	174 ± 158	310	56	Croatian food consumption survey on adults	2002	2011
Estonia	226 ± 231	168 ± 159	394	43	National Dietary Survey among 11-74 years old individuals in Estonia	2124	2013
Montenegro	120 ± 110	163 ± 121	326	50	Montenegrin National Dietary Survey on the general population	697	2017
Netherlands	115 ± 128	163 ± 117	279	59	Dutch National Food Consumption Survey 2012-2016 (DNFCS)	1487	2012
Romania	163 ± 186	154 ± 110	317	49	Romanian national food consumption survey for adolescents, adults, and elderly	740	2019
Slovenia	158 ± 151	153 ± 105	311	50	Slovenian national food consumption survey	385	2017
Ireland	91 ± 95	151 ± 84	242	62	North/South Ireland Food Consumption Survey	958	1997
Bosnia and Herzegovina	130 ± 144	149 ± 122	279	53	Bosnia-Herzegovinian Dietary Survey of adolescents, adults, and pregnant women	850	2017
Spain	155 ± 129	144 ± 102	299	48	Spanish National dietary survey in adults, elderly, and pregnant women	536	2013
France	133 ± 131	144 ± 90	277	52	Individual and national study on food consumption 2	2276	2007
Finland	171 ± 178	126 ± 102	297	43	National Findiet Surveys	1575	2007
Sweden	91 ± 104	126 ± 99	218	59	National Diet and Nutrition Survey - Years 1-3	1266	2008
Belgium	118 ± 122	125 ± 100	244	52	Diet National 2004	1292	2004
Czech Republic	124 ± 123	117 ± 91	241	49	Czech National Food Consumption Survey	1666	2003
Germany	165 ± 175	95 ± 94	260	37	National Nutrition Survey II	10,419	2007
Austria	162 ± 153	90 ± 96	252	36	Austrian Study on Nutritional Status 2010-12 - Adults	308	2010
European average ± STD	147 ± 42	164 ± 41	312 ± 64	53 ± 8	Calculated sum of the 24 listed surveys	37026	

Countries were sorted from top to bottom, based on the average vegetable intake.

[†] European Food consumption data [16]). Age range from 18-64 years.

* median and interquartile range

Table 3: Summarised data from the EPIC cohort based on the data from the year 2002 [14].

men		fruits (F)	vegetables (V)	F + V	leafy V	% V / F+V	% LV / V
Country	n	mean	mean		mean		
Greece	1312	273	270	543	30	50	11
Spain	1777	372	224	596	44	38	19
Italy	1444	403	211	614	35	34	17
Germany	2268	207	160	368	14	44	9
Netherlands	1024	168	137	305	28	45	20
UK *	518	206	193	400	10	48	5
Denmark	1923	160	141	301	10	47	7
Sweden	2765	122	112	234	9	48	8
sum # / av ##	13031	239	181	420	23	44	12
STD		102	53	146	13	5	6
women		fruits (F)	vegetables (V)	F + V	leafy V	% V / F+V	% LV / V
Country	n	mean	mean		mean		
Greece	1374	242	207	449	29	46	14
Spain	1443	355	196	551	34	36	17
Italy	2512	343	181	524	28	34	16
Germany	2150	236	166	403	17	41	10
Netherlands	2960	192	130	322	23	40	18
UK *	768	223	192	415	15	46	8
Denmark	1995	206	150	356	11	42	7
Sweden	3285	164	127	290	11	44	9
France **	4639	251	226	476	45	47	20
Norway **	1798	168	125	293	7	43	5
sum # / av ##	22924	238	170	408	22	42	12
STD		66	36	93	12	4	5
women and men		fruits (F)	vegetables (V)	F + V	leafy V	% V / F+V	% LV / V
Country	n	mean	mean		mean		
Greece	2686	257	238	496	29	48	12
Spain	3220	363	210	573	39	37	18
Italy	3956	373	196	569	32	34	16
UK *	1286	215	193	407	13	47	6
Germany	4418	222	163	385	15	42	9
Denmark	3918	183	145	328	10	44	7
Netherlands	3984	180	133	313	25	43	19
Sweden	6050	143	119	262	10	46	8
sum # / av ##	29518	242	175	417	22	43	12
STD		85	41	118	11	5	5

The countries are sorted from top to bottom, based on the vegetable intake of women and men, n – number of included persons, av – average, * just health aware people were included, ** just woman were examined for this country, # - the sum is relevant for the number (n) of included persons, ## - average of fruits intake, vegetable intake, F + V intake, leafy V intake, % V / F+V and % LV / V.

Figures:

Figure 1: Schematic summary of a dietary pattern focusing on fruit and vegetable intakes in different European countries. *The dashed line indicates the recommended “5 A DAY” recommendation of 5 portions of 80 g fruits and / or vegetables for daily intake.*

Figure 2: Schematic summary of the calculated vitamin A5 intake levels in different European countries. Used abbreviations; V - vegetables

Figure 3: Calculated individual daily 9CBC intake levels based from the data of the a) the EPIC cohort [14], b) summarized national surveys and c) age depending intake, both based on EFSA's Food Consumption database [16] : All estimated values are based on average values for 9CBC in fruits (0.2 µg/g) and vegetables (5.2 µg/g). *The dashed line indicates the recommended intake of 1.1 mg 9CBC (provitamin A5) /day [1].*

Figure 4: Calculated daily 9CBC intake levels of individuals with low, medium and high daily average fruit and vegetable intake from a French survey [14] with an average F+V intake of 295 g F+V / day. *The dashed line indicates the recommended intake of 1.1 mg 9CBC (provitamin A5) /day [1].*

Figure 1:




Figure 2:






Figure 3:

A. EPIC cohort database



B. EFSA - national survey database



C. Comparison age





Figure 4:

Comparison for low, medium and high intake of fruits and vegetables



average intake of F+V portions per day	2	4,2	6,9	3,7
representing % of the French population	54	21	25	

References:

- [1]. Bohn T, Hellmann-Regen J, de Lera Á R, Böhm V, Rühl R. Human nutritional relevance and suggested nutritional guidelines for Vitamin A5/X and Provitamin A5/X. submitted to Nutrition Reviews. 2022.
- [2]. Krężel W, Rivas A, Szklenar M, Ciancia M, Alvarez R, de Lera AR, Rühl R. Vitamin A5/X, a new food to lipid hormone concept for a nutritional ligand to control RXR-mediated signaling. *Nutrients*. 2021; 13(3): 925.
- [3]. Rühl R, Krężel W, de Lera AR. 9-Cis-13, 14-dihydroretinoic acid, a new endogenous mammalian ligand of retinoid X receptor and the active ligand of a potential new vitamin A category: vitamin A5. *Nutrition reviews*. 2018; 76(12): 929-41.
- [4]. Additives EPoF, Nutrient Sources added to F. Scientific Opinion on the re-evaluation of mixed carotenes (E 160a (i)) and beta-carotene (E 160a (ii)) as a food additive. *EFSA Journal*. 2012; 10(3): 2593.
- [5]. Organization WH. Vitamin and mineral requirements in human nutrition: World Health Organization; 2004.
- [6]. Scott JM, Molloy AM. The discovery of vitamin B(12). *Ann Nutr Metab*. 2012; 61(3): 239-45.
- [7]. van der Worp HB, Howells DW, Sena ES, Porritt MJ, Rewell S, O'Collins V, Macleod MR. Can animal models of disease reliably inform human studies? *PLoS Med*. 2010; 7(3): e1000245.
- [8]. Rucker RB, Rucker MR. Nutrition: ethical issues and challenges. *Nutr Res*. 2016; 36(11): 1183-92.
- [9]. Semba RD. The discovery of the vitamins. *Int J Vitam Nutr Res*. 2012; 82(5): 310-5.
- [10]. Hardman TC, Aitchison R, Scaife R, Edwards J, Slater G. The future of clinical trials and drug development: 2050. *Drugs Context*. 2023; 12.
- [11]. Roe DA. A plague of corn. The social history of pellagra. 1973.
- [12]. van Het Hof KH, West CE, Weststrate JA, Hautvast JG. Dietary factors that affect the bioavailability of carotenoids. *J Nutr*. 2000; 130(3): 503-6.
- [13]. Cifuentes M, Vahid F, Devaux Y, Bohn T. Biomarkers of food intake and their relevance to metabolic syndrome. *Food & Function*. 2024.
- [14]. Agudo A, Slimani N, Ocke M, Naska A, Miller A, Kroke A, et al. Consumption of vegetables, fruit and other plant foods in the European Prospective Investigation into Cancer and Nutrition (EPIC) cohorts from 10 European countries. *Public health nutrition*. 2002; 5(6b): 1179-96.
- [15]. EFSA. The EFSA Comprehensive European Food Consumption Database. EFSA; 2022.
- [16]. EFSA. EFSA Food Consumption Database. <https://www.efsa.europa.eu/en/data-report/food-consumption-data>. 2022; accessed 27.02.2024.
- [17]. Tavoularis G, Hébel P. Fruits et légumes : les Français suivent de moins en moins la recommandation. *Credoc - Consommation et modes de vie*. 2017; 292.
- [18]. Castetbon K, Vernay M, Malon A, Salanave B, Deschamps V, Roudier C, et al. Dietary intake, physical activity and nutritional status in adults: the French nutrition and health survey (ENNS, 2006–2007). *British Journal of Nutrition*. 2009; 102(5): 733-43.
- [19]. Francou A, Hebel P, Braesco V, Drewnowski A. Consumption Patterns of Fruit and Vegetable Juices and Dietary Nutrient Density among French Children and Adults. *Nutrients*. 2015; 7(8): 6073-87.
- [20]. Bánáti D, Hellman-Regen J, Mack I, Young HA, Benton D, Eggersdorfer M, et al. Defining a vitamin A5/X specific deficiency - vitamin A5/X as a critical dietary factor for mental health. *Int J Vitam Nutr Res*. 2024; 94(5-6): 443-75.
- [21]. Stea TH, Nordheim O, Bere E, Stornes P, Eikemo TA. Fruit and vegetable consumption in Europe according to gender, educational attainment and regional affiliation-A cross-sectional study in 21 European countries. *PLoS One*. 2020; 15(5): e0232521.

- [22]. Stea TH, Nordheim O, Bere E, Stornes P, Eikemo TA. Fruit and vegetable consumption in Europe according to gender, educational attainment and regional affiliation—A cross-sectional study in 21 European countries. *Plos one*. 2020; 15(5): e0232521.
- [23]. da Costa GG, da Conceição Nepomuceno G, da Silva Pereira A, Simões BFT. Worldwide dietary patterns and their association with socioeconomic data: an ecological exploratory study. *Global Health*. 2022; 18(1): 31.
- [24]. OECD/European Union. Health at a Glance: Europe 2022: State of Health in the EU Cycle. Paris: OECD Publishing; 2022.
- [25]. Głąbska D, Guzek D, Groele B, Gutkowska K. Fruit and Vegetable Intake and Mental Health in Adults: A Systematic Review. *Nutrients*. 2020; 12(1).
- [26]. Morris MC, Wang Y, Barnes LL, Bennett DA, Dawson-Hughes B, Booth SL. Nutrients and bioactives in green leafy vegetables and cognitive decline: Prospective study. *Neurology*. 2018; 90(3): e214-e22.
- [27]. Ben-Amotz A, Fishier R. Analysis of carotenoids with emphasis on 9-cis β -carotene in vegetables and fruits commonly consumed in Israel. *Food Chem*. 1998; 62(4): 515-20.
- [28]. Khoo H-E, Prasad KN, Kong K-W, Jiang Y, Ismail A. Carotenoids and Their Isomers: Color Pigments in Fruits and Vegetables. *Molecules* [Internet]. 2011; 16(2):[1710-38 pp.].
- [29]. Vahid F, Brito A, Le Coroller G, Vaillant M, Samouda H, Bohn T, Group oboOW. Dietary Intake of Adult Residents in Luxembourg Taking Part in Two Cross-Sectional Studies—ORISCAV-LUX (2007–2008) and ORISCAV-LUX 2 (2016–2017). *Nutrients*. 2021; 13(12): 4382.