

## **Some markers of placental inflammation in pregnant women with gestational diabetes**

Thiago Paes de Barros De Luccia

Department of Obstetrics, Paulista School of Medicine, Federal University of São Paulo (EPM-UNIFESP)

### **Abstract**

**Background** analysis of fetal/placental components of women with gestational diabetes presented a slightly inflammatory profile, compared with non-diabetic pregnant women in previous studies by our group. Leptin, resistin and IL-6 are involved in the inflammatory process while adiponectin can act in anti-inflammatory processes.

**Purpose** Since both obesity and gestational diabetes are associated with inflammatory conditions, through these mediators we seek to evaluate systemic patterns in pregnant women and fetal patterns of this possible inflammation.

**Materials and Methods** Three adipokines (leptin, resistin and adiponectin) and one cytokine (IL-6) were studied in three different compartments (maternal serum, fetal serum and amniotic membrane culture supernatant). Four pregnant groups were analyzed (eutrophic, overweight, obese and gestational diabetics (GDM)). Maternal and fetal serum and amnion membrane biopsies were collected from 20 GDM and 28 normoglycemic subjects (Controls) who underwent elective cesarean sections.

**Results** We found a higher production of IL-6 in the culture supernatant of the amniotic membrane of obese pregnant women and more significantly in pregnant women with gestational diabetes. We did not observe correlation between the levels of mediators detected in the three compartments (mother serum, fetal serum and culture supernatant of amniotic membrane).

**Conclusion** Mainly in the amniotic membrane of pregnant women with GDM, a slight increase in inflammatory markers was observed.

## **Introduction:**

In previous studies carried out by our group, analysis of fetal/placental components of women with gestational diabetes presented a slightly inflammatory profile, compared with non-diabetic pregnant women<sup>1,2</sup>. In this article we will present supplementary data obtained after the aforementioned publications. In this new data, we added a more detailed analysis of the behavior of markers (leptin, adiponectin, resistin and IL-6) according to the pregnant women's BMI. Since both obesity and gestational diabetes are associated with inflammatory conditions, through these mediators we seek to evaluate systemic patterns in pregnant women and fetal patterns of this possible inflammation.

## **Methods:**

Three adipokines (leptin, resistin and adiponectin) and one cytokine (IL-6) were studied in three different compartments (maternal serum, fetal serum and amniotic membrane culture supernatant). Four pregnant groups were analyzed (eutrophic, overweight, obese and gestational diabetics). Maternal and fetal serum and amnion membrane biopsies were collected from 20 GDM and 28 normoglycemic subjects (control) who underwent elective cesarean sections during the years 2017 to 2020. The amnion membrane was manually peeled from the fetal membranes and placed in a 12-well culture for 24 hours at 37°C in a 5% CO<sub>2</sub> atmosphere and 95% air humidity (details about amniotic membrane culture in Menon et al., 2011<sup>3</sup>). The cytokine and adipokines were evaluated in serum and amnion culture supernatant samples (Luminex and ELISA). The detailed methodology of the work can be consulted in a previously published article<sup>1</sup>. The participants were recruited at the Obstetric Center of Hospital São Paulo-Brazil and of the Hospital do Servidor Público Municipal do Estado de São Paulo-Brazil (Ethics and Research Committee of UNIFESP CAAE 80728417.8.0000.5505).

## **Results:**

Here we will only present the most consistent results, as in some compartments (MS, FS and CS) measuring all markers was not possible. IL-6 measurements, for example, were better in the fetal compartment. We were unable to achieve good results

in detecting this cytokine in maternal and fetal serum. Figure 1 shows these results of this cytokine in amniotic membrane culture. In the graph (**Figure 1**) we see a greater production of IL-6 in the membrane of obese pregnant women and those with gestational diabetes.

In table 1 we have the most consistent results in all groups of pregnant women. In **Figure 2-A** and **Figure 2-B** we present the leptin/adiponectin ratio in pregnant controls and pregnant women with gestational diabetes.

## Discussion:

Ayina et al. showed that leptin to adiponectin ratio was significantly and positively correlated with the body mass index ( $r = 0.669$ ,  $p < 0.0001$ )<sup>4</sup>. Higher levels of this ratio are associated with greater cardiovascular risks<sup>5-7</sup>. In our study, in the GDM group, the positive correlation did not happen, maybe because the variation of BMI was smaller in this group. In the GDM group, the BMI was more similar to the overweight BMI group.

In relation to adipokines, there was a certain similarity in dosages between overweight pregnant women and gestational diabetics, suggesting that BMI variation is an important factor controlling adipokines. If we look at table 1, we see that there was no significant difference in dosages between the GDM group and overweight controls. More studies with a larger number of participants need to be carried out to understand the role of gestational diabetes in changing these markers (with BMI matching).

In studies on intra-amniotic inflammation, IL-6 was shown to be a good marker of inflammation<sup>8</sup>. The increase in this marker in amniotic fluid and cervical fluid has also been associated with preterm birth<sup>9</sup>. In this scenario, the amniotic membrane may play a role in the production of this cytokine. Our data showed greater amounts of IL-6 produced by the amniotic membrane in obese women and more significantly in gestational diabetics. However, there is no clear association between obesity/gestational diabetes and premature birth, but rather with advanced maternal age and premature birth<sup>10</sup>.

In our previous studies, we did not observe correlation between the levels of mediators detected in the three compartments (mother serum, fetal serum and culture supernatant of amniotic membrane). Mainly in the amniotic membrane of pregnant women with GDM, a slight increase in inflammatory markers was observed.

## References:

- 1- De Luccia TPB, Ono E, Menon R, Borbely AU, Mattar R, Richardson L, da Silva ALM, Botelho RM, da Rocha MLTLF, Daher S. The effect of Gestational Diabetes Mellitus on the fetal compartment. *J Reprod Immunol* 2021, 145:103314. doi: 10.1016/j.jri.2021.103314.
- 2- De Luccia TPB. Functional characterization of the amniotic membrane of pregnant women with gestational diabetes mellitus. 2021. Thesis presented to the Federal University of São Paulo – Escola Paulista de Medicina. Doctoral advisor: Silvia Daher; Co-advisor: Rosiane Mattar.
- 3- Menon, R., Fortunato, S.J., Yu, J., Milne, G.L., Sanchez, S., Drobek, C.O., Lappas, M., Taylor, R.N. Cigarette smoke induces oxidative stress and apoptosis in normal term fetal membranes. *Placenta* 2011, 32:317–322.
- 4- Ayina CNA, Endomba FTA, Mandengue SH, Noubiap JJN, Ngoa LSE, Boudou P, et al. Association of the leptin-to-adiponectin ratio with metabolic syndrome in a sub-Saharan African population. *Diabetol Metab Syndr* 2017, 5;9(1):66.
- 5- Frühbeck G, Catalán V, Rodríguez A, Gómez-Ambrosi J. Adiponectin-leptin ratio: A promising index to estimate adipose tissue dysfunction. relation with obesity-associated cardiometabolic risk. *Adipocyte* 2018, 7:57–62. doi: 10.1080/21623945.2017.1402151
- 6- Larsen MA, Isaksen VT, Moen OS, Wilsgaard L, Remijn M, Paulssen EJ, Florholmen J, Goll R. Leptin to adiponectin ratio - A surrogate biomarker for early detection of metabolic disturbances in obesity. *Nutr Metab Cardiovasc Dis* 2018, 28(11):1114-1121.
- 7- Senkus, K.E., Crowe-White, K.M., Bolland, A.C. et al. Changes in adiponectin:leptin ratio among older adults with obesity following a 12-month exercise and diet intervention. *Nutr Diabetes* 2022, 12:30 .
- 8- Leañós-Miranda A, Nolasco-Leañós AG, Carrillo-Juárez RIz, et al. Interleukin-6 in Amniotic Fluid: A Reliable Marker for Adverse Outcomes in Women in Preterm Labor and Intact Membranes. *Fetal Diagn Ther* 2021, 48 (4): 313–320.

9- Chang Y, Li W, Shen Y, Li S, Chen X. Association between interleukin-6 and preterm birth: a meta-analysis. *Annals of Medicine* 2023, 55.

10- Preda A, Iliescu DG, Comănescu A, et al. Gestational Diabetes and Preterm Birth: What Do We Know? Our Experience and Mini-Review of the Literature. *J Clin Med* 2023, 12(14):4572.

Figure captions:

**Figure 1-** IL-6 (culture supernatant - CS). Eut=eutrophic, Over=overweight, Ob=obese and GDM=gestational diabetics.

**Figure 2 - A** - correlation between leptin to adiponectin ratio and BMI in Controls.

**Figure 2 - B** - correlation between leptin to adiponectin ratio and BMI in GDM.

Table captions

**Table 1-** Adipokines and cytokines: median (interquartile range) - Mann Whitney test. BMI: mean (standard deviation) – t test. \* significant difference. GDM: gestational diabetes *mellitus*. MS: mother serum; FS: fetal serum; CS: culture supernatant of amniotic membrane.

Variable	Eutrophic (14)	Overweight (8)	Obese (6)	GDM (20)	Eut x Over	Eut x Ob	Eut x GDM	Over x Ob	Over x GDM	Ob x GDM
mean BMI (kg/m <sup>2</sup> )	22,3 (2,2)	26,3 (0,9)	32,8 (3,6)	28,4 (5,1)	<b>&lt;0.0001*</b>	<b>&lt;0.0001*</b>	<b>&lt;0.0001*</b>	<b>&lt;0.0001*</b>	0,0780	<b>0,0091*</b>
Adiponectin MS (µg/mL)	3,7 (1,9-4,2)	2,2 (1,2-3,3)	1,1 (1,0-1,5)	2,8 (1,2-4,2)	0,1051	<b>0,0023*</b>	0,6430	<b>0,0281*</b>	0,3493	<b>0,0061*</b>
Adiponectin FS (µg/mL)	3,4 (1,1-6,4)	5,0 (1,7-6,5)	1,1 (0,9-2,1)	8,1 (3,5-14,5)	0,5832	0,0564	<b>0,0090*</b>	<b>0,0152*</b>	<b>0,0491*</b>	<b>0,0007*</b>
Leptin MS (ng/mL)	0,2 (0,1-0,6)	40,4 (11,1-63,1)	89,1 (65,3-182,5)	39,7 (19,5-56,2)	<b>0,0003*</b>	<b>0,0003*</b>	<b>&lt;0.0001*</b>	<b>0,0041*</b>	1,0000	<b>0,0010*</b>
Leptin FS (ng/mL)	6,1 (4,0-17,0)	12,5 (5,5-18,5)	16,6 (13,3-29,8)	17,5 (7,8-30,4)	0,3040	<b>0,0324*</b>	<b>0,0261*</b>	0,1605	0,3423	0,8365
Leptin CS (ng/mL)	0,25 (0,1-0,6)	0,44 (0,08-1,4)	0,46 (0,2-0,7)	0,41 (0,1-0,7)	0,9109	0,4333	0,6132	0,7308	0,9702	0,5919
Resistin MS (ng/mL)	9,1 (6,3-12,3)	8,6 (8,0-12,1)	16,6 (14,1-17,7)	6,6 (4,3-10,5)	0,8255	<b>0,0044*</b>	0,2048	<b>0,0256*</b>	0,0579	<b>0,0036*</b>
Resistin FS (ng/mL)	11,6 (9,3-18,4)	19,6 (11,7-27,0)	17,2 (14,8-21,9)	13,4 (12,0-15,3)	0,1148	0,0872	0,9078	1,0000	0,2573	<b>0,0307*</b>
Resistin CS (ng/mL)	1,3 (0,7-2,1)	0,5 (0,2-1,8)	0,8 (0,7-1,0)	1,6 (0,9-2,1)	0,0998	0,2055	0,8178	0,5237	0,1803	0,0611
IL-6 CS (pg/mL)	612,8 (153-1364)	881,0 (136-2381)	3424,0 (999-22071)	2016,0 (1416-2515)	0,2423	<b>0,0426*</b>	<b>0,0053*</b>	0,2343	0,3595	0,6128

**Table 1**









