

Title: Longitudinal monitoring of respiratory syncytial virus, SARS-CoV-2 and influenza in wastewater of a Southern Indian city

Apoorva Venkatesh^{1,2}, Sreelekshmi RS^{1,3}, Parishuddha Babu Movva^{1,2}, Manas K Madhukar^{1,3}, Aruna Panda⁴, S. Venkata Mohan⁵, Rakesh Mishra,^{1,3,4,*} and Shivranjani C Moharir^{1,3,4,*}

Affiliations:

¹ Tata Institute for Genetics and Society (TIGS), Bengaluru- 560065, India

² National Centre for Biological Sciences (NCBS), Bengaluru- 560065, India

³ Academy of Scientific and Innovative Research (AcSIR), Ghaziabad- 201002, India

⁴ Centre for Cellular and Molecular Biology (CCMB), Hyderabad- 500007, India

⁵ Indian Institute of Chemical Technology (IICT), Hyderabad- 500007, India

Details for correspondence

Dr Shivranjani C Moharir

Email ID: shivranjani.c@tigs.res.in

Contact No. +91-8790295507, +91-080-61948156

Address: Tata Institute for Genetics and Society, Gandhi Krishi Vignana Kendra (GKVK), Rajiv Gandhi Nagar, Kodigehalli, Bengaluru, Karnataka 560065

Dr Rakesh K Mishra

Email ID: rakesh.mishra@tigs.res.in

Address: Tata Institute for Genetics and Society, Gandhi Krishi Vignana Kendra (GKVK), Rajiv Gandhi Nagar, Kodigehalli, Bengaluru, Karnataka 560065

Funding sources: This study was supported by the funding from the Rockefeller Foundation, Grant Number 2021 HTH 018 provided to the Tata Institute for Genetics and Society (TIGS), Centre for Cellular and Molecular Biology (CCMB) and the National Centre for Biological Sciences (NCBS). The study is also supported by the institutional core funding to Tata Institute for Genetics and Society by the Tata Trusts.

Abstract: The detection of diverse pathogens and chemical compounds in community wastewater facilitates the monitoring of public health trends of emerging diseases and health conditions. During the COVID-19 pandemic, detection of viral RNA in wastewater assisted in monitoring the infection rates in different geographies and this drew attention to the method of wastewater-based epidemiology (WBE). In contrast to individual clinical testing, WBE offers an affordable, population-wide overview of the infection status of a community including asymptomatic carriers and those without access to healthcare facilities. To understand the community status of the infections caused by respiratory syncytial virus, SARS-CoV-2 and influenza virus subtypes in the monsoon season in Vijayawada city in the state of Andhra Pradesh in India, we longitudinally analyzed wastewater samples once a week between July-August 2024 from 7 sewage treatment plants in the city. The data obtained from the multiplexed RT-qPCR was used to calculate the viral loads in the wastewater samples at the city level and the proportion of population shedding the virus was estimated. All three viruses RSV A+B, SARS-CoV-2 and Influenza A+B were detected in the wastewater during our sampling period. Amongst the three viruses, the city had the largest load of SARS-CoV-2 RNA in the wastewater followed by RSV A+B and Influenza A+B. The study demonstrates the potential of utilizing wastewater surveillance system coupled to multiplexed RT-PCR to understand the population level dynamics of co-existence of multiple pathogens during the monsoon season in the urban settings.

Key words: respiratory syncytial virus, influenza, wastewater surveillance, SARS-CoV-2, multiplex RT-PCR

Abbreviations: COVID-19- Corona Virus Disease 2019, SARS-CoV-2- sever acute respiratory syndrome coronavirus -2, WBE- wastewater based epidemiology, RSV- respiratory syncytial virus, STPs- sewage treatment plants, RT-PCR - real time polymerase chain reaction, MLD- Million Liters per Day

1 Introduction:

Wastewater monitoring is a method traditionally employed to track the spread of pathogens, including viruses such as poliovirus and norovirus, in a community (Nelson et al., 1967; Sinclair et al., 2008). It has recently emerged as a valuable tool for community-wide surveillance of Corona Virus Disease 2019 (COVID-19) and antimicrobial resistance (Ahmed et al., 2020; Madhukar et al., 2024; Munk et al., 2022; Tharak et al., 2022). The COVID-19 pandemic has exposed numerous unforeseen realities. Within months of severe acute respiratory syndrome coronavirus- 2 (SARS-CoV-2) infections spreading globally, most governments found themselves grappling with the fundamental task of determining the extent and geographical distribution of the infected populations. The presence of SARS-CoV-2 in the fecal matter has led several research groups worldwide to advocate for wastewater based epidemiology (WBE) as a means to evaluate the viral circulation among populations (Agrawal et al., 2021; Mallapaty, 2020). Studies have consistently revealed correlations between wastewater monitoring and clinical case data for various viruses including SARS-CoV-2, respiratory syncytial virus (RSV) and influenza, thus reinforcing the value of wastewater surveillance as an epidemiological instrument (Hughes et al., 2022; Wolfe et al., 2022). Numerous investigations have identified other respiratory and enteric viruses in wastewater samples, including enteroviruses A- D, rhinovirus, parainfluenza and metapneumovirus (Boehm et al., 2023; Brinkman et al., 2017). These detections have also aligned with clinical disease patterns, offering real-time information on the viral circulation.

The virtue of wastewater sample to serve as a composite sample for a large population, and to provide health status of the entire community gives it a unique advantage, particularly when individual testing may be restricted by logistical or resource limitations. The detection of viral RNA in wastewater can also function as an early warning sign of the infection surges in communities, often preceding clinical caseloads (Ahmed et al., 2021). The COVID-19 pandemic has highlighted the critical need for more accurate and versatile methods for estimating and understanding the spread of the virus at community level that can complement the public health surveillance systems. Since COVID-19 was a global health threat during the pandemic, the focus was on monitoring SARS-CoV-2, the causative virus. However, attention must now be shifted on including the surveillance of the traditional seasonal respiratory viruses that have reemerged after being temporarily suppressed by pandemic control measures and efforts should be directed towards understanding their patterns of emergence along with the cropping patterns of the other dormant viruses.

Influenza A and B viruses and RSV are particularly noteworthy, which account for the majority of respiratory infections along with SARS-CoV-2 (García-Arroyo et al., 2022). Influenza A and B viruses are primarily linked to seasonal influenza outbreaks, with different subtypes of influenza A causing worldwide influenza pandemics (Hutchinson, 2018). The persistence of the viral nucleic acids in wastewater and the associated patterns of viral shedding of influenza A and B suggest that wastewater surveillance can be a useful method for identifying and tracking seasonal flu outbreaks and pandemics (Minodier et al., 2015). Research indicates that people carrying these viruses can release around millions copies of the virus per gram of fecal matter (Arena et al., 2012). Given that, influenza viruses and RSV present symptoms similar to SARS-CoV-2, it is challenging to differentiate between these infections based on only the clinical presentations (Chung et al., 2021). Using molecular approaches, wastewater surveillance that involves the detection of viral RNA in sewage, can be utilized as a crucial tool for distinguishing between COVID-19 surges and seasonal outbreaks of other respiratory viruses at community level, allowing researchers and public health authorities to monitor multiple respiratory pathogens simultaneously. This could facilitate a judicious resource relocation to manage the infectious diseases effectively.

This study aims to evaluate, monitor, and compare the longitudinal RNA levels of the three respiratory viruses- influenza, respiratory syncytial virus, and SARS-CoV-2, simultaneously, in wastewater samples collected from seven sewage treatment plants (STPs) in the southern Indian city of Vijayawada, in the state of Andhra Pradesh. The study demonstrates the potential of using a multiplexed real time polymerase chain reaction (RT-PCR) based approach to simultaneously monitor the longitudinal patterns of these viruses and differentiate their burden at community level through wastewater-based epidemiology. It also demonstrates how wastewater surveillance of these viral pathogens can be used to get an estimate of the proportion of the population shedding the virus and can be utilized as a complementary approach to strengthen the public health surveillance system.

2 Methods:

2.1 Sample collection and processing:

The wastewater samples were collected once a week in between 20th July 2024 to 24th August 2024 from 7 sewage treatment plants (STPs) of Vijayawada city that has a population of around 1,467,000 as of September 2024 (Population, 2024). The city generates around 132 million liters per day (MLD) sewage and most of it is treated through the sewage treatment plants in the city (Corporation, 2024). The details of the sampling sites that receive majorly

human influx, are given in Supplementary Table 1. Around 1 liter wastewater sample was collected in sterile bottle during the peak flow from each STP by grab sampling method, as per the procedures described earlier (Kopperi et al., 2021; Madhukar et al., 2024; Tharak et al., 2022). The samples were inactivated using sodium hypochlorite (20 ml of 0.1% for 1 liter sample) (Hemalatha et al., 2021). Following thorough mixing, 50 ml of the wastewater sample was aliquoted, and 4 g of polyethylene glycol 8000 (Sigma -Aldrich, Catalogue No. P5413) and 0.9 g of sodium chloride (SRL, Catalogue No. 33205) were added to the sample and then incubated at 4°C overnight. The samples were centrifuged for 30 minutes at 10,000 rpm at 4°C. The pellet was either utilized directly for viral RNA extraction with the QIAamp Viral RNA Mini Kit (Qiagen, Catalogue No. 52906) or stored at – 80° C till further processing. Samples handling and processing was carried out in a biosafety cabinet following appropriate biosafety measures.

2.2 Detection of respiratory virus by RT-PCR:

The RNA extracted from the wastewater samples were used for multiplex RT-PCR for the detection of the respiratory viruses using the GenePath Dx RespiRFC + Influenza subtyping + PMMoV Detection and Quantitation RT-qPCR Reagents kit (Catalog number: GPDXRFC-3T-PM-01-100). Each sample was run in triplicate and, a 3 tubes 12 plex (4 plex X 3 tubes) reaction was set up for every sample. Tube 1 contained RSV A+B (FAM channel), SARS-CoV2 (HEX/ VIC channel), Influenza A+B (ROX channel), and an internal control PMMoV (Cy5 channel). Tube 2 contained targets for Influenza A (ROX channel), Influenza A H1N1 pdm09 (FAM channel), Influenza A H3N2 (HEX/ VIC channel), and an internal control PMMoV (Cy5 channel). And tube 3 contained Influenza A H7 (FAM channel), Influenza A H5 (HEX/ VIC channel), Influenza B (ROX channel), and the internal control PMMoV (Cy5 channel) as targets. RT-PCR was set with control templates of known concentrations of all the targets for plotting the standard curves for the respective targets. 5ul of 0.5 copies/ ul, 5 copies/ ul, 50 copies/ ul, 500 copies/ ul and 5000 copies/ ul of the control templates were used in triplicate for setting up RT-PCR reaction for obtaining the standard curves. The Ct values were plotted on Y axes against viral copies/ ul on the X axes. The equations obtained from the standard curves of each of the targets were used to calculate the respective viral RNA copies in the wastewater samples.

2.3 Statistical analysis: Based on the known concentrations of the controls and their respective Ct values, standard curves were plotted for the RNA concentrations for each of the viruses. An equation in the form of ' $y = mx + c$ ', where $y =$ Ct value, $m =$ gradient or slope, $x =$ viral copies and $c =$ y intercept, was obtained for each of the viruses. Pearson's correlation co-efficient (R^2) was calculated between the trend line and the line obtained by joining the datapoints. A correlation coefficient of ≥ 0.9 indicates a very strong positive correlation between the trend line and the line obtained by joining the datapoints.

2.4 Calculations of the estimated number of individuals shedding the virus:

The total viral RNA load in the STP was estimated by multiplying the viral RNA copies per liter of wastewater by the total flow (MLD: Million Liters per Day) of wastewater processed in the treatment plant on that particular day (Supplementary Table 2 and 3).

*Total viral RNA load in the STP = viral RNA load per liter of wastewater * the volume of the STP*

Once the total viral RNA load in the STP is known then the estimated number of individuals contributing to this viral load can be calculated if the viral copies shed by an individual in feces is known (Ahmed et al., 2020; Hemalatha et al., 2021).

Based on literature for SARS-CoV-2, a single infected individual sheds 10^7 copies of the viral RNA per ml of feces. Volume of feces excreted per day is 120 ml (Foladori et al., 2020; Hemalatha et al., 2021).

For influenza, literature reports that an infected individual sheds on an average 844700 viral RNA copies per gram of stool (Supplementary Table 4). This value is obtained by averaging the viral copies shed per gram of feces across different samples (Arena et al., 2012). On an average one person is known to excrete 128 grams of feces per day (Rose et al., 2015)

The number of infected individuals contributing to the viral load in the STP is estimated by dividing the total viral load in the STP by the viral load shed by one infected individual per day:

*Number of individuals shedding the virus = Total viral load in STP / (viral copies per gram or liter stool * gram or liter of stool excreted per day)*

3 Results:

3.1 Simultaneous detection of RNA of respiratory viruses RSV A + B, SARS-CoV-2 and influenza in wastewater samples:

Seven STPs in Vijayawada city were monitored once a week for the respiratory viruses- RSV A + B, SARS-CoV-2 and Influenza A + B between 20th July 2024 to 24th August 2024. Out of the 42 wastewater samples analyzed, 26 had Ct values ≤ 35 for RSV A + B, 41 had Ct ≤ 35 for SARS-CoV-2 and 32 had Ct ≤ 35 for Influenza A + B. PMMoV was used as an internal control and was detected in all the samples at Ct ≤ 35 . 24 samples had Ct ≤ 35 for all the three viruses- RSV A + B, SARS-CoV-2 and Influenza A + B together (Table 1).

3.2 Longitudinal monitoring of RSV A+B, SARS-CoV-2 and Influenza A+B viral RNA in wastewater:

RSV A+B, SARS-CoV-2 and Influenza A+B viral RNA were detected by RT-PCR in most of the samples analyzed during the monsoon period of 20th July 2024 to 24th August 2024. In order to calculate the viral RNA concentrations in the wastewater samples from the Ct values, we plotted the standard curves for each of the viruses using control templates of known concentrations (Figure 2 A- J). Using the Ct values obtained for each of the known concentrations, an equation was derived for the respective viruses. We then calculated the concentration of the viral RNA in the respective STPs on the dates of sample collection using the Ct values obtained from the RT-PCR of the RNA extracted from the wastewater samples and fitting them in the standard curve equations for the respective virus. To understand the longitudinal patterns of these viruses, the viral RNA concentrations were plotted against the date of sampling (Figure 3- 5). Amongst the three respiratory viruses- RSV A+B, SARS-CoV-2 and Influenza A+B, the RNA load was maximum for SARS-CoV-2 in general and on the peak dates as compared to the other two.

For RSV A+B, there was a clear rise in the viral RNA load in the city in the third sampling week, i.e., on 3rd August 2024 (Figure 3A). This increase in RSV A+B RNA load on 3rd August was also observed in the individual sampling locations ASN1, ASN2 and JKP, while for the location RLN1, the RNA load peaked on 10th August 2024 (Figure 3 B, C, F and H). The location ATN1 showed the rise in RSV A+B RNA load on 17th August 2024, which declined in the subsequent weeks (Figure 3D). The RSV A+B RNA was not detected in the location ATN2 on 20th July and 27th July 2024 and the location JKP on 20th July 2024 (Figure 3E and H). In the location RLN2, the RSV A+B RNA load was higher in the first week of sampling, i.e., on 20th July 2024 and declined in the subsequent weeks (Figure 3G).

SARS-CoV-2 RNA load peaked in the second week of sampling i.e., on 27th July 2024 during the sampling window in the city (Figure 4A). The peak of SARS-CoV-2 RNA load was also seen in the individual locations ASN1, ASN2, ATN1, RLN1 and JKP on 27th July 2024 (Figure 4 B, C, D, F and H). In the location ATN2, the peak was seen on 10th August 2024, while in RLN2 it was seen on 17th August 2024 (Figure 4E and G).

The RNA load of Influenza A+B virus was highest in the city on 20th July 2024 and had maximum contribution from the site JKP, ATN2 and RLN2 on that day (Figure 5A, E, G and H). The locations ASN2 and ATN1 showed the peak on 3rd August 2024 (Figure 5C and D). The location ASN1 showed two peaks- one on 3rd August and the other on 24th August 2024 during the sampling window, while the location RLN1 showed the peak on 10th August 2024 (Figure 5B and F).

3.3 Detection of subtypes of influenza through wastewater surveillance: Since the Influenza viral RNA was detected in all the locations in the city during the sampling period, we investigated the presence of the RNA of different subtypes of Influenza virus viz Influenza A virus and its subtypes - Influenza A H1N1, Influenza A H3N2, Influenza A H5 and Influenza A H7, as well as Influenza B virus. 12 out of 32 Influenza A + B samples with ≤ 35 had Ct value ≤ 35 for primer- probes specific to Influenza A as well, of which 4 samples showed Ct ≤ 35 for Influenza A H1N1. Influenza A H3N2, Influenza A H5 and Influenza A H7 were not picked up in any of the samples. 27 samples showed Ct value ≤ 35 for Influenza B (Table 1). The longitudinal trend plots for the RNA load of Influenza A, Influenza A H1N1 and Influenza B are shown in supplementary figures 1-3.

3.4 Estimation of number of infected individuals: The viral RNA copies excreted in feces by an individual infected with SARS-CoV-2 and Influenza virus is known in literature. Combining this information with the viral load that observed in the individual STPs in our study, we estimated the number of individuals shedding the viral RNA into the wastewater in the respective individual STPs and the whole city (Table 2 and 3). We observed that during our sampling window, on 20th July 2024, 0.21% population of the city was shedding the SARS-CoV-2 RNA in the wastewater (Figure 6A). The percentage of individuals shedding the SARS-CoV-2 RNA in wastewater increased to 0.94% in the next week, followed by a decline in subsequent weeks. A more granular analysis at individual location level suggested that the location ASN1 had the maximum SARS-CoV-2 RNA shedding in wastewater, by

0.344 % of the population, on 27th July 2024, followed by 0.251% at the location ASN2 and 0.194% at the location JKP on the same day (Figure 6C).

During our sampling window, we observed the maximum RNA shedding of Influenza virus in the wastewater in the city, corresponding to shedding by 0.2% of the population, on 20th July 2024 (Figure 6 B). The location JKP had the maximum percentage of individuals, 0.126%, contributing to the shedding of viral RNA on that day (Figure 6D). There was a decrease in the shedding of viral RNA in the wastewater in the subsequent weeks.

Except for the location JKP on 2th July 2024, at any point of time, the percentage of population shedding SARS-CoV-2 in wastewater was higher as compared to those shedding Influenza A + B at city level as well as individual location level. The concentration of the RSV RNA shed by an infected individual in the feces is not known. Hence, we could not estimate the number of individuals shedding the virus in the STPs wastewater.

Discussion:

Though the prominent symptoms shown by the individuals infected with the respiratory viruses RSV, SARS-CoV-2 and Influenza RNA are related to the respiratory systems, the RNA of these viruses are shed in the wastewater through the fecal route as well. Earlier studies have demonstrated that there is a strong correlation between the shedding of viral RNA by infected individuals in wastewater and the respective number of clinically confirmed individuals (Tisza et al., 2023; Toribio-Avedillo et al., 2023). In this study, we investigated the levels of these three viruses together in wastewater in Vijayawada. All the three viruses could be detected in the wastewater in the city. The load of SARS-CoV-2 RNA was the highest in the city throughout our sampling window, followed by RSV A+ B and the Influenza A + B.

The influenza virus consists of 4 distinct types of viruses- influenza A, B, C and D. While influenza A primarily affects the humans as well as other hosts including aquatic birds, pigs, horses and poultry, influenza B and C mainly infects humans, and influenza D is known to infect cattle, pigs and goats (Francis, 1940; Toribio-Avedillo et al., 2023). It is well understood that influenza A and B viruses co-occur in humans across different age groups and exhibit varying degrees of severity (He et al., 2022). Annually, influenza A and B viruses contribute to between 291,243 and 645,832 respiratory-related deaths, with 3 to 5 million cases of severe illness occurring worldwide (Iuliano et al., 2018).

In India, a nation of 1.428 billion people, the surveillance capacity for influenza A H1N1 virus is limited, with only few laboratories equipped to conduct clinical testing according to National Centre for Disease Control (NCDC, 2019). In such scenario, wastewater and sewage surveillance can offer considerable potential for the detection of influenza viruses and their subtypes. A recent pan-India study that investigated 34,260 cases of influenza-like illness (ILI) and severe acute respiratory infection found that during the 2021 monsoon season in India, influenza A(H3) and B/Victoria strains predominated. However, in the 2022 monsoon season, influenza A (H1N1) pdm09 became the dominant strain (Potdar et al., 2023). Concordantly, in our study, we could detect the H1N1 subtype in wastewater in the monsoon season of July and August 2024. The avian influenza A subtypes H3N2, H5, and H7 were not detected in any of the samples during the sampling window. This could be possible due to four reasons- i) in this study, we sampled the sewage treatment plants that receive influx from human catchments and not from poultry, so these subtypes if present, could not find a way in the sampled wastewater, ii) due to absence of these infectious during our sampling window, iii) inefficient or improbable detection of these subtypes through wastewater surveillance or iv) probably these subtypes are not shed in the feces. H5N1 caused significant outbreaks of avian flu in Asian countries in 2004, H7N3 in Pakistan, and H5N2 in Taiwan, leading to the culling of hundreds of millions of poultry (Sturm-Ramirez et al., 2004). Further investigations are necessary towards detection of these subtypes in wastewater surveillance and to optimize the protocols.

Interestingly, in our wastewater samples, influenza B virus was detected in 27 of the 42 samples (Ct value ≤ 35), representing a relatively higher proportion compared to influenza A. Unlike influenza A, which can infect both humans and animals, influenza B is a strictly human pathogen, having evolved stably to infect humans. This likely explains its higher detection rate in sewage samples (Chen and Holmes, 2008).

The presence of SARS-CoV-2 in wastewater has been documented in numerous studies and has garnered significant attention within the WBE community (Zhang et al., 2022). While much attention has been focused on WBE for tracking SARS-CoV-2, this approach has also demonstrated its effectiveness in detecting and monitoring a variety of enteric viruses like influenza, RSV, poliovirus, norovirus, rotaviruses, human astroviruses, and hepatitis A and E viruses. To date, there has been minimal research on the concurrent detection of RSV,

Influenza, and SARS-CoV-2 in wastewater in India. Our study represents the comprehensive investigation in India to simultaneously monitor the presence of SARS-CoV-2, RSV, and influenza viruses in wastewater and provides valuable insights into the co-circulation of these pathogens within the community.

Figures:




Figure 1: Co-ordinates of the sampling sewage treatment plants plotted on the map of Vijayawada city.




Figure 2: Standard curves of viruses, - A) RSV A+B, B) SARS CoV-2, C) INF A+B, D) INF A, E) INF A H1N1, F) INF A H3N2, G) INF A H5, H) INF A H7, I) INF B, and J) PMMoV.




Figure 3: Longitudinal plots of RNA load of RSV in wastewater samples A) At the city level, and the 7 sampling locations, - B) at the location ASN1 -, C) at the location ASN2, D) at the location ATN1, E) at the location ATN2, F) at the location RLN1, G) at the location RLN2 and H) at the location JKP.




Figure 4: Longitudinal plots of RNA load of SARS-CoV-2 in wastewater samples A) At the city level, and the 7 sampling locations, B) at the location ASN1 -, C) at the location ASN2, D) at the location ATN1, E) at the location ATN2, F) at the location RLN1, G) at the location RLN2 and H) at the location JKP.




Figure 5: Longitudinal plots of RNA load of Influenza in wastewater samples A) At the city level, and the 7 sampling locations, B) at the location ASN1 -, C) at the location ASN2, D) at the location ATN1, E) at the location ATN2, F) at the location RLN1, G) at the location RLN2 and H) at the location JKP.




Figure 6: Estimated percentage of virus shedding population A) Estimated percentage of SARS-CoV-2 shedding population at the city level, B) Estimated percentage of Influenza A+B shedding population at the city level, C) Estimated percentage of SARS-CoV-2 shedding population at the 7 individual STPs and A) Estimated percentage of Influenza A+B shedding population at the 7 individual STPs.

Table 1: Table showing the Ct values obtained by RT-PCR of wastewater samples for RSV A+B, SARS-CoV-2, influenza A+B, influenza A, influenza A H1N1, influenza A H3N2, influenza A H5, influenza A H7, influenza B and PMMoV. The Ct values in red font are ≤ 35 . UD= Undetermined

Collection date	Sample ID	RSV A+B	SARS-CoV2	INF A+B	INF A	INF A H1N1	INF A H3N2	INF A H5	INF A H7	INF B	PMMoV
20/07/24	ASN1	36.70	32.21	35.23	38.57	38.18	UD	UD	UD	UD	26.15
	ASN2	37.02	32.69	36.70	37.20	37.74	UD	UD	UD	UD	26.92
	ATN1	35.21	31.51	33.86	36.70	37.16	UD	UD	UD	UD	25.80
	ATN2	UD	31.48	32.35	34.35	38.14	UD	UD	UD	33.77	25.16
	RLN1	35.10	31.55	34.03	36.58	38.05	UD	UD	UD	34.91	25.60
	RLN2	32.76	30.71	33.20	35.66	36.14	UD	UD	UD	34.97	25.12
	JKP	UD	30.37	30.96	37.04	37.53	UD	UD	UD	37.02	25.31
27/07/24	ASN1	34.60	28.44	34.19	37.07	36.91	UD	UD	UD	36.43	24.93
	ASN2	35.63	27.77	33.49	35.55	37.14	UD	UD	UD	36.21	24.24
	ATN1	36.30	29.29	36.31	38.91	37.76	UD	UD	UD	35.41	27.74
	ATN2	UD	32.16	35.84	38.39	38.18	UD	UD	UD	UD	28.00
	RLN1	36.25	29.62	34.28	36.04	38.21	UD	UD	UD	36.10	26.87
	RLN2	36.32	30.90	34.46	37.42	37.73	UD	UD	UD	37.28	27.62
	JKP	37.64	28.23	36.66	37.19	37.71	UD	UD	UD	36.14	25.52
03/08/24	ASN1	32.45	30.47	33.42	34.99	35.66	UD	UD	UD	34.08	25.13
	ASN2	32.41	29.78	32.71	34.01	36.36	UD	UD	UD	32.95	25.24
	ATN1	33.77	30.29	33.28	33.91	34.35	UD	UD	UD	33.57	24.83
	ATN2	34.15	31.12	35.37	35.76	UD	UD	UD	UD	34.65	26.22
	RLN1	34.74	31.39	34.28	35.62	37.51	UD	UD	UD	34.24	27.00
	RLN2	34.70	30.27	33.58	34.99	37.98	UD	UD	UD	34.61	25.75
	JKP	32.54	30.21	33.10	36.32	37.69	UD	UD	UD	33.16	24.73
10/08/24	ASN1	32.97	32.23	34.68	36.38	37.03	UD	UD	UD	35.36	25.44
	ASN2	35.41	33.17	35.62	36.44	38.53	UD	UD	UD	36.09	27.34
	ATN1	36.09	31.61	34.38	36.10	37.49	UD	UD	UD	34.88	24.42
	ATN2	33.50	29.64	33.86	35.15	37.58	UD	UD	UD	35.77	24.63
	RLN1	34.23	31.00	33.78	35.27	38.65	UD	UD	UD	34.60	25.03
	RLN2	34.92	32.98	35.13	37.43	38.78	UD	UD	UD	34.77	27.26
	JKP	UD	36.59	UD	40.00	UD	UD	UD	UD	UD	32.82
17/08/24	ASN1	34.65	30.57	33.76	33.92	34.60	UD	UD	UD	34.16	24.49
	ASN2	33.47	30.17	33.36	33.13	34.60	UD	UD	UD	33.63	24.33
	ATN1	32.69	30.64	34.30	35.19	36.83	UD	UD	UD	34.02	24.58
	ATN2	34.42	30.38	34.88	33.77	36.43	UD	UD	UD	34.07	23.54
	RLN1	35.14	30.55	35.51	35.31	35.16	UD	UD	UD	34.95	23.96
	RLN2	34.70	29.92	33.96	34.40	35.84	UD	UD	UD	33.29	24.82
	JKP	34.71	30.32	34.62	35.74	36.30	UD	UD	UD	33.61	24.61
24/08/24	ASN1	34.27	31.96	33.32	34.15	37.05	UD	UD	UD	31.40	24.41

	ASN2	34.88	32.80	34.62	35.86	36.90	UD	UD	UD	33.28	26.21
	ATN1	34.78	32.52	34.76	35.47	36.13	UD	UD	UD	34.29	25.89
	ATN2	33.39	31.29	34.13	34.64	36.62	UD	UD	UD	33.43	24.83
	RLN1	34.34	31.43	34.29	35.53	37.04	UD	UD	UD	34.28	25.73
	RLN2	34.91	32.59	34.21	35.88	35.74	UD	UD	UD	33.55	26.39
	JKP	34.51	31.01	32.84	33.06	34.13	UD	UD	UD	33.19	24.53

Table 2: Table showing the SARS-CoV-2 RNA load in wastewater samples, the estimated number of infected individuals and the estimated percentage of infected individuals in the city

Collection date	STP	SARS-CoV-2 RNA in wastewater sample (copies/l)	Estimated number of infected individuals	Estimated percentage of infected individuals in the city
20/07/24	ASN1	18098.16	603	0.041
	ASN2	13753.78	229	0.016
	ATN1	26855.92	224	0.015
	ATN2	27242.88	227	0.015
	RLN1	26234.19	219	0.015
	RLN2	41990.41	700	0.048
	JKP	51060.08	851	0.058
	Total	205235.42	3053	0.208
27/07/24	ASN1	151472.89	5049	0.344
	ASN2	221066.78	3684	0.251
	ATN1	93899.71	782	0.053
	ATN2	18574.85	155	0.011
	RLN1	77723.00	648	0.044
	RLN2	37824.57	630	0.043
	JKP	170908.98	2848	0.194
	Total	771470.78	13797	0.941
03/08/24	ASN1	48269.27	1609	0.110
	ASN2	71260.30	1188	0.081
	ATN1	53458.68	445	0.030
	ATN2	33431.84	279	0.019
	RLN1	28652.69	239	0.016
	RLN2	53942.23	899	0.061
	JKP	55647.88	927	0.063
	Total	344662.90	5586	0.381
10/08/24	ASN1	17824.67	594	0.041
	ASN2	10481.96	175	0.012
	ATN1	25400.46	212	0.014
	ATN2	76792.06	640	0.044
	RLN1	35788.12	298	0.020
	RLN2	11690.55	195	0.013
	JKP	1523.79	25	0.002
	Total	179501.61	2139	0.146
17/08/24	ASN1	45651.60	1522	0.104
	ASN2	57042.80	951	0.065


	ATN1	43884.52	366	0.025
	ATN2	50829.28	424	0.029
	RLN1	45962.46	383	0.026
	RLN2	65526.86	1092	0.074
	JKP	52457.21	874	0.060
	Total	361354.71	5611	0.382
24/08/24	ASN1	20783.28	693	0.047
	ASN2	12968.10	216	0.015
	ATN1	15202.12	127	0.009
	ATN2	30315.02	253	0.017
	RLN1	28042.03	234	0.016
	RLN2	14568.96	243	0.017
	JKP	35551.75	593	0.040
	Total	157431.26	2357	0.161

Table 3: Table showing the Influenza RNA load in wastewater samples, the estimated number of infected individuals and the estimated percentage of infected individuals in the city

Sample collection date	STP	Influenza RNA in wastewater sample (copies/l)	Estimated number of infected individuals	Estimated percentage of infected individuals in the city
20/07/24	ASN1	435.94	161	0.011
	ASN2	148.32	27	0.002
	ATN1	1196.58	110	0.008
	ATN2	3603.65	332	0.023
	RLN1	1055.97	97	0.007
	RLN2	1944.42	358	0.024
	JKP	10017.57	1846	0.126
	Total	18402.45	2933	0.200
27/07/24	ASN1	937.01	345	0.024
	ASN2	1563.21	288	0.020
	ATN1	198.09	18	0.001
	ATN2	279.28	26	0.002
	RLN1	880.60	81	0.006
	RLN2	770.90	142	0.010
	JKP	153.60	28	0.002
	Total	4782.69	929	0.063
03/08/24	ASN1	1650.42	608	0.041
	ASN2	2778.36	512	0.035
	ATN1	1830.73	169	0.012
	ATN2	395.30	36	0.002
	RLN1	878.19	81	0.006
	RLN2	1463.59	270	0.018
	JKP	2083.85	384	0.026
	Total	11080.44	2060	0.140
10/08/24	ASN1	653.33	241	0.016
	ASN2	328.74	61	0.004
	ATN1	815.76	75	0.005
	ATN2	1194.42	110	0.008
	RLN1	1270.80	117	0.008
	RLN2	469.92	87	0.006
	JKP		0	0.000
	Total	4732.97	690	0.047
17/08/24	ASN1	1282.85	473	0.032
	ASN2	1726.08	318	0.022
	ATN1	862.89	80	0.005

	ATN2	564.21	52	0.004
	RLN1	356.91	33	0.002
	RLN2	1110.39	205	0.014
	JKP	686.01	126	0.009
	Total	6589.35	1287	0.088
24/08/24	ASN1	1775.40	654	0.045
	ASN2	686.34	127	0.009
	ATN1	618.34	57	0.004
	ATN2	981.26	90	0.006
	RLN1	872.69	80	0.005
	RLN2	926.46	171	0.012
	JKP	2523.79	465	0.032
	Total	8384.27	1645	0.112

Graphical abstract



Author contributions: SCM, RKM, VM and AP conceptualized and planned the study. RKM and SCM received the funding for the study. AV, SRS, PBM, MKM performed the experiments. SCM, AV, SRS, MKM performed the data analysis and wrote the first draft of the manuscript. All the authors contributed to the refinement of the manuscript draft.

Acknowledgements: The authors acknowledge the generous intellectual support of Dr. L S Shashidhara, Director, NCBS, Bengaluru and appreciate his inputs on the study.

References:

- Agrawal S, Orschler L, Lackner S. Long-term monitoring of SARS-CoV-2 RNA in wastewater of the Frankfurt metropolitan area in Southern Germany. *Sci Rep* 2021; 11: 5372.
- Ahmed W, Angel N, Edson J, Bibby K, Bivins A, O'Brien JW, et al. First confirmed detection of SARS-CoV-2 in untreated wastewater in Australia: A proof of concept for the wastewater surveillance of COVID-19 in the community. *Sci Total Environ* 2020; 728: 138764.
- Ahmed W, Tschärke B, Bertsch PM, Bibby K, Bivins A, Choi P, et al. SARS-CoV-2 RNA monitoring in wastewater as a potential early warning system for COVID-19 transmission in the community: A temporal case study. *Science of The Total Environment* 2021; 761: 144216.
- Arena C, Amoros JP, Vaillant V, Balay K, Chikhi-Brachet R, Varesi L, et al. Simultaneous investigation of influenza and enteric viruses in the stools of adult patients consulting in general practice for acute diarrhea. *Virology* 2012; 9: 116.
- Boehm AB, Hughes B, Duong D, Chan-Herur V, Buchman A, Wolfe MK, et al. Wastewater concentrations of human influenza, metapneumovirus, parainfluenza, respiratory syncytial virus, rhinovirus, and seasonal coronavirus nucleic-acids during the COVID-19 pandemic: a surveillance study. *Lancet Microbe* 2023; 4: e340-e348.
- Brinkman NE, Fout GS, Keely SP. Retrospective Surveillance of Wastewater To Examine Seasonal Dynamics of Enterovirus Infections. *mSphere* 2017; 2: 10.1128/msphere.00099-17.
- Chen R, Holmes EC. The evolutionary dynamics of human influenza B virus. *J Mol Evol* 2008; 66: 655-63.
- Chung HY, Jian MJ, Chang CK, Lin JC, Yeh KM, Chen CW, et al. Novel dual multiplex real-time RT-PCR assays for the rapid detection of SARS-CoV-2, influenza A/B, and respiratory syncytial virus using the BD MAX open system. *Emerg Microbes Infect* 2021; 10: 161-166.
- Corporation VM. Vijayawada's clean drive towards swachhata. 2024.
- Foladori P, Cutrupi F, Segata N, Manara S, Pinto F, Malpei F, et al. SARS-CoV-2 from faeces to wastewater treatment: What do we know? A review. *Sci Total Environ* 2020; 743: 140444.
- Francis T, Jr. A NEW TYPE OF VIRUS FROM EPIDEMIC INFLUENZA. *Science* 1940; 92: 405-8.
- García-Arroyo L, Prim N, Del Cuerpo M, Marín P, Roig MC, Esteban M, et al. Prevalence and seasonality of viral respiratory infections in a temperate climate region: A 24-year study (1997-2020). *Influenza Other Respir Viruses* 2022; 16: 756-766.
- He J, Hou S, Chen Y, Yu JL, Chen QQ, He L, et al. The Epidemiological Pattern and Co-infection of Influenza A and B by Surveillance Network From 2009 to 2014 in Anhui Province, China. *Front Public Health* 2022; 10: 825645.

- Hemalatha M, Kiran U, Kuncha SK, Kopperi H, Gokulan CG, Mohan SV, et al. Surveillance of SARS-CoV-2 spread using wastewater-based epidemiology: Comprehensive study. *Sci Total Environ* 2021; 768: 144704.
- Hughes B, Duong D, White BJ, Wigginton KR, Chan EMG, Wolfe MK, et al. Respiratory Syncytial Virus (RSV) RNA in Wastewater Settled Solids Reflects RSV Clinical Positivity Rates. *Environmental Science & Technology Letters* 2022; 9: 173-178.
- Hutchinson EC. *Influenza Virus*. *Trends Microbiol* 2018; 26: 809-810.
- Iuliano AD, Roguski KM, Chang HH, Muscatello DJ, Palekar R, Tempia S, et al. Estimates of global seasonal influenza-associated respiratory mortality: a modelling study. *The Lancet* 2018; 391: 1285-1300.
- Kopperi H, Tharak A, Hemalatha M, Kiran U, Gokulan CG, Mishra RK, et al. Defining the methodological approach for wastewater-based epidemiological studies- Surveillance of SARS-CoV-2. *Environ Technol Innov* 2021; 23: 101696.
- Madhukar MK, Singh N, Iyer VR, Sowpati DT, Tallapaka KB, Mishra RK, et al. Antimicrobial resistance landscape in a metropolitan city context using open drain wastewater-based metagenomic analysis. *Environ Res* 2024; 252: 118556.
- Mallapaty S. How sewage could reveal true scale of coronavirus outbreak. *Nature* 2020; 580: 176-177.
- Minodier L, Charrel RN, Ceccaldi PE, van der Werf S, Blanchon T, Hanslik T, et al. Prevalence of gastrointestinal symptoms in patients with influenza, clinical significance, and pathophysiology of human influenza viruses in faecal samples: what do we know? *Virology* 2015; 12: 215.
- Munk P, Brinch C, Møller FD, Petersen TN, Hendriksen RS, Seyfarth AM, et al. Genomic analysis of sewage from 101 countries reveals global landscape of antimicrobial resistance. *Nature Communications* 2022; 13: 7251.
- NCDC. List of labs testing the clinical samples for Influenza A (H1N1) in the States/UTs of India. <https://ncdc.mohfw.gov.in/wp-content/uploads/2024/02/90384949221554289446.pdf>. 2019.
- Nelson DB, Circo R, Evans AS. Strategic viral surveillance of sewage during and following an oral poliovirus vaccine campaign. *Am J Epidemiol* 1967; 86: 641-52.
- Population V. Vijayawada Population, 2024, <https://www.census2011.co.in/census/city/408-vijayawada.html>, 2024.
- Potdar V, Vijay N, Mukhopadhyay L, Aggarwal N, Bhardwaj SD, Choudhary ML, et al. Pan-India influenza-like illness (ILI) and Severe acute respiratory infection (SARI) surveillance: epidemiological, clinical and genomic analysis. *Front Public Health* 2023; 11: 1218292.
- Rose C, Parker A, Jefferson B, Cartmell E. The Characterization of Feces and Urine: A Review of the Literature to Inform Advanced Treatment Technology. *Crit Rev Environ Sci Technol* 2015; 45: 1827-1879.
- Sinclair RG, Choi CY, Riley MR, Gerba CP. Pathogen surveillance through monitoring of sewer systems. *Adv Appl Microbiol* 2008; 65: 249-69.
- Sturm-Ramirez KM, Ellis T, Bousfield B, Bissett L, Dyrting K, Rehg JE, et al. Reemerging H5N1 influenza viruses in Hong Kong in 2002 are highly pathogenic to ducks. *J Virol* 2004; 78: 4892-901.
- Tharak A, Kopperi H, Hemalatha M, Kiran U, C GG, Moharir S, et al. Longitudinal and Long-Term Wastewater Surveillance for COVID-19: Infection Dynamics and Zoning of Urban Community. *Int J Environ Res Public Health* 2022; 19.

- Tisza M, Javornik Cregeen S, Avadhanula V, Zhang P, Ayvaz T, Feliz K, et al. Wastewater sequencing reveals community and variant dynamics of the collective human virome. *Nature Communications* 2023; 14: 6878.
- Toribio-Avedillo D, Gómez-Gómez C, Sala-Comorera L, Rodríguez-Rubio L, Carcereny A, García-Pedemonte D, et al. Monitoring influenza and respiratory syncytial virus in wastewater. *Beyond COVID-19. Sci Total Environ* 2023; 892: 164495.
- Wolfe MK, Duong D, Bakker KM, Ammerman M, Mortenson L, Hughes B, et al. Wastewater-Based Detection of Two Influenza Outbreaks. *Environmental Science & Technology Letters* 2022; 9: 687-692.
- Zhang D, Duran SSF, Lim WYS, Tan CKI, Cheong WCD, Suwardi A, et al. SARS-CoV-2 in wastewater: From detection to evaluation. *Mater Today Adv* 2022; 13: 100211.