

1 **Knowledge and Willingness towards Human Papillomavirus**
2 **Vaccination among the Parents and School Teachers of**
3 **Eligible Girls in Dhaka, Bangladesh: A School-Based Cross-**
4 **Sectional Study**

5
6 Samina Sultana ^{¶1}, MD Nahid Hassan Nishan ^{¶1}, Aklima Akter ², Dalia Rahman ²,
7 Fowzia Yesmin ², Mohammad Delwer Hossain Hawlader ^{*1}

8
9 1 Department of Public Health, North South University, Dhaka-1229, Bangladesh.
10 2 Department of Obstetrics & Gynecology, Dhaka Medical College Hospital, Dhaka-1000,
11 Bangladesh
12

13 ***Corresponding Author**

14 Mohammad Delwer Hossain Hawlader

15 MBBS, MPH, PhD

16 Department of Public Health, School of Health and Life Sciences, North South University,
17 Dhaka 1229, Bangladesh

18 E-mail address: mohammad.hawlader@northsouth.edu

19 [¶]These authors contributed equally to this work.

20

21

22 **Abstracts**

23 **Background**

24 Cervical cancer ranks as the common prevalent cancer, among women worldwide especially
25 impacting low-resource countries. In Bangladesh, this accounts for 12% of all cancer cases. The
26 development of cancer is closely linked to Human Papillomavirus (HPV) infection. Despite the
27 availability of HPV vaccines, their uptake remains limited in Bangladesh. Thus, this research
28 aims to assess the knowledge and willingness of parents and school teachers regarding HPV
29 vaccination for eligible girls in Bangladesh.

30 **Methodology**

31 This study involved 406 parents and school teachers of girls aged 9-14 years from Dhaka city. A
32 cross-sectional study design was used. Data collection was done through a questionnaire
33 administered by interviewers after pre-testing and refinement for clarity and reliability. Analysis
34 was carried out using Stata 17 software. Chi-square tests and logistic regression were used to
35 uncover associations and predictors related to knowledge levels and willingness.

36 **Results**

37 Findings revealed that a majority of participants (64.04%) exhibited an understanding of HPV
38 and cervical cancer yet a high percentage (98.28%) expressed willingness to engage in HPV
39 vaccination initiatives. participants with primary (AOR=3.306, $p<0.005$), secondary
40 (AOR=8.806, $p<0.001$), and higher education (AOR=5.059, $p<0.001$), as well as those from
41 upper-middle-income groups (AOR=3.038, $p<0.001$), had significantly higher knowledge of
42 HPV and cervical cancer.

43 **Conclusion**

44 The research emphasizes lack of knowledge regarding HPV and its vaccination among parents
45 and educators in Bangladesh despite a willingness to vaccinate. These results emphasize the

46 importance of tailored initiatives and better access, to health information to increase HPV
47 vaccine acceptance and lower the incidence of cervical cancer.

48 **Keywords**

49 *Adolescent girls, Bangladesh, Cervical cancer, HPV vaccination, Parents, School teachers*

50 **Introduction**

51 Cervical cancer ranks as the fourth most common cancer among women globally, with a
52 significant burden in low-resource countries [1]. In 2022, 660,000 women were diagnosed and
53 350,000 died from this disease worldwide [1]. In Bangladesh, cervical cancer represents 12% of
54 female cancers, following breast cancer (19%) and esophageal cancer (11.1%) [2]. In 2020, there
55 were 8,268 new cases and 4,971 deaths in Bangladesh [2]. Infection with HPV, a non-enveloped,
56 double-stranded DNA virus, is a critical cause of cervical cancer [2]. Among the approximately
57 100 fully sequenced HPV types, high-risk types 16 and 18 are the main contributors to cervical
58 cancer [3,4], with HPV type 16 responsible for over 50% of squamous cell carcinomas and most
59 adenocarcinoma cases [5–8].

60 HPV infections typically occur in late teens or early 20s in unvaccinated populations [6,9]. The
61 Immature metaplastic cells in the cervix, especially during puberty and early pregnancy, are
62 highly susceptible to HPV infection [10]. However, the HPV vaccine played a major public
63 health role in preventing HPV-related cancers [11,12]. Prophylactic vaccines, such as bivalent,
64 quadrivalent, and nonavalent, are produced using virus-like particle (VLP) technology
65 [11,13,14], inducing type-specific antibodies that prevent persistent HPV infection and
66 precancerous lesions [15,16].

67 In 2008, the US Food and Drug Administration approved the first HPV vaccine, which prevents
68 70% of cervical cancers and other HPV-related diseases [17]. By 2017, 71 countries had
69 introduced the vaccine for girls aged 9 to 13 years [18]. Bangladesh introduced the HPV vaccine
70 in 2016, in the Gazipur district, funded by the Global Alliance for Vaccines and Immunization
71 (GAVI) [19]. However, successful implementation requires addressing knowledge gaps and
72 societal attitudes toward the vaccine [20].

73 Studies highlight the importance of evaluating vaccine-related knowledge among adolescents,
74 parents, and school teachers. For instance, in Kampong Speu, Cambodia, 35% of women knew
75 about cervical cancer prevention through vaccination, and 62% were willing to receive the
76 vaccine [21]. Another study revealed high willingness among both urban and rural respondents
77 to vaccinate themselves or their daughters [22]. These findings emphasize the influence of
78 parents and school teachers on adolescent vaccination decisions.

79 Bangladesh's National Cervical Cancer Control Program aims to integrate the HPV vaccine into
80 the Expanded Program on Immunization (EPI) and expand cervical cancer screening and
81 treatment [23]. The high public health burden of cervical cancer in Bangladesh makes achieving
82 widespread vaccination essential. The WHO's goal of vaccinating 90% of girls under 15 by 2030
83 further underscores this necessity [24]. However, adolescents themselves cannot make
84 vaccination decisions; this responsibility falls to their parents, with school teachers playing a
85 crucial role in influencing awareness and dispelling myths about vaccination [25,26].

86 While global studies have assessed HPV vaccine awareness, significant gaps remain, particularly
87 in low-resource countries like Bangladesh. Therefore, this study aims to address these gaps by
88 evaluating the knowledge and willingness of parents and school teachers towards HPV
89 vaccination for adolescent girls in Bangladesh. This study helps to generate actionable insights
90 that can inform future public health interventions, ultimately improving vaccine coverage and
91 reducing cervical cancer incidence. The large sample size and focus on socio-demographic
92 variations make this study highly relevant and generalizable.

93 **Materials and Methods**

94 **Study Design**

95 This study employed a cross-sectional design. The study involved quantitative data collection
96 using a structured, interviewer-administered questionnaire. The structured questionnaire was
97 designed based on a thorough review of existing literature and consultations with three experts in
98 the field. It was pre-tested and refined to ensure clarity, reliability, and validity. **Table 1** shows
99 that in the questionnaire, there were 8 sets of questions for knowledge and 8 sets of questions for
100 willingness. The reliability and internal consistency of the questions were assessed through

101 Cronbach's alpha (α) which showed the coefficient value is higher than 0.80 indicating very
102 strong internal consistency and reliability were present on it. Data were collected from a diverse
103 sample of parents and school teachers from different types of schools (public, private) and
104 various geographical locations within Dhaka (urban, and rural areas).

105 **Insert Table 1 here**

106 **Study Population and Place of Study**

107 This study focused on parents and school teachers of adolescent girls aged 9-14 years attending
108 primary and secondary schools in Dhaka, Bangladesh. Dhaka is the capital and largest city of
109 Bangladesh, and more than 10 million people live here [27]. This provides a diverse
110 demographic profile which was one of the suitable sites for conducting the study. The research
111 was conducted across various schools within this city to ensure a representative sample reflective
112 of the broader population.

113 **Study Period and Inclusion/Exclusion Criteria**

114 The research was carried out over five months, from August 2023 to December 2023, following
115 the approval of the research proposal. Participants included parents of adolescent girls aged 9-14
116 years and school teachers who directly interacted with girls within the same age range. To
117 maintain the study's integrity and relevance, several exclusion criteria were applied: participants
118 residing outside Dhaka, individuals with major psychological or psychiatric conditions, parents
119 of non-school-going adolescent girls, and individuals unwilling to provide information. These
120 criteria ensured the inclusion of a relevant and targeted participant pool.

121 **Sampling Technique and Sample Size**

122 A convenient sampling technique was employed in this study. This approach was chosen due to
123 practical considerations such as time constraints and the accessibility of participants within the
124 study period. Convenient sampling allowed for the efficient collection of data from readily
125 available and willing participants, which was crucial given the logistical challenges of accessing
126 a large and diverse population within Dhaka.

127 Although convenient sampling may limit the generalizability of the findings, it provided a
128 practical solution for obtaining a sufficient sample size within the specified timeframe. Schools
129 were selected based on their willingness to participate and their accessibility to the research
130 team. Within these schools, parents and teachers who met the inclusion criteria and were willing
131 to participate were approached.

132 The sample size was determined using a standard formula for prevalence studies:

133
$$N = z^2 y \frac{1 - y}{w^2}$$

134 In this formula, N represents the desired sample size, Z the standard normal deviation at 5%
135 type-1 errors ($p < 0.05$) with a 95% confidence interval, which is 1.96. The proportion of the
136 population exhibiting the characteristic of interest, based on previous studies, was set at 59.2% (y
137 = 0.592) [28]. The allowable error or precision (w) was set at 0.05. Substituting these values into
138 the formula, the initial sample size calculated was 371.15. To account for a potential 5% dropout
139 rate, the minimum sample size was needed to 390 participants. In this study, a total of 435
140 samples were collected. Upon applying exclusion criteria, removing missing values, and
141 duplicate entries finally 406 samples were kept for the study.

142 **Data Collection**

143 Data collection involved a structured, face-to-face, interviewer-administered questionnaire.
144 Initially, the questionnaire was developed in English and then translated into Bengali to ensure
145 comprehensibility among all participants. The development process of the questionnaire
146 involved several steps: drafting an initial version based on a thorough literature review and
147 expert consultations, translating the draft into Bengali by bilingual experts, pre-testing the
148 questionnaire with a small group of parents and teachers to identify any language or
149 comprehension issues, refining the questionnaire based on feedback from the pre-test, and
150 conducting a field trial in two schools to finalize the questionnaire before the main data
151 collection phase.

152 **Ethical Consideration**

153 The study received approval from the Institutional Review Board (IRB) of North South
154 University (IRB #2024/OR-NSU/IRB/0803), which was mandated by regulatory requirements.
155 The ethical guidelines specified in the 1964 Declaration of Helsinki and its later amendments
156 were adhered to wherever applicable. Prior to participation, all eligible respondents were
157 provided with a detailed explanation of the study's purpose, procedures, potential risks, and
158 benefits. Written informed consent was obtained from each participant. The respondents were
159 assured that their privacy and confidentiality would be strictly protected, and no part of their
160 responses would be disclosed to unauthorized individuals under any circumstances. This
161 commitment to ethical standards ensured the protection and respect of all participants throughout
162 the study.

163 **Data analysis**

164 The collected data were meticulously coded and entered into Stata version 17 for analysis. After
165 applying inclusion and exclusion criteria and removal of any missing and duplicate data, the final
166 dataset was prepared. This was checked by the supervisor as to see if the integrity and data
167 quality were assured properly or not. Descriptive statistics were used initially to summarize the
168 demographic characteristics of the study population, providing a clear overview of the study
169 sample, including the distribution of demographics such as age, gender, educational level, and
170 occupation.

171 For the knowledge assessment, eight specific questions were used, each with a binary response
172 format of 'Yes' or 'No.' These questions covered key aspects of HPV and its vaccination.
173 Participants were asked whether having multiple sexual partners is a risk factor for HPV
174 infection, whether engaging in sexual activity at a young age increases the risk of HPV infection,
175 whether being a smoker increases the risk of HPV infection, whether sexual contact is the
176 primary route of transmission for HPV infection, whether the primary cause of cervical
177 carcinoma is HPV infection, whether people can transmit HPV to their partners even without
178 showing any symptoms, whether receiving the HPV vaccine before sexual intercourse can
179 prevent cervical carcinoma, and whether the recommended age for receiving the HPV vaccine is
180 nine to fourteen years old. Responses were coded as 1 for 'Yes' and 0 for 'No'. A composite
181 knowledge score was calculated for each participant, ranging from 0 to 8. The cut point was set

182 to 4 as if the participants scoring equal and less than 4 were categorized as having poor
183 knowledge, while those scoring higher than 4 were categorized as having good knowledge.

184 Willingness to vaccinate was assessed using eight Likert-scale questions, with responses scored
185 from 1 (Strongly Disagree) to 5 (Strongly Agree). Participants were asked whether having only
186 one sexual partner can protect an individual from HPV infection, whether HPV vaccine
187 education should be provided to adolescents in schools, whether cervical cancer is a significant
188 health issue for women, whether cervical cancer can cause death in women, whether men's
189 involvement is important in the prevention of cervical carcinoma, whether girls should receive
190 the HPV vaccine before their first sexual intercourse, whether health information about the HPV
191 vaccine is necessary for adolescents, and whether the HPV vaccine is effective in preventing
192 cervical cancer. A total willingness score was calculated for each participant, ranging from 8 to
193 40. Participants with a total score above 24 were classified as willing to vaccinate, while those
194 with a score of 24 or below were classified as not willing.

195 Bivariate analysis was performed to explore the association between demographic variables and
196 the categorized knowledge and willingness level. Chi-square tests were used to identify
197 significant associations between categorical variables. Variables with a p-value ≤ 0.2 in the
198 bivariate analysis were included in the logistic regression model to ensure that potentially
199 relevant factors were not excluded prematurely.

200 To adjust for potential confounders and to determine the independent effect of each variable on
201 the outcomes of interest, multivariate logistic regression was employed. In this step, all
202 independent variables that showed a p-value ≤ 0.2 in the chi-square were included in the
203 multivariate model. The final model was refined by sequentially removing non-significant
204 variables and reassessing the fit of the model. The significance level was shown as the level of
205 strength as ψ for a p-value less than 0.05, $\psi\psi$ for a *P-value* less than 0.01, and $\psi\psi\psi$ for a p-value
206 less than 0.001. Adjusted odds ratios (AOR) with 95% confidence intervals (CI) were also shown
207 to identify significant predictors. The results of the analysis were presented using text, tables,
208 and figures to provide a comprehensive depiction of the findings.

209 **Results**

210 In this study, a total of 406 respondents were included encompassing a diverse population in
211 terms of age, participant type, residence, education level, and socioeconomic status. **Table 2**
212 shows the demographic characteristics of the participants showing that the majority were aged
213 between 31-40 years (52.96%). Most respondents were parents (69.21%), while the rest were
214 school teachers (30.79%). The sample was nearly evenly split between urban (49.51%) and rural
215 (50.49%) residents. Education levels varied, with higher education being the most common
216 (47.78%) among the participants. Socioeconomic status was also balanced, with 50.25% of
217 participants classified as lower middle income and 49.75% as upper middle income.

218 **Insert Table 2 here**

219 **Table 3** provides a detailed breakdown of the responses to knowledge and willingness-related
220 statements. There were a significant proportion of respondents (60.84%) recognized that having
221 multiple sexual partners increases the risk of HPV infection, and (55.17%) understood that
222 receiving the HPV vaccine before sexual intercourse can prevent cervical carcinoma.
223 Additionally, a substantial majority (87.37%) believed that providing HPV vaccine education in
224 schools is essential, and (83.98%) agreed that men's involvement is important for the prevention
225 of cervical carcinoma.

226 **Insert Table 3 here**

227 Among the participants, a majority (72.17%) had heard about cervical cancer, and (88.42%) had
228 no history of cervical cancer in their family. Most respondents (92.12%) indicated no history of
229 sexual activity among their girls, and (67.00%) did not express fear of sexual transmission of
230 HPV. Regarding HPV vaccination, (29.56%) had received the vaccine. These findings highlight
231 a high level of awareness about cervical cancer and a substantial portion of participants taking
232 preventive measures such as HPV vaccination.

233 **Insert Table 4 here**

234 After properly scoring and labeling the knowledge and willingness category the results indicated
235 that the majority of participants had poor knowledge about the HPV vaccine and cervical cancer,
236 with (64.04%) demonstrating poor knowledge and only (35.96%) showing good knowledge.
237 However, the willingness to participate in HPV vaccination programs was notably high, with
238 (98.28%) of participants expressing their willingness.

239 **Insert Fig 1 here:** Bar chart showing the prevalence (%) of knowledge and Willingness level
240 according to their category.

241 The association analysis revealed significant relationships between knowledge levels and various
242 demographic characteristics. The age group was significantly associated with knowledge levels
243 ($X^2=10.721$, $p=0.013$), with the highest knowledge observed in the 31-40 years age group.
244 Participant type also showed a significant association ($X^2=7.287$, $p=0.007$). Residence was
245 another significant factor ($X^2=18.367$, $p<0.001$), with urban residents having higher knowledge
246 levels than rural residents. Moreover, Education level ($X^2=37.068$, $p<0.001$) and socioeconomic
247 status ($X^2=42.073$, $p<0.001$) were both significantly associated with knowledge levels,
248 highlighting that higher education and better socioeconomic status correlate with better
249 knowledge about HPV vaccine and cervical cancer.

250 However, willingness to participate in HPV vaccination programs didn't show any significant
251 association.

252 **Insert Table 5 here**

253 Through the logistic regression model, the study analyzed the demographic factors influencing
254 knowledge levels about HPV and cervical cancer. Significant findings indicated that participants
255 with primary education (AOR=3.306, $p<0.005$), secondary education (AOR=8.806, $p<0.001$)
256 and higher education (AOR=5.059, $p<0.001$) had considerably higher likelihood of knowledge
257 levels compared to those with no education. Additionally, individuals from upper-middle-income
258 groups were significantly more knowledgeable than those from lower-middle-income groups
259 (AOR=3.038, $p<0.001$). These results highlight the critical role of higher education and better
260 socioeconomic status in enhancing awareness and knowledge about HPV and cervical cancer.

261 **Insert Table 6 here**

262 **Discussion**

263 The results of this study highlight several important aspects regarding the knowledge and
264 willingness of parents and school teachers of adolescent girls toward the HPV vaccination in
265 Bangladesh. The demographic characteristics revealed that the majority of participants were aged
266 between 31-40 years, with a significant proportion being parents rather than school teachers.

267 This demographic distribution is important as it suggests that the primary caregivers, who are
268 directly responsible for the health decisions of adolescent girls, form the bulk of the respondents.

269 The knowledge-related findings showed that a significant portion of respondents recognized that
270 having multiple sexual partners increases the risk of HPV infection and that receiving the HPV
271 vaccine before sexual intercourse can prevent cervical carcinoma. However, there were notable
272 gaps in knowledge, such as the misunderstanding that HPV infection is not the primary cause of
273 cervical carcinoma and the lack of awareness that HPV can be transmitted without symptoms.
274 These findings align with previous studies where awareness about HPV and its link to cervical
275 cancer is generally low. For instance, similar gaps in knowledge, suggest that the lack of
276 comprehensive sexual health education and cultural taboos surrounding discussions of sexual
277 health contribute to these misconceptions [29,30].

278 Furthermore, the study found that urban residents and those with higher education levels had
279 significantly better knowledge about HPV and cervical cancer. Urban residents typically have
280 better access to health information and services, which likely contributes to their higher
281 knowledge levels [31]. Additionally, higher education is often correlated with improved health
282 literacy, enabling individuals to better understand and utilize health information [32].

283 Despite the gaps in knowledge, the study revealed a high willingness among respondents to
284 participate in HPV vaccination programs. The overwhelming majority of participants expressed
285 support for providing HPV vaccine education in schools and recognized the importance of men's
286 involvement in preventing cervical carcinoma. This high level of willingness can be attributed to
287 the general recognition of cervical cancer as a significant health issue and the perceived benefits
288 of vaccination. Previous research supports this finding, indicating that when individuals are
289 aware of the benefits of vaccination, they are more likely to support and participate in
290 vaccination programs [33].

291 The logistic regression analysis provided further insights into the key factors influencing
292 knowledge levels about HPV and cervical cancer. Participants with primary, secondary, and
293 higher education had a significantly higher likelihood of having better knowledge levels
294 compared to those with no education. Additionally, individuals from upper-middle-income
295 groups were more knowledgeable than those from lower-middle-income groups. These results
296 emphasize the critical role of education and socioeconomic status in enhancing awareness and

297 knowledge about HPV. Education equips individuals with the necessary skills to seek out,
298 understand, and apply health information, while higher socioeconomic status provides better
299 access to healthcare resources and information [34].

300 **Limitation**

301 Although every research has limitations, one issue could be this research relies on information
302 provided by the participants themselves which could lead to biases like wanting to appear as
303 memory-related biases. People might exaggerate behaviors that are seen positively or downplay
304 those that are not affecting the accuracy of the results. Moreover, because the study only captures
305 a moment in time through its cross-sectional design it can't fully establish cause-and-effect
306 relationships or track changes over time. Additionally, although the research involves individuals
307 from both urban and rural areas using a convenience sampling method might limit how broadly
308 we can apply the findings to the population. Finally, even though steps were taken to make sure
309 the survey was reliable and valid difference, in culture and language might still influence how
310 participants interpret questions and provide responses potentially impacting what conclusions we
311 draw from the study.

312 **Conclusion**

313 The findings reveal significant knowledge gaps about HPV and its vaccination among parents
314 and school teachers in Bangladesh, despite a strong willingness to support vaccination programs.
315 Factors such as education level and socioeconomic status were found to significantly influence
316 knowledge levels. The high willingness to participate in HPV vaccination programs highlights
317 the potential for targeted educational interventions and improved vaccine access, driven by a
318 general awareness of cervical cancer as a major health issue and the perceived benefits of
319 vaccination. Enhancing access to accurate health information and making vaccines more
320 available, especially in rural and lower socioeconomic areas, can substantially reduce HPV-
321 related cervical cancer in Bangladesh. These efforts are crucial for improving the health
322 outcomes of adolescent girls and supporting the broader public health objectives of the country.

323 **Abbreviation**

| Abbreviation | Full Term |
|----------------|---|
| AOR | Adjusted Odds Ratio |
| CI | Confidence Interval |
| EPI | Expanded Program on Immunization |
| GAVI | Global Alliance for Vaccines and Immunization |
| HPV | Human Papillomavirus |
| P-value | Probability Value |
| VLP | Virus-Like Particle |
| WHO | World Health Organization |
| X ² | Chi-Square |
| α | Cronbach's Alpha |

324

325 Acknowledgments

326 We would like to express our gratitude to all the parents and teachers for their participation.
327 Without their support, it would be difficult to get the data.

328 Reference

- 329 1. Cervical cancer. [cited 26 Jun 2024]. Available: [https://www.who.int/news-room/fact-](https://www.who.int/news-room/fact-sheets/detail/cervical-cancer)
330 [sheets/detail/cervical-cancer](https://www.who.int/news-room/fact-sheets/detail/cervical-cancer)
- 331 2. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global
332 Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for
333 36 Cancers in 185 Countries. *CA A Cancer J Clinicians*. 2021;71: 209–249.
334 doi:10.3322/caac.21660
- 335 3. Schiffman M, Clifford G, Buonaguro FM. Classification of weakly carcinogenic human
336 papillomavirus types: addressing the limits of epidemiology at the borderline. *Infect Agent*
337 *Cancer*. 2009;4: 8. doi:10.1186/1750-9378-4-8
- 338 4. International Agency for Research on Cancer, editor. Human papillomaviruses: this
339 publication represents the views and expert opinions of an IARC Working Group on the
340 Evaluation of Carcinogenic Risks to Humans, which met in Lyon, 15 - 22 February 2005.
341 Lyon: IARC; 2007.

- 342 5. Muñoz N, Castellsagué X, De González AB, Gissmann L. Chapter 1: HPV in the etiology
343 of human cancer. *Vaccine*. 2006;24: S1–S10. doi:10.1016/j.vaccine.2006.05.115
- 344 6. Lo KWK, Wong YF, Chan MKM, Li JCB, Poon JS, Wang VW, et al. Prevalence of human
345 papillomavirus in cervical cancer: A multicenter study in China. *Intl Journal of Cancer*.
346 2002;100: 327–331. doi:10.1002/ijc.10506
- 347 7. Bosch FX, de Sanjosé S. The epidemiology of human papillomavirus infection and cervical
348 cancer. *Dis Markers*. 2007;23: 213–227. doi:10.1155/2007/914823
- 349 8. Castellsagué X, Díaz M, De Sanjosé S, Muñoz N, Herrero R, Franceschi S, et al.
350 Worldwide Human Papillomavirus Etiology of Cervical Adenocarcinoma and Its Cofactors:
351 Implications for Screening and Prevention. *JNCI: Journal of the National Cancer Institute*.
352 2006;98: 303–315. doi:10.1093/jnci/djj067
- 353 9. Bosch FX, De Sanjose S. Chapter 1: Human Papillomavirus and Cervical Cancer--Burden
354 and Assessment of Causality. *JNCI Monographs*. 2003;2003: 3–13.
355 doi:10.1093/oxfordjournals.jncimonographs.a003479
- 356 10. Okunade KS. Human papillomavirus and cervical cancer. *J Obstet Gynaecol*. 2020;40:
357 602–608. doi:10.1080/01443615.2019.1634030
- 358 11. Frazer IH. CHAPTER 8 HPV vaccines. *Intl J Gynecology & Obste*. 2006;94.
359 doi:10.1016/S0020-7292(07)60014-9
- 360 12. Smith MA, Lew J-B, Walker RJ, Brotherton JML, Nickson C, Canfell K. The predicted
361 impact of HPV vaccination on male infections and male HPV-related cancers in Australia.
362 *Vaccine*. 2011;29: 9112–9122. doi:10.1016/j.vaccine.2011.02.091
- 363 13. Zhou J, Sun XY, Stenzel DJ, Frazer IH. Expression of vaccinia recombinant HPV 16 L1
364 and L2 ORF proteins in epithelial cells is sufficient for assembly of HPV virion-like
365 particles. *Virology*. 1991;185: 251–257. doi:10.1016/0042-6822(91)90772-4

- 366 14. Kirnbauer R, Taub J, Greenstone H, Roden R, Dürst M, Gissmann L, et al. Efficient self-
367 assembly of human papillomavirus type 16 L1 and L1-L2 into virus-like particles. *J Virol.*
368 1993;67: 6929–6936. doi:10.1128/jvi.67.12.6929-6936.1993
- 369 15. Giuliano AR, Palefsky JM, Goldstone S, Moreira ED, Penny ME, Aranda C, et al. Efficacy
370 of Quadrivalent HPV Vaccine against HPV Infection and Disease in Males. *N Engl J Med.*
371 2011;364: 401–411. doi:10.1056/NEJMoa0909537
- 372 16. Palefsky JM, Giuliano AR, Goldstone S, Moreira ED, Aranda C, Jessen H, et al. HPV
373 Vaccine against Anal HPV Infection and Anal Intraepithelial Neoplasia. *N Engl J Med.*
374 2011;365: 1576–1585. doi:10.1056/NEJMoa1010971
- 375 17. Gardasil 9 (human papillomavirus 9-valent vaccine, recombinant) FDA Approval History.
376 In: Drugs.com [Internet]. [cited 26 Jun 2024]. Available:
377 <https://www.drugs.com/history/gardasil-9.html>
- 378 18. Vaccines and immunization: What is vaccination? [cited 26 Jun 2024]. Available:
379 [https://www.who.int/news-room/questions-and-answers/item/vaccines-and-immunization-](https://www.who.int/news-room/questions-and-answers/item/vaccines-and-immunization-what-is-vaccination)
380 [what-is-vaccination](https://www.who.int/news-room/questions-and-answers/item/vaccines-and-immunization-what-is-vaccination)
- 381 19. Salwa M, Abdullah Al-Munim T. Ethical issues related to human papillomavirus
382 vaccination programs: an example from Bangladesh. *BMC Med Ethics.* 2018;19: 39.
383 doi:10.1186/s12910-018-0287-0
- 384 20. Kessels SJM, Marshall HS, Watson M, Braunack-Mayer AJ, Reuzel R, Tooher RL. Factors
385 associated with HPV vaccine uptake in teenage girls: A systematic review. *Vaccine.*
386 2012;30: 3546–3556. doi:10.1016/j.vaccine.2012.03.063
- 387 21. Touch S, Oh J-K. Knowledge, attitudes, and practices toward cervical cancer prevention
388 among women in Kampong Speu Province, Cambodia. *BMC Cancer.* 2018;18: 294.
389 doi:10.1186/s12885-018-4198-8
- 390 22. Islam JY, Khatun F, Alam A, Sultana F, Bhuiyan A, Alam N, et al. Knowledge of cervical
391 cancer and HPV vaccine in Bangladeshi women: a population based, cross-sectional study.
392 *BMC Women’s Health.* 2018;18: 15. doi:10.1186/s12905-018-0510-7

- 393 23. National Strategy for Cervical Cancer Prevention and Control in Bangladesh, 2017-2022.
394 [cited 26 Jun 2024]. Available: [https://www.who.int/bangladesh/news/detail/24-09-2017-](https://www.who.int/bangladesh/news/detail/24-09-2017-national-strategy-for-cervical-cancer-prevention-and-control-in-bangladesh-2017-2022)
395 [national-strategy-for-cervical-cancer-prevention-and-control-in-bangladesh-2017-2022](https://www.who.int/bangladesh/news/detail/24-09-2017-national-strategy-for-cervical-cancer-prevention-and-control-in-bangladesh-2017-2022)
- 396 24. Bruni L, Saura-Lázaro A, Montoliu A, Brotons M, Alemany L, Diallo MS, et al. HPV
397 vaccination introduction worldwide and WHO and UNICEF estimates of national HPV
398 immunization coverage 2010–2019. *Preventive Medicine*. 2021;144: 106399.
399 doi:10.1016/j.ypmed.2020.106399
- 400 25. Smailbegovic MS, Laing GJ, Bedford H. Why do parents decide against immunization?
401 The effect of health beliefs and health professionals. *Child*. 2003;29: 303–311.
402 doi:10.1046/j.1365-2214.2003.00347.x
- 403 26. Gowda C, Schaffer SE, Dombkowski KJ, Dempsey AF. Understanding attitudes toward
404 adolescent vaccination and the decision-making dynamic among adolescents, parents and
405 providers. *BMC Public Health*. 2012;12: 509. doi:10.1186/1471-2458-12-509
- 406 27. Population of Cities in Bangladesh 2024. [cited 26 Jun 2024]. Available:
407 <https://worldpopulationreview.com/countries/cities/bangladesh>
- 408 28. Mihretie GN, Liyeh TM, Ayele AD, Belay HG, Yimer TS, Miskr AD, et al. Female
409 adolescents' knowledge and acceptability of human papillomavirus vaccine in Debre Tabor
410 Town, Ethiopia: a cross-sectional study. *BMJ Open*. 2023;13: e061813.
411 doi:10.1136/bmjopen-2022-061813
- 412 29. Lukumay GG, Mgopa LR, Mushy SE, Rosser BRS, Massae AF, Mkonyi E, et al.
413 Community myths and misconceptions about sexual health in Tanzania: Stakeholders'
414 views from a qualitative study in Dar es Salaam Tanzania. Hodges MH, editor. *PLoS ONE*.
415 2023;18: e0264706. doi:10.1371/journal.pone.0264706
- 416 30. Latifnejad Roudsari R, Javadnoori M, Hasanpour M, Hazavehei SMM, Taghipour A.
417 Socio-cultural challenges to sexual health education for female adolescents in Iran. *Iran J*
418 *Reprod Med*. 2013;11: 101–110.

- 419 31. Chen X, Orom H, Hay JL, Waters EA, Schofield E, Li Y, et al. Differences in Rural and
420 Urban Health Information Access and Use. *J Rural Health*. 2019;35: 405–417.
421 doi:10.1111/jrh.12335
- 422 32. Bayati T, Dehghan A, Bonyadi F, Bazrafkan L. Investigating the effect of education on
423 health literacy and its relation to health-promoting behaviors in health center. *J Educ Health*
424 *Promot*. 2018;7: 127. doi:10.4103/jehp.jehp_65_18
- 425 33. Lechuga J, Swain G, Weinhardt LS. Perceived need of a parental decision aid for the HPV
426 vaccine: content and format preferences. *Health Promot Pract*. 2012;13: 214–221.
427 doi:10.1177/1524839910388622
- 428 34. Protheroe J, Nutbeam D, Rowlands G. Health literacy: a necessity for increasing
429 participation in health care. *Br J Gen Pract*. 2009;59: 721–723.
430 doi:10.3399/bjgp09X472584

431

432

433 *Table 1: Scale Reliability and Covariance for Knowledge and Attitude Statements*

| Scale Type | Average inter-item Covariance | Number of Items | Scale Reliability Coefficient |
|-------------------------------------|-------------------------------|-----------------|-------------------------------|
| Knowledge Related statements | 0.0893693 | 8 | 0.8358 |
| Attitude Related statements | 0.2062145 | 8 | 0.8370 |

434

435 *Table 2: Demographic characteristics of the participants*

| Characteristics | Frequency (N) | Percentage (%) |
|------------------------------|---------------|----------------|
| Age group | | |
| 20-30 y | 65 | 16.01 |
| 31-40 y | 215 | 52.96 |
| 41-50 y | 114 | 28.08 |
| Above 50 y | 12 | 2.96 |
| Participant type | | |
| Parents | 281 | 69.21 |
| School Teacher | 125 | 30.79 |
| Resident | | |
| Urban | 201 | 49.51 |
| Rural | 205 | 50.49 |
| Education Level | | |
| No Education | 55 | 13.55 |
| Primary | 64 | 15.76 |
| Secondary | 93 | 22.91 |
| Higher | 194 | 47.78 |
| Socio Economic Status | | |
| Lower Middle Income | 204 | 50.25 |
| Upper Middle Income | 202 | 49.75 |

436

437 *Table 3: Distribution of Knowledge and Willingness Related Statement Among Respondents*

| Statements | Freq. | Percent |
|---|-------|---------|
| Knowledge Related | | |
| Multiple sexual partners increase HPV infection risk | | |
| No | 159 | 39.16 |
| Yes | 247 | 60.84 |
| Early sexual activity increases HPV infection risk | | |
| No | 330 | 81.28 |
| Yes | 76 | 18.72 |
| Smoking increases HPV infection risk | | |
| No | 202 | 49.75 |
| Yes | 204 | 50.25 |
| Sexual contact is the primary HPV transmission route | | |
| No | 219 | 53.94 |
| Yes | 187 | 46.06 |
| HPV infection is the primary cause of cervical carcinoma | | |
| No | 260 | 64.04 |
| Yes | 146 | 35.96 |
| HPV can be transmitted without symptoms | | |
| No | 269 | 66.26 |

| | | |
|---|-----|-------|
| Yes | 137 | 33.74 |
| HPV vaccine before sex prevents cervical carcinoma | | |
| No | 182 | 44.83 |
| Yes | 224 | 55.17 |
| The recommended age for the HPV vaccine is 9-14 years | | |
| No | 235 | 57.88 |
| Yes | 171 | 42.12 |
| Willingness Related | | |
| I believe having one sexual partner protects from HPV | | |
| Strongly Disagree | 10 | 2.46 |
| Disagree | 14 | 3.45 |
| Neutral | 56 | 13.79 |
| Agree | 190 | 46.80 |
| Strongly Agree | 136 | 33.50 |
| I support providing HPV vaccine education in schools | | |
| Strongly Disagree | 1 | 0.25 |
| Disagree | 11 | 2.71 |
| Neutral | 19 | 4.68 |
| Agree | 133 | 32.76 |
| Strongly Agree | 242 | 59.61 |
| I consider cervical cancer a significant issue for women | | |
| Strongly Disagree | 1 | 0.25 |
| Disagree | 2 | 0.49 |
| Neutral | 14 | 3.45 |
| Agree | 140 | 34.48 |
| Strongly Agree | 249 | 61.33 |
| I believe cervical cancer can cause death | | |
| Strongly Disagree | 2 | 0.49 |
| Disagree | 5 | 1.23 |
| Neutral | 10 | 2.46 |
| Agree | 154 | 37.93 |
| Strongly Agree | 235 | 57.88 |
| I think men's involvement is important for prevention | | |
| Strongly Disagree | 1 | 0.25 |
| Disagree | 10 | 2.46 |
| Neutral | 43 | 10.59 |
| Agree | 169 | 41.63 |
| Strongly Agree | 183 | 45.07 |
| support girls getting the HPV vaccine before their first sex | | |
| Strongly Disagree | 2 | 0.49 |
| Disagree | 7 | 1.72 |
| Neutral | 48 | 11.82 |
| Agree | 199 | 49.01 |
| Strongly Agree | 150 | 36.95 |
| I believe HPV vaccine information is necessary for adolescents | | |
| Strongly Disagree | 0 | 0.00 |
| Disagree | 4 | 0.99 |
| Neutral | 10 | 2.46 |
| Agree | 187 | 46.06 |
| Strongly Agree | 205 | 50.49 |
| I agree that the HPV vaccine prevents cervical cancer | | |
| Strongly Disagree | 3 | 0.74 |
| Disagree | 5 | 1.23 |
| Neutral | 32 | 7.88 |
| Agree | 196 | 48.28 |
| Strongly Agree | 170 | 41.87 |

439

440 *Table 4: Reproductive Health Factors vs. Human Papillomavirus Vaccine Awareness Among Participants*

| Characteristics | Response | Frequency | Percent |
|------------------------------------|----------|-----------|---------|
| Heard about Cervical Cancer | Yes | 293 | 72.17 |
| | No | 113 | 27.83 |
| History of Cervical Cancer | No | 359 | 88.42 |
| | Yes | 47 | 11.58 |
| History of Sexual Activity | No | 374 | 92.12 |
| | Yes | 32 | 7.88 |
| Fear of Sexual Transmission | No | 272 | 67.00 |
| | Yes | 134 | 33.00 |
| Taken HPV Vaccine | No | 286 | 70.44 |
| | Yes | 120 | 29.56 |

441 *Table 5: Association between Knowledge and Willingness Levels across Demographic Characteristics*

| Characteristics | Good Knowledge N (%) | X ² Value | P-value | Have Willingness N (%) | X ² Value | P-value |
|-----------------------------|-------------------------|----------------------|-----------------------|---------------------------|----------------------|---------|
| Age Group | | 10.721 | 0.013 ^ψ | | 4.461 | 0.216 |
| 20-30 y | 15 (10.27) | | | 64 (16.04) | | |
| 31-40 y | 73 (50.00) | | | 213 (53.38) | | |
| 41-50 y | 53 (36.30) | | | 111 (27.82) | | |
| Above 50 y | 5 (3.42) | | | 11 (2.76) | | |
| Participant Type | | 7.287 | 0.007 ^{ψψ} | | 0.016 | 0.898 |
| Parents | 89 (60.96) | | | 276 (69.17) | | |
| School Teacher | 57 (39.04) | | | 123 (30.60) | | |
| Residence | | 18.367 | <0.001 ^{ψψψ} | | 1.369 | 0.242 |
| Urban | 93 (63.70) | | | 196 (49.12) | | |
| Rural | 53 (36.30) | | | 203 (50.88) | | |
| Education Level | | 37.068 | <0.001 ^{ψψψ} | | 2.959 | 0.398 |
| No Education | 4 (2.74) | | | 53 (13.28) | | |
| Primary | 13 (8.90) | | | 63 (15.79) | | |
| Secondary | 39 (26.71) | | | 93 (23.31) | | |
| Higher | 90 (61.64) | | | 190 (47.62) | | |
| Socioeconomic Status | | 42.073 | <0.001 ^{ψψψ} | | 1.338 | 0.247 |
| Lower Middle Income | 42 (28.77) | | | 202 (50.63) | | |
| Upper Middle Income | 104 (71.23) | | | 197 (49.37) | | |

442

443 Denote: (Chi² significance: ^ψ = p-value<0.05, ^{ψψ} = p-value=<0.01, ^{ψψψ} = p-value<0.001)

444

445 *Table 6: Demographic factors associated with the Knowledge level*

| Characteristics | AOR | St. Error | [95% Conf. Interval] |
|-------------------------|-------|-----------|----------------------|
| Age Group | | | |
| 20-30 y | Ref | | |
| 31-40 y | 1.598 | 0.602 | 0.763 - 3.346 |
| 41-50 y | 1.615 | 0.649 | 0.735 - 3.549 |
| Above 50 y | 2.548 | 1.883 | 0.598 - 10.847 |
| Participant Type | | | |


| | | | |
|-------------------------------|----------|-------|----------------|
| <i>Parents</i> | Ref | | |
| <i>School Teachers</i> | 1.278 | 0.396 | 0.697 - 2.344 |
| <i>Residence</i> | | | |
| <i>Urban</i> | Ref | | |
| <i>Rural</i> | 0.674 | 0.180 | 0.400 - 1.138 |
| <i>Education Level</i> | | | |
| <i>No Education</i> | Ref | | |
| <i>Primary</i> | 3.306* | 2.032 | 0.991 - 11.030 |
| <i>Secondary</i> | 8.806*** | 5.186 | 2.776 - 27.928 |
| <i>Higher</i> | 5.059*** | 3.082 | 1.533 - 16.698 |
| <i>Monthly Income</i> | | | |
| <i>Lower Middle Income</i> | Ref | | |
| <i>Upper Middle Income</i> | 3.038*** | 0.838 | 1.769 - 5.217 |

446

447 Denote: (P-value indication: * = P-value<0.05, ** = P-value=<0.01, *** = P-value<0.001, AOR=adjusted odds ratio)

448

Prevalence of Knowledge and Willingness Levels



Figure