

Oral hygiene practice, status, and periodontitis among dental outpatient elderly in tertiary hospitals: a cross-sectional study in Dhaka, Bangladesh

Sadia Mehjabeen Kanta¹, Sifat Sharmin², Md. Golam Sharower³, Zaki Farhana⁴, Shuvojit Kumar Kundu⁵, Sadhan Kumar Das⁶, Anton Abdulbasah Kamil⁷, Mohammad Meshbahur Rahman^{8*}

¹Department of Health Education, National Institute of Preventive and Social Medicine, Mohakhlai, Dhaka-1212. Email: sadia.shd3@gmail.com

²Department of Hospital Management, National Institute of Preventive and Social Medicine, Mohakhlai, Dhaka-1212. Email: limutanvir7889@gmail.com

³Department of Entomology, National Institute of Preventive and Social Medicine, Mohakhlai, Dhaka-1212. Email: sharower_bau@yahoo.com

⁴Division of Statistics, Bangladesh Bank-The Central Bank of Bangladesh, Bangladesh. Email: zfarhanaesha@gmail.com

⁵Directorate General of Health Services, Ministry of Health & Family Welfare, Government of the People's Republic of Bangladesh. Email: shuvojit495@gmail.com

⁶Department of Health Education, National Institute of Preventive and Social Medicine, Mohakhlai, Dhaka-1212. Email: sadhondas81@gmail.com

⁷Department of Business Administration, Istanbul Gelisim University, Istanbul, Turkey. Email: akamil@gelisim.edu.tr

⁸Department of Biostatistics, National Institute of Preventive and Social Medicine, Mohakhlai, Dhaka-1212. Email: meshbah037@gmail.com

Abstract

Background

Oral health significantly affects older people's quality of life, overall health, and well-being. The study aimed to assess oral hygiene practice, status, and periodontitis among elderly patients.

Methods

A descriptive cross-sectional study was conducted among elderly patients attending dental outpatient departments of tertiary hospitals in Dhaka city from January to December 2022. Data from two hundred and twenty-seven elderly patients were collected by face-to-face interview through a pretested, semi-structured questionnaire and oral examination using the Oral Hygiene Index Simplified (OHI-S), and Community Periodontal Index component of CPITN. Statistics were presented in frequency, mean, percentage, chi-square test, fisher's exact test, and binary logistic regression.

Results

Among 227 elderly patients, the mean (\pm SD) age was 55.75 (\pm 8.20) years; 51% were female, and 49% were male. Among respondents, 70.48% cleaned their teeth regularly, 61.23% cleaned their teeth once daily, 70.5% used toothbrushes and toothpaste for cleaning their teeth, and only 2.2% used interdental aids. More than half of the respondents, 54%, had poor oral hygiene status, and 55.5% had periodontitis. A significant association existed between periodontitis, age, gender, educational status, occupation, and monthly family income. Older adults aged 66 years and above were 7.2 times more likely to have periodontitis (OR_{adj} : 12.637, 95% CI: 3.371-47.371, $p < 0.0001$). Respondents who engaged in labour work were more likely to have periodontitis than the service holders (OR_{adj} :5.855, 95% CI: .594-57.75, $p < 0.130$). Periodontitis was significantly associated with oral hygiene practices and oral hygiene status. Elderly who did not clean their teeth regularly were roughly 15 times more likely to develop periodontitis (OR_{adj} : 9.474, 95% CI: 3.636-24.689, $p < 0.0001$). Older adults who did not use toothpaste and toothbrush were 5.1 times more likely to develop periodontitis (OR_{adj} : 3.792, 95% CI: 1.758- 8.178, $p < 0.001$).

Conclusion

More than half of the elderly individuals in the study had periodontitis and poor oral hygiene. The research found that the increase in periodontitis was linked to respondents' socio-demographic characteristics and oral hygiene practices. These findings underscore the importance of maintaining good oral hygiene and getting regular dental checkups for older adults.

Keywords: Periodontitis, Simplified Oral Hygiene Index (OHI-S), CPITN, older adults, Oral hygiene practice

Introduction

Bangladesh is currently experiencing an epidemiological transition characterized by a significant decline in mortality rates from acute, infectious, and parasitic diseases [1–6]. The acceleration of population ageing and an increase in the prevalence of oral health problems and non-communicable diseases will considerably challenge health and social policy planners [7–10].

The World Health Organization (WHO) identified the oral health of older adults as one of its primary concerns due to the growing population of elderly people [11]. Poor oral health in elderly persons is prevalent around the world, with high rates of tooth loss, dental caries, and the prevalence of periodontal disease, xerostomia, and oral precancer/cancer being particularly noticeable [12–14]. Individuals' physical and psychological well-being, as well as their appearance, pronunciation, ability to chew, taste of food, and sociability, are all impacted by their oral health [15]. In addition to affecting diet, nutrition, and phonetics, poor oral health and tooth loss also endanger general health. Numerous systemic disorders, including cardiovascular disease, bacterial pneumonia, diabetes mellitus, and low birth weight, are affected by oral infection, particularly periodontitis, on their course and aetiology [16]. Thus, good oral health is essential for improved systemic health and healthy dental status.

Proper oral hygiene is the cornerstone of good oral health, preventing eighty percent of oral health problems [17]. Effective oral hygiene and plaque removal are the most crucial methods in preventing periodontal disease since dental plaque is the principal etiological factor in the pathogenesis of this disease [18]. Periodontal disease is a group of inflammatory conditions affecting the gingiva, bone, and periodontal ligament, which can result in tooth loss and spread inflammation throughout the body [19]. Gingivitis, a reversible inflammation, is the first stage of periodontal disease, which can progress to periodontitis, an irreversible infection [20]. One-third of the general population suffers from periodontitis, which is more prevalent in older persons than in younger and middle-aged people [21].

A patient's dietary habits, quality of life, and sense of self-worth are all impacted by severe periodontal disease, which also raises the chance of multiple tooth loss, masticatory dysfunction, and edentulousness. Beyond its effects on the orofacial structures, severe periodontal disease has

been associated with several systemic conditions, including diabetes mellitus, cardiovascular disease, respiratory illnesses, renal disorders, and an elevated risk of head and neck cancers [22]. Therefore, it is imperative to address periodontitis in the elderly population. Nevertheless, there is a shortage of information on the oral health of older adults in Bangladesh [23]. Therefore, it is vital to identify the extent of oral hygiene practice and assess the periodontal health status among older adults. Furthermore, the study's findings may help the responsible stakeholders take necessary interventions to raise healthier oral hygiene and oral health status among older adults concerning public health activities.

Methods

Ethical consent and permission for data collection

This study was approved by the institutional review board of the National Institute of Preventive and Social Medicine (NIPSOM), Bangladesh (Ref No: NIPSOM/IRB/2017/09). The declaration of Helsinki (revised version) was followed to conduct the study. Both written and verbal consent was obtained before initiating the interview. A brief on the aims and objectives was given to the participants. Participants who agreed to give consent were finally included in the study.

Study setting and participants

The study was a descriptive cross-sectional study conducted among elderly patients who attended the dental outpatient departments of selected hospitals in Dhaka city within one year, from January to December 2022. The elderly patients aged 46years and above attending dental OPD of the selected hospitals (1. Shaheed Suhrawardy Medical College & Hospital, Sher-E-Bangla Nagar, Dhaka 2. Dhaka Dental College Hospital, Mirpur-14, Dhaka 3. Probin Hospital, Agargoan, Dhaka) were included in our study.

Sampling technique and sample size

A convenient type of non-probability sampling technique was followed in the study. The sample size of the study was calculated by using the formula below:

$$n = z^2pq/d^2$$

Here, n=assumed/desired sample size; z=the standard normal deviation, usually set at 1.96 at a 95% confidence level; p=proportion of periodontitis among elderly people is 82%=0.82, obtained by the literature search [24]; q=1-p; and d=Margin of error (5%) = 0.05.

The sample size for the study is

$$n = \frac{(1.96)^2 \times 0.82 \times 0.18}{(0.05)^2} = 226.8 = 227$$

Data were collected from 227 patients after obtaining informed written consent.

Selection criteria

The inclusion criteria were: (1) Elderly patients aged 46 years and above attending dental OPD (2) Willing to participate and given informed written consent. And the exclusion criteria were Patients with mental or physical disabilities, those who were edentulous, and those who did not have at least two teeth in each sextant were all excluded from this study.

Data collection tool and procedures

A pretested semi-structured questionnaire was used to collect the data through face-to-face interviews. The questionnaire contained a checklist for oral examination. The simplified Oral Hygiene Index (OHI-S) was used to assess individuals' oral hygiene status. The Community periodontal index (CPI) component of the Community Periodontal Index of Treatment Need (CPITN) was used to measure the periodontal disease condition of the respondents. Oral examinations were performed in the outpatient department of the hospital. The individuals to be examined remained seated on a chair, and the examiner stood. The examiner was vested with a mask and gloves and used a WHO community periodontal index probe, mirror, and caries probe (dental explorer).

Simplified Oral Hygiene Index (OHI-S)

A simplified oral hygiene index with debris (plaque) and calculus components was first described by J. Greene and J.R. Vermillion in 1964 [25]. OHI-S is an index that measures the current oral hygiene status based on the number of debris and calculus on six tooth surfaces in the mouth. Six tooth surfaces were scored, four posterior and two anterior. The index values were calculated from the recordings of the calculus and debris scores. The debris scores were added and divided by the number of surfaces scored for each person. A score for a group of individuals was obtained, and the average of the individual scores was calculated. The average group or individual score was

defined as the Simplified Debris Index (DI-S). The same process was used to obtain the calculus scores or the Simplified Calculus Index (CI-S). To obtain the Simplified Oral Hygiene Index (OHI-S), the average group or individual calculus and debris scores were added. Range of scores for oral hygiene index: Good: 0-1.2, Fair: 1.3-3, Poor: 3.1-6. A clinical examination of plaque and calculus was done using a caries probe (dental explorer) and mouth mirror.

Community Periodontal Index of Treatment Needs (CPITN)

The Community Periodontal Index of Treatment Needs (CPITN), introduced by the World Health Organization (WHO) in 1977, included periodontal probing on ten index teeth [26]. CPITN uses a periodontal probe to assess the presence or absence of gingival bleeding on probing, supra or sub-gingival calculus, and periodontal pockets. The mouth was divided into six sextants, and the worst score in each sextant was recorded. The 10 index teeth were examined at mesial and distal proximal sites on the buccal and lingual/palatal sides. Periodontal status was evaluated using the Community Periodontal Index (CPI) component of the CPITN index. About periodontitis ($CPI \geq 3$) and non-periodontitis ($CPI \leq 2$), all research variables were examined [24].

The scoring criteria of CPITN are as follows Score 0: Healthy periodontal conditions or no periodontal disease; Score 1: Bleeding on probing; Score 2: Calculus with plaque felt by probing, Score 3: Shallow periodontal pockets (4 to 5 millimeters), Score 4: Deep periodontal pockets (6 millimeters or more). Pocket depths were measured with a graduated WHO CPI probe.

Data quality control and statistical analysis

To ensure the reliability and validity of the study results, we used several techniques: (1) conducted pretesting of our study and revised the questionnaire (2) checked the data and fixed errors by observing descriptive statistics; (3) employed an updated version of statistical software in data analysis.

In data analysis, descriptive statistics, such as frequency, percentage, mean, and SD (standard deviation), were used to describe the respondents' socio-demographic characteristics and selective attributes. Inferential statistics including Chi-square test, Fisher's exact test, and Binary logistic regression were also done. A p-value of 5% was considered significant at the 95% confidence interval. After data collection and processing, this study compiled and analyzed data according to the objectives and variables by IBM Software – SPSS (Statistical Package for Social Science) version-26.

Result

Socio-demographic profile of the respondents

Table 1 shows the socio-demographic characteristics of the patients who participated in our study. According to the distribution of respondents by age group, the 46-55-year-old age group had the highest frequency, and the mean age was 55.72 years. Among 227 respondents, 51% were female, and 49% were male. It was observed that 53 (23.36%) respondents had no formal education (Table 1).

Table 1: Socio-demographic characteristics of the respondents

Variables	f (%)
Age group(years)	
46-55	110 (48.5)
56-65	89 (39.2)
66 and above	28 (12.3)
Gender	
Male	111 (49)
Female	116 (51)
Educational Status	
No formal education	53 (23.3)
Primary	46 (20.3)
Secondary	63 (27.8)
Higher secondary and above	65 (28.6)
Occupation	
Service holder	38 (16.7)
Businessman	24 (10.6)
Labourer	16 (7.0)
Homemaker	108 (47.6)
Unemployed/ Retired	41 (18.1)
Monthly family income	
5000-24000	146 (64.32)
24001-44000	38 (16.74)
44001 and above	43 (18.9)

About 46 (20.3%), 63 (27.85%), and 65 (28.6%) had completed primary, secondary, higher secondary and above, respectively. Almost half of the respondents were homemakers, 108

(47.6%), and the percentage of labourer respondents was the lowest (7%). It was also observed that the highest proportion of respondents, 146 (64.32%), had monthly family income within 5000–24000 BDT (Table-1).

Oral hygiene practices of respondents:

It was found that most respondents, 160 (70.48%), cleaned their teeth regularly, and 67 (29.52%) did not clean their teeth regularly. Most of them, 222 (97.8%), did not use interdental aids, 219(96.5%) did not use mouthwash, 200 (88.1%) did not clean their tongue, whereas 59 (26%) respondents rinsed their mouth after a meal. It was found that the majority of the respondents, 222 (97.8%), visited the dentist only when they had a problem. Among 227 respondents, 139 (61.23%) cleaned their teeth once daily, and 88 (38.77%) cleaned their teeth twice or more daily. Most of the respondents, 160 (70.5%), used toothbrushes and toothpaste for cleaning their teeth, whereas 67 (29.5%) used other materials (tooth powder, charcoal, miswak, or chew stick) (Table- 2).

Table 2: Oral hygiene practices of the respondents

Practices	f (%)
Regularity of tooth cleaning	
Yes	160 (70.48)
No	67 (29.52)
Frequency of tooth cleaning	
Once daily	139 (61.23)
Twice or more daily	88 (38.77)
Materials used for tooth cleaning	
Toothbrush and toothpaste	160 (70.5)
Others	67 (29.5)
Use of interdental aids	
Yes	5 (2.2)
No	222 (97.8)
Tongue cleaning	
Yes	27 (11.9)
No	200 (88.1)
Rinsing the mouth after a meal	
Yes	59 (26)
No	168 (74)
Use of mouthwash	
Yes	8 (3.5)
No	219 (96.5)
Visit to dentist	
Only in problem	222 (97.8)

Oral Hygiene status, Periodontal Health status, Periodontitis of Respondents of respondents

Figure 1A illustrates the distribution of respondents based on oral hygiene status. Out of 227 participants, 123 (54%) exhibited poor oral hygiene, 69 (30%) demonstrated fair hygiene, and the remaining 35 (16%) had good oral hygiene.

Figure 1B shows the distribution of respondents by periodontal health status. Of the total, 31 respondents (13.66%) were free from periodontal disease, 17 (7.49%) exhibited bleeding on probing, 53 (23.35%) presented with calculus detectable by probing, 62 (27.31%) had pathological pockets of 4–5 mm, and 64 (28.19%) exhibited pockets measuring 6 mm or more.

Figure 1C depicts the presence of periodontitis among respondents. Out of the 227 individuals, 126 (55.5%) were diagnosed with periodontitis, while 101 (44.5%) did not present with the condition.






Figure 1. A. Distribution of respondents by oral hygiene status; B. Distribution of respondents by periodontal health status; C. Distribution of respondents by presence of periodontitis.

Association of respondent's socio-demographic characteristics with periodontitis

Table 3 shows the distribution of periodontitis concerning socio-demographic characteristics. Age group, sex, educational status, occupation, and monthly family income were significantly associated with periodontitis.

Variable	Periodontitis f (%)	Non-periodontitis f (%)	P value
Age group			
46-55 years	50 (45.50)	60 (54.50)	
56-65 years	52 (58.40)	37 (41.60)	0.001
66 years and above	24 (85.7)	4 (14.3)	
Gender			
Male	69 (62.20)	42 (37.80)	0.048
Female	57 (49.10)	59 (50.90)	
Educational Status			
No formal education	45 (84.90)	8 (15.10)	
Primary	31 (67.40)	15 (32.60)	0.0001
Secondary	26 (41.30)	37 (58.70)	
Higher Secondary and above	24 (36.9)	41 (63.1)	
Occupation			
Service holder	17 (44.70)	21 (55.30)	
Businessman	13 (54.20)	11 (45.80)	0.019
Laborer	15 (93.80)	1 (6.30)	
Homemaker	57 (52.80)	51 (47.20)	
Unemployed/ Retired	24 (58.6)	17 (41.5)	

Monthly family income (in taka)			
5000-24000	95 (65.10)	51 (34.90)	
24001-44000	16 (42.10)	22 (57.90)	0.0001
44001 and above	15 (34.9)	28 (65.1)	

Table 3: Distribution of periodontitis concerning socio-demographic characteristics

Association of respondent's oral hygiene practices, status with periodontitis

Table 4 shows the distribution of periodontitis regarding oral hygiene practices and status. Regularity, frequency, materials used for tooth cleaning, use of interdental aids, tongue cleaning, rinsing mouth after meal, use of mouthwash, and visits to the dentist were significantly associated with periodontitis. Regarding oral hygiene status, it was found that 100% (35) of respondents with good oral hygiene status had not periodontitis, and 97.60% (120) of respondents with poor oral hygiene status had periodontitis. The association was found statistically significant ($p < 0.05$).

Table 4: Association between oral hygiene practices, status, and periodontitis

Variable	Periodontitis f (%)	Non-periodontitis f (%)	P value
Regularity of tooth cleaning			
Yes	65 (40.60)	95 (59.40)	0.0001
No	61 (91)	6 (9)	
Frequency of tooth cleaning			
Once	98 (70.50)	41 (29.50)	0.0001
Twice or more daily	28 (31.80)	60 (68.20)	
Materials used for tooth cleaning			
Toothbrush and toothpaste	72 (45.00)	88 (55.00)	0.0001
Others	54 (80.6)	13 (19.4)	
Use of interdental aids			
Yes	0 (0)	5 (100)	0.016
No	126 (56.80)	96 (43.20)	
Tongue Cleaning			
Yes	7 (25.90)	20 (74.10)	0.001
No	119 (59.50)	81 (40.50)	
Rinsing the mouth after meal			
Yes	17 (28.80)	42 (71.20)	0.0001
No	109 (64.90)	59 (35.10)	
Use of mouthwash			
Yes	0 (0.00)	8 (100.00)	0.001

No	126 (57.50)	93 (42.50)	
Visit to dentist			
Only in problems	126 (56.8)	96 (43.2)	0.012
Once or twice a year	0 (0.00)	5 (100.00)	
Oral hygiene status			
Good	0 (0)	35 (100)	0.0001
Fair	6 (8.7)	63 (91.3)	
Poor	120 (97.60)	3 (2.40)	

Associated factors of periodontitis

Socio-demographic characteristics associated with periodontitis were identified by unadjusted odds ratio and adjusted odds ratio with 95% confidence intervals after adjusting for several important covariates (Table 5). Older adults aged 66 years and above were 7.2 times more likely to have periodontitis than the 46-55 years age group (OR_{un-adj} : 7.2, 95% CI: 2.342-22.135, $p < 0.001$; OR_{adj} : 12.637, 95% CI: 3.371-47.371, $p < 0.0001$). Male older adults were more likely to develop periodontitis than female older adults (OR_{un-adj} : 1.7, 95% CI: 1.002-2.886, $p < 0.049$; OR_{adj} : 3.309, 95% CI: .859-12.747, $p < 0.082$). Education had a significant impact on periodontitis. Older adults having higher secondary or above education had less chance of developing periodontitis (OR_{un-adj} : .104, 95% CI: .042-.257, $p < 0.0001$; OR_{adj} : .100, 95% CI: .032-.307, $p < 0.0001$). Regarding the impact of occupation on periodontitis, laborers were more likely to have periodontitis than the service holders (OR_{un-adj} : 18.529, 95% CI: 2.218-154.810, $p < 0.007$; OR_{adj} : 5.855, 95% CI: .594-57.75, $p < 0.130$). Monthly family income also significantly impacted periodontitis, where older people had higher incomes (Income. \geq 44001 BDT) family had a lower chance of developing periodontitis (OR_{un-adj} : .288, 95% CI: .141-.587, $p < 0.001$; OR_{adj} : .515, 95% CI: .216-1.227, $p < 0.134$) (Table 5).

Table 5: Multiple logistic regression analysis of the association between periodontitis and socio-demographic characteristics

Characteristics	Unadjusted OR	CI	P-value	Adjusted OR	CI	P-value
Age						
46-55	Ref			Ref		
56-65	1.686	0.96-2.97	0.070	1.73	0.85-3.53	0.13
66 and above	7.200	2.34-22.14	0.001	12.64	3.37-47.37	0.0001
Gender						
Female	Ref			Ref		
Male	1.701	1.00-2.89	0.049	3.31	0.86- 12.75	0.082
Education						
No formal education	Ref			Ref		
Primary	0.367	0.14-0.97	0.044	0.39	0.14-1.09	0.074
Secondary	0.125	0.05-0.31	0.0001	0.14	0.05-.38	0.000
Higher secondary and above	0.104	0.04-0.26	0.0001	0.10	0.03-.31	0.0001
Occupation						
Service holder	Ref			Ref		
Businessman	1.46	0.52-4.07	0.470	1.118	.332- 3.768	0.857
Laborer	18.53	2.22-154.81	0.007	5.855	.594-57.75	0.130
Homemaker	1.38	0.66-2.90	0.395	1.788	.453-7.055	0.407
Unemployed/ Retired	1.74	0.72-4.25	0.221	0.747	.233-2.397	0.624
Monthly family income						
5000-24000	Ref			Ref		
24001-44000	.390	0.19-0.81	0.011	0.502	0.20-1.25	0.140
44001 and above	.288	0.14-0.59	0.001	0.515	0.22-1.23	0.134

Regularity of tooth cleaning, frequency of tooth cleaning, the material used for tooth cleaning, and rinsing the mouth after the meal significantly impacted periodontitis (Table 6). Elderly who did not clean their teeth regularly were roughly 15 times more likely to develop periodontitis (OR_{un-}

adj: 14.85, 95% CI: 6.066- 36.400, $p < 0.0001$; OR_{adj}: 9.474, 95% CI: 3.636- 24.689, $p < 0.0001$). Similarly, respondents who did not use toothpaste and toothbrush were 5.077 times more likely to develop periodontitis (OR_{un-adj}: 5.077, 95% CI: 2.570- 10.030, $p < 0.0001$; OR_{adj}: 3.792, 95% CI: 1.758- 8.178, $p < 0.001$). Older adults who cleaned their teeth twice or more daily were less likely to develop periodontitis (OR_{un-adj}: .195, 95% CI: .000- .348, $p < 0.0001$; OR_{adj}: .403, 95% CI: .216- .856, $p < 0.016$). Rinsing the mouth after meals also had a significant impact on periodontitis, where older people who did not rinse their mouth after meal had 4.56 times more chance of developing periodontitis (OR_{un-adj}: 4.564, 95% CI: 2.391- 8.711, $p < 0.002$; OR_{adj}: 3.14, 95% CI: 1.272- 7.755, $p < 0.013$) (Table 6).

Table 6: Multivariate analysis of the association between periodontitis and Oral Hygiene practices

Characteristics	Unadjusted OR	CI	P-value	Adjusted OR	CI	P-value
Regularity of tooth cleaning						
Yes	Ref			Ref		
No	14.86	6.07- 36.40	0.0001	9.47	3.64- 24.69	0.0001
Frequency of tooth cleaning						
Once daily	Ref			Ref		
Twice or more daily	.19	0.00- 0.35	0.0001	0.40	0.22-0.86	0.016
Materials used for tooth cleaning						
Toothbrush and toothpaste	Ref			Ref		
Others	5.08	2.57- 10.03	0.0001	3.79	1.76- 8.18	0.001
Tongue cleaning						
Yes	Ref			Ref		
No	4.19	1.69- 10.38	0.002	0.83	0.23- 3.08	0.637
Rinsing the mouth after meal						
Yes	Ref			Ref		
No	4.56	2.39- 8.71	0.0001	3.14	1.27-7.75	0.013

Discussion

In developing countries like Bangladesh, where the elderly population is increasing significantly, dental health is especially important but frequently overlooked in terms of an individual's overall health and well-being. People are more prone to oral health issues as they become older, particularly periodontitis, which can have a serious effect on their general health and quality of life. This study highlighted the lack of oral hygiene practices, increased poor oral hygiene status and the association of periodontitis with several factors.

The present study was conducted among 227 elderly patients attending the dental outpatient department of three hospitals in Dhaka city. Most of the respondents, 110 (48.5%), were aged between 46 and 55 years, which is inconsistent with the study among the older adults in Dhaka, Bangladesh, reporting that among the elderly patients, the highest proportion of the respondents (49.10%) was in the age group of 60-69 years (1). Regarding educational qualification, 63 (27.8%) respondents had completed a secondary educational level, which is the highest frequency, and 53 (23.3%) had no formal education. This result is incompatible with the study among older adults in Bangladesh, which reported that most respondents had no schooling [27]. In this study, 108 (47.6%) of the respondents were homemakers, which is comparable with another study where most were homemakers [28]. According to our study findings, the highest proportion, 146 (64.32%) of respondents, had a monthly family income within 5000 – 24000 Taka, which is in line with the study conducted in the Narayanganj district of Bangladesh, reported the monthly family income of most of the respondents ranged from 4000 to 20000 taka per month [27].

The risk of developing periodontitis was higher in older adults who did not regularly brush their teeth. Brushing teeth once daily is enough to keep your mouth healthy and prevent periodontal and dental cavities [29]. Although the majority of the respondents, 160 (70.48%), cleaned their teeth regularly, the percentage of respondents brushing their teeth twice or more daily was 38.77%, which is similar to a previous study where 151 (90.41%) cleaned their teeth regularly, and 41.91% cleaned their teeth twice daily [27].

Another study conducted in New Delhi, India, revealed that 40 (16.1%) elderly participants brushed twice or more daily, less than the current study's findings [29]. Notably, 160 (70.5%) respondents used toothbrushes and toothpaste for cleaning teeth. This finding agrees with the previous study, where 79.04% of elderly respondents used toothbrushes for tooth cleaning [27].

The most widely used oral hygiene technique worldwide is tooth brushing, while regular brushing seems less prevalent among older people [30]. Similarly, the results of this study indicate that the risk of developing periodontitis was 5.077 times higher for those who did not use a toothbrush and toothpaste. Chahar *et.al* found that 48.2% of respondents used toothbrushes and toothpaste for tooth cleaning, which is less as compared with the current study [30]. The present study revealed that only 5 (2.2%) used interdental aids such as dental floss and interdental brushes. Likewise, the use of dental floss as an interdental cleaning device was almost negligible in the survey conducted in India [31]. Among the 227 respondents, 27 (11.9%) cleaned their tongue, 200 (88.1%) did not clean their tongue, and 59 (26%) respondents rinsed their mouth after meals, whereas the rest of them did not. These findings are in accordance with the study stating that oral hygiene- awareness and practice among patients in Jodhpur, India, found that only 20% of participants cleaned their tongue either with a toothbrush or tongue cleaner, and only 29% of the sample population rinsed their mouth after eating food [32]. According to this study, older adults who did not rinse their mouths after meals had a 4.56 times increased risk of acquiring periodontitis. Rinsing the mouth after meals also substantially affected the development of periodontitis. Again, most of the respondents, 219(96.5%), did not use mouthwash, which is comparable with a cross-sectional study among older Mexican adults, where 16.5 % of the study participants used mouthwash [33]. The current study also showed that the majority of the respondents, 222 (97.8%), visited the dentist only when they had a problem, consistent with the previous study stating that 54% of the study population visited the dentist only in pain [33].

More than half of the respondents, 123 (54%), had poor oral hygiene status, 69 (30%) had fair, and the rest, 35 (16%), had good oral hygiene status. A cross-sectional study conducted among rural dwellers found that one-third (33.8 %) of the participants had poor oral hygiene status, 57.2% had fair, and only 9% had good oral hygiene status [34]. In comparison, the current study had more participants with poor oral hygiene status. In the study from India [35], among 480 workers, 332(81.4%) respondents had poor oral hygiene status, and only 10 (2.5%) had good oral hygiene status, similar to the present study, though the study population is different. In the current study, it was observed that among 227 respondents, 31 (13.66%) had no periodontal disease, followed by 17 (7.49%) respondents who had bleeding on probing, 53 (23.35%) had calculus with plaque felt by probing, 62 (27.31%) had pathological pocket 4-5mm, 64 (28.19%) had pathological pocket

6mm or more. This finding is comparable with the study among the geriatric population in Delhi, India, which revealed that 25.4 % of older people had healthy periodontium, 71.1% had a periodontal pocket of 6 mm or more, and 2.40% had a pocket depth of 4-5 mm [36].

About 126 (55.5%) were found to have periodontitis, while 101 (44.5%) hadn't. The study among patients visiting a dental hospital in Dhaka reported that 15% had periodontitis, which is discordant with the present study [37]. Assessment of the association between periodontitis and socio-demographic characteristics of respondents revealed that periodontitis is significantly associated with age group, sex, educational status, occupation, and monthly family income. A previous study [38] found that the prevalence of periodontitis was higher among older participants and those of lower educational attainment, which is in accordance with the current study's findings. The findings of this study showed that elderly adults with a higher secondary education or above had a lower risk of developing periodontitis. Another study [39] observed that periodontitis was significantly higher in males than females and the old elderly compared to the young elderly, which is consistent with the present study. In the current study, oral hygiene practices were significantly associated with periodontitis. Similarly, earlier research [35] found that periodontal disease was significantly associated with oral hygiene practice. The current study's result supports another study's findings [40], suggesting that a lack of regular oral hygiene care was associated with periodontal disease. In contrast, a study among healthcare professionals showed that oral hygiene practices were not associated with periodontitis [41]. This contrast may be due to differences in the study population. A significant association was noticed between oral hygiene status and periodontitis, which supports the finding of a previous study [40] that reported that oral hygiene was significantly associated with periodontal disease. Subjects with the poorest periodontal health also displayed the poorest oral hygiene. Another study [35] found a significant association between periodontitis and oral hygiene status. We must prioritize the core public health functions of evaluation, policy formulation, and assurance. This involves redefining the impact of periodontal disease, assessing the effectiveness of preventive and control measures, and highlighting periodontal disease as a major public health concern[42].

The study provided information about oral hygiene practice, status, and periodontal status of elderly patients attending dental outpatient departments. The findings indicate the need to improve the oral health of older adults through preventive and curative public health measures. Further

research with a larger sample size could be done to generalize the results and plan for necessary public health interventions accordingly.

Conclusion

The findings of the study provide insight into the oral hygiene practice, status, and periodontal health status of elderly patients. A significant association existed between periodontitis, age, gender, educational status, occupation, and monthly family income. Oral hygiene practices and oral hygiene status were found to be significantly associated with periodontitis. These observations underscore the urgent need for targeted interventions to improve oral health education, promote proper oral hygiene practices and awareness regarding maintaining good oral hygiene and regular dental checkups, and enhance access to dental care services for elderly individuals in Bangladesh. Addressing these factors makes mitigating the burden of periodontitis and associated oral health complications possible, thereby enhancing the ageing population's overall quality of life and well-being.

Acknowledgement

We acknowledge the Department of Biostatistics and the Department of Health Education for their technical support during the study. We are also grateful to all participants included in this study.

Ethical Consent and Permission for data collection

This study was approved by the institutional review board of the National Institute of Preventive and Social Medicine (NIPSOM), Bangladesh (Ref No: NIPSOM/IRB/2017/09). The declaration of Helsinki (revised version) was followed to conduct the study. Both written and verbal consent was obtained before initiating the interview. A brief on the aims and objectives was given to the participants. Participants who agreed to give consent were finally included in the study.

Conflict of Interests

All authors declared no conflicts of interest.

Data availability statement

The datasets generated and analyzed during the current study are available from the

corresponding author on reasonable request to meshbah037@gmail.com.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References:

- 1 Rahman MM, Noman A Al, Islam ANMS, *et al.* Bangladesh striving against double burden: Dengue outbreak surges amid COVID-19 pandemic. *Journal of Global Health Economics and Policy*. 2021;1:e2021015. doi: 10.52872/001c.29657
- 2 Siddik MAB, Munmun MS, Hasan N, *et al.* Physio-Psychosocial Risk of Depression among College-Going Adolescents: A Cross-Sectional Study in Bangladesh. *J Affect Disord Rep*. 2024;16:100768. doi: <https://doi.org/10.1016/j.jadr.2024.100768>
- 3 Al Noman A, Das D, Nesa Z, *et al.* Importance of Wolbachia-mediated biocontrol to reduce dengue in Bangladesh and other dengue-endemic developing countries. *Biosaf Health*. Published Online First: 2023. doi: 10.1016/j.bsheal.2023.03.003
- 4 Mondal M, Islam MN, Ullah A, *et al.* Clinical manifestations and outcomes of COVID-19 in maintenance hemodialysis patients of a high infectious epidemic country: a prospective cross-sectional study. *Annals of Medicine and Surgery*. 2023;85. doi: 10.1097/MS9.0000000000001046
- 5 Noman AA, Joarder T, Islam MS, *et al.* Assessment of home health care seeking behaviors and socio-demographic impact on the recovery time of COVID-19 patients: A cross-sectional study in Bangladesh. *South East Asia Journal of Public Health*. 2021;11:1–13.
- 6 Das SK, Tamannur T, Nesa A, *et al.* Exploring the knowledge and practices on road safety measures among motorbikers in Dhaka, Bangladesh: a cross-sectional study. *Injury Prevention*. 2023;ip-2023-045071. doi: 10.1136/ip-2023-045071
- 7 Allen F, Tsakos G. Challenges in oral health research for older adults. *Gerodontology*. 2024;41:2–8.
- 8 Islam ANMS, Sultana H, Nazmul Hassan Refat Md, *et al.* The global burden of overweight-obesity and its association with economic status, benefiting from STEPs survey of WHO member states: A meta-analysis. *Prev Med Rep*. 2024;46:102882. doi: <https://doi.org/10.1016/j.pmedr.2024.102882>
- 9 Rahman M, Begum M, Uddin M, *et al.* Factors Affecting Health Status of Urban Aged Population: Evidence from Sylhet, Bangladesh. *Indian Journal of Gerontology*. 2018;32:103–18.
- 10 Rahman MM, Farhana Z, Chowdhury TT, *et al.* High Nutritional Vulnerability and Associated Risk Factors of Bangladeshi Wetland Community People Aged 50 Years and Older. *Journal of Ageing and Longevity*. 2022;2:102–12. doi: 10.3390/jal2020010
- 11 Chan AKY, Tsang YC, Jiang CM, *et al.* Integration of Oral Health into General Health Services for Older Adults. *Geriatrics (Switzerland)*. 2023;8. doi: 10.3390/geriatrics8010020

- 12 Petersen PE, Yamamoto T. Improving the oral health of older people: the approach of the WHO Global Oral Health Programme. *Community Dent Oral Epidemiol.* 2005;33:81–92. doi: 10.1111/j.1600-0528.2004.00219.x
- 13 Barua S, Bosu S, Haque MF, *et al.* Oral Health Vulnerability and Associated Risk Factors Among Tribal Elderly People in Bangladesh: A Primary Pilot Initiative Demanding a National Oral Health Survey. *Global Biosecurity.* 2024;6. doi: <https://doi.org/10.31646/gbio.232>
- 14 Tamannur T, Das SK, Nesa A, *et al.* Mothers' knowledge and practices towards oral hygiene of their children aged 5-9 years old: a cross-sectional study in Dhaka. *medRxiv.* 2024;2024.04.05.24305403. doi: 10.1101/2024.04.05.24305403
- 15 Sheiham A, Steele JG, Marcenes W, *et al.* Prevalence of impacts of dental. *Community Dent Oral Epidemiol.* 2001;29:195–203.
- 16 Li X, Kolltveit KM, Tronstad L, *et al.* Systemic Diseases Caused by Oral Infection. 2000.
- 17 Pereira WD. Knowledge, attitude and practice on oral hygiene measures among students in rural areas: Kanchipuram. *Pharma Innov.* 2017;6:382.
- 18 Madiba T, Bhayat A. Periodontal disease - risk factors and treatment options. *South African Dental Journal.* 2018;73. doi: 10.17159/2519-0105/2018/v73no9a5
- 19 Kinane DF, Stathopoulou PG, Papapanou PN. Periodontal diseases. *Nat Rev Dis Primers.* 2017;3.
- 20 Grant WB, van Amerongen BM, Boucher BJ. Periodontal Disease and Other Adverse Health Outcomes Share Risk Factors, including Dietary Factors and Vitamin D Status. *Nutrients.* 2023;15.
- 21 Syrjälä AMH, Ylöstalo P, Knuutila M. Periodontal condition of the elderly in Finland. *Acta Odontol Scand.* 2010;68:278–83. doi: 10.3109/00016357.2010.494619
- 22 Tefera A, Bekele B. Periodontal disease status and associated risk factors in patients attending a tertiary hospital in Northwest Ethiopia. *Clin Cosmet Investig Dent.* 2020;12:485–92. doi: 10.2147/CCIDE.S282727
- 23 Ahmad MS, Al Mamun MA, Islam MS, *et al.* Oral hygiene practice and oral health status of geriatric population in selected area of Rangpur, Bangladesh. *KYAMC Journal.* 2018;9:48–52.
- 24 Bokhari S, Suhail A, Malik A, *et al.* Periodontal disease status and associated risk factors in patients attending a Dental Teaching Hospital in Rawalpindi, Pakistan. *J Indian Soc Periodontol.* 2015;19:678–82. doi: 10.4103/0972-124X.156882
- 25 GREENE JC, VERMILLION JR. THE SIMPLIFIED ORAL HYGIENE INDEX. *J Am Dent Assoc.* 1964;68:7–13. doi: 10.14219/jada.archive.1964.0034
- 26 Benigeri M, Brodeur J-M, Payette M, *et al.* Community periodontal index of. *J Clin Periodontol.* 2000;27:308–12.

- 27 Eusuf Zai SZ, Jamayet N, Alam M. A Study of Teeth Status and Oral Health Related Quality of Life among Elderly in Bangladesh. *International Medical Journal (1994)*. 2013;20:610–4.
- 28 Mahmud SZ, Amin MS, Tarafder MA, *et al*. Measurement of Oral Health Literacy Level Among Bangladeshi Adults Seeking Dental Care and Its Relationship With Socio-demographic Characteristics. *Anwer Khan Modern Medical College Journal*. 2017;7:34–9. doi: 10.3329/akmmcj.v7i1.31611
- 29 Singh A, Purohit BM, Masih N. Geriatric oral health predicaments in New Delhi, India. *Geriatr Gerontol Int*. 2016;16:37–45. doi: 10.1111/ggi.12434
- 30 Chahar P, Mohanty VR, Aswini YB. Oral health-related quality of life among elderly patients visiting special clinics in public hospitals in Delhi, India: A cross-sectional study. *Indian J Public Health*. 2019;63:15–20. doi: 10.4103/ijph.IJPH_316_17
- 31 Goryawala SN, Chavda P, Udhani S, *et al*. A survey on oral hygiene methods practiced by patients attending Dentistry Department at a Tertiary Care Hospital from Central Gujarat. *J Int Soc Prev Community Dent*. 2016;6:115–9. doi: 10.4103/2231-0762.178750
- 32 Jain N, Mitra D, Ashok KP, *et al*. Oral hygiene awareness and practice among patients attending OPD at Vyas Dental College and Hospital, Jodhpur. *J Indian Soc Periodontol*. 2012;16:524–8. doi: 10.4103/0972-124X.106894
- 33 Islas-Granillo H, Casanova-Rosado JF, De La Rosa-Santillana R, *et al*. Self-reported oral hygiene practices with emphasis on frequency of tooth brushing: A cross-sectional study of Mexican older adults aged 60 years or above. *Medicine (United States)*. 2020;99:E21622. doi: 10.1097/MD.00000000000021622
- 34 Azodo CC, Amenaghawon OP. Oral hygiene status and practices among rural dwellers. *European J Gen Dent*. 2013;2:42–5. doi: 10.4103/2278-9626.106806
- 35 Baishya B, Satpathy A, Nayak R, *et al*. Oral hygiene status, oral hygiene practices and periodontal health of brick kiln workers of Odisha. *J Indian Soc Periodontol*. 2019;23:163–7. doi: 10.4103/jisp.jisp_383_18
- 36 Yadav NR, Jain M, Sharma A, *et al*. Periodontal health of the geriatric population in old-age homes of Delhi, India. *J Indian Soc Periodontol*. 2021;25:427–31. doi: 10.4103/jisp.jisp_193_20
- 37 Iqbal MA, Mohol J, Afrin F, *et al*. Prevalence of periodontal diseases among the patient visiting at Periodontology OPD Update Dental College Hospital, Dhaka. *Update Dental College Journal*. 2016;5:23–9. doi: 10.3329/updcj.v5i2.27265
- 38 Azodo C, Umoh A. Prevalence of gingivitis and periodontitis in an adult male population in Nigeria. *Nigerian Journal of Basic and Clinical Sciences*. 2012;9:65. doi: 10.4103/0331-8540.108465

- 39 Braimoh OB, Alade GO. Gingivitis and Periodontitis among the Elderly in Port Harcourt Nigeria: A Population-based Study. *Journal of Oral Health and Community Dentistry*. 2019;13:5–9. doi: 10.5005/jp-journals-10062-0038
- 40 Hopcraft MS, Morgan M V., Satur JG, *et al*. Oral hygiene and periodontal disease in Victorian nursing homes. *Gerodontology*. 2012;29. doi: 10.1111/j.1741-2358.2010.00448.x
- 41 Merchant A, Pitiphat W, Douglass CW, *et al*. Oral Hygiene Practices and Periodontitis in Health Care Professionals. 2002.
- 42 Janakiram C, Dye BA. A public health approach for prevention of periodontal disease. *Periodontol* 2000. 2020;84:202–14.