Depression and Prefrontal-Hippocampal Pathway

# 1 Depression and the Prefrontal-Hippocampal Pathway - A Pilot

# 2 Multimodal Neuroimaging Study in Transgender Women

- 3 Setthanan Jarukasemkit<sup>1,2,3</sup>, MD, Karen M. Tam<sup>1</sup>, Seksan Yoadsanit<sup>1</sup>, Ty Easley<sup>2</sup>, Hailey Modi<sup>2</sup>,
- 4 Lyn Stahl<sup>2</sup>, Adun Kampaengtip<sup>1</sup>, MSc, Thanissara Chansakul<sup>1</sup>, MD, Rena Janamnuaysook<sup>4,5</sup>,
- 5 MBA, Akarin Hiransuthikul, MD<sup>4,6</sup>, Chaipat Chunharas<sup>\*3</sup>, MD, PhD and Janine D.
- 6 Bijsterbosch<sup>\*2</sup>, PhD.
- 7 (1) Faculty of Medicine Ramathibodi Hospital, Mahidol University, Bangkok Thailand.
- 8 (2) Personomics Lab, Department of Radiology, Washington University School of Medicine, Saint Louis, Missouri, USA.
- 9 (3) Cognitive Clinical and Computational Neuroscience (CCCN) Center of Excellence, Chulalongkorn University, Bangkok, Thailand.
- 10 (4) Institute of HIV Research and Innovation (IHRI), Bangkok, Thailand.
- 11 (5) Center of Excellence in Transgender Health, Chulalongkorn University, Bangkok, Thailand.
- 12 (6) Department of Preventive and Social Medicine, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand.
- 13 \* joint last authors.
- 14 Co-correspondence:

#### 15 Chaipat Chunharas, MD, PhD

- 16 Department of Neurology, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand.
- 17 Address: 1873 Ratchadamri Rd., Pathum Wan, Bangkok 10330, Thailand
- 18 Phone: +66 89 777 9493, Email: chaipat.c@chula.ac.th

### 19 Janine D. Bijsterbosch, PhD

- 20 Department of Radiology, Washington University in St. Louis, Missouri, United States
- 21 Address: 4525 Scott Avenue, St. Louis, MO 63110, USA
- 22 Phone: +1 314 629 3344, Email: janine.bijsterbosch@wustl.edu
- 23 Keywords: Depression; Transgender women; Neuroimaging; Spectroscopy; fMRI
- Abstract word counts: 91; Main text word counts: 1827; Number of figures: 3; Number of tables:
- 25 0; Number of supplemental documents: 4.

Depression and Prefrontal-Hippocampal Pathway

## 26 Abstract

27 This pilot study explored associations between gamma-aminobutyric acid (GABA) spectroscopy 28 and functional magnetic resonance imaging in transgender women with depression. Twenty 29 participants completed mental health surveys and imaging between January and March 2024. 30 Depression severity was measured by Patient Health Questionnaire-9 (PHQ-9) scores. Higher 31 PHQ-9 scores were associated with lower GABA levels in the hippocampus and lower fractional 32 amplitude of low-frequency fluctuations (fALFF) in the dorsolateral prefrontal cortex. These 33 findings revealed interactions between neurotransmitter signaling and functional brain activity of 34 the hippocampal-prefrontal circuit in depression. Future large-scale neuroimaging studies are 35 needed.

Depression and Prefrontal-Hippocampal Pathway

### 36 Introduction

The transgender population experiences higher rates of depression compared to the cisgender population, with estimates in Thailand reaching around 22.7% for the transgender community compared to 3.2% in the overall population.<sup>1</sup> Despite this high prevalence, transgender individuals are not typically represented in neuroimaging research projects aimed at elucidating the brain basis of depression. More broadly, there is a problematic lack of diversity in neuroimaging research, which largely focuses on cisgender participants from Western, Educated, Industrialized, Rich, and Democratic (WEIRD) countries.<sup>2</sup>

44

In this pilot study, we focused on two key brain regions implicated in depression, namely the hippocampus and the dorsolateral prefrontal cortex (DLPFC). Self-referential processes such as rumination, a key symptom of depression, have been associated with hippocampal hyperactivity, and hippocampal atrophy is commonly observed in recurrent depression.<sup>3,4</sup> Furthermore, repetitive transcranial magnetic stimulation (TMS) applied to the left DLPFC has shown significant effectiveness in treating depression.<sup>5</sup> As such, the hippocampal-prefrontal circuit is of key interest in the context of depression.

52

53 A hierarchical model proposed by Hasler and Northoff has suggested that non-invasive 54 measurement of cerebral gamma-aminobutyric acid (GABA) concentrations may offer endophenotypic insights into inhibitory pathways relevant to depression.<sup>6</sup> A recent multimodal 55 56 neuroimaging study showed that hippocampal GABA was necessary to recruit DLPFC activation in a task context that required suppression of unwanted thoughts.<sup>7</sup> Despite the importance of the 57 58 prefrontal-hippocampal circuit in cognitive and emotional regulation, its specific relationship 59 with depression measurements remains poorly understood. To address these challenges with 60 limited transgender representation in neuroimaging depression research and to elucidate the role 61 of the prefrontal-hippocampal circuit in depression, the goal of this study was to investigate the 62 prefrontal-hippocampal pathway of depression in a non-WEIRD cohort of transgender women.

#### Depression and Prefrontal-Hippocampal Pathway

### 63 Material and Method

#### 64 *Patients and Data Collection*

65 Twenty transgender women with mild-to-severe depression were recruited from the Tangerine 66 Community Health Clinic, a specialized healthcare center for transgender people in Bangkok, Thailand.<sup>8</sup> Recruitment and data collection took place between January and March 2024. Mild-67 68 to-severe depression was defined based on Patient Health Questionnaire (PHQ-9) scores greater 69 than 5 and a neuroticism score on the Five-Factor Inventory (NEO-FFI) greater than 30.1 which was previously defined as one standard deviation above the population mean.<sup>9,10</sup> Exclusion 70 71 criteria comprised any history of neurological or cognitive disorders and contraindications for 72 magnetic resonance imaging (MRI) scanning. See Supplementary Table S1 for a detailed 73 overview of inclusion and exclusion criteria. Of the twenty subjects recruited, half were 74 currently participating in gender-affirming hormone therapy (GAHT).

75

This study was approved by the Institutional Review Board, Faculty of Medicine Ramathibodi Hospital, Mahidol University (COA. MURA2023/522). After obtaining informed consent, participants completed a series of mental health questionnaire assessments followed by a onehour MRI session at the Advanced Imaging Diagnostic Center, Faculty of Medicine Ramathibodi Hospital, Bangkok, Thailand. For the mental health assessments used in this manuscript, participants performed Thai language versions of the PHQ-9 and NEO-FFI self-report questionnaires on paper prior to the MRI scanning session.<sup>9,10</sup>

83

### 84 Neuroimaging Acquisition and Preprocessing

The MRI session included T1-weighted and T2-weighted structural MRI, resting-state functional MRI (rs-fMRI), and GABA magnetic resonance spectroscopy (MRS) imaging, using a Philips Ingenia Elition 3.0T scanner (software version R5.7.1). We leveraged a state-of the-art imaging protocol informed by the Human Connectome Project Lifespan and Disease protocols, and the MRS utilized the MEGA-PRESS sequence with parameters based on standard guidelines.<sup>11</sup> For more details, the neuroimaging acquisition parameters are listed in Supplementary Table S2.

#### Depression and Prefrontal-Hippocampal Pathway

91

92 Functional imaging data were preprocessed by FMRIB's Software Library (FSL) software. 93 Briefly, the preprocessing steps included motion correction, distortion correction, structural co-94 registration, spatial normalization to Montreal Neurological Institute space using nonlinear 95 registration, smoothing with a 5 millimeter kernel, and noise correction with independent 96 component analysis. The distortion correction of echo planar imaging (EPI) of rs-fMRI was 97 performed utilizing a technique that leverages two different phase encoding directions.<sup>12</sup> 98 Specifically, FSL's topup tool estimated distortion fields from opposite phase-encoded EPI 99 images, and the estimated fields were applied for geometric correction. Please see the 100 Supplementary Methods for further detail. Additional cleanup of rs-fMRI data was performed 101 using independent component analysis noise removal by FSL AROMA.

102

For MRS images, signals obtained from the left DLPFC, and left hippocampus were processed
with baseline initial peak subtraction with 90% Gaussian character for fitting the power
spectrum. MEGA Subtract All (ON-OFF) Spectra was performed. Please see the Supplementary
Methods for further detail.

107

#### 108 Data Analysis

109 The study utilized neuroimaging parameters focused on two regions of interest: the hippocampus 110 and the DLPFC. For the functional analyses, the bilateral hippocampus was defined based on the 111 Harvard/Oxford Subcortical atlas and the bilateral DLPFC was defined based on prior work that utilized a task fMRI localizer (Figure 1).<sup>13</sup> We extracted the mean time series of the left and right 112 113 hippocampus and DLPFC following preprocessing and estimated the fractional amplitude of 114 low-frequency fluctuation (fALFF). This measure captures the normalized strength of 115 spontaneous signal fluctuations in each brain region by calculating the ratio of the power in the 116 low-frequency range (0.01-0.1 Hertz) to the power across the full frequency range.<sup>14</sup> 117 Subsequently, the fALFF estimates were averaged between left and right hippocampus and 118 between left and right DLPFC before further analysis.

#### Depression and Prefrontal-Hippocampal Pathway

119

For the spectroscopy data, GABA concentrations were normalized to creatinine concentrations (GABA/Cr) using creatinine as a stable reference metabolite. Notably, hippocampal spectroscopy can be challenging due to the small size of the hippocampus, which can lead to low spectral resolution and a low signal-to-noise ratio.<sup>15</sup> To address this concern, we followed prior advice on the minimum voxel volume while also preserving spatial specificity for the regions of interest.<sup>11</sup> Furthermore, we controlled for motion-related noise by including the framewise displacement parameter from rs-fMRI as a confound in subsequent analyses.

127

128 Following the extraction of neuroimaging parameters, multiple univariate linear regressions were 129 performed using the ordinary least squares method. In these analyses, multimodal neuroimaging 130 measures served as the independent variables, PHQ-9 scores were included as the dependent 131 variables, and framewise displacement was included as a control variable. Mediation analyses 132 were also conducted to investigate whether DLPFC fALFF mediated the relationship between 133 hippocampal GABA and depression severity as measured with PHQ-9 scores. Additionally, an 134 unpaired t-test was performed to investigate differences in neuroimaging measures and PHQ-9 135 scores between the GAHT and non-GAHT groups using permutation testing. Equations for all 136 regression and mediation models can be found in the Supplementary Material.

137

#### 138 Results

139 The study consisted of 20 transgender women with mean age (standard deviation) of  $30.1 (\pm 6.5)$ 140 years. The mean neuroticism score (NEO-FFI) was 45.55 ( $\pm$  7.5), and the mean PHQ-9 score was 141 14.6 ( $\pm$  5.73), indicating the mental health status of the sample. Nearly half of the participants 142 (45%) were undergoing gender-affirming hormone therapy at the time of the study. The linear 143 regression models revealed a significant negative association between hippocampal GABA and 144 PHQ-9 scores ( $\beta = -0.48$ , p = 0.039; Figure 2A), such that higher depression severity was 145 associated with decreased hippocampal GABA. Furthermore, our findings revealed a significant 146 negative association between DLPFC fALFF and PHQ-9 scores ( $\beta = -0.54$ , p = 0.015; Figure 147 2B), such that higher depression severity was associated with decreased fALFF in the DLPFC.

Depression and Prefrontal-Hippocampal Pathway

148 No significant associations were observed between PHQ-9 scores and either hippocampal fALFF149 or DLPFC GABA.

150

Figure 3 illustrates the results from the mediation analysis model. We observed a trend-level mediating effect of DLPFC fALFF on the relationship between hippocampal GABA and PHQ-9 scores. Specifically, the results showed a significant relationship between hippocampal GABA and DLPFC fALFF ( $\beta = 0.48$ , p = 0.037) with a trend-level mediation effect ( $\beta = -0.20$ , p = 0.060).

156

We did not observe a significant difference between GAHT groups on PHQ-9 scores or on any of the neuroimaging measurements. Although prior work has shown a decrease in mental health symptomatology following GAHT, the lack of group differences in our study is explained by our inclusion criteria which required the presence of depressive symptoms to enable the investigation of the neural correlations of depression.<sup>16</sup>

Depression and Prefrontal-Hippocampal Pathway

### 162 Discussion

163 This pilot study aimed to investigate the multimodal prefrontal-hippocampal circuit in relation to 164 depression in a transgender cohort. Our findings revealed negative associations between 165 depression severity and hippocampal GABA, and negative associations between depression 166 severity and the amplitude of functional fluctuations in the DLPFC. These findings on the fronto-167 hippocampal relationship in depression are consistent with previous research in cisgender 168 subjects, where higher hippocampal GABA was shown to recruit the DLPFC in an unwanted 169 thought suppression task, indicating that DLPFC top-down signaling may depend on 170 hippocampal GABA.<sup>7</sup> Moreover, lower hippocampal GABA may be indicative of impaired inhibitory signaling, which could contribute to hippocampal hyperactivity.<sup>3,17</sup> 171

172

173 Hypoactivity of the DLPFC has previously been observed in major depressive disorder patients and in peripartum and post-stroke depression.<sup>18,19,20</sup> Given the DLPFC's pivotal role in cognitive 174 175 control and emotion regulation, prior work has suggested that left DLPFC fALFF could serve as a biomarker to predict treatment responses to escitalopram and TMS intervention.<sup>21,22</sup> Although 176 177 GABA depletion is commonly observed in depression, GABA also plays critical roles in 178 regulating neurochemical processes, including the hypothalamus-pituitary-adrenal axis and neurosteroid homeostasis.<sup>23</sup> Disruption of GABA signaling in postpartum depression and 179 premenstrual dysphoria may suggest abnormal neurochemical responses to hormonal 180 fluctuations, particularly those driven by estradiol.<sup>24,25</sup> As such, this link between estradiol and 181 182 GABA may be of interest for future research on interactions between GAHT and depression in 183 the transgender community.

184

185 Additionally, our study supports the Hasler & Northoff depression model, which links 186 neurotransmitter imbalances. particularly in GABAergic signaling, to functional dysconnectivity.<sup>6</sup> However, further research is required to explore all potential pathways within 187 188 this model, including other circuitry like limbic and salience networks and other relevant 189 neurotransmitters such as serotonin and dopamine. The findings of this pilot study provide

Depression and Prefrontal-Hippocampal Pathway

preliminary validation for the potential use of GABA MRS in tracking symptom severity in thisunderrepresented population.

192

Although we were underpowered to detect differences linked to GAHT, GAHT has been shown to reduce the risk of depression in the transgender population and has previously been associated with lower odds of both depression and suicidality.<sup>16</sup> These prior GAHT findings underscore the potential mediating role of hormone therapy in mitigating depression risk in the transgender population. Large-scale neuroimaging studies in transgender cohorts will be crucial to further understand the relationship between GAHT (and sex hormones more broadly) and neural circuitry dysfunction in depression.

200

#### 201 *Limitation*

The pilot study was limited in statistical power due to the relatively small sample size. As a result, we performed focused analyses on two brain regions (the hippocampus and DLPFC). Future large-scale multimodal neuroimaging studies that are sufficiently powered to interrogate a larger number of relationships in a broader neural circuitry are essential to address the complexities inherent in hierarchical models of depression.

207

#### 208 Conclusion

We investigated the prefrontal-hippocampal circuit in relation to depression in a transgender cohort using multimodal neuroimaging data. Our results revealed that higher depression severity was associated with reduced hippocampal GABA and with reduced DLPFC fALFF. Larger scale neuroimaging studies are needed to further study the brain basis of depression in the transgender community and the potential impact of GAHT on depression-related neural pathways.

Depression and Prefrontal-Hippocampal Pathway

# 214 Author's Contributions

- 215 The authors made the following contributions to this manuscript:
- 216 Conceptualization: SJ, TE, RJ, AH, CC, JDB;
- 217 Data curation: SJ, AK, TC;
- 218 Funding acquisition: SJ, TC;
- 219 Investigation: SJ, KMT, SY, TC;
- 220 Project administration: SJ;
- 221 Methodology: SJ, CC, JDB;
- 222 Validation: CC, JDB;
- 223 Software: CC, JDB;
- 224 Writing original draft: SJ, CC, JDB;
- 225 Writing review and editing: SJ, TE, HM, LS, RJ, AH, CC, JDB.
- 226

#### 227 Acknowledgements

We would like to express our gratitude to the Center for High Performance Computing and the Research Computing and Informatics Facility at Washington University in St. Louis for providing computational resources and technical support. We are also grateful to the Prince Mahidol Award Youth Program, Prince Mahidol Award Foundation under the Royal Patronage, Thailand for the financial support that enabled Dr. Jarukasemkit to conduct research at Washington University in St. Louis.

Depression and Prefrontal-Hippocampal Pathway

# 234 Author Disclosure Statement

235 The authors declare no conflict of interest

236

# 237 Funding Information

- 238 This work was supported by Advance Diagnostic Imaging Center Fund (No. 32080009), and
- 239 Faculty of Medicine Ramathibodi Hospital (No. RF\_66138), Mahidol University, Thailand.
- 240 Setthanan Jarukasemkit is supported by the Prince Mahidol Award Youth Program, Prince
- 241 Mahidol Award Foundation under the Royal Patronage.

Depression and Prefrontal-Hippocampal Pathway

## 242 References

- 243 1. Nimitpanya P, Wainipitapong S, Wiwattarangkul T, et al. Evaluation of mental health and
  244 satisfaction following transfeminine gender-affirming surgery in Thailand. Transgend Health
  245 2022;7(1):61-7.
- 246 2. Ricard JA, Parker TC, Dhamala E, et al. Confronting racially exclusionary practices in the
- acquisition and analyses of neuroimaging data. Nat Neurosci 2023;26(1):4-11.
- 248 3. Sheline YI, Barch DM, Price JL, et al. The default mode network and self-referential processes
- in depression. Proc Natl Acad Sci U S A 2009;106(6):1942-7.
- 250 4. Xu X, Zhou Q, Wen F, et al. Meta Analysis of Brain Volumetric Abnormalities in Patients
- with Remitted Major Depressive Disorder. Depress Anxiety 2024;2024(1):6633510.
- 252 5. Kishi T, Ikuta T, Sakuma K, et al. Theta burst stimulation for depression: a systematic review
  253 and network and pairwise meta-analysis. Mol Psychiatry 2024;1-7.
- 254 6. Hasler G, Northoff G. Discovering imaging endophenotypes for major depression. Mol
  255 Psychiatry 2011;16(6):604-19.
- 256 7. Schmitz TW, Correia MM, Ferreira CS, et al. Hippocampal GABA enables inhibitory control
  257 over unwanted thoughts. Nat Commun 2017;8(1):1311.
- 8. Barmania S. Tangerine Clinic: championing transgender health care. Lancet HIV
  2023;10(11):e701.
- 260 9. Lotrakul M, Sumrithe S, Saipanish R. Reliability and validity of the Thai version of the PHQ261 9. BMC Psychiatry 2008;8:1-7.

Depression and Prefrontal-Hippocampal Pathway

- 262 10. Loavattanaving K. A comparative study of personality type (five factor model), emotional
- 263 quotient (Goleman), and work satisfaction of hotel sales person in Bangkok [master's thesis].
- **264** Bangkok:ThammasatUniversity;2004.Availableform:
- 265 https://digital.library.tu.ac.th/tu\_dc/frontend/Info/item/dc:115260 [Last Accessed: 9/16/2024].
- 266 11. Peek AL, Rebbeck TJ, Leaver AM, et al. A comprehensive guide to MEGA-PRESS for
- GABA measurement. Anal Biochem 2023;669:115113.
- 268 12. Schallmo MP, Weldon KB, Burton PC, et al. Assessing methods for geometric distortion
- compensation in 7 T gradient echo functional MRI data. Hum Brain Mapp 2021;42(13):4205-23.
- 270 13. Bijsterbosch J, Smith S, Forster S, et al. Resting state correlates of subdimensions of anxious
- affect. J Cogn Neurosci 2014;26(4):914-26.
- 272 14. Zuo XN, Di Martino A, Kelly C, et al. The oscillating brain: complex and reliable.
  273 Neuroimage 2010;49(2):1432-45.
- 274 15. Allaïli N, Valabrègue R, Auerbach EJ, et al. Single□voxel 1H spectroscopy in the human
  275 hippocampus at 3 T using the LASER sequence: characterization of neurochemical profile and
  276 reproducibility. NMR Biomed. 2015 Oct;28(10):1209-17.
- 277 16. Aldridge Z, Patel S, Guo B, et al. Long term effect of gender affirming hormone treatment
  278 on depression and anxiety symptoms in transgender people: A prospective cohort study.
  279 Andrology 2021;9(6):1808-16.
- 280 17. Campbell S, MacQueen G. The role of the hippocampus in the pathophysiology of major
  281 depression. J Psychiatry Neurosci 2004;29(6):417-26.

Depression and Prefrontal-Hippocampal Pathway

- 282 18. Dong D, Ming Q, Zhong X, et al. State-independent alterations of intrinsic brain network in
- current and remitted depression. Prog Neuropsychopharmacol Biol Psychiatry 2019;89:475-80.
- 284 19. Che K, Mao N, Li Y, et al. Altered spontaneous neural activity in peripartum depression: a
- resting-state functional magnetic resonance imaging study. Front Psychol 2020;11:656.
- 286 20. Egorova N, Veldsman M, Cumming T, et al. Fractional amplitude of low-frequency
- fluctuations (fALFF) in post-stroke depression. NeuroImage Clin 2017;16:116-24.
- 288 21. Kamishikiryo T, Okada G, Itai E, et al. Left DLPFC activity is associated with plasma
  289 kynurenine levels and can predict treatment response to escitalopram in major depressive
- disorder. Psychiatry Clin Neurosci 2022;76(8):367-76.
- 22. Liu J, Shu Y, Wu G, et al. A neuroimaging study of brain activity alterations in treatmentresistant depression after a dual target accelerated transcranial magnetic stimulation. Front
  Psychiatry 2024;14:1321660
- 23. Schür RR, Draisma LW, Wijnen JP, et al. Brain GABA levels across psychiatric disorders: A
  systematic literature review and meta□analysis of 1H□MRS studies. Hum Brain Mapp
  2016;37(9):3337-52.
- 297 24. Schiller CE, Meltzer-Brody S, Rubinow DR. The role of reproductive hormones in298 postpartum depression. CNS Spectr 2015;20(1):48-59.
- 25. Hantsoo L, Payne JL. Towards understanding the biology of premenstrual dysphoric
  disorder: From genes to GABA. Neurosci Biobehav Rev 2023;149:105168.
- 301

302

Depression and Prefrontal-Hippocampal Pathway

- **303** Figure 1: Regions of interest
- 304 Legend: Regions of interest included the hippocampus (green) and DLPFC (red). L indicates the
- 305 left hemisphere, and R indicates the right hemisphere.

306

307 Figure 2: Associations between neuroimaging measures and depression severity

Legend: Scatter plots illustrate the relationship of patient health questionnaire (PHQ-9) scores
with gamma-aminobutyric acid (GABA) in the hippocampal (A) and fractional amplitude of low
frequency fluctuation (fALFF) in the dorsolateral prefrontal cortex (DLPFC) (B) after
controlling for the confounder. 95% Confidence intervals and marginal histograms are also

312 displayed. GABA/Cr indicates GABA concentrations normalized to the creatinine reference.

313

**314** Figure 3: Path diagram of mediation analysis

315 Legend: The right diagram displays separate linear models, while the left diagram illustrates the 316 mediation analysis model. A) Direct associations between gamma-aminobutyric acid (GABA) in 317 the hippocampus and patient health questionnaire (PHQ-9) scores, and between fractional 318 amplitude of low frequency fluctuations (fALFF) in the dorsolateral prefrontal cortex (DLPFC) 319 and PHQ-9 were estimated. B) The mediation analysis revealed a trend-level mediation effect of 320 DLPFC fALFF on the relationship between hippocampal GABA and PHQ-9. The arrows 321 indicate regression coefficients, and asterisks denote significance association (p < 0.05). L 322 indicates the left hemisphere, and R indicates the right hemisphere.







Hippocampal GABA/Cr residuals

![](_page_16_Picture_4.jpeg)

![](_page_16_Figure_6.jpeg)

![](_page_16_Figure_8.jpeg)

(A)

![](_page_17_Picture_1.jpeg)

Hippocampus GABA

![](_page_17_Picture_3.jpeg)

DLPFC fALFF

![](_page_17_Figure_5.jpeg)

![](_page_17_Figure_6.jpeg)

![](_page_17_Picture_7.jpeg)