1 2 3	Intestinal microbiota changes due to <i>Giardia intestinalis</i> infections in a longitudinal Ecuadorian birth cohort and impact on cobalamin biosynthesis.
4 5 6 7	Ashish Damania <sup>1</sup> , Andrea Arévalo-Cortés <sup>2, 3</sup> , Andrea Lopez <sup>2</sup> , Victor Seco-Hidalgo <sup>4</sup> , Diana Garcia-Ramon <sup>2</sup> , Emilie Lefoulon <sup>5</sup> , Courtney Long <sup>5</sup> , Evan Drake <sup>5</sup> , Barton Slatko <sup>5, 6</sup> , Philip J Cooper <sup>2, 4</sup> , Rojelio Mejia <sup>*2, 6</sup>
8 9 10 11 12 13 14 15	<ol> <li>Platform for Innovative Microbiome and Translational Research, Department of Genomic Medicine, The University of Texas MD Anderson Cancer Center, Houston, Texas.</li> <li>School of Medicine, Universidad Internacional del Ecuador, Quito, Ecuador.</li> <li>Faculty of Human Health, Universidad Nacional de Loja, Loja, Ecuador.</li> <li>Institute of Infection and Immunity, St George's University of London, London, UK.</li> <li>Genome Biology Division, New England Biolabs, Inc, Ipswich, MA, USA.</li> <li>Department of Pediatrics – Tropical Medicine, Baylor College of Medicine, Houston, Texas.</li> </ol>
16	"Corresponding author: Rojello Mejia, mejia@bcm.edu
18 19	Keywords: microhiome, cobalamin, vitamin B12, <i>Giardia intestinalis</i>
20	
21	
22	
24	
25	
26	
27	
28	
30	
31	
32	
33	
34	
33 36	
37	
38	
39	
40	
41	
42	
44	
45	
46	
47	
48 40	
<del>1</del> 2 50	
51	

### 52 Abstract

53	Background: Giardia intestinalis is a protozoal parasite infecting the gastrointestinal tract
54	worldwide. Chronic infections/reinfections are common, with adverse nutritional consequences
55	for critical growth during the first five years of life. Vitamin B12 (cobalamin) is absorbed primarily
56	in the ileum of the small intestine, where Giardia trophozoites attach and replicate. Bacteria
57	activate bioavailable vitamin B12, which is essential for human DNA synthesis and
58	development. A disturbance in cobalamin biosynthesis caused by giardiasis may contribute to
59	impairment of childhood development and growth.
60	Methodology/Principal Findings: We performed a longitudinal analysis on 61 Ecuadorian
61	children using multi-parallel real-time quantitative PCR and whole genome sequencing.
62	Children had increased <i>Giardia</i> DNA burden from ages 3 to 5 (p = 0.0012) and 7.58 times more
63	frequent <i>Giardia</i> infections (1.31 to 34.33, P = 0.0176). There was an increased alpha diversity/
64	Giardia fg/µl in three-year-olds compared to age-matched non-infected (30.20 vs 4.37, p =
65	0.050), but decreased alpha diversity/ Giardia fg/µl in five-year-olds compared to age-matched
66	non-infected children (0.21 vs 4.31, p = 0.021). Alpha diversity/ Giardia fg/µl was also
67	decreased in samples collected longitudinally from the same children at five compared to 3
68	years (p = 0.031). Cobalt transport protein ( <i>Cbi</i> N) (FDR < 0.003) and IPR011822 (Cobalamin-
69	dependent methionine synthase) sequences were decreased in infected children (p < 0.002)
70	and among those with greatest <i>Giardia</i> burdens (p < 0.001).
71	Conclusion/Significance: Giardia intestinalis infection may affect bacterial diversity in the
72	ileum where vitamin B12 is activated, as suggested by a reduced proportion of Cobalt transport
73	protein component (CbiN) gene sequences in the gut microbiome of infected children. These
74	findings are potentially important to our understanding of how Giardia infections may affect
75	childhood growth.
-	

76

# 77 Background

78 Giardiasis is a common parasitic disease in humans in low-income settings. It is caused by the 79 protozoan Giardia intestinalis (syn. G. duodenalis or G. lamblia) that infects humans through the 80 ingestion of cysts present in a fecally-contaminated environment. Trophozoites, released in the 81 stomach, adhere to and replicate in the small intestine, while in the jejunum, trophozoites 82 transform back into cysts that are excreted in feces. Infections cause clinical manifestations 83 ranging from an asymptomatic state to acute or chronic diarrhea, malabsorption, abdominal 84 pain, vomiting, and nausea. Differences in clinical presentation may vary according to the 85 infecting Giardia assemblage, infectious dose, age and nutritional status of the host, and 86 specific effects of *Giardia* on gut microbiome composition (1,2) (3-6).

*G. intestinalis* is a common infection in children living in poorly hygienic conditions, and chronic infections impair growth and cognition (2). The prevalence of giardiasis among children in some of Ecuador's peri-urban and rural regions has been estimated between 20% and 31.5% (7–9). In contrast, in samples of children from Brazil and Colombia, prevalence was 44% and 60%, respectively (10,11).

92 The vital role of maternal and early-life nutrition in physical and neural development (12) 93 has been well described. Vitamin B12, or cobalamin, is a critical micronutrient exclusively 94 biosynthesized by specific microbes in the gastrointestinal tract. Vitamin B12 is required during 95 infancy and childhood to develop higher brain functions such as language and cognition. A 96 deficiency can result in megaloblastic anemia and neurological complications such as sensory 97 and motor disturbances and cognitive declines (13,14) (15). Vitamin B12 deficiency has been 98 linked to decreased intrinsic factors, bacterial overgrowth, and parasitic infections, including 99 Giardia (13,16). Cobalamin deficiency can also cause bacteria to shift elsewhere in the 100 gastrointestinal tract and could disrupt microbial interactions (17).

101Few studies have examined the interactions between the gut microbiome and vitamin102B12. Degnan et al. (2014) described how most of the known human gut microbial species103required B12 but lacked the genes necessary to synthesize B12 *de novo* - less than 25% of

104 intestinal bacterial species can synthesize vitamin B12(18). Vitamin B12 does not appear to 105 affect the intestinal microbiome: an analysis of infants aged 4-6 months did not show alterations 106 in the intestinal microbiota following vitamin B12 supplementation of vitamin B12 deficient infants 107 (19). In contrast, vitamin B12 supplementation in patients with homocystinuria did not affect the 108 alpha or beta diversity of the gut microbiome (20). 109 Giardia infections could affect vitamin B12 through its effects on the intestinal 110 microbiome. Mejia et al. (2020) evaluated the potential impact of giardiasis on the interaction 111 between the gut microbiome and vitamin B12, showing that G. intestinalis infected children had 112 a higher abundance of the genus *Prevotella*, and reduced presence of *Cbi*M, a gene involved in 113 vitamin B12 biosynthesis, compared to uninfected children (21). Other protozoas, including 114 Giardia, also show shifts in microbial communities(22).

115 In the present study, we hypothesized that *Giardia* infections alter the gut microbiome, 116 causing changes in bacterial species that activate vitamin B12 and may deplete bioavailable 117 vitamin B12, contributing to growth impairment in young children. To test this hypothesis, we did 118 a taxonomic and metagenomic analysis of the bacterial microbiome from fecal samples 119 collected longitudinally from Ecuadorian children at 3 and 5 years of age who were infected (or 120 not) with G. intestinalis as determined using multi-parallel real-time quantitative PCR (qPCR). 121 Our analyses of the bacterial microbiome of the gut focused on the diversity of bacteria and their 122 vitamin B12 activating sequences.

123

#### 124 Methods

125 Sample selection and study procedures

For the present study, we used a nested sample of children that were randomly selected from a birth cohort of 2,404 newborns followed up from birth to 8 years of age and living in the rural District of Quininde, Esmeraldas Province, in tropical coastal Ecuador (23). For the present study, we selected 61 stool samples from children collected at 3 and 5 years of age with 1) a

130 known positive or negative result for G. intestinalis DNA using gPCR and 2) who were negative 131 for the presence of DNA from other gastrointestinal parasites (Ascaris 132 *lumbricoides, Ancylostoma duodenale, Necator americanus, Strongyloides stercoralis,* 133 Trichuris trichiura, Cryptosporidium species, and Entamoeba histolytica) (9). None of the 134 children had received antibiotics within three months of stool samples being collected. After the 135 study sample collection, all children with positive gastrointestinal parasite results were treated 136 with appropriate anti-parasitic medication. Paired stool samples at 3 and 5 years were available 137 for 27 children, while unpaired samples from 7 children were available at 3 or 5 years of age. 138 Data on sociodemographic, individual, and household factors were collected by maternal 139 questionnaire around the time of birth of the child. A trained member of the study team 140 administered the questionnaire. 141 142 Ethics statement 143 Informed written consent was obtained from a parent or guardian of each child. The study 144 protocols were approved by the Bioethics Committee of the Universidad Internacional De 145 Ecuador, Quito, Ecuador. From the parent study, the results of stool microscopy were given to 146 the parent/guardian and anti-parasite treatment was provided where appropriate. The study was 147 also approved by the Baylor College of Medicine, institutional review board H-33219. 148 149 Parasite sample preparation 150 151 Stool samples were stored in ice and air-tight containers immediately after collection for later 152 DNA extraction using MP FastDNA Spin Kits for Soil (MP Biomedicals, Irvine, CA). DNA from 153 samples stored at -20° C and processed by multi-parallel gPCR (G. intestinalis, A. 154 lumbricoides, A. duodenale, N. americanus, S. stercoralis, T. trichiura, Cryptosporidium species, 155 and E. histolytica) as described (9,24). 156 157 Microbiome sequencing

158	Microbiome sequencing was performed at New England Biolabs, Inc. Preparation included
159	removing methylated DNA with NEBNext® Microbiome enrichment kit with no size selection. All
160	samples used one $\mu g$ of DNA with eight cycles of PCR enrichment (NEBNext Ultra DNA Library
161	Prep Kit for Illumina, Version 5.1, 9/17). Illumina NextSeq (Illumina, San Diego, CA) or Illumina
162	MiSeq (Illumina, San Diego, CA) was used to produce 151bp reads (single and paired-end);
163	however, only NextSeq sequencer reads were retained for downstream analysis.
164	
165	Bioinformatics analysis
166	Paired ends were interleaved using the BBMap program (version 38.73) to convert all
167	fastq reads to single-end reads for downstream processing (25). All single-end fastq reads were
168	filtered using a fastp program set at a minimum Phred score of 15 and a minimum length set at
169	50bp (26). MultiQC was used to visualize and summarize the filtered fastq reads (27). The NR
170	database was downloaded in fasta format from <a href="http://ftp.ncbi.nlm.nih.gov/blast/db/FASTA/">http://ftp.ncbi.nlm.nih.gov/blast/db/FASTA/</a> on
171	19th December 2019 as the reference database (28). Filtered fastq reads were processed using
172	Diamond (Version v0.9.29) with blastx mode at the default setting and maximum target
173	sequences set to one (29). Output from Diamond was processed to extract taxonomy and
174	InterPro protein database results using the Megan (community edition version 6.20.17) program
175	in STAMP format (30–32).
176	

### 177 Taxonomic and metagenomic analysis

The resultant taxonomy text file was imported into Phyloseq for calculating Shannon alpha-diversity (species richness) and beta-diversity (species variability) (33). The "dplyr" package was used for data transformation within the R programming language environment (34,35).

Taxonomic data were filtered to remove reads less than 500 reads in 20% (12 of 61
 samples) of the total samples. Any reads from phylum chordate were removed to filter for

possible human reads. Batch effects were investigated using an ordinate function from the
Phyloseq package with the "bray" distance method and Principal Coordinate Analysis (PCoA).
The InterPro database was searched for using the keyword "cobalamin" to manually extract all
vitamin B12 biosynthesis and transporter genes using InterPro entry ID and description, which
resulted in 142 proteins<sup>[1]</sup>.

189 The taxonomic and InterPro results from individual samples were exported in tab-190 delimited file format and subsequently merged using the "merge\_metaphlan\_table.py" Python 191 script, which was included with metaphlan software version 2.2.0 (36). Vegan package was 192 used to determine beta-diversity metrics, including PERMANOVA and beta-dispersion among 193 the *Giardia* guartile (zero to three), reflecting the intensity of *Giardia* DNA found via gPCR. 194 Giardia infection status and anthropometric scores (37). MaAsLin2 was used to test for 195 differentially abundant taxa with minimum prevalence of 25% and minimum abundance of 100 196 reads (38). Giardia parasite burden was measured in 4 ordinal groups that included uninfected 197 children and the 3 of 4 quartiles of increasing concentrations of Giardia DNA (1st (0.01 - 0.15), 198  $2^{nd}$  (0.17-24.50) and  $3^{rd}$  (118.58 – 29946.58). Differential analysis for genes from metagenomic 199 data was performed using the corncob package (39) with "differentialTest" function. Statistically 200 significant results were further run through "bbdml" function to obtain an effect size estimate 201 using maximum likelihood. Lastly, "plot bbdml" function on the bbdml output with 10,000 202 bootstrap simulations was used to obtain a 95% confidence interval of individual relative 203 abundance values.

204

205 Statistical analysis

The Kruskal-Wallis test was used to compare three or more independent groups. Pairwise Wilcox tests were used as post-hoc tests with a false discovery rate as a p-value correction method. Statistical significance was set at p < 0.05, and the false-discovery rate was set at 0.05 for all tests. Linear mixed effect models were conducted using the ImerTest R package (40).

- 210 Graphs were created using Prism Version 10.3.0 (Graphpad Software, Boston, MA) and the R
- 211 ggplot2 package (41).
- 212
- 213 Results
- 214 Comparing Giardia DNA burden and assemblage among infected children
- 215 There was a significant increase in the *Giardia* DNA burden at five compared to three years
- among samples from all children with evidence of infection at both ages (p=0.0012) (Figure
- 1A), and this was also seen in paired samples (p=0.031) (Figure 1B).
- 218



219 Figure 1: A. Higher levels of *Giardia* DNA in 5-year-old infected children compared to three-

year-old infected children (0.145 vs 18.76 fg/µl, P = 0.0012). B. In individually *Giardia-infected* 

221 children, there was a significant increase in Giardia DNA burden in paired samples (P =

222 **0.0312**).

223 If a 3-year-old was infected with *Giardia*, there were 7.58 greater odds (95% CI 1.31-

34.33, P = 0.0176) of having giardiasis at five years old (Figure 2). Using A and B assemblage





and B assemblages. Because of the large percentage of mixed assemblages in this cohort,

assemblages were not considered for the analysis.

228

Figure 2. Children infected with *Giardia* at three years had a 7.58 greater odds of being infected

230 at five years (1.31 to 34.33, P = 0.0176).

231 Alpha diversity of infected and non-infected children

Alpha diversity, measuring gut microbiome species diversity using the Shannon index, did not

233 differ significantly by *Giardia* parasite burden between these two groups; there were no

significant differences relative to *Giardia* DNA abundance (Supplemental Figure 1) or between

age or sex (Supplemental Figure 2).

236 To incorporate the impact of *Giardia* DNA burden on bacterial alpha diversity, a ratio of

237 alpha diversity was divided by *Giardia* DNA fg/µl was calculated for each sample. There was

some evidence of increased alpha diversity/ *Giardia* fg/µl in three-year-olds compared to age-

matched non-infected (30.20 vs. 4.37, p = 0.050), but decreased alpha diversity/ Giardia fg/µl in

five-year-olds compared to age-matched non-infected children (0.21 vs 4.31, p = 0.021).

241 Individual children that were *Giardia* positive at both time points at three and five years showed

a decrease in the ratio of alpha diversity relative to *Giardia* DNA concentration  $fg/\mu l$  (p = 0.022).

243 In the presence of *Giardia* infection, alpha diversity was greater in relation to *Giardia* DNA (fg/µl)



in three compared to five-year-olds and also compared to *Giardia*-negative children (Kruskal-

246

Figure 3. Alpha diversity normalized to concentration (fg/µl) of *Giardia intestinalis* DNA in infected children aged three-year-olds was higher than in non-infected three-year-olds (30.20 vs 4.37, p = 0.050). The opposite was seen in five-year-olds infected and non-infected (0.21 vs 4.31, p = 0.021). There was also a decrease in alpha diversity/ *Gi* fg/µl in the five-year-olds and an overall difference between all cohorts (p = 0.0053). For children without *Giardia*, their results are on the Y2 axis and are labeled Alpha Diversity. In paired samples, there was a similar decrease in Alpha diversity/ *Gi* fg/µl in the five-

year-olds (p = 0.031) with no differences in the other paired groups (Figure 4).





256

Figure 4. Paired results of Alpha diversity normalized to *Giardia* DNA concentrations (fg/ $\mu$ I) between infected and non-infected children. There was a decrease in Alpha diversity/ *Giardia* fg/ $\mu$ I in five-year-old children compared to three-year-olds (p = 0.031).

260

261

262 Beta Diversity in Giardia infected and non-infected children.

For three-year-old children, there was no difference in beta diversity between *Giardia, which was* positive and negative (R<sup>2</sup> = 0.032, p = 0.401). For five-year-old children, there was no difference in Beta diversity between *Giardia* positive and negative (R<sup>2</sup> = 0.0262, p = 0.837) (**Supplemental Figures 3**). Exploration of independent effects of *Giardia* infection status or *Giardia* quartile, age, and sex on distance matrix calculated using PERMANOVA did not reveal any statistically significant terms to explain variance (**Supplemental Table 1**). Variance in beta diversity between groups did not differ by *Giardia* infection status or across ordinal groups of

- 270 parasite burden. Beta dispersion plots, ANOVA, and permutation tests with 10,000 permutations
- 271 confirmed no significant difference in variance by *Giardia* infection quartiles for three-year-olds
- (p = 0.842) or five-year-olds (p = 0.796) (Supplemental Figure 2).
- 273
- 274 Differential Taxonomic differences
- 275 Differential abundance analysis was conducted to see possible changes at the species and
- 276 genus level with respect to infection status, the latter allowing comparisons with other 16S rRNA
- 277 gene sequencing studies. MaAsLin2 analysis (differential test function) was done to identify
- 278 OTUs that were differentially abundant relative to *Giardia* infection. Species were differentially
- abundant and positively enriched in uninfected compared to infected children in 3-year-olds
- 280 (Figure 5A) and 5-year-olds (Figure 5B). All genus and species OTUs shown have FDR < 0.05.
- 281 At the species level, we saw bacteria belonging to the genera Veillonella, Clostridium, and
- 282 Enterobacter more abundant in infected children. Uninfected children had abundant
- 283 Ruminococcus, Bacteroides, and Eubacterium species compared to Giardia-infected children of
- the same age (Figure 5)





- 287 Figure 5: Top differentially abundant OTUs at species level comparison of *Giardia* infected
- versus uninfected children controlling for three-year-olds (A) and five-year-olds (B).

289 A comparison of paired samples infected with Giardia DNA with uninfected children 290 showed 33 OTUs at genus and 132 at species levels to be differentially abundant. Differentially 291 abundant taxa between these groups included the genera Bacteroides sp. Ruminococcus sp. 292 and Mycoplasma sp. were also observed among the top 20 OTUs at the species level (Figure 293 6A). Children who were initially not infected at three years old and then positive at 5-years-old 294 had several Prevotella species (Figure 6B). Three-year-old infected children had more 295 Bacteroides sp., Paraprevotella sp., and Acinetobacter sp. compared to non-infected five-year-296 olds had more Ruminococcus sp, Bifidobacterium longum, and Eubacterium sp. (Figure 6C). Children of both ages had fewer bacterial differences than the other cohorts (Figure 6D). 297

Giardia Positive at Age 5 vs Age 3 А В Giardia -+ Positive at Age 5 vs Negative Age 3 onas sp. CAG:777 KT\_(Shigella) votella sp. CAG:5226 coccus vestibularis votella sp. MGM2 hiaella flexneri reptococcus salivarius votella sp. CAG:891 npylobacter concisus votellamassilia timonensis ibacterium sp. OM08–24 ntella sp. CAG:386 cillus cereus andidatus Melainabacteria bacterium MEL.A1 nicutes bacterium CAG:238 . KT\_(Prevotellaceae) Species autia sp. BCRC 81119 ella sp. CAG:755 Species **3** 5 **3** 5 Dysgonomonas gad Murihaculaceae bacterium Isolate-013 (NC Sporobacter term Roseburia sp. AF25-25L Bacteroides sp. D2 Ruminococcus sp. CAG:35 Bacteroides sp. CAG:18 Sutterella sp. AM11-3 Mycoplasma sp. CAG:877 Roseburia sp. AF42 Bacteroides sp. CAG:54 Prevotella sp. CAG:87 Bacteroides sp. CAG:70 , Roseburia sp. AF12-17L cutes bacterium CAG:10 Roseburia sp. AF15-2 Ruminococcus sp. CAG:48 Roseburia sp. CAG: Clostridium sp. CAG:3 Maaslin2 Coefficient Maaslin2 Coefficient wn are the top 10 sp cies with the highest and lowest coeffi Shown are the top 10 vith the highest and lo Giardia +- Negative at Age 5 vs Positive Age 3 D Giardia Negative at Age 5 vs Age 3 С ospirillum sp. 51\_20 roides sp. AR29 Eubacterium sp. CAG:76 cteroides faecis CAG:32 aprevotella clara CAG:116 achnospira pectinoschiza KT (Paraprevotella) cinetobacter sp. CAG:196 eillonella sp. CAG:933 raprevotella clara roides clarus achvspira sp. CAG:484 nococcus sp. AF21–3 Age pecies es sp. AF27-10BH Species 35 Eubacterium sp. TM06-4 **3** 5 Clostridium sp. AF15-4 Ruminococcus broi Clostridium sp. CAG:5 Prevotella sp. MGM2 ninococcaceae bacterium FB201. Ruminococcus sp. AM28-29I F , revotella sp. Marseille-P411. Bifidobacterium longu Ruminococcus sp. CAG:35 I KT (Bifidobacterii Actinomyces gra Firmicutes bacterium CAG:88 Prevotella sp. CAG:87 Maaslin2 Coefficient Maaslin2 Coefficient are the top 10 species with the highest and lowest coefficients Shown are the top 10 spe ies with the highest and lowest coefficients

Age

Figure 6. Top 20 differentially abundant OTUs at genus and species level for *Giardia* positives at both ages (A), the non-infected at three years old and then infected at five years old (B), and those infected at three years old and then non-infected at five years old (C). Only 8 OTUs were significantly different in abundance for those children not infected at both ages (D).

304

#### 305 Differential abundant cobalamin genes

306 Out of 143 vitamin B12 biosynthesis genes, 32 genes were found to be present among the 61 307 samples. When conducting a differential abundance test from the Corncob package by Giardia 308 infection status using beta-binomial regression with sex and age as covariates, IPR003705 – 309 Cobalt transport protein (*Cbi*N) was found to be significantly in low abundance in *Giardia*-310 positive children with FDR < 0.001. CbiN was further tested individually with sex and age as 311 covariates. It was found to be less prominent in the *Giardia-positive* group with p = 0.002. The 312 resultant graph shows the predicted confidence interval of the *Cbi*N gene in samples after 313 10,000 bootstrap simulations (Figure 7A).

Notably, the differential abundant test with *Giardia* quartile instead of *Giardia* infection status indicated genes coding for IPR003705 – Cobalt transport protein (*Cbi*N) (FDR < 0.003) and IPR011822 (Cobalamin-dependent methionine synthase) with FDR = 0.003 were found to be statistically significantly differentially abundant in the *Giardia* negative group vs. *Giardia* positive group, with sex and age as covariates.

Differential analysis of these genes was individually analyzed to get estimates by the Giardia quartile. For the *CbiN* gene, the remaining *Giardia* quartiles (2 and 3), as compared to the no *Giardia* quartile, had statistically significant negative abundances with p = 0.01 and p < 0.001, respectively (**Figure 7B**). The predicted intervals from the differential abundance testing for *Cbi*N alone are noticeably higher in the *Giardia-positive* group.

324



Figure 7. Relative abundance of the CbiN gene was decreased in children infected with *Giardia* compared to non-infected children (A). Relative abundance of CbiN gene among all the samples by *Giardia* concentration quartiles, Neg (0), and increasing concentrations (1, 2, 3). The prediction intervals are calculated using 10,000 bootstrap simulations.

329

#### 330 Discussion

331 Giardia infections are estimated to be present in about one-third of developing countries' 332 children and adult populations (42). Microbiome perturbation due to Giardia infection was 333 minimal compared to Giardia-negative children, as there were no significant changes in the 334 alpha-diversity and beta-diversity between the groups. Our results also show a non-significant 335 increase in the alpha diversity with age. A study conducted on 417 Swedish children from 336 newborn to five years of age found that the compositional variation was the least between three 337 and five years (0.9%) (43). This indicates that microbiome changes between these time points 338 are leveling out as they may be approaching an adult-like state. Such a decline in change may 339 have flattened Giardia's impact on diversity changes. Mice studies have shown an increase in 340 Shannon alpha-diversity or richness in species following Giardia infection, while other studies 341 have found no alterations (3,5,44). However, these studies were conducted with 16S rRNA 342 sequencing and murine models. Microbiome studies in humans show that Giardia infection can

reduce alpha diversity (6,21) or distinctive microbial profiles in individuals with asymptomatic
versus symptomatic giardiasis(45). Toro-Londono et al. found significant changes in beta
diversity in the *Giardia* group vs. the control group(46).

346 It is speculated that the amount of *Giardia*, or its burden, is related to disease 347 progression, including changes in the intestinal microbiota(47). Therefore, we used the amount 348 of *Giardia* DNA, in fg/µl, to modulate the microbial alpha diversity by mathematically dividing 349 them and creating a ratio of alpha diversity/ *Giardia* fg/µl of DNA. This ratio explores how 350 *Giardia's* DNA burden affects microbial alpha diversity per infected child.

351 This study revealed a number of differentially abundant taxa, which may help understand 352 the microbiome changes caused by the *Giardia* infection. It was not possible to distinguish the 353 stool sample bacteria from the various intestinal regions. However, the small intestine has a 354 higher oxygen level and less bacteria density than the colon due to the antimicrobial activity of 355 bile acids (48). Samples with the highest *Giardia* infection showed abundant anaerobic species 356 from *Clostridium* and *Veillonella* (Figure 5), which reside in the duodenum and cause biofilms in 357 vitro (49,50). Biofilms can lead to pathogenesis, antimicrobial resistance, virulence, and 358 restructure the microbiome (51). Interestingly, V. atypica from genus Veillonella is a lactate 359 metabolizer in a low pH environment such as the small intestine and may negate some of the 360 probiotic effects of lactic acid-producing bacteria Lactobacillus and Bifidobacterium on Giardia 361 (52, 53).

Cobalamin is absorbed by Intrinsic factor (IF) at the cobalamin complex at the receptors in the ileum. It is known that cobalamin absorption can be impacted by small intestine bacterial overgrowth (SIBO) due to competition with IF (54). Some microbes, such as *Bacteroides thetaiotamicron,* can remove cobalamin from IF with BtuG2 protein and compete directly with the human host (55). This study found bacteria containing *Btu*G homologs, such as *B. coprophilus,* with lower abundance in the highest *Giardia* quartile than the *Giardia* negative group.

369 Specific cobalamin biosynthesis sequences were decreased in the Giardia-positive 370 groups, showing that the microbiome was deficient in cobalamin generation genes. Changes in 371 microbiome diversity and the low presence of BtuG homologs bacteria may have been due to 372 the effects of Giardia infection in these children. Corncob analysis limited to cobalamin-related 373 genes revealed the lack of CbiN – a cobalt transporter protein among the Giardia positive 374 groups, suggesting an indirect impact on activating cobalamin and discordance at the individual 375 vital amino acid level. In a larger study, children with giardiasis had broad amino acid 376 deficiencies (56). The available data in this study cannot determine whether the lack of the CbiN 377 gene impacts the overall cobalamin availability across the gut microbiome. 378 Limitations of this study include the lack of statistical power across relatively small 379 samples of children sampled at 3 and 5 years of age. Also, serum levels of cobalamin were 380 unavailable, and metabolomics data could not correlate with microbiome findings or 381 anthropometric measures. Finally, a gap of a two-year timeframe between sample collection 382 may be comparatively long to detect microbiome changes. 383

#### 384 Conclusion

385 This longitudinal analysis observed a decrease in alpha diversity normalized to Giardia DNA 386 (fg/µl) concentration as children increased in age from 3 to 5 years old, revealing the impact on 387 microbiome diversity changes as children age. Differential taxa abundance analysis showed 388 microbiome changes at genus and species levels, which could allow biofilm formation and 389 reduction in potentially beneficial butyrate-producing bacteria. There were also fewer cobalamin 390 synthesis genes among the microbiomes of the children with a higher infectious burden of 391 Giardia. These findings may help understand the mechanism of how Giardia may lead to growth 392 delays and stunting.

393

Funding: Internal funds from New England Biolabs for DNA sequencing. Research funding was
 provided by the US Department of Health and Human Services, Health Resources and Services
 Administration for Baylor College of Medicine Center of Excellence in Health Equity, Training,

397 and Research (Grant No: D34HP31024). Also, the Baylor College of Medicine Junior Faculty

398 Seed Award and BCM Center for Globalization Pilot Project were used for research supplies.

- 399 Also, Wellcome Trust grant 088862/Z/09/Z.
- 400

401 Authors' contributions: RM and AD were responsible for bioinformatics and statistical

- 402 analysis. RM, AA, AL performed DNA extraction, qPCR, demethylation, and library preparation.
- 403  $\,$  EL, CL, ED, BS were involved in the DNA sequencing sample reactions
- 404 and preliminary analysis. RM, AD, VSH, DGR, BS, PJC wrote the manuscript. All authors read 405 and approved the final manuscript.
- 406

407 408	References			
409 410 411	1.	Certad G, Viscogliosi E, Chabé M, Cacciò SM. Pathogenic Mechanisms of Cryptosporidium and Giardia. Trends Parasitol. 2017 Jul 1;33(7):561–76.		
412 413	2.	Halliez MC, Buret AG. Extra-intestinal and long term consequences of Giardia duodenalis infections. World J Gastroenterol WJG. 2013 Dec 21;19(47):8974–85.		
414 415 416	3.	Beatty JK, Akierman SV, Motta JP, Muise S, Workentine ML, Harrison JJ, et al. Giardia duodenalis induces pathogenic dysbiosis of human intestinal microbiota biofilms. Int J Parasitol. 2017 May 1;47(6):311–26.		
417 418 419	4.	Huth S von, Thingholm LB, Kofoed PE, Bang C, Rühlemann MC, Franke A, et al. Intestinal protozoan infections shape fecal bacterial microbiota in children from Guinea-Bissau. PLoS Negl Trop Dis. 2021 Mar 3;15(3):e0009232.		
420 421 422 423	5.	Riba A, Hassani K, Walker A, Best N van, Zeschwitz D von, Anslinger T, et al. Disturbed gut microbiota and bile homeostasis in Giardia-infected mice contributes to metabolic dysregulation and growth impairment. Sci Transl Med [Internet]. 2020 Oct 14 [cited 2021 Mar 9];12(565). Available from: https://stm.sciencemag.org/content/12/565/eaay7019		
424 425 426	6.	Toro-Londono MA, Bedoya-Urrego K, Garcia-Montoya GM, Galvan-Diaz AL, Alzate JF. Intestinal parasitic infection alters bacterial gut microbiota in children. PeerJ. 2019 Jan 7;7:e6200.		
427 428 429	7.	Jacobsen KH, Ribeiro PS, Quist BK, Rydbeck BV. Prevalence of intestinal parasites in young Quichua children in the highlands of rural Ecuador. J Health Popul Nutr. 2007 Dec;25(4):399–405.		
430 431 432	8.	Lowenstein C, Vasco K, Sarzosa S, Salinas L, Torres A, Perry MJ, et al. Determinants of Childhood Zoonotic Enteric Infections in a Semirural Community of Quito, Ecuador. Am J Trop Med Hyg. 2020 Mar 30;102(6):1269–78.		

433 9. Mejia R, Vicuña Y, Broncano N, Sandoval C, Vaca M, Chico M, et al. A Novel, Multi-434 Parallel, Real-Time Polymerase Chain Reaction Approach for Eight Gastrointestinal 435 Parasites Provides Improved Diagnostic Capabilities to Resource-Limited At-Risk 436 Populations. Am J Trop Med Hyg. 2013 Jun 5;88(6):1041-7. 437 10. Bryan PE, Romero M, Sánchez M, Torres G, Gómez W, Restrepo M, et al. Urban versus 438 Rural Prevalence of Intestinal Parasites Using Multi-Parallel qPCR in Colombia. Am J 439 Trop Med Hyg. 2020 Dec 14; 440 11. Fantinatti M, Lopes-Oliveira LAP, Cascais-Figueredo T, Austriaco-Teixeira P, Verissimo E, 441 Bello AR, et al. Recirculation of Giardia lamblia Assemblage A After Metronidazole 442 Treatment in an Area With Assemblages A, B, and E Sympatric Circulation. Front 443 Microbiol [Internet]. 2020 Oct 22 [cited 2021 Mar 6];11. Available from: 444 https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7642054/ 445 12. Forgie AJ, Drall KM, Bourque SL, Field CJ, Kozyrskyj AL, Willing BP. The impact of 446 maternal and early life malnutrition on health: a diet-microbe perspective. BMC Med 447 [Internet]. 2020 May 12 [cited 2021 Mar 26];18. Available from: 448 https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7216331/ 449 13. Green R, Allen LH, Bjørke-Monsen AL, Brito A, Guéant JL, Miller JW, et al. Vitamin B 12 450 deficiency. Nat Rev Dis Primer. 2017 Jun 29;3(1):1–20. 451 14. Langan RC, Goodbred AJ. Vitamin B12 Deficiency: Recognition and Management. Am Fam 452 Physician. 2017 Sep 15;96(6):384-9. 453 15. Goraya JS. Vitamin B12 deficiency in Indian infants and children. Paediatr Int Child Health. 454 2020 May;40(2):75-7. 455 16. Handbook of Vitamins [Internet]. CRC Press; 2013 [cited 2021 Mar 9]. Available from: 456 https://www.taylorfrancis.com/books/handbook-vitamins-janos-zempleni-john-suttie-jesse-457 gregory-iii-patrick-stover/10.1201/b15413 458 17. Mach N, Clark A. Micronutrient Deficiencies and the Human Gut Microbiota. Trends 459 Microbiol. 2017 Aug 1;25(8):607–10. 460 18. Degnan PH, Taga ME, Goodman AL. Vitamin B 12 as a Modulator of Gut Microbial 461 Ecology. Cell Metab. 2014 Nov;20(5):769-78. 462 19. Boran P, Baris HE, Kepenekli E, Erzik C, Soysal A, Dinh DM. The impact of vitamin B12 463 deficiency on infant gut microbiota. Eur J Pediatr. 2020 Mar 1;179(3):385–93. 464 20. Rizowy GM, Poloni S, Colonetti K, Donis KC, Dobbler PT, Leistner-Segal S, et al. Is the gut 465 microbiota dysbiotic in patients with classical homocystinuria? Biochimie. 2020 Jun;173:3– 466 11.

- 467 21. Mejia R, Damania A, Jeun R, Bryan PE, Vargas P, Juarez M, et al. Impact of intestinal
  468 parasites on microbiota and cobalamin gene sequences: a pilot study. Parasit Vectors. 2020
  469 Apr 19;13(1):200.
- 470 22. Von Huth S, Thingholm LB, Kofoed PE, Bang C, Rühlemann MC, Franke A, et al. Intestinal
  471 protozoan infections shape fecal bacterial microbiota in children from Guinea-Bissau.
  472 Bartelt LA, editor. PLoS Negl Trop Dis. 2021 Mar 3;15(3):e0009232.
- 473 23. Cooper PJ, Chico ME, Platts-Mills TA, Rodrigues LC, Strachan DP, Barreto ML. Cohort
  474 Profile: The Ecuador Life (ECUAVIDA) study in Esmeraldas Province, Ecuador. Int J
  475 Epidemiol. 2015 Oct;44(5):1517–27.
- 476 24. Cimino RO, Jeun R, Juarez M, Cajal PS, Vargas P, Echazú A, et al. Identification of human
  477 intestinal parasites affecting an asymptomatic peri-urban Argentinian population using
  478 multi-parallel quantitative real-time polymerase chain reaction. Parasit Vectors. 2015 Jul
  479 17;8:380.
- 480
  480
  481
  481
  481
  481
  482
  482
  483
  483
  484
  484
  485
  485
  485
  486
  486
  487
  487
  488
  488
  489
  480
  480
  480
  480
  480
  480
  481
  481
  482
  483
  483
  483
  484
  484
  485
  485
  485
  486
  486
  487
  487
  488
  488
  488
  489
  480
  480
  480
  480
  480
  481
  481
  481
  482
  483
  483
  484
  484
  484
  485
  485
  486
  486
  487
  487
  487
  488
  488
  488
  488
  489
  480
  480
  480
  480
  480
  481
  481
  481
  482
  482
  483
  484
  484
  484
  485
  485
  486
  486
  487
  487
  488
  488
  488
  488
  488
  489
  489
  480
  480
  480
  480
  480
  480
  480
  480
  480
  480
  480
  480
  480
  481
  481
  481
  482
  482
  483
  483
  484
  484
  484
  484
  484
  485
  485
  486
  486
  487
  487
  488
  488
  488
  488
  488
  488
  488
  488
  488
  488
  488
  488
  488
  488
  488
  488
  488
  488
  488
  488
  488
  488
  488
  488
  488
  488
- 484 26. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
  485 Bioinformatics. 2018 Sep 1;34(17):i884–90.
- 486 27. Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results for
  487 multiple tools and samples in a single report. Bioinformatics. 2016 Oct 1;32(19):3047–8.
- 488 28. O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, et al. Reference
  489 sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional
  490 annotation. Nucleic Acids Res. 2016 Jan 4;44(D1):D733-745.
- 491 29. Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND. Nat
  492 Methods. 2015 Jan;12(1):59–60.
- 30. Blum M, Chang HY, Chuguransky S, Grego T, Kandasaamy S, Mitchell A, et al. The
  InterPro protein families and domains database: 20 years on. Nucleic Acids Res. 2021 Jan
  8;49(D1):D344–54.
- 496 31. Huson DH, Beier S, Flade I, Górska A, El-Hadidi M, Mitra S, et al. MEGAN Community
  497 Edition Interactive Exploration and Analysis of Large-Scale Microbiome Sequencing
  498 Data. PLOS Comput Biol. 2016 Jun 21;12(6):e1004957.
- 499 32. Parks DH, Tyson GW, Hugenholtz P, Beiko RG. STAMP: statistical analysis of taxonomic
  500 and functional profiles. Bioinforma Oxf Engl. 2014 Nov 1;30(21):3123–4.
- 33. McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis and
   Graphics of Microbiome Census Data. PLOS ONE. 2013 Apr 22;8(4):e61217.

- 34. R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna,
   Austria: R Foundation for Statistical Computing; 2020. Available from: https://www.R project.org/
- 35. Wickham H, François R, Henry L, Müller K. dplyr: A Grammar of Data Manipulation
   [Internet]. 2021. Available from: https://CRAN.R-project.org/package=dplyr
- Segata N, Waldron L, Ballarini A, Narasimhan V, Jousson O, Huttenhower C. Metagenomic
   microbial community profiling using unique clade-specific marker genes. Nat Methods.
   2012 Aug;9(8):811–4.
- 511 37. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan:
  512 Community Ecology Package [Internet]. 2020. Available from: https://CRAN.R513 project.org/package=vegan
- 38. Mallick H, Rahnavard A, McIver LJ, Ma S, Zhang Y, Nguyen LH, et al. Multivariable
  association discovery in population-scale meta-omics studies. PLOS Comput Biol. 2021
  Nov 16;17(11):e1009442.
- 39. Martin BD, Witten D, Willis AD. Modeling microbial abundances and dysbiosis with beta binomial regression. Ann Appl Stat. 2020 Mar;14(1):94–115.
- 40. Kuznetsova A, Brockhoff PB, Christensen RHB. ImerTest Package: Tests in Linear Mixed
  Effects Models. J Stat Softw. 2017;82(13):1–26.
- 41. Wickham H. ggplot2: elegant graphics for data analysis. Second edition. Cham: Springer;
   2016. 260 p. (Use R!).
- 42. Dunn N, Juergens AL. Giardiasis. In: StatPearls [Internet]. Treasure Island (FL): StatPearls
  Publishing; 2021 [cited 2021 Apr 16]. Available from: http://www.ncbi.nlm.nih.gov/books/NBK513239/
- 43. Roswall J, Olsson LM, Kovatcheva-Datchary P, Nilsson S, Tremaroli V, Simon MC, et al.
  Developmental trajectory of the healthy human gut microbiota during the first 5 years of
  life. Cell Host Microbe [Internet]. 2021 Mar 31 [cited 2021 Apr 24];0(0). Available from:
  https://www.cell.com/cell-host-microbe/abstract/S1931-3128(21)00100-1
- 44. Barash NR, Maloney JG, Singer SM, Dawson SC. Giardia Alters Commensal Microbial
  Diversity throughout the Murine Gut. Infect Immun [Internet]. 2017 May 23 [cited 2021
  Apr 17];85(6). Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5442636/
- 45. McGregor BA, Razmjou E, Hooshyar H, Seeger DR, Golovko SA, Golovko MY, et al. A
  shotgun metagenomic analysis of the fecal microbiome in humans infected with Giardia
  duodenalis. Parasit Vectors. 2023 Jul 18;16(1):239.
- 46. Toro-Londono MA, Bedoya-Urrego K, Garcia-Montoya GM, Galvan-Diaz AL, Alzate JF.
   Intestinal parasitic infection alters bacterial gut microbiota in children. PeerJ. 2019 Jan
   7;7:e6200.

- 47. Mejia R, Damania A, Jeun R, Bryan PE, Vargas P, Juarez M, et al. Impact of intestinal
  parasites on microbiota and cobalamin gene sequences: a pilot study. Parasit Vectors. 2020
  Dec;13(1):200.
- 542 48. Donaldson GP, Lee SM, Mazmanian SK. Gut biogeography of the bacterial microbiota. Nat
  543 Rev Microbiol. 2016 Jan;14(1):20–32.
- 49. Martinez-Guryn K, Leone V, Chang EB. Regional Diversity of the Gastrointestinal
  Microbiome. Cell Host Microbe. 2019 Sep 11;26(3):314–24.
- 546 50. Motta JP, Wallace JL, Buret AG, Deraison C, Vergnolle N. Gastrointestinal biofilms in
  547 health and disease. Nat Rev Gastroenterol Hepatol. 2021 Jan 28;1–21.
- 548 51. Tytgat HLP, Nobrega FL, van der Oost J, de Vos WM. Bowel Biofilms: Tipping Points
  549 between a Healthy and Compromised Gut? Trends Microbiol. 2019 Jan 1;27(1):17–25.
- 550 52. Fekete E, Allain T, Siddiq A, Sosnowski O, Buret AG. Giardia spp. and the Gut Microbiota:
  551 Dangerous Liaisons. Front Microbiol [Internet]. 2021 Jan 12 [cited 2021 Feb 23];11.
  552 Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7835142/
- 53. Scheiman J, Luber JM, Chavkin TA, MacDonald T, Tung A, Pham LD, et al. Meta-omics
  analysis of elite athletes identifies a performance-enhancing microbe that functions via
  lactate metabolism. Nat Med. 2019 Jul;25(7):1104–9.
- 556 54. Rowley CA, Kendall MM. To B12 or not to B12: Five questions on the role of cobalamin in
   557 host-microbial interactions. PLOS Pathog. 2019 Jan 3;15(1):e1007479.
- 55. Wexler AG, Schofield WB, Degnan PH, Folta-Stogniew E, Barry NA, Goodman AL.
  Human gut Bacteroides capture vitamin B12 via cell surface-exposed lipoproteins. eLife.
  2018 Sep 18;7:e37138.
- 56. Giallourou N, Arnold J, McQuade ETR, Awoniyi M, Becket RVT, Walsh K, et al. Giardia
  hinders growth by disrupting nutrient metabolism independent of inflammatory
  enteropathy. Nat Commun. 2023 May 18;14(1):2840.
- 564