| $\frac{1}{2}$ | A decade of dengue disease burden in Africa (2013–2023): a systematic review |
|---------------|--|
| 3 | <u>Gaspary O. Mwanyika</u> ^{a,b} , <u>Monika Moir^a, Abdualmoniem O. Musa^c, Jenicca</u> |
| 4 | Poongavanan ^a , <u>Graeme Dor^a, Eduan Wilkinson^a, Cheryl Baxter^a, <u>Tulio de Oliveira</u>^{a,d}</u> |
| 5 | and <u>Houriiyah Tegally</u> ^a |
| 6 | |
| 7 | ^a Centre for Epidemic Response and Innovation (CERI), School for Data Science and |
| 8 | Computational Thinking, Stellenbosch University, Stellenbosch, South Africa |
| 9 | ^b Department of Applied Sciences, Mbeya University of Science and Technology, |
| 10 | Mbeya, Tanzania |
| 11 | ^c Kassala University & Ministry of Health, Kassala State, Sudan |
| 12 | ^d KwaZulu-Natal Research Innovation and Sequencing Platform, University of |
| 13 | KwaZulu- Natal, Durban, South Africa |
| 14 | |
| 15 | |
| 16 | |
| 17 | |
| 18 | |
| 19 | Corresponding authors: Houriiyah Tegally (houriiyah@sun.ac.za) |
| 20 | |
| 21 | |
| 22 | |
| 23 | |
| 24 | |
| 25 | |

Travel Medicine and Infectious Diseases

26 Abstract

27 Background

- 28 Dengue is a major mosquito-borne disease worldwide. The epidemiological trends of the
- 29 disease in Africa over the past decade remain unclear. This review aims to provide insight
- 30 into the epidemiological trends of dengue in Africa from 2013–2023.

31 Methods

- 32 We systematically searched PubMed/MEDLINE and Scopus for studies published
- 33 between January 2013 and December 2023. Additionally, we collected official
- 34 records from the World Health Organization for Africa and African Centre for Disease
- 35 Control. We included studies that reported dengue cases in humans in Africa and
- 36 excluded publications prior to 2013, review articles and non-human studies. For
- 37 specific countries, the suspected cases per 100,000 population and fatality rates
- 38 were estimated and the trend predicted using a negative binomial model. The
- 39 statistical analyses and visualisations were performed using R programming.

40 **Results**

- 41 Of the 453 reports screened, 87 from 25 African countries were selected for
- 42 systematic review. Between 2013 and 2023, approximately 200,000 suspected
- 43 dengue cases, 90,000 confirmed cases and 900 deaths were reported in Africa.
- 44 Over 80% of confirmed cases originated from West Africa, with Burkina Faso
- 45 reporting over 500 cases per 100,000 population. DENV1 and DENV2 predominating
- 46 at different times with transmission closely linked to rainy seasons.

47 Conclusions

48 The rising dengue cases across Africa, highlight the need to strengthen surveillance

49 and implement effective regional-specific interventions against future dengue

Travel Medicine and Infectious Diseases

- 50 outbreaks. Further research is necessary to improve our understanding on dengue
- 51 transmission dynamics and suitability of regions in Africa.
- 52 Key words: dengue, disease burden, Africa, systematic review
- 54 **1.** Introduction

53

55 Dengue is a major mosquito-borne viral disease transmitted by infected female 56 mosquitoes of the Aedes genus [1]. The disease is caused by dengue virus (DENV) 57 of the Flaviviridae family in the genus Orthoflavivirus. The virus consists of four 58 antigenically distinct serotypes: DENV1, DENV2, DENV3 and DENV4, which are 59 capable of inducing mild to severe illnesses in humans [2]. Globally, approximately 60 390 million dengue infections, 500,000 hospitalizations and more than 20,000 deaths 61 are reported annually [3]. In recent decades, the virus has transcended its traditional 62 boundaries, extending into temperate regions including Europe and North America [4,5]. The disease has been reported in Africa since the late 19th and early 20th 63 64 centuries, with reported cases in Zanzibar (1870), Burkina Faso (1925), Egypt 65 (1887), South Africa (1926–1927), and Senegal (1927–1928). From the 1960s to 66 2010, laboratory confirmed cases were reported in 15 African countries, with recent 67 endemicity established in more than 34 countries [6]. Although the prevalence of 68 dengue virus across Africa has been previously documented [7,8], epidemiological 69 trends in the context of morbidity and mortality, geographical distribution, seasonality 70 and transmission suitability remain unclear. In this systematic review, we collate a 71 decade of dengue epidemiological data from the African continent to synthesise and 72 analyse epidemiological trends between 2013 and 2023. We highlight regional and 73 country-specific epidemiological trends to inform public health response against 74 future dengue outbreaks in Africa.

Travel Medicine and Infectious Diseases

75 **2.** Methods

76 **2.1 Search strategy and selection criteria**

77 This review analysed studies and reports describing dengue cases in African

78 countries and territories. Medical Subject Headings (MeSH) terms such as: dengue

virus, prevalence, epidemiology, serotype, in conjunction with a compilation of

80 specific African countries and territories were used in the search. We searched

81 PubMed/MEDLINE and Scopus databases for relevant English articles using an

82 advanced search strategy (Appendix S1). Additional epidemiological reports were

83 obtained from the World Health Organization for Africa (WHO AFRO) and the African

84 Centre for Disease Control (AFRICA CDC) between January 2013 and December

85 2023. We included articles that reported dengue cases in human studies in Africa

and official records from the WHO AFRO and AFRICA CDC surveillance reports. We

87 excluded publications prior to 2013, review articles, and non-human studies.

88 **2.2 Screening and quality assessment**

89 Two reviewers (GOM and AM) conducted an initial screening of the titles and

90 abstracts from the search results using Rayyan application software accessible at

91 <u>https://rayyan.ai/</u> to identify relevant articles. Reasons for exclusion of irrelevant

92 articles were documented. Three reviewers (GOM, AM and HT) evaluated the quality

93 of included reports. The selection of reports for inclusion in the systematic review

94 followed the Preferred Reporting Items for Systematic Reviews and Meta-analysis

95 guidelines (PRISMA) [9].

96

97 2.3 Key definitions

Travel Medicine and Infectious Diseases

- 98 The following terms have been used in the analysis to signify different metrics used
 99 to describe dengue burden in Africa.
- 100 1. A suspected case was defined as any individual residing in or having travelled 101 to areas with dengue transmission within the past 14 days and presents with 102 acute fever, typically lasting from two to seven days' duration with two or more 103 of the following symptoms: nausea/vomiting, abdominal pain, chills, rash, 104 headache/retro-orbital pain, myalgia and arthralgia; may exhibit petechial or 105 positive tourniquet test (+ >10 pinpoint-sized spots of bleeding under the skin 106 (petechiae) per square inch, low platelet and white blood cell counts even 107 without any warning sign. 108 2. A confirmed case was defined as a suspected dengue case with laboratory 109 confirmation of infection which may include polymerase chain reaction (PCR), 110 virus culture, IgM seroconversion in paired sera (acute and convalescent 111 samples), IgG seroconversion in paired sera or fourfold IgG titre in paired 112 sera.
- 1133. A severe case was defined as a suspected/probable/confirmed dengue case
- 114 presenting with one or more of the following symptoms: severe plasma
- 115 leakage leading to dengue shock syndrome fluid accumulation with
- 116 respiratory distress, severe bleeding, severe damage of organs such as liver
- 117 (aspartate aminotransferase (ASAT) or alanine aminotransferase (ALT)
- 118 elevation \geq 1000) and central nervous system.
- 4. Case fatality rate was defined as the proportion of deaths within a specified
 population that are attributable to the total number of suspected dengue cases
 over a specific period.

Travel Medicine and Infectious Diseases

| 122 | 5. | Transmission potential (index P) was defined as a measure that quantifies the |
|-----|----|--|
| 123 | | risk of dengue virus transmission in a specific region, taking into account |
| 124 | | climate-based factors such as temperature and humidity, which directly |
| 125 | | influence the breeding, survival, and biting behaviour of the mosquito vectors |
| 126 | | responsible for spreading the virus. This index provides insight into how |
| 127 | | favourable environmental conditions are for dengue circulation in a given |
| 128 | | area. The index was developed by Nakase et. al (2023) [10]. The spatio- |
| 129 | | temporal estimates of transmission potential were then compiled for African |
| 130 | | countries and territories. A threshold of 1.0 was selected to compare with the |
| 131 | | basic reproductive rate. A value of 1.0 indicates that in a population where the |
| 132 | | average number of adult female mosquitoes per host is 1.0 corresponds to a |
| 133 | | reproduction number of 1. The period of transmission suitability is defined as |
| 134 | | a month in which the transmission potential is greater than 1.0 [10]. |

135

2.4 Data synthesis and analysis

136 Data on the year of outbreak, country of origin, number of suspected and confirmed 137 cases, deaths and serotype counts were extracted from included reports and 138 compiled into an Excel spreadsheet (Miscrosoft Corp., 2016 Redmond, WA, USA). 139 The estimation of dengue burden was based on the number of suspected cases per 140 country's population during the respective years. The population data for the 141 respective years were obtained from the Worldometer web-source, accessible at 142 https://www.worldometers.info/population/africa/. Case fatality rate (CFR) was 143 computed based on the number of reported deaths per total number of suspected 144 dengue cases in the specific country. A negative binomial model was used to predict 145 the growth of suspected cases using the year as a predictor variable. The model was

Travel Medicine and Infectious Diseases

- 146 selected to suit count data, such as dengue case numbers and accounts for
- 147 overdispersion. The model was presented by the following equation;

$$\log(y) = \beta_0 + \beta_1 \cdot Year_t$$

149 Where;

150 y = Expected number of suspected dengue cases

151 β_0 = Intercept of the model

152 β_1 = Coefficient for the year t

153 $Year_t$ = The year variable for which the number of dengue cases was predicted

154 $e^{\beta 0 + \beta 1} = A$ multiplicative factor for the growth in dengue cases. The best fit was

- 155 compared with a Poisson regression model using likelihood ratio test. Statistical
- analysis and visualizations were conducted using R version 4.3.2 with primary
- 157 packages ggplot2, dplyr and MASS.
- 158

168

159 **3.** Results

160 **3.1 Literature search**

The review protocol was registered in the PROSPERO International prospective register for systematic reviews of human studies under CRD42023480486. The search yielded a total of 453 results, including 297 from academic databases and 156 official sources. After removing 151 duplicates, 302 unique records were screened by titles and abstracts. 173 reports were excluded from the screening process, and 129 were evaluation for eligibility. Finally, 42 records were excluded for specific reasons (Appendix S2), and 87 included in the systematic review (**Figure 1**).

Travel Medicine and Infectious Diseases



169

170 Figure 1.Selection process of included reports according to PRISMA guideline.171

172 3.2 An overview of included reports

- 173 Table 1 presents the epidemiological data extracted from the included reports,
- 174 comprising 35 published articles and 52 official records from the WHO AFRO and
- 175 AFRICA CDC surveillance data. Between January 2013 and December 2023, data
- 176 on dengue were available from 25 African countries. Seven countries, including,

- 177 Burkina Faso (7 records, n = 14,7286), Cameroon (6, n = 2513), Cote d'Ivoire (5, n =
- 178 8,250), Kenya (8, n = 7,828), Senegal (8, n = 4,606), Sudan (7, n = 2,696) and
- 179 Tanzania (6, n = 10,369) accounted for more than 50% of all records (46/87) and
- 180 80% of cumulative suspected cases (19,7707/24,4352). A comparison year-on-year
- reveals that the highest number of reports (21.8%, 19/87) and suspected cases
- 182 (60.3%, 174,888/245,143) were documented in 2023. In this year, Burkina Faso
- reported the most lethal outbreak with a total of 147,286/174,888 (84.2%) cumulative
- 184 suspected cases, 68,402/72677 (94.1%) confirmed cases, and 688/765 (89.9%).

| Reference | Country | Population | Year of | Suspected | Confirmed | Deaths | | Serc | otype | |
|-------------|--------------|------------|----------|-----------|-----------|--------|---------|-----------|-------|-------|
| | | | outbreak | cases | cases | | DENV1 | DENV2 | DENV3 | DENV4 |
| [11] | Tanzania | 49253643 | 2013 | 431 | 431 | 2 | 0 | 431 | 0 | 0 |
| [12] | Kenya | 44790000 | 2013 | 267 | 101 | 1 | 51 | 48 | 0 | 0 |
| [13] | Tanzania | 50814552 | 2014 | 2121 | 1017 | 4 | 0 | 0 | 0 | 0 |
| [14] | South Sudan | 37003245 | 2014 | 155 | 35 | 8 | 35 | 0 | 0 | 0 |
| [15] | Tanzania | 50814552 | 2014 | 483 | 101 | 0 | 0 | 101 | 0 | 0 |
| [16] | Kenya | 45831863 | 2014 | 1022 | 361 | 0 | 36 | 105 | 10 | 11 |
| [17] | Nigeria | 179E+08 | 2014 | 526 | 24 | 0 | 0 | 0 | 0 | 0 |
| [18] | Kenya | 45831863 | 2014 | 489 | 43 | 0 | 0 | 0 | 0 | 0 |
| [19] | Burkina Faso | 18106000 | 2015 | 399 | 21 | 0 | 0 | 0 | 0 | 0 |
| [20] | Cameroon | 23012646 | 2015 | 349 | 21 | 0 | 0 | 0 | 0 | 0 |
| [21] | Egypt | 97720000 | 2015 | 253 | 28 | 0 | 28 | 0 | 0 | 0 |
| [22] | Senegal | 14360000 | 2015 | 104 | 3 | 0 | 3 | 0 | 0 | 0 |
| [23] | Burkina Faso | 19280000 | 2016 | 1327 | 19 | 0 | 0 | 11 | 6 | 0 |
| [24] | Democratic | 81430000 | 2016 | 253 | 14 | 0 | 12 | 2 | 0 | 0 |
| | Republic of | | | | | | | | | |
| | Congo | | | | | | | | | |
| [25] | Sudan | 39380000 | 2016 | 106 | 4 | 0 | 0 | 4 | 0 | 0 |
| [26] | Seychelles | 94677 | 2016 | 1062 | 422 | 0 | 0 | 0 | 0 | 0 |
| [27] | Burkina Faso | 19280000 | 2016 | 2929 | 317 | 0 | 6 | 191 | 104 | 0 |
| [28] | Kenya | 47894670 | 2016 | 560 | 5 | 0 | 0 | 4 | 0 | 0 |
| [00] | Burkina Faso | 19280000 | 2016 | 1266 | 29 | 15 | 0 | 29 | 0 | 0 |
| [29] | Angola | 21000000 | 2017 | 401 | 00 | 0 | 1 | 62 | 0 | 0 |
| [30] | Burkina Faso | 19280000 | 2017 | 9029 | 141 | 18 | ۲ 12 | 00 101 | 12 | 0 |
| [3] [22] | Equat | 24213022 | 2017 | 1421 | 322 | 2 | 13 | 07 | 70 | 0 |
| [32] | Konva | 101500000 | 2017 | 1537 | 806 | 1 | 0 | 97 | 0 | 0 |
| [34] | Sudan | 40680000 | 2017 | 90 | 90 | 2 | 0 | 0 | 0 | 0 |
| [35] | Cameroon | 24390000 | 2017 | 114 | 8 | 0 | 5 | 0 | 0 | 0 |
| [36] | Ethiopia | 108E+08 | 2017 | 101 | 101 | 1 | 0 | 15 | Õ | 0 |
| [37] | Cameroon | 24390000 | 2017 | 791 | 86 | 0 | 5 | 16 | 8 | C C |
| [38] | Mali | 19310000 | 2017 | 429 | 33 | 0 | 0 | 0 | 0 | 0 |
| [39] | Cameroon | 24390000 | 2017 | 629 | 2 | 0 | 2 | 0 | 0 | 0 |
| [40] | Mauritania | 4160000 | 2017 | 307 | 165 | 0 | 0 | 104 | 0 | 0 |
| [41] | Seychelles | 95843 | 2017 | 4068 | 1429 | 0 | 0 | 0 | 0 | 0 |

 Table 1. Epidemiological trends of dengue cases and deaths in Africa over the last decade (2013–2023, n= 87)

| Reference | Country | Population | Year of | Suspected | Confirmed | Deaths | Serotype | | | |
|-----------|--------------------------|------------|----------|-----------|-----------|--------|----------|-------|-------|-------|
| | | | outbreak | cases | cases | | DENV1 | DENV2 | DENV3 | DENV4 |
| [42] | Senegal | 15570000 | 2018 | 2981 | 342 | 0 | 0 | 0 | 0 | 0 |
| [43] | Mauritania | 42710000 | 2018 | 322 | 28 | 0 | 0 | 28 | 0 | 0 |
| [44] | Tanzania | 58090000 | 2018 | 226 | 37 | 0 | 0 | 27 | 0 | 0 |
| [45] | Ghana | 30637585 | 2018 | 150 | 4 | 0 | 1 | 0 | 3 | 0 |
| [46] | Nigeria | 198E+08 | 2018 | 130 | 11 | 0 | 5 | 0 | 6 | 0 |
| [47] | Reunion | 856942 | 2018 | 6770 | 951 | 6 | 0 | 951 | 0 | 0 |
| [48] | Senegal | 15570000 | 2018 | 198 | 17 | 0 | 0 | 4 | 11 | 0 |
| [49] | Senegal | 15570000 | 2018 | 832 | 224 | 0 | 6 | 35 | 103 | 0 |
| [50] | Seychelles | 96762 | 2018 | 6120 | 1511 | 0 | 0 | 0 | 0 | 0 |
| [51] | Kenya | 48950000 | 2019 | 660 | 286 | 0 | 0 | 0 | 0 | 0 |
| [52] | Benin | 12290444 | 2019 | 26 | 14 | 2 | 0 | 0 | 0 | 0 |
| [53] | Côte d'Ivoire | 26150000 | 2019 | 2919 | 302 | 2 | 95 | 28 | 0 | 0 |
| [54] | Tanzania | 59870000 | 2019 | 6917 | 5286 | 13 | 0 | 0 | 0 | 0 |
| [52] | Ethiopia | 114E+08 | 2019 | 1251 | 6 | 0 | 0 | 0 | 0 | 0 |
| [52] | Mali | 20570000 | 2019 | 20 | 9 | 0 | 0 | 0 | 0 | 0 |
| [55] | Senegal | 16000000 | 2019 | 6 | 1 | 0 | 0 | 0 | 0 | 0 |
| [56] | Sudan | 43230000 | 2019 | 265 | 145 | 0 | 35 | 10 | 10 | 0 |
| [57] | Sudan | 43230000 | 2019 | 100 | 23 | 0 | 0 | 0 | 23 | 0 |
| [58] | Reunion | 861200 | 2019 | 30 | 16 | 0 | 16 | 0 | 0 | 0 |
| [59] | Sudan | 43230000 | 2019 | 76 | 17 | 2 | 0 | 0 | 0 | 0 |
| [60] | Tanzania | 59870000 | 2019 | 191 | 20 | 0 | 20 | 0 | 0 | 0 |
| [61] | Mauritius | 1296279 | 2019 | 265 | 141 | 0 | 136 | 5 | 0 | 0 |
| [62] | Cameroon | 25780000 | 2019 | 310 | 14 | 0 | 0 | 0 | 0 | 0 |
| [63] | Sudan | 43230000 | 2019 | 395 | 67 | 0 | 0 | 0 | 0 | 0 |
| [64] | Comoros | 806166 | 2020 | 696 | 4 | 0 | 4 | 0 | 0 | 0 |
| [65] | Mauritania | 4615000 | 2020 | 7 | 2 | 0 | 0 | 0 | 0 | 0 |
| [66] | Cameroon | 27200000 | 2020 | 320 | 41 | 0 | 2 | 8 | 28 | 0 |
| [67] | Nigeria | 213996181 | 2020 | 82 | 11 | 0 | 11 | 0 | 0 | 0 |
| [68] | Senegal | 17763163 | 2021 | 86 | 27 | 0 | 0 | 0 | 0 | 0 |
| [69] | Kenya | 53542175 | 2021 | 867 | 36 | 2 | 0 | 0 | 0 | 0 |
| [70] | Angola | 34500000 | 2021 | 86 | 38 | 0 | 0 | 0 | 0 | 0 |
| [71] | Côte d'Ivoire | 27480000 | 2021 | 4 | 4 | 0 | 0 | 0 | 0 | 0 |
| [72] | Côte d'Ivoire | 28200000 | 2022 | 11 | 11 | 1 | 0 | 0 | 0 | 0 |
| [73] | Kenya | 54027487 | 2022 | 2426 | 68 | 2 | 0 | 0 | 0 | 0 |
| [74] | Sao Tome and Principe | 227380 | 2022 | 1150 | 1150 | 8 | 0 | 0 | 0 | 0 |

| Reference | Country | Population | Year of | Suspected | Confirmed | Deaths | | Serot | уре | |
|-----------|--------------------------|------------|----------|-----------|-----------|--------|-------|-------|-------|-------|
| | | | outbreak | cases | cases | | DENV1 | DENV2 | DENV3 | DENV4 |
| [75] | Niger | 26207977 | 2022 | 1 | 1 | 0 | 0 | 0 | 0 | 0 |
| [74] | Senegal | 17763163 | 2022 | 196 | 169 | 0 | 0 | 0 | 0 | 0 |
| [76] | Chad | 18278568 | 2023 | 1581 | 41 | 1 | 0 | 0 | 0 | 0 |
| [77] | Côte d'Ivoire | 28873034 | 2023 | 3895 | 321 | 27 | 0 | 0 | 0 | 0 |
| [78] | Burkina Faso | 23251485 | 2023 | 146878 | 68346 | 688 | 0 | 0 | 0 | 0 |
| [78] | Senegal | 17763163 | 2023 | 203 | 203 | 0 | 0 | 0 | 0 | 0 |
| [78] | Mali | 23293698 | 2023 | 4427 | 629 | 29 | 0 | 0 | 0 | 0 |
| [78] | Cape Verde | 573007 | 2023 | 410 | 193 | 0 | 0 | 0 | 0 | 0 |
| [76] | Angola | 37174067 | 2023 | 3 | 3 | 0 | 0 | 0 | 0 | 0 |
| [76] | Sudan | 48667653 | 2023 | 1664 | 1664 | 7 | 0 | 0 | 0 | 0 |
| [78] | Ethiopia | 127E+08 | 2023 | 14249 | 127 | 7 | 0 | 0 | 0 | 0 |
| [76] | Sao Tome and Principe | 233931 | 2023 | 69 | 69 | 3 | 0 | 0 | 0 | 0 |
| [79] | Togo | 9053799 | 2023 | 8 | 2 | 1 | 0 | 0 | 0 | 0 |
| [76] | Egypt | 113E+08 | 2023 | 578 | 578 | 0 | 0 | 0 | 0 | 0 |
| [78] | Mauritius | 1300557 | 2023 | 265 | 265 | 0 | 0 | 0 | 0 | 0 |
| [79] | Guinea | 14190612 | 2023 | 6 | 6 | 1 | 0 | 0 | 0 | 0 |
| [78] | Niger | 27202843 | 2023 | 148 | 148 | 0 | 0 | 0 | 0 | 0 |
| [78] | Nigeria | 224E+08 | 2023 | 72 | 14 | 0 | 0 | 0 | 0 | 0 |
| [78] | Ghana | 34121985 | 2023 | 18 | 9 | 0 | 0 | 0 | 0 | 0 |
| [78] | Benin | 13712828 | 2023 | 6 | 3 | 1 | 0 | 0 | 0 | 0 |
| [80] | Burkina Faso | 23251485 | 2023 | 408 | 56 | 0 | 18 | 0 | 38 | 0 |
| Total | | | | 245,143 | 90,053 | 857 | 542 | 2,551 | 440 | 11 |

Travel Medicine and Infectious Diseases

1 3.3 The spatial distribution of suspected dengue cases in Africa

- 2 Spatial distribution analysis indicates differences in number and distribution of
- 3 suspected dengue cases across African countries and territories. From 2013-2023,
- 4 Burkina Faso, Ethiopia and Tanzania reported the highest number of cases,
- 5 surpassing 10,000. Reunion and Seychelles reported the highest number of cases
- 6 (5,001—10,000) among territories (**Figure 2**).





9 Figure 2. A map of Africa illustrating the geographical distribution of suspected
10 dengue cases in various African countries and territories based on data reported
11 from 2013-2023. The map was developed using QGIS open-source software version
12 3.38 accessed at https://qgis.org/download/

13

14 **3.4** A rise in the number of confirmed dengue cases in West Africa

Travel Medicine and Infectious Diseases

- 15 Over the past decade, dengue disease burden has increased in Africa, with a 5-fold
- 16 increase in West Africa, from approximately 14,000 in 2014 to 70,000 cumulative
- 17 confirmed cases in 2023. The region was responsible for approximately 80% of the
- 18 confirmed cases (71, 793/89,967) (**Figure 3**).
- 19



20

Figure 3: A. The magnitude and trend of cumulative confirmed dengue cases across Africa. B. The number of suspected dengue cases per 100,000 population and fatality rates (%) for specific countries based on data reported from 2013 to 2023. Burkina Faso, and Sao Tome and Principe recorded over 500 cases per 100,000 population surpassing all other countries (Fig 3B). Sao Tome and Principe, Sudan and Mali reported higher fatality rates of 0.9%, 0.7% and 0.6% respectively.

27 **3.5** The occurrence of multiple DENV serotypes and severe dengue

Since 2013, all four serotypes of DENV, DENV1, DENV2, DENV3, and DENV4)

- 29 have been reported in Africa. Continentally, DENV1 and DENV2 dominated at
- 30 different time (Figure 4). From 2019 to 2020, DENV1 was predominate serotype in

Travel Medicine and Infectious Diseases

31 both Eastern and Western Africa. In 2023, DENV3 dominated Western Africa



- 32 whereas DENV2 prevailed in Eastern Africa.
- 33





Since 2013, a few cases of severe dengue, specifically Dengue Haemorrhagic Fever (DHF) and Dengue Shock Syndrome (DSS), have been reported in Africa. Twenty cases were reported in the United Republic of Tanzania (2014), nine in Burkina Faso (2015), five in Ethiopia (2017), two in Benin (2019) and 40 in Sudan (2019). Severe outcomes were associated with diabetes in Tanzania [11], pregnant women in Burkina Faso [19], male gender in Ethiopia [36] and malaria-dengue co-infection in Sudan [81].

Travel Medicine and Infectious Diseases

45 **3.6 Regional differences in transmission seasonality and suitability**

| 46 | Central (Figure 5A) and Eastern (Figure 5B) Africa experienced prolonged dengue |
|----|--|
| 47 | transmission seasons from April to November. In Central Africa, high peaks were |
| 48 | observed in June (> 400 cases), September (> 1,000) and November (> 700) with a |
| 49 | monthly median of 94 cases. Eastern Africa exhibited a dynamic transmission |
| 50 | pattern with the highest peaks in May and June (> 4,000), September and November |
| 51 | (> 8,000, respectively) with a monthly median of 88 cases. Northern Africa (Figure |
| 52 | 5C exhibited sporadic transmission patterns with high peaks in February, July, |
| 53 | August and November (>200, respectively) with a monthly median of 93 cases. |
| 54 | Western Africa (Figure 5D) had distinct high transmission seasons with high peaks |
| 55 | in October (> 17,000), November (> 50,000) and December (30,000) and a monthly |
| 56 | median of 98 cases while Southern Africa reported less than 100 cases. |
| 57 | Western Africa reported the highest number of deaths (> 500), followed by Eastern |
| 58 | Africa (> 40). In Western Africa, deaths were mostly reported during the high |
| 59 | transmission season between October and December, whereas in Eastern and |
| 60 | Central Africa, deaths were reported during both the low and high transmission |
| 61 | seasons. There have been no reported fatalities in Northern Africa. |

Travel Medicine and Infectious Diseases

62



Figure 5. The seasonality of dengue transmission across regions based on data
available from 2017–2023. A. Central Africa, B. Eastern Africa, C. Northern Africa
and D. Western Africa.

66 Central and Western Africa experience persistent suitability (index P > 1) for dengue 67 transmission between April and November. Eastern Africa exhibits two phases of 68 transmission suitability (**Figure 6B**), that coincide with short and long rainy seasons 69 from October to December and March to May, respectively, whereas Northern Africa 70 exhibits transmission suitability between August to November (**Figure 6C**). 71

Travel Medicine and Infectious Diseases



72

Figure 6. Dengue transmission potential (TP) across regions based on data
available from 2017–2023. A. Central Africa B. Eastern Africa C. Northern Africa
and D. Western Africa.

76

3.7 Increasing trend in the number of predicted dengue cases across Africa
The negative binomial model predicted a rising trend in the number of dengue cases
in all African regions for each passing year (Figure 7) with the growth rate exceeding
30% in West Africa (Table 2). There were limited cases from Southern Africa that
could be included in the model.

Travel Medicine and Infectious Diseases



83

84 Figure 7. Predicted dengue cases across regions based on a negative binomial

85 model using data reported from 2013–2023. A. Central Africa B. Eastern Africa, C.

- 86 Northern Africa and D. Western Africa.
- 87

Table 2. Growth of predicted dengue cases in Africa regions according to negative binomial model using data available from 2013–2023.

| Region | Year coefficient () | Multiplicative factor (|) | Growth rate (%)/year |
|-----------------|----------------------|-------------------------|---|----------------------|
| Western Africa | 0.33 | 1.39 | | 39 |
| Eastern Africa | 0.19 | 1.21 | | 21 |
| Central Africa | 0.14 | 1.15 | | 15 |
| Northern Africa | 0.10 | 1.11 | | 11 |
| | | | | |

90

91 **4. Discussion**

92 Dengue represents a significant public health threat in Africa. However, the non-

93 specific clinical presentation of the disease, which resembles malaria and other

94 febrile illnesses such as yellow fever and chikungunya, limits better detection,

95 reporting and understanding of the disease burden. Additionally, there are limited

Travel Medicine and Infectious Diseases

| 96 | health resources for surveillance and timely detection [82]. This review presents the |
|-----|--|
| 97 | first systematic analysis to define the epidemiological trends of dengue disease |
| 98 | burden in Africa and associated territories from 2013 to 2023. The findings of this |
| 99 | review can inform the strengthening of intervention strategies to reduce morbidity |
| 100 | and mortality of dengue in Africa. |
| 101 | The spatial analysis reveals disparities in the quantity and distribution of suspected |
| 102 | dengue cases across Africa (Figure 2). This observation may indicate gaps in |
| 103 | epidemiological surveillance and case reporting, given that tropical regions in Africa |
| 104 | shares similar vector ecologies and transmission indices. Overall, West Africa was |
| 105 | responsible for more than two-thirds of confirmed dengue cases and one-third of |
| 106 | surveillance reports, indicating increasing transmission activities and improved case |
| 107 | reporting in this region. |
| 108 | Burkina Faso recorded the highest burden of cases per 100,000 population (Figure |
| 109 | 3). In 2023, the country accounted for more than 80% of confirmed cases and |
| 110 | deaths. These estimates are consistent with the World Health Organization's |
| 111 | surveillance report on health emergency situations [83]. The impact of climate |
| 112 | hazards on the distribution of vectors and extensive international travels of infected |
| 113 | individuals from endemic countries are likely to exacerbate dengue transmission |
| 114 | through spillover events and introductions, respectively [84,85]. |
| 115 | There are several probable factors that may have contributed to dengue |
| 116 | transmission in Western Africa over the past decade. Several studies have |
| 117 | demonstrated that the abundance of Aedes aegypti breeding habitats, particularly |
| 118 | waste tyres, was a significant factor for transmission, particularly in peri urban |
| 119 | centres [86,87]. Other studies have revealed that dengue outbreaks in Western |
| | |

- 120 Africa are driven by the presence of abundant infected Aedes vectors [88]. In 2023,
- 121 Ouédraogo and others reported the presence of a high number of immature Aedes
- 122 *aegypti* vectors in the handwashing stations that were constructed in public areas
- 123 during the COVID-19 pandemic in Burkina Faso [89].
- 124 Despite a significant increase in the number of dengue cases, case fatality rate
- remains below 1% in Africa (Figure 3), compared to 3%–10% in Asia [90]. Given the
- 126 continuous circulation of multiple DENV serotypes within the same region (Figure 4),
- 127 more cases of severe dengue were expected due to lack of cross immunity.
- 128 However, it is possible that limited diagnostic capabilities, and under-reporting due to
- 129 misdiagnosis with other febrile illnesses, such as malaria are contributing to low
- 130 prevalence [91]. Further, genetic evidence from global ancestral analysis suggests
- 131 that African descendants may be protective against dengue haemorrhagic
- 132 phenotype [92].
- 133 The seasonality of dengue transmission in Africa shows regional differences (Figure
- 134 **5**), with Central and Eastern Africa experiencing long transmission seasons that
- 135 coinciding with rainy seasons from May to November in Central Africa (Figure 5A)
- and from April to May and November to December in Eastern Africa (Figure 5B).
- 137 The erratic transmission pattern observed in Northern Africa (Figure 5C), may be
- 138 attributed to various factors including the storage of water in open containers [93,94]
- and heavy rainfall [95], that attract Aedes mosquitoes. Moreover, there are inter-
- 140 regional migrations of people from endemic countries and extensive intra-regional
- 141 trade activities that play a significant role in the transmission of dengue [93]. Western
- 142 Africa exhibits a distinct high transmission season between October and December.

Travel Medicine and Infectious Diseases

143 These observations are consistent with the results from previous studies conducted144 in this region [96,97].

145 From April to November, Central and Western Africa (Figure 6A and 6D), exhibit 146 long transmission suitability periods (Index P > 1.0) that correlate with the annual 147 rainfall seasons [98,99]. In contrast, high transmission potential in Eastern Africa 148 (Figure 6B), correlates with two rainy seasons from September to December and 149 from March to May [100]. The low number of dengue cases reported in Northern 150 Africa could be due to a short transmission suitability period between August to 151 October. In general, West Africa experiences the highest transmission suitability 152 (Index P > 2.0) between July and November in comparison to other regions. The 153 prolonged period of transmission suitability may have contributed to the rise of 154 Western Africa as a hotspot of dengue transmission. 155 The negative binomial model predicted an increasing trend of suspected dengue

cases across Central, Eastern and Western Africa (Figure 7), with a growth rate
exceeding 30% in West Africa (Table 2). Results from previously described climate
suitability models indicate that these regions will experience significant growth in
dengue incidences over the coming decades [101]. These findings will aid to inform
healthcare policy and practices in Africa to enhance surveillance and implement
effective interventions to prevent ongoing dengue transmission.

162 Limitations

The results of this review are subject to several limitations. First, confirmed case
counts may be underestimated due to the application of different case definitions.
Second, limited dengue surveillance and case reporting between 2020–2022 period
due to COVID-19 pandemic introduces bias in estimating dengue burden. Third,

Travel Medicine and Infectious Diseases

- 167 confirmed cases among travellers returning from African countries and territories
- 168 were not included. Therefore, dengue burden estimates reported in this report should
- 169 be interpreted with caution.
- 170

171 **5.** Conclusions

- 172 Over the past decade, there has been a rise in the number of confirmed dengue
- 173 cases, particularly in West Africa. The persistent presence of multiple DENV
- 174 serotypes within the same region increases the likelihood of severe dengue due to
- the lack of cross-immunity. It is important to strengthen surveillance and implement
- 176 region-specific interventions to prevent future dengue outbreaks. We advocate
- 177 further research for understanding the evolution and transmission dynamics of the
- 178 specific dengue virus lineages in Africa.
- 179

180 **Funding**

181 Research activities at KRISP and CERI are supported in part by grants from the 182 Rockefeller Foundation (HTH 017), the Abbott Pandemic Defense Coalition (APDC), 183 the National Institute of Health USA (U01 AI151698) for the United World Antivirus 184 Research Network (UWARN), the SAMRC South African mRNA Vaccine Consortium 185 (SAMVAC), Global Health EDCTP3 Joint Undertaking and its members as well as 186 Bill & Melinda Gates Foundation (101103171), the Health Emergency Preparedness 187 and Response Umbrella Program (HEPR Program), managed by the World Bank 188 Group (TF0B8412), the UK's Medical Research Foundation (MRF-RG-ICCH-2022-189 100069), and the Wellcome Trust for the Global health project (228186/Z/23/Z). The 190 content and findings reported herein are the sole deduction, view and responsibility

Travel Medicine and Infectious Diseases

- 191 of the researcher/s and do not reflect the official position and sentiments of the
- 192 funding agencies.
- 193 Institutional review board statement
- 194 No ethical clearance because this is a review of published research.
- 195 Informed consent statement
- 196 Not applicable
- 197
- 198

Data availability statement

- 200 All the relevant data are contained within the manuscript. Any additional data is
- 201 made available through Mendeley data repository accessible at
- 202 https://data.mendeley.com/my-data/.
- 203 Authors' contributions
- 204 Gaspary O. Mwanyika: Conceptualization, data curation, formal analysis,
- 205 methodology, validation, writing-original draft, writing-reviewing & editing.
- Abdualmoniem O. Musa: Data curation and writing-reviewing & editing. Jenicca
- 207 **Poongavanan**: Methodology, writing-reviewing & editing. **Monika Moir**:
- 208 Methodology, Writing-reviewing & editing. **Graeme Dor**: Writing-review & editing.
- 209 Eduan Wilkinson: Writing-reviewing & editing. Cheryl Baxter: Methodology, project
- administration, writing-original draft, writing-reviewing & editing. Tulio de Oliveira:
- 211 Resources, funding acquisition, project administration, validation, writing-reviewing &
- editing. Houriiyah Tegally: Methodology, data curation, validation, writing-original
- 213 draft and writing-reviewing & editing.
- 214 Competing interest

Travel Medicine and Infectious Diseases

215 The authors declare no competing interest in this work.

216 Acknowledgements

- 217 We acknowledge the World Health Organization for Africa (#WHOAfro) and Africa
- 218 Centers for Disease Control and Prevention (#AfricaCDC) that made epidemiological
- 219 surveillance reports for different countries accessible. We also express our gratitude
- to all the members of the Climate Amplified Disease Epidemics (#CLIMADE)
- 221 consortium for their valuable contributions in this review.

222

223 Supplementary materials

- 224 Mwanyika, Gaspary (2024), "S1.Advanced search strategy for PubMed/Medline
- databases", Mendeley Data, V1, <u>https://data.mendeley.com/datasets/zf7x7wy7xg/1</u>
 226
- Mwanyika, Gaspary (2024), "S2. Excluded articles for specific reasons.", Mendeley
 Data, V1, <u>https://data.mendeley.com/datasets/3gjypt9yvs/1</u>
- 229

Mwanyika, Gaspary (2024), "S3. PRISMA 2020 checklist", Mendeley Data, V1,
 <u>https://data.mendeley.com/datasets/hx4r4jywmb/1</u>

- 232
- Mwanyika, Gaspary (2024), "S4. R programming codes ", Mendeley Data, V1, https://data.mendeley.com/datasets/d9kyhgsx2m/1
- 235
- 236 Mwanyika, Gaspary (2024), "S5. Transmission potential (Index P) data", Mendeley
- 237 Data, V1, <u>https://data.mendeley.com/datasets/cv9c4gzdy9/1</u>
- 238

239 References

- [1] Weaver SC, Reisen WK. Present and future arboviral threats. Antiviral Res
 2010;85:328–45. doi: 10.1016/j.antiviral.2009.10.008.
- [2] Murray NEA, Quam MB, Wilder-Smith A. Epidemiology of dengue: past, present
 and future prospects. Clin Epidemiol 2013;5:299–309.
- 244 https://doi.org/10.2147/CLEP.S34440.
- [3] Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, et al. The
 global distribution and burden of dengue. Nature 2013;496:504–7.
 https://doi.org/10.1038/pature12060
- 247 https://doi.org/10.1038/nature12060.
- [4] Riccò M, Peruzzi S, Balzarini F, Zaniboni A, Ranzieri S. Dengue Fever in Italy:
 The "Eternal Return" of an Emerging Arboviral Disease. Trop Med Infect Dis
 249 Density of the set of the set
- 250 2022;7:10. https://doi.org/10.3390/tropicalmed7010010.

| 251 | [5] Ryan SJ, Carlson CJ, Mordecai EA, Johnson LR. Global expansion and |
|-----|---|
| 252 | redistribution of Aedes-borne virus transmission risk with climate change. PLoS |
| 253 | Negl Trop Dis 2019;13:e0007213. https://doi.org/10.1371/journal.pntd.0007213. |
| 254 | [6] Amarasinghe A, Kuritsk JN, Letson GW, Margolis HS. Dengue virus infection in |
| 255 | Africa. Emerg Infect Dis 2011;17:1349–54. |
| 256 | https://doi.org/10.3201/eid1/08.101515. |
| 257 | [7] Eltom K, Enan K, El Hussein ARM, Elkhidir IM. Dengue Virus Infection in Sub- |
| 258 | Sanaran Africa Between 2010 and 2020: A Systematic Review and Meta- |
| 259 | Analysis. Front Cell Infect Microbiol 2021;11:678945. |
| 200 | 1111ps.//doi.org/10.3309/101110.2021.070945. |
| 201 | al Dengue virus infection and associated risk factors in Africa: a systematic |
| 262 | al. Deligue virus infection and associated fisk factors in Africa. a systematic review and meta-analysis. Viruses 2021:13:536 |
| 203 | [9] Selcuk AA A Guide for Systematic Reviews: PRISMA Turk Arch |
| 265 | Otorhinolaryngol 2019:57:57–8. https://doi.org/10.5152/tao.2019.4058 |
| 265 | [10] Nakase T. Giovanetti M. Obolski U. Lourenco J. Global transmission suitability |
| 267 | maps for dengue virus transmitted by Aedes aegypti from 1981 to 2019. Sci Data |
| 268 | 2023: 10:275. https://doi.org/10.1038/s41597-023-02170-7. |
| 269 | [11] Boillat-Blanco N, Klaassen B, Mbarack Z, Samaka J, Mlaganile T, Masimba J, |
| 270 | et al. Dengue fever in Dar es Salaam, Tanzania: clinical features and outcome in |
| 271 | populations of black and non-black racial category. BMC Infect Dis 2018;18:644. |
| 272 | https://doi.org/10.1186/s12879-018-3549-z |
| 273 | [12] Ellis EM, Neatherlin JC, Delorey M, Ochieng M, Mohamed AH, Mogeni DO, et |
| 274 | al. A household serosurvey to estimate the magnitude of a dengue outbreak in |
| 275 | Mombasa, Kenya, 2013. PLoS Negl Trop Dis 2015;9:e0003733. |
| 276 | https://doi.org/10.1371/journal.pntd.0003733. |
| 277 | [13] World Health Organization (WHO). Dengue outbreak in the United Republic of |
| 278 | Tanzania (Situation as of 30 May 2014). United Republic of Tanzania: World |
| 279 | Health Organization, https://reliefweb.int/report/united-republic-tanzania/dengue- |
| 280 | outbreak-united-republic-tanzania-situation-30-may-2014. ; 2014 [accessed 29 |
| 281 | November 2023j. |
| 282 | [14] Anmed A, All Y, Elmagboul B, Monamed O, Elduma A, Bashab H, et al. Dengue |
| 283 | https://doi.org/10.2201/oid2511.191766 |
| 204 | 11105.//001.019/10.3201/elu2311.101700. |
| 205 | [15] Valid F, Modela LE, De Naldo F, Onyo Nivi, Meschi S, Rumisha SF, et al. Clinical virologic and epidemiologic characteristics of dengue outbreak. Dar es |
| 280 | Salaam Tanzania 2014 Emerg Infect Dis 2016:22:895 doi: |
| 288 | 10.3201/eid2205.151462 |
| 289 | [16] Shah MM, Ndenga BA, Mutuku FM, Vu DM, Grossi-Soyster FN, Okuta V, et al. |
| 290 | High dengue burden and circulation of 4 virus serotypes among children with |
| 291 | undifferentiated fever, Kenya, 2014–2017. Emerg Infect Dis 2020;26:2638. |
| 292 | doi: 10.3201/eid2611.200960. |
| 293 | [17] Onyedibe K. A cross sectional study of dengue virus infection in febrile patients |
| 294 | presumptively diagnosed of malaria in Maiduguri and Jos plateau, Nigeria. |
| 295 | Malawi Med J 2018;30:276–82. doi: 10.4314/mmj.v30i4.11. |
| 296 | [18] Ngoi CN, Price MA, Fields B, Bonventure J, Ochieng C, Mwashigadi G, et al. |
| 297 | Dengue and chikungunya virus infections among young febrile adults evaluated |

for acute HIV-1 infection in coastal Kenya. PLoS One 2016;11:e0167508. doi:

[19] Sondo KA, Ouattara A, Diendéré EA, Diallo I, Zoungrana J, Zémané G, et al.

Travel Medicine and Infectious Diseases

10.1371/journal.pone.0167508.

298

299

300

301 Dengue infection during pregnancy in Burkina Faso: a cross-sectional study. 302 BMC Infect Dis 2019;19:997. https://doi.org/10.1186/s12879-019-4587-x. 303 [20] Nkenfou CN, Fainguem N, Dongmo-Nguefack F, Yatchou LG, Kameni JJK, 304 Elong EL, et al. Enhanced passive surveillance dengue infection among febrile 305 children: Prevalence, co-infections and associated factors in Cameroon. PLoS Neal Trop Dis 2021;15:e0009316. doi: 10.1371/journal.pntd.0009316. 306 307 [21] World Health Organization (WHO). Dengue– Egypt. Disease Outbreak News. 308 World Health Organization (WHO), https://www.who.int/emergencies/disease-309 outbreak-news/item/12-november-2015-dengue-en; 2015 [accessed 29 November 310 20231. 311 [22] Dieng I, Hedible BG, Diagne MM, El Wahed AA, Diagne CT, Fall C, et al. Mobile 312 Laboratory Reveals the Circulation of Dengue Virus Serotype I of Asian Origin in 313 Medina Gounass (Guediawaye), Senegal. Diagn Basel Switz 2020;10:408. 314 https://doi.org/10.3390/diagnostics10060408. 315 [23] Tarnagda Z, Cissé A, Bicaba BW, Diagbouga S, Sagna T, Ilboudo AK, et al. 316 Dengue Fever in Burkina Faso, 2016. Emerg Infect Dis 2018;24:170-2. 317 https://doi.org/10.3201/eid2401.170973. 318 [24] Proesmans S, Katshongo F, Milambu J, Fungula B, Muhindo Mavoko H, Ahuka-319 Mundeke S, et al. Dengue and chikungunya among outpatients with acute 320 undifferentiated fever in Kinshasa, Democratic Republic of Congo: A crosssectional study. PLoS Negl Trop Dis 2019;13:e0007047. 321 322 https://doi.org/10.1371/journal.pntd.0007047. 323 [25] Hamid Z, Hamid T, Alsedig K, Abdallah T, Elaagip A, Ahmed A, et al. Molecular 324 Investigation of Dengue Virus Serotype 2 Circulation in Kassala State, Sudan. 325 Jpn J Infect Dis 2019;72:58–61. https://doi.org/10.7883/yoken.JJID.2018.267. 326 [26] International Federation of Red Cross and Red Crescent Societies (IFRC). 327 Seychelles: Dengue Outbreak Emergency Plan of Action Final Report DREF 328 Operation n° MDRSC004. International Federation of Red Cross and Red 329 Crescent Societies (IFRC), https://reliefweb.int/report/seychelles/seychelles-dengue-330 outbreak-emergency-plan-action-final-report-dref-operation-n; 2016 [accessed 29 331 November 2023]. 332 [27] World Health Organization (WHO). Dengue Fever – Burkina Faso. Disease 333 Outbreak News. World Health Organization (WHO), 334 https://www.who.int/emergencies/disease-outbreak-news/item/18-november-2016-335 dengue-burkina-faso-en; 2016 [accessed 29 November 2023]. 336 [28] Masika MM, Korhonen EM, Smura T, Uusitalo R, Vapalahti K, Mwaengo D, et 337 al. Detection of dengue virus type 2 of Indian origin in acute febrile patients in 338 rural Kenya. PLoS Negl Trop Dis 2020;14:e0008099. 339 https://doi.org/10.1371/journal.pntd.0008099. 340 [29] Neto Z, Martinez PA, Hill SC, Jandondo D, Thézé J, Mirandela M, et al. 341 Molecular and genomic investigation of an urban outbreak of dengue virus 342 serotype 2 in Angola, 2017–2019. PLoS Negl Trop Dis 2022;16:e0010255. 343 https://doi.org/10.1371/journal.pntd.0010255.

| 344 | [30] World Health Organization (WHO), Dengue – Burkina Faso, Disease Outbreak |
|-----|---|
| 345 | News. World Health Organization (WHO), https://www.who.int/emergencies/disease- |
| 346 | outbreak-news/item/6-november-2017-dengue-burkina-faso-en; 2017 [accessed 29 |
| 347 | November 2023]. |
| 348 | [31] World Health Organization (WHO) Regional Office for Africa. Weekly bulletins |
| 349 | on outbreaks and other emergencies. World Health Organization (WHO), |
| 350 | https://iris.who.int/bitstream/handle/10665/258888/OEW35-268192017.pdf?sequence=1; |
| 351 | 2017 [accessed 29 November 2023]. |
| 352 | [32] El-Kady AM, Osman HA, Alemam MF, Marghani D, Shanawaz MA, Wakid MH, |
| 353 | et al. Circulation of Dengue Virus Serotype 2 in Humans and Mosquitoes During |
| 354 | an Outbreak in El Quseir City, Egypt. Infect Drug Resist 2022;15:2713-21. |
| 355 | https://doi.org/10.2147/IDR.\$360675. |
| 356 | [33] World Health Organization (WHO). Weekly bulletins on outbreaks and other |
| 357 | emergencies. World Health Organization (WHO), |
| 358 | https://iris.who.int/bitstream/handle/10665/258888/OEW35-268192017.pdf?sequence=1; |
| 359 | 2017 [accessed 29 November 2023]. |
| 360 | [34] World Health Organization (WHO). Weekly epidemiological monitor. World |
| 361 | Health Organization (WHO), |
| 362 | https://applications.emro.who.int/docs/epi/2017/Epi_Monitor_2017_10_49.pdf; 2017 [|
| 363 | accessed 29 November 2023]. |
| 364 | [35] Mouiche MMM, Ntumvi NF, Maptue VT, Tamoufe U, Albert B, Ngum Ndze V, et |
| 365 | al. Evidence of Low-Level Dengue Virus Circulation in the South Region of |
| 366 | Cameroon in 2018. Vector Borne Zoonotic Dis Larchmt N 2020;20:314-7. |
| 367 | https://doi.org/10.1089/vbz.2019.2531. |
| 368 | [36] Gutu MA, Bekele A, Seid Y, Mohammed Y, Gemechu F, Woyessa AB, et al. |
| 369 | Another dengue fever outbreak in Eastern Ethiopia-An emerging public health |
| 370 | threat. PLoS Negl Trop Dis 2021;15:e0008992. |
| 371 | https://doi.org/10.1371/journal.pntd.0008992. |
| 372 | [37] Letizia AG, Pratt CB, Wiley MR, Fox AT, Mosore M, Agbodzi B, et al. |
| 373 | Retrospective Genomic Characterization of a 2017 Dengue Virus Outbreak, |
| 374 | Burkina Faso. Emerg Infect Dis 2022;28:1198–210. |
| 375 | https://doi.org/10.3201/eid2806.212491. |
| 376 | [38] World Health Organization (WHO). Weekly bulletins on outbreaks and other |
| 377 | emergencies. In: WHO Regional Office for Africa. Mali, |
| 378 | https://iris.who.int/bitstream/handle/10665/259809/OEW1-2018.pdf?sequence=1; 2018 |
| 379 | [accessed 29 November 2023]. |
| 380 | [39] Mouiche MMM, Ntumvi NF, Maptue VT, Tamoufe U, Albert B, Ngum Ndze V, et |
| 381 | al. Evidence of Low-Level Dengue Virus Circulation in the South Region of |
| 382 | Cameroon in 2018. Vector Borne Zoonotic Dis Larchmt N 2020;20:314–7. |
| 383 | https://doi.org/10.1089/vbz.2019.2531. |
| 384 | [40] World Health Organization (WHO). Outbreaks and other emergencies updates. |
| 385 | In: WHO Regional Office for Africa, https://www.afro.who.int/health-topics/disease- |
| 386 | outbreaks/outbreaks-and-other-emergencies-updates?page=14; 2017 [accessed 29 |
| 387 | November 2023]. |
| 388 | [41] World Health Organization (WHO). World Health Organization. Weekly bulletins |
| 389 | on outbreaks and other emergencies. In: WHO Regional Office for Africa, |

Travel Medicine and Infectious Diseases

| 390 | https://iris.who.int/bitstream/handle/10665/259809/OEW1-2018.pdf?sequence=1; 2017 |
|-----|--|
| 391 | [accessed 29 November 2023]. |
| 392 | [42] World Health Organization (WHO). Weekly bulletins on outbreaks and other |
| 393 | emergencies. In: WHO Regional Office for Africa, |
| 394 | https://iris.who.int/bitstream/handle/10665/277423/OEW52-2228122018.pdf; 2018 |
| 395 | [accessed 29 November 2023]. |
| 396 | [43] World Health Organization (WHO). Weekly bulletins on outbreaks and other |
| 397 | emergencies. In: WHO Regional Office for Africa, |
| 398 | https://iris.who.int/bitstream/handle/10665/278952/OEW01-29122018-\04012019.pdf; |
| 399 | 2018 [accessed 29 November 2023]. |
| 400 | [44] World Health Organization (WHO). Weekly bulletins on outbreaks and other |
| 401 | emergencies. In: WHO Regional Office for Africa, |
| 402 | https://iris.who.int/bitstream/handle/10665/278952/OEW01-29122018-\04012019.pdf; |
| 403 | 2018 [accessed November 29, 2023]. |
| 404 | [45] Bonney JHK, Hayashi T, Dadzie S, Agbosu E, Pratt D, Nyarko S, et al. |
| 405 | Molecular detection of dengue virus in patients suspected of Ebola virus disease |
| 406 | in Ghana. PloS One 2018;13:e0208907. |
| 407 | https://doi.org/10.1371/journal.pone.0208907. |
| 408 | [46] Ayolabi CI, Olusola BA, Ibemgbo SA, Okonkwo GO. Detection of Dengue |
| 409 | viruses among febrile patients in Lagos, Nigeria and phylogenetics of circulating |
| 410 | Dengue serotypes in Africa. Infect Genet Evol J Mol Epidemiol Evol Genet Infect |
| 411 | Dis 2019;75:103947. https://doi.org/10.1016/j.meegid.2019.103947. |
| 412 | [47] Vincent M, Larrieu S, Vilain P, Etienne A, Solet J-L, François C, et al. From the |
| 413 | threat to the large outbreak: dengue on Reunion Island, 2015 to 2018. |
| 414 | Eurosurveillance 2019;24:1900346. https://doi.org/10.2807/1560- |
| 415 | 7917.ES.2019.24.47.1900346. |
| 416 | [48] Gaye A, Ndiaye T, Sy M, Deme AB, Thiaw AB, Sene A, et al. Genomic |
| 417 | investigation of a dengue virus outbreak in Thiès, Senegal, in 2018. Sci Rep |
| 418 | 2021;11:10321. https://doi.org/10.1038/s41598-021-89070-1. |
| 419 | [49] Dieng I, Fall C, Barry MA, Gaye A, Dia N, Ndione MHD, et al. Re-emergence of |
| 420 | dengue serotype 3 in the context of a large religious gathering event in Touba, |
| 421 | Senegal. Int J Environ Res Public Health 2022;19:16912. |
| 422 | [50] World Health Organization (WHO). Weekly bulletins on outbreaks and other |
| 423 | emergencies. In: WHO Regional Office for Africa, |
| 424 | https://iris.who.int/bitstream/handle/10665/27/423/OEW52-2228122018.pdf; 2018 |
| 425 | [accessed 29 November 2023]. |
| 426 | [51] World Health Organization (WHO) Regional Office for Africa. Weekly bulletins |
| 427 | on outbreaks and other emergencies. In: WHO Regional Office for Africa, |
| 428 | https://www.who.int/docs/default-source/who-afro-outbreaks-and-emergencies- |
| 429 | updates/oew15-0814042019.pdf; 2019 [accessed 29 November 2023]. |
| 430 | [52] VVorid Health Organization (VVHO). Weekly bulletins on outbreaks and other |
| 451 | emergencies. In: WHO Regional Office for Affica, |
| 432 | <u>https://fils.wno.int/bitstream/nandie/10005/550555/OEW01-05012020.pdf</u> ; 2020 |
| 455 | [accessed 29 NOVEITIDEF 2023]. |
| 434 | [53] WORD HEARD ORGANIZATION (WHO). WEEKIY DUILETINS ON OUTDREAKS and other |

435 emergencies. In: WHO | Regional Office for Africa,

| 436 | https://iris.who.int/bitstream/handle/10665/326596/OEW34-1925082019.pdf; 2019 |
|-----|---|
| 437 | [accessed 29 November 2023]. |
| 438 | [54] World Health Organization (WHO). Weekly bulletins on outbreaks and other |
| 439 | emergencies. In: WHO Regional Office for Africa, |
| 440 | https://iris.who.int/bitstream/handle/10665/328778/OEW40-300906102019.pdf; 2019 |
| 441 | [accessed 29 November 2023]. |
| 442 | [55] World Health Organization (WHO). Weekly bulletins on outbreaks and other |
| 443 | emergencies. In: WHO Regional Office for Africa, |
| 444 | https://iris.who.int/bitstream/handle/10665/330353/OEW01-05012020.pdf; 2020 |
| 445 | [accessed 29 November 2023]. |
| 446 | [56] Desogi M, Ali M, Gindeel N, Khalid F, Abdelraheem M, Alnaby A, et al. |
| 447 | Detection of dengue virus serotype 4 in Sudan. East Mediterr Health J Rev Sante |
| 448 | Mediterr Orient Al-Majallah Al-Sihhiyah Li-Sharq Al-Mutawassit 2023;29:436–41. |
| 449 | https://doi.org/10.26719/emhj.23.041. |
| 450 | [57] Eldigail MH, Abubaker HA, Khalid FA, Abdallah TM, Musa HH, Ahmed ME, et |
| 451 | al. Association of genotype III of dengue virus serotype 3 with disease outbreak |
| 452 | in Eastern Sudan, 2019. Virol J 2020;17:118. https://doi.org/10.1186/s12985- |
| 453 | 020-01389-9. |
| 454 | [58] Hafsia S, Barbar T, Wilkinson DA, Atyame C, Biscornet L, Bibi J, et al. Genetic |
| 455 | characterization of dengue virus serotype 1 circulating in Reunion Island, 2019- |
| 456 | 2021, and the Seychelles, 2015-2016. BMC Infect Dis 2023;23:294. |
| 457 | https://doi.org/10.1186/s12879-023-08125-y. |
| 458 | [59] Ahmed A, Eldigail M, Elduma A, Breima T, Dietrich I, Ali Y, et al. First report of |
| 459 | epidemic dengue fever and malaria co-infections among internally displaced |
| 460 | persons in humanitarian camps of North Darfur, Sudan. Int J Infect Dis IJID Off |
| 461 | Publ Int Soc Infect Dis 2021;108:513–6. https://doi.org/10.1016/j.ijid.2021.05.052. |
| 462 | [60] Mwanyika GO, Mboera LEG, Rugarabamu S, Makange M, Sindato C, Lutwama |
| 463 | JJ, et al. Circulation of dengue serotype 1 viruses during the 2019 outbreak in |
| 464 | Dar es Salaam, Tanzania. Pathog Glob Health 2021;115:467–75. |
| 465 | https://doi.org/10.1080/20477724.2021.1905302. |
| 466 | [61] World Health Organization (WHO). Weekly bulletins on outbreaks and other |
| 467 | emergencies. In: WHO Regional Office for Africa, |
| 468 | https://iris.who.int/bitstream/handle/10665/326403/OEW21-2026052019.pdf; 2019 |
| 469 | [accessed 29 November 2023]. |
| 470 | [62] Tchuandom SB, Lissom A, Ateba GHM, Tchouangueu TF, Tchakounte C, Ayuk |
| 471 | AR, et al. Dengue virus serological markers among potential blood donors: an |
| 472 | evidence of asymptomatic dengue virus transmission in Cameroon. Pan Afr Med |
| 473 | J 2020;36. doi: 10.11604/pamj.2020.36.185.22128. |
| 474 | [63] World Health Organization (WHO). Weekly bulletins on outbreaks and other |
| 475 | emergencies. In: WHO Regional Office for Africa, |
| 476 | https://iris.who.int/bitstream/handle/10665/332246/OEW22-2531052020.pdf; 2020 |
| 477 | [accessed 30 November 2023]. |
| 478 | [64] World Health Organization (WHO). Weekly bulletins on outbreaks and other |
| 479 | emergencies: Week 22, https://iris.who.int/bitstream/handle/10665/332246/OEW22- |
| 480 | 2531052020.pdf; 2020 [accessed 29 November 2023]. |
| 481 | [65] World Health Organization (WHO) Regional Office for Africa. Weekly bulletins |
| 482 | on outbreaks and other emergencies: Week 26 In: WHO Regional Office for |

Travel Medicine and Infectious Diseases

| 483 | Africa, https://iris.who.int/bitstream/handle/10665/344522/OEW26-2127062021.pdf; |
|-----|--|
| 484 | 2020 [accessed 30 November 2023]. |
| 485 | [66] Simo Tchetgna H. Sado Yousseu F. Kamgang B. Tediou A. McCall PJ. Wondii |
| 486 | CS. Concurrent circulation of dengue serotype 1, 2 and 3 among acute febrile |
| 487 | patients in Cameroon, PLoS Neol Trop Dis 2021:15:e0009860. |
| 488 | https://doi.org/10.1371/journal.pntd.0009860. |
| 489 | [67] Tizhe DT, Kwaga JKP, Nok Kia GS, Serological and Molecular Survey for |
| 490 | Dengue Virus Infection in Suspected Febrile Patients in Selected Local |
| 491 | Government Areas in Adamawa State, Nigeria, Vaccines 2022:10:1407 |
| 492 | https://doi.org/10.3390/vaccines10091407 |
| 493 | [68] World Health Organization (WHO). Weekly bulletins on outbreaks and other |
| 494 | emergencies. In: WHO Regional Office for Africa. |
| 495 | https://iris.who.int/bitstream/handle/10665/350967/OEW01-271202012022.pdf: 2022 |
| 496 | [accessed 30 November 2023]. |
| 497 | [69] World Health Organization (WHO). Weekly bulletins on outbreaks and other |
| 498 | emergencies. In: WHO Regional Office for Africa. |
| 499 | https://iris.who.int/bitstream/handle/10665/342386/OEW27-280604072021.pdf: 2021 |
| 500 | Jaccessed 30 November 2023 |
| 501 | [70] World Health Organization (WHO). Weekly bulletins on outbreaks and other |
| 502 | emergencies. In: WHO Regional Office for Africa. |
| 503 | https://iris.who.int/bitstream/handle/10665/343406/OEW31-260701082021.pdf: 2021 |
| 504 | [accessed 30 November 2023]. |
| 505 | [71] World Health Organization (WHO). Weekly bulletins on outbreaks and other |
| 506 | emergencies. In: WHO Regional Office for Africa. |
| 507 | https://iris.who.int/bitstream/handle/10665/342123/OEW18-2602052021.pdf; 2021 |
| 508 | [accessed 30 November 2023]. |
| 509 | [72] Weekly bulletins on outbreaks and other emergencies. In: WHO Regional |
| 510 | Office for Africa, https://iris.who.int/bitstream/handle/10665/356076/OEW14- |
| 511 | 280303042022.pdf; 2022 [accessed 30 November 2023]. |
| 512 | [73] World Health Organization (WHO). Weekly bulletins on outbreaks and other |
| 513 | emergencies. In: WHO Regional Office for Africa, |
| 514 | https://iris.who.int/bitstream/handle/10665/355799/OEW23-300505062022.pdf; 2022 |
| 515 | [accessed 30 November 2023]. |
| 516 | [74] World Health Organization (WHO). Weekly bulletins on outbreaks and other |
| 517 | emergencies. In: WHO Regional Office for Africa, |
| 518 | https://iris.who.int/bitstream/handle/10665/365480/OEW52-1925120222.pdf; 2022 |
| 519 | [accessed 30 November 2023]. |
| 520 | [75] World Health Organization (WHO) Regional Office for Africa. Weekly bulletins |
| 521 | on outbreaks and other emergencies. In: WHO Regional Office for Africa. Niger, |
| 522 | https://iris.who.int/bitstream/handle/10665/364200/OEW36-290804092022.pdf; 2022 [|
| 523 | accessed 30 November 2023]. |
| 524 | [76] Africa Centres for Disease Control and Prevention. Weekly Event-Based |
| 525 | Surveillance Report. Africa Centres for Disease Control and Prevention (CDC), |
| 526 | https://africacdc.org/download/africa-cdc-weekly-event-based-surveillance-report- |
| 527 | december-2023/; 2023 [accessed 30 November 2023]. |
| 528 | [77] World Health Organization (WHO). Dengue Situation Report 001. Côte d'Ivoire: |
| 500 | |

529 World Health Organization,

| 530 | https://iris.who.int/bitstream/handle/10665/375392/AFRO.Dengue.Sitrep001- |
|-----|---|
| 531 | 20231219.pdf; 2023 [accessed 30 November 2023]. |
| 532 | [78] World Health Organization (WHO). AFRO Dengue Situation Report 001. World |
| 533 | Health Organization, |
| 534 | https://iris.who.int/bitstream/handle/10665/375392/AFRO.Dengue.Sitrep001- |
| 535 | <u>20231219.pdf;</u> 2023 [accessed 22 January 2024]. |
| 536 | [79] Africa Centres for Disease Control and Prevention (CDC). Weekly Event-Based |
| 537 | Surveillance Report. Africa Centres for Disease Control and Prevention, |
| 538 | https://iris.who.int/bitstream/handle/10665/375392/AFRO.Dengue.Sitrep001- |
| 539 | <u>20231219.pdf;</u> 2023 [accessed 22 January 2024]. |
| 540 | [80] Gomgnimbou MK, Belem LRW, Some K, Diallo M, Barro B, Kaboré A, et al. |
| 541 | Utilization of novel molecular multiplex methods for the detection and, |
| 542 | epidemiological surveillance of dengue virus serotypes and chikungunya virus in |
| 543 | Burkina Faso, West Africa. Mol Biol Rep 2024;51:906. |
| 544 | https://doi.org/10.1007/s11033-024-09847-1. |
| 545 | [81] Ahmed A, Eldigail M, Elduma A, Breima T, Dietrich I, Ali Y, et al. First report of |
| 546 | epidemic dengue fever and malaria co-infections among internally displaced |
| 547 | persons in humanitarian camps of North Darfur, Sudan. Int J Infect Dis IJID Off |
| 548 | Publ Int Soc Infect Dis 2021;108:513–6. https://doi.org/10.1016/j.ijid.2021.05.052. |
| 549 | [82] World Health Organization (WHO). Disease Outbreak News; Dengue – Global |
| 550 | situation. World Health Organization, https://www.who.int/emergencies/disease- |
| 551 | outbreak-news/item/2023- |
| 552 | DON498#:~:text=Between%201%20January%202023%20and,of%20the%20Americas% |
| 553 | 2C%20with%2015; 2023 [accessed 05 January 2024]. |
| 554 | [83] World Health Organization (WHO). Weekly bulletins on outbreaks and other |
| 555 | emergencies. World Health Organization, |
| 556 | https://iris.who.int/bitstream/handle/10665/375392/AFRO.Dengue.Sitrep001- |
| 557 | <u>20231219.pdf;</u> 2023 [accessed 05 January 2024]. |
| 558 | [84] Aliaga-Samanez A, Romero D, Murray K, Segura M, Real R, Olivero J. Potential |
| 559 | climate change effects on the distribution of urban and sylvatic dengue and |
| 560 | yellow fever vectors. Pathog Glob Health 2024;118:397–407. |
| 561 | https://doi.org/10.1080/2047724.2024.2369377. |
| 562 | [85] Carlson CJ, Albery GF, Merow C, Trisos CH, Zipfel CM, Eskew EA, et al. |
| 563 | Climate change increases cross-species viral transmission risk. Nature |
| 564 | 2022;607:555–62. doi: 10.1038/s41586-022-04788-w. |
| 565 | [86] Gyasi P, Bright Yakass M, Quaye O. Analysis of dengue fever disease in West |
| 566 | Africa. Exp Biol Med Maywood NJ 2023;248:1850–63. |
| 567 | https://doi.org/10.11/7/15353702231181356. |
| 568 | [87] Badolo A, Somble A, Yameogo F, Wangrawa DW, Sanon A, Pignatelli PM, et al. |
| 569 | First comprehensive analysis of Aedes aegypti bionomics during an arbovirus |
| 570 | outbreak in west Africa: Dengue in Ouagadougou, Burkina Faso, 2016-2017. |
| 571 | PLOS Negl Trop Dis 2022;16:e0010059. |
| 572 | nttps://doi.org/10.13/1/journal.pntd.0010059. |
| 573 | [88] Pappoe-Ashong P, Otosu-Appian L, Mingle J, Jassoy C. Seroprevalence of |
| 5/4 | aengue virus intections in Gnana. East Atr Med J 2018;95:2132–40. |
| 575 | [89] Ouedraogo WM, Zanre N, Rose NH, Zahouli JZB, Djogbenou LS, Viana M, et |
| 5/6 | ai. Dengue vector nabitats in Ouagadougou, Burkina Faso, 2020: an unintended |

- 577 consequence of the installation of public handwashing stations for COVID-19
- 578 prevention. Lancet Glob Health 2024;12:e199–200.
- 579 https://doi.org/10.1016/S2214-109X(23)00565-X.
- [90] Yeh C-Y, Lu B-Z, Liang W-J, Shu Y-C, Chuang K-T, Chen P-L, et al.
 Trajectories of hepatic and coagulation dysfunctions related to a rapidly fatal
 outcome among hospitalized patients with dengue fever in Tainan, 2015. PLoS
 Negl Trop Dis 2019;13:e0007817. doi: 10.1371/journal.pntd.0007817.
- [91] Waggoner JJ, Gresh L, Vargas MJ, Ballesteros G, Tellez Y, Soda KJ, et al.
 Viremia and clinical presentation in Nicaraguan patients infected with Zika virus, chikungunya virus, and dengue virus. Clin Infect Dis 2016:ciw589. https://doi:
 10.1093/cid/ciw589.
- [92] Sierra B, Triska P, Soares P, Garcia G, Perez AB, Aguirre E, et al. OSBPL10,
 RXRA and lipid metabolism confer African-ancestry protection against dengue
 haemorrhagic fever in admixed Cubans. PLoS Pathog 2017;13:e1006220.
- 591 https://doi.org/10.1371/journal.ppat.1006220.
- [93] Seidahmed OME, Hassan SA, Soghaier MA, Siam HAM, Ahmed FTA,
 Elkarsany MM, et al. Spatial and temporal patterns of dengue transmission along
 a Red Sea coastline: a longitudinal entomological and serological survey in Port
 Sudan city. PLoS Negl Trop Dis 2012;6:e1821.
- 596 https://doi.org/10.1371/journal.pntd.0001821.
- 597 [94] Andayi F, Charrel RN, Kieffer A, Richet H, Pastorino B, Leparc-Goffart I, et al. A
 598 sero-epidemiological study of arboviral fevers in Djibouti, Horn of Africa. PLoS
 599 Negl Trop Dis 2014;8:e3299. https://doi.org/10.1371/journal.pntd.0003299.
- [95] Seidahmed OME, Siam H a. M, Soghaier MA, Abubakr M, Osman HA, Abd
 Elrhman LS, et al. Dengue vector control and surveillance during a major
 outbreak in a coastal Red Sea area in Sudan. East Mediterr Health J Rev Sante
 Mediterr Orient Al-Majallah Al-Sihhiyah Li-Sharq Al-Mutawassit 2012;18:1217–
 24. https://iris.who.int/handle/10665/118472.
- [96] Donatien K, Hien YE, Salam S, Yacouba NK, Denise IP, Nikièma AR, et al.
 Seroepidemiological Study of Dengue Virus Infection Suspected Cases in
 Burkina Faso. J Biosci Med 2023;11:47–56. doi: 10.4236/jbm.2023.111006.
- [97] Ouattara CA, Traore S, Sangare I, Traore TI, Meda ZC, Savadogo LGB.
 Spatiotemporal analysis of dengue fever in Burkina Faso from 2016 to 2019.
 BMC Public Health 2022;22:462. https://doi.org/10.1186/s12889-022-12820-x.
- BMC Public Health 2022;22:462. https://doi.org/10.1186/s12889-022-12820-x.
 [98] Klassou KS, Komi K. Analysis of extreme rainfall in Oti river basin (West Africa).
- J Water Clim Change 2021;12:1997–2009. https://doi.org/10.2166/wcc.2021.154.
 [99] Fotso-Nguemo TC, Diallo I, Diakhaté M, Vondou DA, Mbaye ML, Haensler A, et
- al. Projected changes in the seasonal cycle of extreme rainfall events from
- 615 CORDEX simulations over Central Africa. Clim Change 2019;155:339–57. 616 https://doi.org/10.1007/s10584-019-02492-9.
- 617 [100] Mwangi E, MacLeod D, Kniveton D, Todd MC. Variability of rainy season 618 onsets over East Africa. Int J Climatol 2024;44:3357–79.
- 619 https://doi.org/10.1002/joc.8528.
- [101] Sintayehu DW, Tassie N, De Boer WF. Present and future climatic suitability
 for dengue fever in Africa. Infect Ecol Epidemiol 2020;10:1782042.
- 622 https://doi.org/10.1080/20008686.2020.1782042.
- 623