

1 **Variation in Nutritional Composition of Anatomical Parts and Taxonomic Classes of**
2 **Wild Animals: A Systematic Review Using Data Imputation with Artificial Intelligence**

3 Short title: Nutritional Composition of Wildmeat: A Systematic Review.

4 Ana Luisa dos Santos Medeiros^a, Amanda Letícia Bezerra de Oliveira^b, Maria Fernanda
5 Araújo de Medeiros^c, Daniel Tregidgo^d, Eliana Bistriche Giuntini^e, Elias Jacob de Menezes
6 Neto^f, Juliana Kelly da Silva Maia^g, Michelle Cristine Medeiros Jacob^h

7 ^a *Universidade Federal do Rio Grande do Norte, Avenida Senador Salgado Filho, s/n, Natal,*
8 *RN, Brazil, 59078-970, nutri.analuisa97@gmail.com, ORCID: 0009-0003-3238-5240*

9 ^b *Universidade Federal do Rio Grande do Norte, Avenida Senador Salgado Filho, s/n, Natal,*
10 *RN, Brazil, 59078-970, amanda.Oliveira.102@ufrn.edu.br, ORCID: [0009-0008-5356-5810](https://orcid.org/0009-0008-5356-5810)*

11 ^c *Universidade Federal do Rio Grande do Norte, Avenida Senador Salgado Filho, s/n, Natal,*
12 *RN, Brazil, 59078-970, maria.fernanda.059@ufrn.edu.br, ORCID: 0000-0001-7674-9307*

13 ^d *Grupo de Pesquisa em Ecologia de Vertebrados Terrestres, Instituto de Desenvolvimento*
14 *Sustentável Mamirauá, Tefé, Amazonas, Brazil, dantregidgo@gmail.com, ORCID: 0000-*
15 *0002-5468-831X*

16 ^e *Food Research Center ((FoRC) -Universidade de São Paulo / USP, São Paulo, Brazil,*
17 *elibi@alumni.usp.br, ORCID: 0000-0002-8386-5418*

18 ^f *Metropole Digital Institute, Universidade Federal do Rio Grande do Norte, Avenida*
19 *Senador Salgado Filho, s/n, Natal, RN, Brazil, 59078-970, elias.jacob@ufrn.br, ORCID:*
20 *0000-0002-1153-8899*

21 *§Programa de Pós-graduação em Nutrição / UFRN, Universidade Federal do Rio Grande do*
22 *Norte, Avenida Senador Salgado Filho, s/n, Natal, RN, Brazil, 59078-970,*
23 *juliana.maia@ufrn.br, ORCID:0000-0002-7970-534X*

24 *^hLaboratory of Biodiversity and Nutrition, Universidade Federal do Rio Grande do Norte,*
25 *Avenida Senador Salgado Filho, s/n, Natal, RN, Brazil, 59078-970, michelle.jacob@ufrn.br,*
26 *ORCID: [0000-0002-4881-7285](https://orcid.org/0000-0002-4881-7285), corresponding author*

27

28

29 **ABSTRACT**

30 Wildmeat is crucial for the food security of Indigenous Peoples and Local Communities,
31 yet information about its nutritional profile remains scarce. This study systematically
32 reviewed the impact of anatomical part and taxonomic class (mammals, birds, reptiles)
33 on the nutritional composition of wildmeat. Using the PRISMA protocol, we selected
34 articles from Web of Science, Scopus, and Medline/PubMed databases, with criteria
35 including original articles on wild animal meat composition consumed by humans,
36 excluding studies presenting secondary data or lacking detailed methodologies. We
37 employed a quality questionnaire and concordance analysis (Fleiss' Kappa = 1.00) for
38 robustness. Artificial intelligence techniques (eg., K-Nearest Neighbors) estimated
39 missing nutritional values in all 21 articles included in our study, covering 26 species and
40 10 nutrients. Results show statistically significant nutritional variations between
41 anatomical parts and animal classes. Reptile viscera have over twice the fat content and
42 triple the iron compared to muscles. Mammal viscera contain five times more omega-6
43 and selenium, four times more iron and manganese, and almost double the zinc compared
44 to muscles. Among classes, bird muscles have over 90% higher fat content than mammal

45 muscles and 20% higher than reptile muscles. Mammals have over 100% higher zinc
46 levels than birds, and reptiles have over 400% more selenium than birds. No significant
47 difference in iron content between mammals and birds was noted, likely due to bird
48 slaughter methods. This study highlights the importance of wildmeat for food security.
49 Importantly, we demonstrate an enormous variation in nutritional composition,
50 underscoring how different anatomical parts and taxonomic classes can contribute to
51 tackling different nutritional deficiencies. Additionally, it introduces a novel
52 methodology for handling missing nutritional composition data, providing a
53 comprehensive approach to understanding the nutritional value of wildmeat. Our findings
54 can inform food security policies and wildlife management, balancing conservation and
55 subsistence.

56 Keywords: Wildmeat, Nutritional Composition, Food Security, Artificial Intelligence,
57 Data Imputation, Systematic Review

58 **1. INTRODUCTION**

59 Meat, as a fundamental component of human diets, stands out for its diversity and
60 nutritional value, offering an abundant source of essential nutrients for health. In this context,
61 meat from wild animals, known as wildmeat, bushmeat or game meat, emerges as a
62 fundamental resource for various populations around the world, especially Indigenous Peoples
63 and Local Communities (IPLC). In this study, we consider wildmeat to be derived from wild,
64 non-domesticated animals, (1) particularly vertebrates, excluding insects, crustaceans, larvae,
65 mollusks, and fish. Wildmeat is a crucial resource for ensuring food access for populations
66 within traditional food systems (2–4). It is estimated that approximately 230 to 833 million
67 people rely on wild animal meat as a source of protein worldwide (5). Furthermore, its
68 consumption is associated with improved nutritional status and cognitive development (6). For

69 instance, an investigation in the Amazon region found a correlation between wildmeat
70 consumption and children's health, showing an average increase in hemoglobin concentration
71 of 0.25 g/dL in children from vulnerable rural areas (7). Similarly, a study in Madagascar
72 revealed that families most reliant on wildmeat, who are also the most economically
73 disadvantaged, have a four-fold higher likelihood of developing anemia if they lose access to
74 wildlife (8) .

75 Despite progress in recognizing the importance of wildmeat for food security and
76 health, significant gaps remain in understanding its nutritional composition, particularly
77 regarding micronutrients. Few studies have specifically evaluated the nutritional composition
78 of wild species consumed (9), and existing nutritional composition tables, such as the Brazilian
79 Food Composition Table, (10) often do not distinguish between wild and domesticated species.
80 Researchers frequently use food matching to estimate the nutritional composition of foods with
81 unavailable direct data, (11) a method critical for assessing products derived from biodiversity.
82 However, this approach may introduce inaccuracies if the data is based on domestic animals,
83 potentially misrepresenting wild species' nutritional profiles (12).

84 Current literature provides limited and scattered data on the nutritional composition
85 of wild animals, (9,13,14) hindering a comprehensive understanding of nutritional variations.
86 Therefore, our systematic review aims to identify patterns in the nutritional composition of
87 wildmeat, compiling data and applying artificial intelligence techniques to address missing
88 values (15). This approach offers a detailed analysis of wild animal meat's nutritional profile,
89 addressing ethical challenges in studying wild species.

90 Currently, the scientific literature presents, albeit limited, data on the nutritional
91 composition of wild animals (9,13,14). However, this information is scattered across various
92 articles and diverse fields of knowledge. This dispersion and scarcity of data significantly

93 hinder our comprehensive understanding of nutritional variations in wild animals. In light of
94 this, we propose the systematic review we conducted to identify patterns in the nutritional
95 composition of wildmeat. By compiling this data and applying artificial intelligence techniques
96 to address missing values, (15) we provide a detailed analysis of the nutritional profile of
97 wildmeat. This approach represents a viable strategy in the face of the ethical challenges
98 involved in analyzing wild species, considering that our database contained many gaps due to
99 the scarcity of data in the literature, likely caused by legal and ethical limitations in sample
100 collection (16).

101 Different tissues, such as viscera and muscles, have distinct nutritional profiles due
102 to their specific physiological functions (17). Additionally, animal class may determine
103 nutritional composition variations influenced by physiological and environmental factors (2,9).
104 In this sense, to identify nutritional patterns in wildmeat composition, we hypothesized that the
105 anatomical part (viscera vs. muscles) and the animal class (mammals, birds, reptiles) influence
106 meat's nutritional composition.

107 **2. MATERIALS AND METHODS**

108 The data used to create the *corpus* for this project were gathered through a systematic
109 review conducted according to the Preferred Reporting Items for Systematic Reviews and
110 Meta-Analyses (PRISMA) guidelines, [(18)] see Supplementary Material 1. The protocol for
111 this review was not registered in advance, as the research does not directly analyze any health-
112 related outcomes.

113 **2.1 Selection criteria and search**

114 To address the research question posed in this project, "How does the nutritional
115 composition of anatomical parts and taxonomic classes of wild animals vary?", we selected
116 articles based on the following eligibility criteria: (i) original articles published in any language,

117 with no date restrictions, that (ii) provided data on the composition of wild animal meat
118 consumed by human populations. We also included articles recommended by experts in the
119 field, provided they met the eligibility criteria. We excluded studies that (i) did not provide
120 nutritional composition data of wild animals, (ii) presented secondary data, or (iii) did not detail
121 the methodology for food composition analysis.

122 In June 2022, we conducted searches in the Web of Science, Scopus, and
123 Medline/PubMed databases (via the National Library of Medicine). The search involved
124 applying specific descriptors in each database. The following strategy guided the search:
125 (BUSHMEAT OR “WILD MEAT” OR “GAME MEAT” OR “INDIGENOUS MEAT” OR
126 “WILD ANIMALS” OR "HUNTING ANIMALS MEAT") AND (“FOOD CONSUMPTION”
127 OR “FOOD INTAKE” OR DIETARY) AND (MICRONUTRIENT OR NUTRIENT OR
128 NUTRITION OR “FOOD COMPOSITION” OR MINERAL).

129 **2.2 Study selection**

130 We used the Rayyan tool to organize records and remove duplicates identified during
131 the search. Three authors (ALSM, MFAM, ALBO) independently screened the articles,
132 applying the eligibility criteria. During the initial screening, titles and abstracts that did not
133 meet the inclusion criteria were excluded. In the next stage, full-text reviews were conducted,
134 and studies presenting secondary data, incomplete works, or not meeting the selection criteria
135 were excluded.

136 In cases of discrepancies or uncertainties, two authors (MCMJ, JKSM) were consulted.
137 Subsequently, potentially eligible articles were read in full. After the reading and selection
138 process, the chosen articles were stored in the Zotero reference management software.

139 **2.3 Data extraction**

140 Three authors (ALSM, MFAM, ALBO) independently extracted data from the selected
141 articles to ensure transcription accuracy. Initially, we organized article identification data and
142 information on the method of food composition analysis used in each study to determine the
143 comparability of the available nutritional information. Additionally, we extracted data on
144 research variables of interest, namely: taxonomy, animal class, analyzed part, and levels of
145 macronutrients and micronutrients.

146 The nutrients analyzed, given the scope of the study (e.g., excluding heavy metal
147 analysis) and data availability, include the following: iron, selenium, zinc, potassium,
148 magnesium, manganese, sodium, proteins, fat, and omega-6. Nutritional composition data were
149 converted into grams per 100 grams, and all levels were standardized on a wet weight basis.
150 Articles presenting data on a dry weight basis were individually converted and calculated using
151 moisture information described in the methodology of the respective articles. It is noteworthy
152 to mention that we did not have access to the raw data from the original analyses conducted by
153 the authors of the papers reviewed, as this information was not provided.

154 **2.4 Quality analysis**

155 To date, there are no specific quality questionnaires for research on the nutritional
156 composition of wild animals in the scientific literature. Creating such a tool is essential,
157 considering the ethical and legal complexities associated with studying wildmeat (19). These
158 complexities include the use of opportunistic sampling and the need for rigorous control over
159 potential conflicts of interest in the studies analyzed. To address this gap, we developed a
160 quality questionnaire aimed at assessing the methodological robustness of these studies.

161 The assessment tool consisted of a checklist with nine items. The development of this
162 instrument followed an integrative approach, incorporating elements from existing protocols
163 such as LatinFoods/FAO, "QUADAS Tool/Timmer's Analysis Tool," OHAT, Cochrane,

164 QUADAS, Timmers, and STROBE, and adapting them to meet the specific needs of our
165 research. For each item, we crafted a corresponding question that a well-reported study should
166 address, covering aspects such as objectives, sample size, study design, methods, results,
167 statistical analysis, funding sources, conflicts of interest, and incomplete data. Structured
168 responses and scoring were developed for each question: 0 points indicate that information is
169 not available, 0.5 points denote ambiguity or unclear information, and 1.0 point signifies that
170 the information is clearly presented or not applicable (see Supplementary Material 2).

171 The tool was tested by three trained evaluators (ALSM, MFAM, and ALBO), who used
172 the instrument to independently assess the quality of an article. Subsequently, the results from
173 each evaluator were discussed with the research team to refine the clarity of the instrument.
174 With the tool finalized, three trained evaluators independently assessed the articles using the
175 checklist. The checklist data were computed individually, and an average score was calculated
176 from the evaluators' assessments. The agreement between the results was measured using Fleiss'
177 Kappa. Articles were classified as low quality if they received between 0 to 2.9 points, medium
178 quality if they received between 3.0 to 5.9 points, and high quality if they received more than
179 6.0 points.

180 **2.5 Data preparation**

181 Table 1 lists the amount of missing data for each nutrient column, considering the total
182 of 76 observations in the dataset.

183 **Table 1.** Overview of available and missing data for each nutrient analyzed

Nutrients	Fe	Mn	Se	Zn	K	Mg	Na	Protein	Fat	w6
Missing	51	54	63	46	61	61	63	22	26	57

Available	25	22	13	30	15	15	13	54	50	19
Total	76	76	76	76	76	76	76	76	76	76

184

185 To address these gaps, we used machine learning to impute missing data. The goal
186 is to frame each missing value as a regression problem, estimating missing values to minimize
187 their impact on analysis results. This method expands the use of the original data, by allowing
188 its use on modeling steps that would not be possible with the missing value (20,21).

189 Generally, there are three main types of missing data: Missing Completely at
190 Random (MCAR), Missing at Random (MAR), and Missing Not at Random (MNAR). MCAR
191 denotes cases where the probability of a value being missing does not depend on the data itself
192 or any other observed variables. In contrast, MAR means that the probability of missingness is
193 related to other observed attributes. Finally, MNAR arises when the probability of a value being
194 missing is explicitly dependent on the missing value itself (22).

195 In the context of our dataset, we consider the absence of values in the nutritional
196 data as Missing Completely at Random (MCAR), indicating that the missing data are unrelated
197 to the dataset itself. This assumption seems reasonable, as our literature review revealed that
198 the scarcity of data is primarily attributed to the limited availability of studies on game meat
199 analysis. This is likely justified by the legal and ethical limitations in collecting samples (16),
200 rather than any specific bias or underlying structural cause. The missing values are likely a
201 result of general research limitations such as funding constraints or legal and ethical constraints
202 on sample collection. We tested several methods to impute our data, including:

203 1. *Multiple Imputation by Chained Equations* (MICE): This method
204 operates iteratively, imputing missing values with estimates derived from regression
205 models using observed data.

206 2. *K-Nearest Neighbors* (KNN): This technique estimates missing values
207 based on the average of the k nearest neighbors, considering the distance between
208 variables in the dataset.

209 3. *Soft Impute*: In this approach, missing values are estimated through a
210 low-rank matrix approximation using a soft thresholding algorithm.

211 4. *Matrix Factorization*: This technique decomposes the matrix into low-
212 dimensional components, which are then used to approximate missing values within the
213 matrix.

214 5. *Nuclear Norm Minimization*: This method involves minimizing the
215 nuclear norm of the matrix, resulting in a low-rank approximation of the original matrix
216 with complete data.

217 6. *Iterative Singular Value Decomposition* (SVD): In this approach, an
218 iterative algorithm is employed to estimate missing values, progressively refining the
219 approximation using SVD.

220 We evaluated the best technique among these mentioned using a cross-validation
221 approach for data imputation methods. This approach involves intentionally removing known
222 values from the dataset, treating them as missing, and then attempting to impute them using
223 various techniques. It is important to note that our imputation process considered the entire
224 dataset, not just individual columns. For example, when imputing missing values in the iron
225 (Fe) column, the process took into account all available data across all nutrient columns. This

226 multivariate approach allows the imputation methods to leverage relationships and patterns
227 between different nutrients, potentially improving the accuracy of the estimates. This way we
228 can capture complex interactions between nutrients that might be missed if we only looked at
229 each nutrient in isolation. This process allows us to compare the performance of each
230 imputation method by estimating the reliability of the imputed values against the actual known
231 values. The steps to execute this evaluation method are as follows:

- 232 1. Selecting a known value from the dataset and masking it, treating it as a missing value.
- 233 2. Apply each imputation technique separately to estimate the masked value.
- 234 3. Store the imputed values obtained from each method for further analyses.
- 235 4. Repeat steps 1-3 for all available values in all columns of the dataset.
- 236 5. Once the imputation process is complete for all values and methods, calculate the
237 chosen evaluation metrics by comparing the original values with those imputed using
238 each of the different imputation methods.

239 For the final step, the assessment of imputation, we used three main metrics:
240 Symmetric Mean Absolute Percentage Error (SMAPE) and Mean Absolute Error (MAE). MAE
241 is useful for assessing the overall performance of the imputation technique, providing a
242 measure of how closely imputed values approximate the actual values. SMAPE quantifies the
243 relative accuracy between observed and imputed values by calculating the average of absolute
244 differences normalized by the average absolute values, subsequently expressed as a percentage.
245 The formula for SMAPE is as follows:

$$246 \quad SMAPE = \frac{1}{n} \sum_{i=1}^n \frac{|y_i - \hat{y}_i|}{(|y_i| + |\hat{y}_i|)} \times 100\%$$

247 Where:

- 248 • y_i is the actual value of the variable

- 249 • \hat{y}_i is the predicted value of the variable
- 250 • n is the number of observations

251 Finally, the MAE (Mean Absolute Error) calculates the average of the absolute
252 differences between observed and imputed values, providing a direct and interpretable
253 representation of the average error magnitude. The formula for MAE is given by:

254

$$255 \quad MAE = \frac{1}{n} \sum_{i=1}^n |y_i - \hat{y}_i|$$

256 In summary, our methodology includes a thorough analysis of missing data
257 characteristics. We classified the pattern of absence and employed a systematic execution of
258 multiple imputation techniques to address the limitation caused by missing values. The scarcity
259 of nutritional information for wildmeat, identified in our literature review, presents a
260 compelling case to treat missing nutrition data as MCAR. This assumption allows for the use
261 of imputation methods, which, as shown in previous research, can enhance the accuracy and
262 reliability of estimates derived from incomplete datasets.

263 All data imputation analyses were conducted using the Python programming
264 language. Upon completion of the imputation process, nutrient values were expressed in units
265 of measure as per the Dietary Reference Intakes. [(23)] The code required to replicate our results
266 is publicly available on GitHub. [(24)]

267 **2.6 Variables related to hypothesis testing**

268 The nutritional composition data of 10 nutrients served as our dependent variable. For
269 hypothesis 1, where we tested the influence of the anatomical part on nutritional composition,

270 the independent variables were muscle and viscera. For hypothesis 2, the independent variables
271 were the animal classes: mammals, birds, and reptiles.

272 We recognize that factors such as animal anatomy (25) cause of death, (13) geographical
273 location and seasonality, (26,27) physical activity level, (28) and physiological and metabolic
274 characteristics (9) can affect meat's nutritional composition. However, we lack access to these
275 data for our research because they are inconsistently reported in the analyzed studies and are
276 only occasionally mentioned when investigating these specific influences.

277 **2.7 Data Analysis**


278 We used descriptive statistics to summarize our findings and assessed the normality
279 of numerical data using the Shapiro test. Based on the data characteristics, we conducted either
280 ANOVA or Kruskal-Wallis tests, setting a significance level of 0.05 (p-value) to determine the
281 significance of the results. In cases of significant differences, we performed post hoc tests such
282 as Tukey or Bonferroni. All statistical analyses were conducted using the R programming
283 language through the RStudio interface.

284 **3. RESULTS**

285 **3.1 Study Selection**

286 After searching the database, we obtained a total of 565 articles (Web of Science:
287 68, Medline/PubMed: 325, Scopus: 172). After excluding 142 duplicates, 423 articles
288 proceeded to title and abstract screening. Following this step, we selected 75 articles for full-
289 text review. At this stage, 62 articles were excluded for presenting secondary data (5 articles),
290 being incomplete (56 articles), or lacking nutrient composition data relevant to our selection (1
291 article). Ultimately, we included 13 articles for review. Additionally, we incorporated 8 external

292 articles, recommended by experts, which met all our criteria. Therefore, a total of 21 articles
293 constitute our study. Figure 1 illustrates the selection process and flowchart.



*Consider, if feasible to do so, reporting the number of records identified from each database or register searched (rather than the total number across all databases/registers).

**If automation tools were used, indicate how many records were excluded by a human and how many were excluded by automation tools.

From: Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71. doi: 10.1136/bmj.n71. For more information, visit: <http://www.prisma-statement.org/>

294

295 **Figure 1.** Flowchart of the study selection process.

296 **3.2. Characteristics of the studies**

297 The systematic review analyzed 26 distinct animal species, including 13 mammalian
298 species, six bird species, and seven reptile species (see Supplementary Material 4). The studies
299 covered sample collections from 11 countries: Italy, Germany, South Africa, Poland, the United
300 States of America, New Zealand, Spain, Canada, Latvia, Peru, and Brazil. Italy, Spain, Poland,
301 and South Africa were prominently featured for their recurring analyses of wildmeat. For a
302 comprehensive understanding of the studies included in this review, Table 2 presents their main
303 characteristics.

304 **Table 2.** Characterization of studies on the nutritional composition of wild animals. Names of
305 animals collected are presented exactly as provided in the papers. Scientific names were
306 updated for the analysis; for updated names, see Supplementary Material 4.

n	Publication	Study objective	Key findings	Country of collection	Wild animals	Samples	Analyses performed	Study quality(29)
1	Aguiar JPL. Tabela de composição de alimentos da Amazônia. ACTA AMAZONICA 26. 1996;121–6.	Analyze samples of wild animal meat obtained from public markets in the Amazon.	Analyses of the samples' composition were conducted. The results were compiled into a centesimal composition table, and a summary of the findings is available for consultation in this article.	Brazil	Cabeçudo (<i>Peltocephalus dumerilianus</i>); catitu (<i>Pecari tajacu</i>); cutia (<i>Dasyprocta leporina</i>); Iaçá (<i>Podocnemis sextuberculat</i>); jacaretinga (<i>Caiman crocodilus</i>); paca (<i>Agouti paca</i>); tartaruga (<i>Podocnemis expansa</i>); tracajá	31 (muscle) + 4 (viscera)	Protein: Method Kjeldahl - AOAC (1984); Lipid: Soxhlet Method	High (6.50)(30)

					<i>(Podocnemis unifilis);</i> <i>veado (Mazama americana)</i>			
2	Amici A, Cifuni GF, Contò M, Esposito L, Failla S. Hunting area affects chemical and physical characteristics and fatty acid composition of wild boar (<i>Sus scrofa</i>) meat. Rendiconti Lincei. 1o de novembro de 2015;26(3):527–34.	To examine how different geographic hunting areas influence the chemical composition and quality indices of wild boar meat.	The characteristics of wild animal meat varied due to environmental factors, including diet. Dietary diversity appears to influence aspects of the meat, such as color parameters, cooking losses, dry matter content, protein levels, and fatty acid profiles. These differences were observed in relation to	Italy	Wild boar (<i>Sus scrofa</i>)	48 (Longissimus thoracis - muscle)	Protein: Method Kjeldahl - AOAC (1995); Lipids: Folch et al. (1957); Lipid Fraction: Method Soxhlet - IUPAC (1992).	Medium (5.50)(31)

			the hunting area.					
3	Milczarek A, Janocha A, Niedziałek G, Zowczak-Romanowicz M, Horoszewicz E, Piotrowski S. Health-Promoting Properties of the Wild-Harvested Meat of Roe Deer (<i>Capreolus capreolus</i> L.) and Red Deer (<i>Cervus elaphus</i> L.) [Internet]. 2021 [citado 5 de setembro de 2024]. Disponível em:	Compare the proximate nutrient composition of roe deer and wild deer meat.	Regarding the macronutrient profile, no differences were observed with respect to sex, except for the fat content in females, which was higher than in males regardless of the species. As for micronutrients, the levels found are in accordance with the recommended values.	Poland	Roe deer (<i>Capreolus capreolus</i>), Red deer (<i>Cervus elaphus</i>)	60 (semimembranosus muscles)	Protein: Kjeldahl (990.03); Lipids: Method Soxhlet (991.36); Lipid fractions: GC-FID; Methodology; Minerals: (Cd, Pb, Fe) PN-EN 14084	High (8.33)

	https://www.mdpi.com/2076-2615/11/7/2108							
4	Dannenberger D, Nuernberg G, Nuernberg K, Hagemann E. The effects of gender, age and region on macro- and micronutrient contents and fatty acid profiles in the muscles of roe deer and wild boar in Mecklenburg-Western Pomerania (Germany). <i>Meat Sci.</i> 2013;94(1):39–46.	Investigate the effects of gender, age, and region on the macronutrients, micronutrients, and fatty acid profiles of wild boar and deer muscles.	The protein content in deer remained constant regardless of region, age, or sex; however, wild boar meat varied across these factors. Fat in deer varied by region, age, and sex, while in wild boar, they remained stable. Selenium and iron were not influenced by region, age, or sex, except for higher iron levels in older females.	Germany	Roe deer (<i>Capreolus capreolus</i>), Wild boar (<i>Sus scrofa</i>)	203 (longissimus muscle)	Proteins and lipids: Meat Analyzer FoodScan™ (FOSS Analytic, Hillerod, Denmark); Lipid fractions: Gas chromatography analysis; Minerals: ICP-MS (Se, Cu, Fe, and Zn)	High (6.50)

			Zinc concentrations varied according to region, age, and sex.					
5(3 2)	Fernandes HR, Deliza R, Neto OC, Silva CM, Albuquerque NI de, Martins TR, et al. Effect of high hydrostatic pressure on the meat of collared peccaries (<i>Tayassu tajacu</i>) with different ages. Afr J Food Sci. 30 de setembro de 2022;16(9):215–25.	Evaluate the effects of high hydrostatic pressure on the physicochemical parameters of collared peccary meat.	High hydrostatic pressure tenderized the meat of adult collared peccaries and influenced meat quality. The results also showed changes in the protein profiles of animals of different ages subjected to different pressure levels.	Brazil	Peccary (<i>Pecari tajacu</i>)	40 (muscle)	Proteins: Method Kjeldahl - AOAC (1995); Lipids: Method Soxhlet - AOAC (1995); Lipid fraction: Bligh and Dyer (1959)	High (6.50)(33)

6	<p>Gálvez C. H, Arbaiza T, Carcelén C. F, Lucas A. O. Revista de Investigaciones Veterinarias del Perú. 1999 [citado 5 de setembre de 2024].</p> <p>VALOR NUTRITIVO DE LAS CARNES DE SAJINO (<i>Tayassu tajacu</i>), VENADO COLORADO (<i>Mazama americana</i>), MAJAZ (<i>Agouti paca</i>) Y MOTELO (<i>Geochelone denticulata</i>).</p>	<p>Determine the nutritional characteristics of wild animal meat.</p>	<p>Wild meats exhibited high protein content and low fat levels.</p>	<p>Peru</p>	<p>Sajino (<i>Tayassu pecari</i>), Motelo (<i>Chelonoidis denticulata</i>), Majaz (<i>Cuniculus paca</i>), Venado colorado (<i>Mazama americana</i>)</p>	<p>20 (muscle)</p>	<p>Macronutrients and micronutrients</p> <p>Proteins : Method Kjeldahl - AOAC, 1990;</p> <p>Lipids: Method Soxhlet - .AOAC, 1990</p>	<p>Medium (5.50)</p>
---	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------	----------------------------------------------------------------------	-------------	----------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------

	Disponível em: https://revistasinvestigacion.unmsm.edu.pe/index.php/veterinaria/article/view/6707							
7(34)	Jarzyńska G, Falandysz J. Selenium and 17 other largely essential and toxic metals in muscle and organ meats of Red Deer (<i>Cervus elaphus</i>) — Consequences to human health. <i>Environ Int.</i> 1o de julho de 2011;37(5):882–8.	To determine the content and composition of micronutrients in red deer, relating them to meat intake rates, human nutritional needs, and associated health impacts.	The meats from muscles and organs of red deer exhibited high concentrations of copper, chromium, cobalt, manganese, selenium, and zinc. Additionally, they contained toxic trace elements such as mercury, lead, cadmium, and titanium.	Poland	Red Deer (<i>Cervus elaphus</i>)	60 (20 muscle; 20 liver 20 de kidney)	Minerals: ICP-AES (Se) , ICP-MS (Ag, Ba, Cd, Co, Cr, Cs, Cu, Ga, Mn, Mo, Pb, Rb, Sb, Sr, Tl, V e Zn)	Medium (4.50)(35)

			The cadmium levels found in the meats exceed tolerance limits.					
8	Johnson HE, Bleich VC, Krausman PR. Mineral deficiencies in tule elk, Owens Valley, California. J Wildl Dis. janeiro de 2007;43(1):61–74.	Assess the mineral content of horn, liver and forage to measure deficiencies and toxicities.	The levels of minerals found in the liver and horn varied between animals from two different locations. It was found that antler breakage in elk at one of the locations was related to a copper and phosphorus deficiency. The levels of micronutrients found in forage varied according to location and season.	United States of America	Tule elk (<i>Cervus elaphus</i>)	240 (liver)	Minerals: Flame atomic absorption spectrophotometry (Cu e Mo); ICP-MS (Ca, Fe, Mg, Mn, S, P, Zn, Cu e Mo)	Medium (5.50)(36)

9	<p>Lima AT.</p> <p>Caracterização Fisico Quimica da Tartaruga da Amazônia de água proveniente de cativeiro e de habitat natural do estado do Amazonas.</p> <p>Universidade Federal do Amazonas; 2009.</p>	<p>To evaluate the yield and chemical composition of <i>Podocnemis expansa</i> raised in commercial captivity and from natural habitats.</p>	<p>The meat evaluated in both systems showed a low fat content. And the free-living ones had a higher protein content.</p>	Brazil	Tartaruga-da-amazônia (<i>Podocnemis expansa</i>)	45 (muscle) + 15 (viscera)	<p>Proteins : Method Kjeldahl ; Lipid (Bligh e Dyer,1959)</p>	High (8.50)(37)
10	<p>Lorenzo JM, Maggiolino A, Gallego L, Pateiro M, Serrano MP, Domínguez R, et al.</p> <p>Effect of age on nutritional properties</p>	<p>Investigating the effects of slaughter age on the levels of macro and microminerals in wild deer meat.</p>	<p>Total fat and fraction contents were affected by the slaughter age, showing higher values in older animals. Regarding cholesterol, its levels decreased with</p>	Spain	Red deer (<i>Cervus elaphus</i>)	150 (muscle)	<p>Lipid: Bligh e Dyer; Fraction lipid analysis: Gas chromatography; Minerals: ICP-OES, Ca,</p>	High (8.50)

	of Iberian wild red deer meat - Journal of the Science of Food and Agriculture [Internet]. 2019 [citado 5 de setembro de 2024]. Disponível em: https://scijournals.onlinelibrary.wiley.com/doi/abs/10.1002/jsfa.9334		increasing slaughter age.				K, Mg, Na, P, Fe, Mn, Zn e Cu.	
11	Cawthorn DM, Hoffman LC. The bushmeat and food security nexus: A global account of the	Determine the proximate and fatty acid composition of zebra meat.	The muscles had high protein content and low fat content. In the fat group, high concentrations of	South Africa	Zebra (<i>Equus quagga</i>)	20 (longissimus lumborum muscle, longissimus thoracis et	Protein: Method Dumas - AOAC 992.15; Lipid: LEE;	High (7.33)(38)

	contributions, conundrums and ethical collisions. Food Res Int. 1o de outubro de 2015;76:906–25.		saturated fatty acids were found.			lumborum)	Fraction lipid analysis: Gas chromatography;	
12	Rudman M, Leslie AJ, van der Rijst M, Hoffman LC. Quality characteristics of Warthog (<i>Phacochoerus africanus</i>) meat. Meat Sci. 1o de novembro de 2018;145:266–72.	Evaluate the sensorial, physical and chemical attributes of wild boar meat.	Analysis of the sensory profile demonstrated that this meat has an undesirable aroma and flavor, described as sour, which was found in the meat of wild boars of all ages. Regarding macronutrient recommendations, these	South Africa	Warthogs (<i>Phacochoerus africanus</i>)	31 (Longissimus lumborum)	Protein: Method Dumas - 992.15; Lipids: LEE; Ashes (AOAC, 1992; AOAC, 2002a, 2002b)	High (8.0)

			meats were considered healthy sources of protein and polyunsaturated fatty acids.					
13(39)	Neto JV, Bressan MC, Faria PB, Vieira JO e, Santana MTA, Kloster M. Composição centesimal e colesterol da carne de jacaré-do-pantanal (Caiman yacare Daudin 1802) oriundo de zoológico e habitat natural. Ciênc E	Determine the proximate composition and cholesterol of meat from the tail and back of alligators, from zoo breeding facilities and natural habitats.	Animals bred in captivity had a lower amount of fat and higher protein levels when compared to animals in their natural habitat.	Brazil	Alligator-swampland (<i>Caiman yacare</i>)	12 (muscle)	Proteins : Method Kjeldahl - AOAC,1990; Lipids: Method Soxhlet - AOAC, 1990.	High (8.50)

	Agrotecnologia. agosto de 2006;30:701–6.							
14	Landi N, Ragucci S, Di Giuseppe AM, Russo R, Poerio E, Severino V, et al. Nutritional profiling of Eurasian woodcock meat: chemical composition and myoglobin characterization. J Sci Food Agric. 2018;98(13):5120–8.	Evaluate the nutritional aspects of Eurasian chicken meat.	Urian chicken meat contains high concentrations of high- quality proteins, due to its amino acid composition and low fat content.	Italy	Eurasian woodcock (<i>Scolopax rusticola</i>)	10 (entire chest and leg muscle, without skin)	Proteins : Method Kjeldahl; lipids: Method Soxhlet - AOAC 991.36; Lipid fractions: GC- FID	High (7.00)(41)

	(40)							
15	Pérez-Peña PE, Riveros-Montalván MS, Vargas-Arana G, Soria FD, Chumbe JV, Baca YB. Consumo, microbiología y bromatología de la carne silvestre durante la COVID-19 en Iquitos, Perú. Cienc Amaz Iquitos.	Know the microbiological and bromatological status of bush meat during the COVID-19 pandemic from two public markets in the Amazon.	Game meat had a low fat content and a high percentage of protein. The calcium content found in wild meat was higher than in domesticated meat. Phosphorus content was also significant in wild meat.	Peru	<i>Pecari tajacu</i> , <i>Tayassu pecari</i> , <i>Cuniculus paca</i> , <i>Chelonoidis denticulatus</i>	16 (muscle)	Protein:Kjeldahl; Fat: Soxhlet; Vitamins: Chromatography high pressure liquid (HPLC); Minerals: (zinc, sodium, potassium and	High (6.50)(26)

	2021;9(2):51–68.						iron) by Absorption Spectroscopy Atomic (EAA) and Phosphorus by UV Visible Spectroscopy	
16	Serrano MP, Maggiolino A, Landete-Castillejos T, Pateiro M, Barbería JP, Fierro Y, et al. Quality of main types of hunted red deer meat obtained in Spain compared to farmed	Assess the combined impact of country of origin and type of slaughter and season on the quality and nutritional value of venison.	Wild deer from Spain had a higher protein concentration than meat from animals raised in New Zealand. The calcium, sodium and phosphorus levels found in meat do not differ between countries. New	Spain and New Zealand	Red deer (<i>Cervus elaphus</i>)	24 (Longissimus thoracis et lumborum)	Lipids: LEE; Protein: DUMAS	High (9.0)

	<p>venison from New Zealand. Sci Rep. 22 de julho de 2020;10(1):12157.</p>		<p>Zealand farm deer had higher concentrations of magnesium, iron, manganese, copper and lower concentrations of zinc when compared to wild deer from Spain. The nutrient profile of meats also varied depending on the season. Meat from winter hunts had higher levels of calcium, sodium, iron and zinc and lower levels of magnesium and phosphorus compared to</p>					
--	----------------------------------------------------------------------------	--	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--	--	--	--	--

			meat from summer hunts.					
17	Sevillano-Caño J, Cámara-Martos F, Aguilar-Luque EM, Cejudo-Gómez M, Moreno-Ortega A, Sevillano-Morales JS. Trace Element Concentrations in Migratory Game Bird Meat: Contribution to Reference Intakes Through a	Determine the nutritional and toxicological content of game bird meat.	Iron and chromium contents were similar in all three bird species studied. Meat from thrush had the highest concentrations of copper. Pigeons and doves had the highest concentrations of zinc. The study concluded that the meat of migratory game birds is	Spain	Woodpigeon (<i>Columba palumbus</i>), Common turtledove (<i>Streptopelia turtur</i>), Thrush (<i>Turdus philomelos</i>)	89 (breast; thigh)	Cu, Fe, Zn: flame atomic absorption spectroscopy (FAAS, Varian SpectraAA – model 50B); Cr, Co: Electrothermal Atomic Absorption Spectrometry	High (6.50)

	Probabilistic Assessment. Biol Trace Elem Res. 1o de outubro de 2020;197(2):651–9.		an excellent dietary source of copper and iron, much better than other types of meat, such as non-wild birds and even red meat.				on an Agilent Model 240Z AA	
18	Spiegelaar N, Martin ID, Tsuji LJS. Indigenous Subarctic Food Systems in Transition: Amino Acid Composition (Including Tryptophan) in Wild-Harvested and Processed Meats. Int J	Determine the amino acid and protein composition of wild meats and processed meats	The wild meats analyzed showed high concentrations of proteins and amino acids.	Canada	Moose (<i>Alces alces</i>), Canada Goose (<i>Branta canadensis</i>), Mallard Duck (<i>Anas platyrhynchos</i>)	25 (muscle)	Proteins : Method Kjeldahl - AOAC 981.10; Amino acids: standardized hydrolysis of NaOH and HCl, and ultra-performance liquid	High (7.50)

	Food Sci. 27 de junho de 2019;2019:e7096416. (42)						chromatograph y	
19(43)	Strazdiņa V, Jemeljanovs A, Šterna V. Nutrition Value of Wild Animal Meat. Proc Latv Acad Sci Sect B Nat Exact Appl Sci. 31 de outubro de 2013;67(4-5):373-7.	Compare the nutritional value of the meat of different types of wild animals with analogous domesticated animals.	Wild meat samples had a higher content of proteins, essential fatty acids, iron and zinc compared to non-wild animals. The highest cholesterol levels were found in wild boar (<i>Sus scrofa</i>).	Latvia	Elk (<i>Alces alces</i>), Deer (<i>Cervus elaphus</i>), Wild boar (<i>Sus scrofa</i>), Roe deer (<i>Capreolus capreolus</i>)	54 (muscle)	Proteins : Method Kjeldahl; Lipids: Method Soxhlet - AOAC 991.36; Lipid fractions: Gas- liquid	Medium (5.00)

							chromatograph y; Cholesterol: Colorimetric Blur; Minerals: ICP- AES/ISO 6869-2002	
20	Zimmerman TJ, Jenks JA, Leslie DM, Neiger RD. Hepatic minerals of white-tailed and mule deer in the southern Black Hills, South Dakota. J Wildl Dis. abril de 2008;44(2):341–50.	Assess hepatic mineral concentrations of deer from different habitats.	The liver mineral levels found differed. The differences were associated with diet, location and reproductive stage.	United States of America	White-tailed deer (<i>Odocoileus virginianus</i>), Mule deer (<i>Odocoileus hemionus</i>)	83 (liver)	Minerais: ICP- AES Al, Sb, As, Ba, B, Cd, Ca, Cr, Co, Cu, Fe, Pb, Mg, Mn, Mo, P, K, Se, Na, S, Tl, Zn	Medium (5.50)

<p>21(44)</p>	<p>Webb EC, Ryssen JBJ van, Erasmus MEA, McCrindle CME. Copper, manganese, cobalt and selenium concentrations in liver samples from African buffalo (<i>Syncerus caffer</i>) in the Kruger National Park. J Environ Monit. 29 de novembro de 2001;3(6):583–5.</p>	<p>Evaluate the levels of micronutrients found in buffalo liver and in different regions.</p>	<p>The results demonstrated that there are significant differences in the amounts of micronutrients available in meat depending on the territory in which the buffalo is located.</p>	<p>South Africa</p>	<p>Buffalo (<i>Syncerus caffer</i>)</p>	<p>311 (liver)</p>	<p>Minerals: ICP-AES copper, manganese and cobalt</p>	<p>High (6.67)</p>
---------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------	-----------------------------------------	--------------------	-------------------------------------------------------	--------------------

309 **3.3 Quality Analysis**

310 During the quality analysis process, the agreement among the assessors was
311 excellent (Fleiss' Kappa = 1.00). No articles were rated as low quality, and 72% received a score
312 indicating high quality (> 6.0 points) (see Supplementary Material 5). The highest ratings were
313 given for procedures used in sample processing, description of analysis methods, and
314 presentation of results with measures such as standard deviation or standard error. On the other
315 hand, the criteria where the articles scored lower pertain to the exclusion of the scientific name
316 of the species, as well as the absence of information on conflicts of interest and sources of
317 funding for the studies.

318 **3.4 General Comments on the Dataset**

319 In the topic of data imputation, the techniques that showed the lowest SMAPE values
320 for each nutrient demonstrated less absolute error and greater consistency, and were therefore
321 selected for imputing the data. SMAPE values ranged from approximately 11.06% (protein) to
322 83.67% (selenium). For more details, see Supplementary Material 6.

323 **3.5 General Comments on the Nutritional Profile of the Meats**

324 All methods used in food composition analyses among the different studies are
325 recognized by the AOAC (Association of Official Analytical Chemists), which supports the
326 comparison of the obtained results (see Supplementary Material 7). The Supplementary
327 Material 8 summarizes the mean and standard deviation values of the nutritional composition
328 of wild animals included in this systematic review. Results of the statistical analyses are
329 available in Supplementary Material 9.

330 Table 3 presents the main results, highlighting the pairs of comparisons where statistical
331 differences were significant. We present the main results of this section by highlighting the

332 following findings: differences in the nutritional composition of muscles among classes of wild
333 animals, differences in the nutritional composition of visceral meat from different classes of
334 wild animals, and differences in nutritional composition between muscle and visceral meats,
335 including a comparison of nutrient profiles between these anatomical parts of wild animals.

336 **Table 3.** Nutritional composition in muscle and viscera of wildmeat, comprising mammals, birds, and reptiles, including original and imputed data.

337 Pairs of comparisons with statistically significant differences are highlighted in black.

Nutrient	Mammal Viscera x Mammal Muscle	Mammal Viscera x Bird Muscle	Mammal Viscera x Reptile Viscera	Mammal Viscera x Reptile Muscle	Muscle Mammal x Muscle Bird	Mammal Muscle x Reptile Viscera	Mammal Muscle x Reptile Muscle	Bird muscle x Reptile Viscera	Bird Muscle x Reptile Muscle	Reptile Viscera x Reptile Muscle
Protein, g	23.26 x 23.15 (>0.05)	23.26 x 23.46 (>0.05)	23.26 x 16.33 (<0.01)	23.26 x 21.34 (<0.01)	23.15 x 23.46 (>0.05)	23.15 x 16.33 (<0.05)	23.15 x 21.34 (>0.05)	23.46 x 16.33 (<0.01)	23.46 x 21.34 (<0.01)	16.33 x 21.34 (>0.05)
Fat, g	1.37 x 1.27 (>0.05)	1.37 x 2.42 (<0.05)	1.37 x 5.7 (<0.05)	1.37 x 2.02 (>0.05)	1.27 x 2.42 (<0.01)	1.27 x 5.7 (<0.01)	1.27 x 2.02 (>0.05)	2.42 x 5.7 (>0.05)	2.42 x 2.02 (<0.05)	5.7 x 2.02 (<0.05)
w6, g	0.24 x 1.37 (<0.01)	1.37 x 0.83 (>0.05)	1.37 x 1.41 (>0.05)	1.37 x 0.5 (<0.01)	0.24 x 0.83 (<0.01)	0.24 x 1.41 (<0.01)	0.24 x 0.5 (>0.05)	0.83 x 1.41 (>0.05)	0.83 x 0.5 (>0.05)	1.41 x 0.5 (>0.05)
Fe, mg	14.51 x 3.21 (<0.01)	4.41 x 14.51 (<0.01)	14.51 x 15.55 (>0.05)	14.51 x 5.39 (<0.01)	3.21 x 4.41 (>0.05)	3.21 x 5.39 (<0.01)	3.21 x 5.39 (<0.01)	3.21 x 15.55 (<0.01)	4.41 x 5.39 (>0.05)	15.55 x 5.39 (<0.01)
K, g	0.13 x 0.21 (<0.05)	0.13 x 0.27 (<0.01)	0.13 x 0.06 (>0.05)	0.13 x 0.21 (>0.05)	0.21 x 0.27 (<0.01)	0.21 x 0.06 (<0.05)	0.21 x 0.21 (>0.05)	0.27 x 0.06 (<0.01)	0.27 x 0.21 (<0.05)	0.06 x 0.21 (>0.05)

Mn, mg	0.39 x 0.08 (<0.01)	0.39 x 0.12 (<0.01)	0.39 x 0.38 (>0.05)	0.39 x 0.14 (>0.05)	0.08 x 0.12 (>0.05)	0.08 x 0.38 (<0.01)	0.08 x 0.14 (<0.01)	0.12 x 0.38 (>0.05)	0.12 x 0.14 (>0.05)	0.38 x 0.14 (>0.05)
Na, g	0.1 x 0.11 (<0.01)	0.11 x 0.1 (>0.05)	0.1 x 0.09 (>0.05)	0.1 x 0.1 (>0.05)	0.11 x 0.1 (>0.05)	0.11 x 0.09 (<0.05)	0.11 x 0.1 (>0.05)	0.1 x 0.09 (>0.05)	0.1 x 0.1 (>0.05)	0.09 x 0.1 (>0.05)
Zn, mg	2.43 x 1.24 (<0.01)	3.33 x 0.86 (<0.01)	3.33 x 2.43 (>0.05)	3.33 x 1.24 (<0.01)	1.76 x 0.86 (<0.01)	1.76 x 2.43 (>0.05)	1.76 x 1.24 (<0.01)	0.86 x 2.43 (<0.01)	0.86 x 1.24 (>0.05)	2.43 x 1.24 (<0.01)
Se, mcg	57.19 x 10.61 (<0.01)	57.19 x 3.45 (<0.01)	57.19 x 81.17 (>0.05)	3.33 x 1.24 (<0.05)	10.61 x 3.45 (>0.05)	10.61 x 81.17 (>0.05)	10.61 x 19.37 (>0.05)	3.45 x 81.17 (<0.01)	3.45 x 19.37 (<0.01)	81.17 x 19.37 (>0.05)

338

339

340

341 **3.6 Nutritional comparison: Nutritional Variation in Muscles of Mammals, Birds, and**
342 **Reptiles**

343 Our results demonstrate significant variations in the nutritional profile of meat
344 depending on the analyzed class (mammal, bird, or reptile).

345 For reptiles, which were all aquatic species, their muscles showed over 60% higher
346 concentrations of iron ($p < 0.01$) and over 75% higher concentrations of manganese ($p < 0.01$)
347 compared to mammals. Additionally, reptile muscles exhibited selenium concentrations that
348 were over 400% higher than those in bird muscles ($p < 0.01$) and nearly 60% higher fat
349 concentrations than those in mammals.

350 Bird muscles stood out for having the highest concentrations of potassium
351 compared to mammals ($p < 0.01$) and reptiles ($p < 0.05$). They also displayed elevated fat
352 concentrations, which were over 90% higher than those found in mammal muscles ($p < 0.01$)
353 and 20% higher than those in reptile muscles ($p < 0.01$). Furthermore, birds showed the highest
354 concentrations of omega-6, surpassing those in mammals by over 200% ($p < 0.01$). There was
355 no statistical difference in iron content between bird and mammal muscles; however, bird
356 muscles had iron concentrations over 30% higher than those in mammals.

357 Mammal muscles exhibited the highest concentrations of zinc, with values over
358 100% higher than those found in bird muscles ($p < 0.01$) and over 40% higher than those
359 recorded in reptile muscles ($p < 0.01$). Mammals also had the lowest fat concentrations, nearly
360 50% lower than in birds and almost 40% lower than in reptiles.

361 **3.7 Nutritional comparison: Nutritional Variation in Viscera of Mammals and Reptiles**

362 When comparing the nutritional content of viscera meat, we found that mammalian
363 viscera had the highest protein concentrations compared to reptiles ($p < 0.01$). Conversely,
364 reptilian viscera stood out for their high fat content, with concentrations over 300% higher than
365 those found in mammals ($p < 0.01$).

366 **3.8 Comparing Anatomical Parts: Nutritional Variation of Macro and Micronutrients in** 367 **Viscera and Muscle of Mammals and Reptiles**

368 Regarding the animal parts, the most frequent variations in macronutrients were related
369 to fat content. We found that reptilian viscera exhibited higher fat content (5.37 g), nearly
370 double ($p < 0.05$), and lower protein content (16.33 g) compared to muscle (2.02 g and 21.34
371 g, respectively). In mammals, the difference was significant only for omega-6, with viscera
372 (1.37 g) showing over 400% higher content than muscles (0.24 g) ($p < 0.01$).

373 Regarding minerals, viscera generally had higher concentrations compared to muscles.
374 Reptilian viscera showed almost 300% more iron ($p < 0.01$) and nearly 200% more zinc ($p <$
375 0.01) than their muscles. In mammals, viscera revealed iron and manganese concentrations over
376 300% higher ($p < 0.01$), selenium levels significantly higher ($p < 0.01$), and zinc concentrations
377 almost 90% higher ($p < 0.01$) than those found in muscles. However, potassium content showed
378 a different pattern, with muscle concentrations over 60% higher than those found in viscera (p
379 < 0.01).

380 **4. DISCUSSION**

381 This study analyzes the nutritional composition patterns found in wild animal meat.
382 Based on these findings, several important conclusions can be drawn: (i) wild animal meats
383 represent significant sources of nutrients, which are often scarce in the diets of populations
384 facing food insecurity; (ii) the viscera of these animals are particularly rich in minerals

385 compared to muscles, similar to what is observed in meats from domesticated animals; and (iii)
386 the nutritional profile, especially in terms of micronutrients, varies among different classes of
387 animals.

388 Considering the interclass variation in both viscera and muscles, it is important to note
389 that, although variations were observed, it is not possible to determine from the available data
390 whether these variations are attributable to intrinsic characteristics of the class or to individual
391 or environmental factors influencing the composition.

392 **4.1 Wildmeat as Source of Nutrients**

393 We compare the nutritional content of wildmeats with the DRI (Dietary Reference
394 Intakes) to provide a reference for understanding the contribution of these wildmeats as a
395 significant source of nutrients for people at different life stages (Supplementary Material 10).
396 Wildmeats can be considered excellent sources of various nutrients. Analyses of the edible
397 portions of muscles and viscera from these animals demonstrate their ability to provide high
398 levels of proteins, as well as micronutrients such as iron, zinc, and selenium (see Supplementary
399 Material 10). These elements are often deficient in the diets of vulnerable groups, such as
400 women and children, particularly in IPLC contexts (45).

401 Nutritional deficiencies have significant health implications. Protein deficiency can lead
402 to growth delays and induce hormonal and immunological changes (46). Iron deficiency is
403 correlated with the development of anemia, (7) while zinc deficiency can lead to immune
404 system dysregulation, compromised cognitive functions, growth deficits, and dermatological
405 disorders (47). Selenium deficiency can compromise the effectiveness of immune system cells
406 (48).

407 Our findings reveal that wild bird meat, in portions equivalent to a breast or thigh piece
408 (100 g), can supply more than half of the daily iron needs for children and women, as well as

409 more than half of the protein needed for women and nearly double the amount needed for
410 children. Meanwhile, mammal meats, in a portion like a medium steak (100 g), provide about
411 half of the protein and 40% of the iron needed for both children and women. Additionally,
412 organs such as the liver and heart are even richer, potentially exceeding 200% of children's iron
413 and selenium needs. For reptiles, a portion of two small pieces of muscle (100 g) can meet over
414 200% of children's iron needs and exceed 400% of their selenium needs. For women, the
415 amounts of iron and selenium provided meet approximately 200% and 150% of their
416 requirements, respectively.

417 These results highlight the potential of wild animal meats as sources of nutrients that
418 are often deficient in the diets of populations facing food insecurity in developing countries.
419 They contribute to improving dietary quality, making their consumption an important
420 nutritional strategy for populations dependent on these resources (49). For this reason, either
421 restricting access to wildmeats due to stringent conservation policies or defaunation of wild
422 animal populations may have adverse nutritional impacts on these vulnerable human
423 populations.

424 **4.2 Viscera of Wild Animals are Important Reservoirs of Nutrients**

425 Our findings indicated that meats derived from viscera have a substantially different
426 nutritional composition compared to those from muscles. Both mammalian and reptilian viscera
427 stood out for their high concentrations of nutrients, surpassing muscular meats, particularly in
428 terms of mineral content.

429 The composition of meat can vary considerably according to the anatomical part and
430 tissue type of the animal (17). While meats derived from viscera are known for their high fat
431 and micronutrient content, muscles are notable for their higher protein concentrations and lower
432 levels of fat and micronutrients.

433 Despite variations in nutrient patterns between animal classes (reptiles and mammals),
434 we observed that reptilian viscera demonstrated a pattern similar to that mentioned by, (17)
435 with higher fat content, lower protein concentrations, and overall mineral concentrations nearly
436 three times higher than in muscles. However, mammalian viscera, while exhibiting higher fat
437 concentrations, particularly omega-6, and high concentrations of micronutrients, had protein
438 concentrations similar to those found in muscles.

439 Furthermore, we observed a distinctive distribution of potassium compared to other
440 minerals present in the viscera. Potassium concentrations in visceral meats were lower than in
441 muscles for both reptiles and mammals. This disparity may be related to the physiological
442 characteristics of these tissues, (17) as muscles require higher concentrations of potassium for
443 muscle contraction.

444 Although wild animal viscera are important nutrient sources, we do not recommend
445 promoting the preferential consumption of specific anatomical parts over others. Utilizing food
446 holistically, a common practice among indigenous peoples and traditional communities,
447 contributes to both sustainability, (45) and nutritional diversification through a
448 complementarity mechanism (50). Furthermore, while viscera may contain high concentrations
449 of certain nutrients, their proportion, both available and consumed, tends to be lower compared
450 to other edible tissues of the animal.

451 Moreover, it is important to note that consuming viscera, despite being a rich source of
452 essential nutrients, may pose risks due to higher levels of heavy metals compared to the muscles
453 of animals. These levels can be higher in wild animal viscera and, in some cases, exceed
454 recommended limits set by health organizations (51). Regular consumption of wildmeat can
455 lead to a significant increase in exposure to heavy metals, such as cadmium and lead, in families

456 consuming these meats (51). Thus, one of the recommendations is to limit the consumption of
457 viscera, such as liver, especially in children.

458 Some studies indicate that certain nutrients can mitigate the toxic effects of heavy metals
459 through chelation. For example, a study conducted in the Brazilian Amazon, where the local
460 population is exposed to a diet rich in selenium and heavy metals such as methylmercury,
461 revealed that selenium plays a chelating role, (52) potentially mitigating the toxic effects of
462 methylmercury. This finding underscores the complexity of nutritional and environmental
463 interactions, highlighting the importance of considering socio-ecological contexts in nutritional
464 recommendations. While high selenium intake can lead to toxicity, (53) the case of the
465 Amazonian population illustrates how excessive consumption of certain nutrients, under certain
466 conditions, can offer protective effects. This highlights the need for a nuanced approach in
467 nutritional guidelines. Selenium was one of the nutrients found in higher concentrations in the
468 organ meats of wild animals, with 100 g of reptile and mammal organ meats providing over
469 100% of daily recommendations for children and adults.

470 **4.3 The Nutritional Composition of Wildmeat Varies Among Classes, Not Necessarily Due** 471 **to Animal Physiology Factors**

472 Our comparison of nutrient content in muscles among species revealed a notable
473 predominance of potassium, fat, and omega-6 in birds compared to other animal classes. Wild
474 birds assessed in this study showed an average fat content of less than 2.5%. Despite being
475 higher than other classes, this is lower than expected for non-wild birds, which typically have
476 fat concentrations around 5%, possibly due to variations in physical activity levels (54). Fat
477 content can also vary considerably depending on the bird species; for example, ostriches
478 generally have lower fat content compared to chickens (55).

479 However, the results obtained with the birds in this study are enlightening in
480 demonstrating that variations in nutrients may not be directly related to the taxonomic class of
481 the animal. We observed that the muscles of wild mammals do not have higher iron
482 concentrations compared to those of wild birds, indicated by a lack of statistically significant
483 difference, a finding that differs from the pattern observed in domesticated animals (10). Based
484 on these findings, we propose two hypotheses to explain the high concentrations of iron and
485 other micronutrients in the meat of wild birds: (i) the distinct anatomy of birds and their
486 migratory lifestyle, and (ii) the slaughter method to which free-ranging birds are subjected.

487 Within bird species, physiological characteristics associated with flight capability
488 influence the nutritional profile of their meat. For flight, birds primarily use the pectoral
489 muscles, responsible for wing movement (56). The continuous use of these muscles at high
490 frequencies requires a high energy cost, leading to physiological adaptations to sustain
491 prolonged flights (56). During migration periods, wild birds exhibit a significant increase in
492 pectoral muscle mass and metabolic rates to sustain flight, allowing them to go long periods
493 without feeding or drinking, primarily relying on fat oxidation for energy (57). This can also
494 affect the low fat concentrations in their meats.

495 When comparing nutrient content in the breast and thigh meat of migratory wild birds,
496 breast muscle has nearly double the iron content found in thigh meat, (13) In contrast,
497 domesticated birds like turkey have lower iron concentrations in breast meat compared to thigh
498 meat. For example, 100 g of turkey breast meat contains approximately 0.73 mg of iron,
499 whereas the thigh contains 1.57 mg per 100g, (10) demonstrating a different nutrient profile
500 compared to migratory wild birds.

501 The high iron concentration in wild bird meat may also be attributed to the slaughter
502 method used. Typically, wild birds are shot in the chest, a highly vascularized area near nutrient-

503 rich organs such as the liver and heart (13). This process can cause the leakage of visceral
504 contents, increasing nutrient content, including iron, in the muscle tissues. The evidence
505 presented on bird behavior and slaughter methods suggests that variations in nutrient levels may
506 not be directly related to the animal's taxonomic class.

507 The anatomy and migratory behavior of wild birds permits them to have a more
508 diverse diet, directly influencing the nutritional composition of their meat (13). We can assume
509 that birds primarily consuming seeds and oilseeds, which are naturally rich in fats, omegas, and
510 minerals, (10) tend to have meat that is richer in these nutrients. In contrast, carnivorous and
511 omnivorous birds have a distinct nutritional profile. These characteristics differ significantly
512 from domesticated birds, which are raised in confinement and tend to feed mainly on
513 agricultural feeds designed for rapid growth (58). Over time, soils used for conventional food
514 production can experience nutritional imbalances and low fertility, especially in terms of
515 micronutrients such as iron, zinc, and manganese (59). Consequently, food produced in
516 conventional systems tends to have lower concentrations of these micronutrients, reflected in
517 lower levels in domesticated bird meats.

518 Regarding reptiles, a significant finding was the high concentration of selenium in
519 their muscles and viscera, much higher than those observed in birds and mammals. These
520 findings may be linked to the physiological characteristics of reptiles, which have the ability to
521 absorb nutrients not only from their diets but also directly from their surrounding aquatic
522 environment (60). Reptiles can take in trace elements through their skin and other permeable
523 membranes while submerged in water. This dual absorption mechanism contributes to the
524 significant presence of trace elements, such as selenium, in their meat (60).

525 The reptiles analyzed in this study are aquatic and were collected mainly in the
526 Brazilian Amazon, known for significant variations in soil and water composition, influenced

527 by different organic matter and soil biota, both geographically (e.g. white, black and transparent
528 rivers) and between dry and rainy periods (61). This region is known for high selenium
529 concentrations in the soil, which play a crucial role in its absorption by local foods and water
530 bodies. For example, Amazonian soils are responsible for the abundant production of Brazil
531 nuts (*Bertholletia excelsa*), a significant source of selenium. Studies show considerable
532 variations in selenium content in these nuts across different Amazonian regions, suggesting
533 significant environmental influence on their concentration. These findings have important
534 implications for understanding the nutritional composition of reptile meat, as selenium may be
535 influenced by the diet of these animals in the Amazon region (62).

536 Our results demonstrated that wild mammals exhibited the highest concentrations of
537 zinc compared to other animal classes, along with elevated levels of iron and potassium. These
538 findings are consistent with, (26) who investigated wild deer meat and found that potassium,
539 zinc, and iron were the most abundant nutrients. However, the content of these nutrients can
540 vary within species depending on the animal's life cycle. For instance, male deer may
541 experience cyclic physiological osteoporosis related to rapid antler growth, decreasing zinc
542 content in the bloodstream and consequently in all other body tissues (26). The results of this
543 study highlight significant variations in the nutritional composition of meat not only between
544 different classes of animals but also among species within the same class.

545 Finally, it is important to note that ecosystems vary considerably between countries, and
546 the wide variations in nutritional profiles are also associated with the specificities from which
547 these species were collected. The study encompassed nutritional composition data from 26
548 animal species, originating from 11 different countries. For this reason, nutrient content in these
549 meats is subject to various factors, as nutrient distribution in the region varies significantly
550 according to soil and water type and collection area (59).

551 Therefore, in our findings, birds stood out for their high concentrations of potassium,
552 lipids, and omega-6 fatty acids, while reptiles exhibited exceptionally high levels of selenium,
553 likely influenced by selenium-rich environments such as the Amazon. Wild mammals, on the
554 other hand, distinguished themselves with high concentrations of zinc, iron, and potassium.
555 These findings suggest that while physiological and metabolic factors may play a role, the
556 nutritional composition of meat is strongly shaped by environmental factors such as diet and
557 habitat. Therefore, it is important to consider these nuances when evaluating the nutritional
558 value of meat from different species, rather than relying solely on taxonomic classifications.

559

560 **4.4 Novelty of the Research**

561 To the best of our knowledge, this is the first study to compile and analyze data
562 concerning a large number of distinct species spanning three different classes in a single
563 analysis. Notably, we have also introduced the technique of data imputation, presumably for
564 the first time in the field of nutritional analysis of wildmeat. While data imputation has been
565 previously employed in areas such as metabolomics for nutritional analyses? (63) its application
566 in this context is innovative. Additionally, during our investigation, we identified a lack of
567 appropriate tools to assess the methodological quality of research related to the nutritional
568 composition of wild animals. Specific ethical and legal considerations in this field exacerbate
569 this need. To address this limitation, we developed our own quality questionnaire (see
570 Supplementary Material 2), which assisted us in evaluating the quality of the studies included
571 in the review, considering the complexity of the resource in question.

572 Several limitations were encountered during the development of this study. One major
573 challenge was the lack of access to original data from the reviewed articles, which may have
574 compromised our ability to capture data variability. Additionally, the lack of specific

575 information on covariates impacting composition in the analyzed studies limited our ability to
576 adjust for these elements during tests and conduct subgroup analyses, preventing us from
577 conducting a traditional meta-analysis. Furthermore, the small sample size of reptile meats
578 demands caution in interpreting results related to this group.

579 We also observed high SMAPE (Symmetric Mean Absolute Percentage Error) values
580 for certain nutrients, indicating reduced quality in the predictions considering the available data.
581 This suggests the need for caution in interpreting these results. Despite this, we proceeded with
582 the imputation process based on three main grounds: (1) the potential for significant natural
583 variability, especially concerning wildmeats; (2) the scarcity of available data for this specific
584 type of resource, which inherently may contribute to greater uncertainty in the forecasts made;
585 and (3) relevant ethical considerations, including the difficulties associated with obtaining wild
586 meat samples.

587 Despite these limitations, our research provides a detailed analysis of nutritional
588 patterns in wild animals, which can underpin and guide future investigations in this field. The
589 data obtained could assist in formulating more precise hypotheses, such as the impact of
590 different slaughter methods on iron content in birds or the influence of the environment on
591 selenium content in reptiles. Furthermore, we emphasize the importance for researchers,
592 especially those focused on wildlife management, to enrich their databases with detailed
593 information on variables influencing nutritional composition. This would allow for more
594 comprehensive analyses, enhancing wildlife conservation efforts and maximizing the benefits
595 derived from these resources, especially for IPLC.

596 **5. CONCLUSION**

597 Wildmeat play a crucial role as nutrient sources for diverse human populations. Findings
598 from this study reveal significant nutrient variations among different animal classes and

599 anatomical parts, underscoring the ability of game meat to provide essential nutrients,
600 especially micronutrients often deficient in the diets of vulnerable groups such as children and
601 women of childbearing age who are particularly susceptible to the effects of malnutrition.
602 Moreover, this study introduces the use of data imputation methodology as an alternative to
603 navigating scarcity of data on wild foods. However, it is important to note that current primary
604 data limitations hinder the precise determination of whether these differences are due to
605 inherent characteristics of animal classes or to individual and environmental factors affecting
606 nutritional composition. Future research that directly analyzes the nutritional composition of
607 wildmeat is needed to clarify the causes of these variations.

608

609

610

611 REFERENCES

612

- 613 1. Ingram DJ, Coad L, Milner-Gulland EJ, Parry L, Wilkie D, Bakarr MI, et al. Wild Meat Is
614 Still on the Menu: Progress in Wild Meat Research, Policy, and Practice from 2002 to
615 2020. *Annu Rev Environ Resour.* 2021;46(Volume 46, 2021):221–54.
- 616 2. Cawthorn DM, Hoffman LC. The bushmeat and food security nexus: A global account of
617 the contributions, conundrums and ethical collisions. *Food Res Int.* 1º de outubro de
618 2015;76:906–25.
- 619 3. Fa JE, Olivero J, Real R, Farfán MA, Márquez AL, Vargas JM, et al. Disentangling the
620 relative effects of bushmeat availability on human nutrition in central Africa. *Sci Rep.* 2
621 de fevereiro de 2015;5(1):8168.
- 622 4. Friant S, Ayambem WA, Alobi AO, Ifebueme NM, Otukpa OM, Ogar DA, et al. Eating
623 Bushmeat Improves Food Security in a Biodiversity and Infectious Disease “Hotspot”.
624 *EcoHealth.* março de 2020;17(1):125–38.
- 625 5. Nielsen MR, Meilby H, Smith-Hall C, Pouliot M, Treue T. The Importance of Wild Meat
626 in the Global South. *Ecol Econ.* 1º de abril de 2018;146:696–705.
- 627 6. Nasi R, Brown, D, Wilkie D, Bennett E, Tutin C, van Tol G, et al. Conservation and use
628 of wildlife-based resources: the bushmeat crisis. Montreal: Secretariat of the Convention
629 on Biological Diversity; 2008. 50 p. (CBD technical series).
- 630 7. Torres PC, Morsello C, Orellana JDY, Almeida O, de Moraes A, Chacón-Montalván EA,
631 et al. Wildmeat consumption and child health in Amazonia. *Sci Rep.* 6 de abril de
632 2022;12(1):5213.

- 633 8. Golden CD, Fernald LCH, Brashares JS, Rasolofoniaina BJR, Kremen C. Benefits of
634 wildlife consumption to child nutrition in a biodiversity hotspot. *Proc Natl Acad Sci U S*
635 *A*. 6 de dezembro de 2011;108(49):19653–6.
- 636 9. Dannenberger D, Nuernberg G, Nuernberg K, Hagemann E. The effects of gender, age
637 and region on macro- and micronutrient contents and fatty acid profiles in the muscles of
638 roe deer and wild boar in Mecklenburg-Western Pomerania (Germany). *Meat Sci*. 1º de
639 maio de 2013;94(1):39–46.
- 640 10. TBCA. TBCA - Tabela Brasileira de Composição de Alimentos. [Internet]. 2023 [citado
641 19 de julho de 2023]. Disponível em: [http://www.tbca.net.br/base-](http://www.tbca.net.br/base-dados/composicao_alimentos.php)
642 [dados/composicao_alimentos.php](http://www.tbca.net.br/base-dados/composicao_alimentos.php)
- 643 11. Masson L. [LATINFOODS and its role in the generation and compilation of data for
644 Latin America]. *Arch Latinoam Nutr*. setembro de 1999;49(3 Suppl 1):89S-91S.
- 645 12. Jacob MCM, Feitosa IS, Albuquerque UP. Animal-based food systems are unsafe: severe
646 acute respiratory syndrome coronavirus 2 (SARS-CoV-2) fosters the debate on meat
647 consumption. *Public Health Nutr*. dezembro de 2020;23(17):3250–5.
- 648 13. Sevillano-Caño J, Cámara-Martos F, Aguilar-Luque EM, Cejudo-Gómez M, Moreno-
649 Ortega A, Sevillano-Morales JS. Trace Element Concentrations in Migratory Game Bird
650 Meat: Contribution to Reference Intakes Through a Probabilistic Assessment. *Biol Trace*
651 *Elem Res*. 1º de outubro de 2020;197(2):651–9.
- 652 14. Zimmerman TJ, Jenks JA, Leslie DM, Neiger RD. Hepatic minerals of white-tailed and
653 mule deer in the southern Black Hills, South Dakota. *J Wildl Dis*. abril de
654 2008;44(2):341–50.
- 655 15. Rizvi STH, Latif MY, Amin MS, Telmoudi AJ, Shah NA. Analysis of Machine Learning
656 Based Imputation of Missing Data. *Cybern Syst*. 2023;0(0):1–15.
- 657 16. Soulsbury CD, Gray HE, Smith LM, Braithwaite V, Cotter SC, Elwood RW, et al. The
658 welfare and ethics of research involving wild animals: A primer. *Methods Ecol Evol*.
659 2020;11(10):1164–81.
- 660 17. Damodaran SD, Parkin KL. *Química de Alimentos de Fennema*. 5º ed. Artmed; 2018.
- 661 18. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred Reporting Items for Systematic
662 Reviews and Meta-Analyses: The PRISMA Statement. *Ann Intern Med*. 18 de agosto de
663 2009;151(4):264–9.
- 664 19. Hayward MW, Callen A, Allen BL, Ballard G, Broekhuis F, Bugir C, et al.
665 Deconstructing compassionate conservation. *Conserv Biol*. 2019;33(4):760–8.
- 666 20. Memon S, Wamala R, Kabano IH. A Comparison of Imputation Methods for Categorical
667 Data [Internet]. Rochester, NY; 2023 [citado 6 de março de 2024]. Disponível em:
668 <https://papers.ssrn.com/abstract=4574180>
- 669 21. Rubin DB. Discussion on Multiple Imputation. *Int Stat Rev Rev Int Stat*. 2003;71(3):619–
670 25.
- 671 22. Myers WR. Handling Missing Data in Clinical Trials: An Overview. *Drug Inf J*. abril de
672 2000;34(2):525–33.
- 673 23. Padovani RM, Amaya-Farfán J, Colugnati FAB, Domene SMÁ. Dietary reference
674 intakes: aplicabilidade das tabelas em estudos nutricionais. *Rev Nutr*. dezembro de
675 2006;19(6):741–60.
- 676 24. Jacob. [eliasjacob/paper_nutritional_composition_wildmeat](https://github.com/eliasjacob/paper_nutritional_composition_wildmeat) [Internet]. 2024. Disponível
677 em: https://github.com/eliasjacob/paper_nutritional_composition_wildmeat
- 678 25. Roça R de O. *COMPOSIÇÃO QUÍMICA DA CARNE*. 2012;
- 679 26. Serrano MP, Maggiolino A, Landete-Castillejos T, Pateiro M, Barbería JP, Fierro Y, et al.
680 Quality of main types of hunted red deer meat obtained in Spain compared to farmed
681 venison from New Zealand. *Sci Rep*. 22 de julho de 2020;10(1):12157.
- 682 27. Soriano A, Murillo P, Perales M, Sánchez-García C, Murillo JA, García Ruiz A.

- 683 Nutritional quality of wild Iberian red deer (*Cervus elaphus hispanicus*) meat: Effects of
684 sex and hunting period. *Meat Sci.* 1º de outubro de 2020;168:108189.
- 685 28. Madeira LA. MORFOLOGIA DAS FIBRAS MUSCULARES ESQUELÉTICAS DE
686 FRANGOS DE CORTE CRIADOS NOS SISTEMAS CONFINADO E SEMI-
687 CONFINADO [Internet]. 2005. Disponível em:
688 [https://repositorio.unesp.br/bitstream/handle/11449/95316/madeira_la_me_botfmvz.pdf?s](https://repositorio.unesp.br/bitstream/handle/11449/95316/madeira_la_me_botfmvz.pdf?squence=1)
689 [equence=1](https://repositorio.unesp.br/bitstream/handle/11449/95316/madeira_la_me_botfmvz.pdf?squence=1)
- 690 29. Aguiar JPL. Tabela de composição de alimentos da Amazônia. *ACTA AMAZONICA* 26.
691 1996;121–6.
- 692 30. Amici A, Cifuni GF, Contò M, Esposito L, Failla S. Hunting area affects chemical and
693 physical characteristics and fatty acid composition of wild boar (*Sus scrofa*) meat.
694 *Rendiconti Lincei.* 1º de novembro de 2015;26(3):527–34.
- 695 31. Milczarek A, Janocha A, Niedziałek G, Zowczak-Romanowicz M, Horoszewicz E,
696 Piotrowski S. Health-Promoting Properties of the Wild-Harvested Meat of Roe Deer
697 (*Capreolus capreolus* L.) and Red Deer (*Cervus elaphus* L.) [Internet]. 2021 [citado 5 de
698 setembro de 2024]. Disponível em: <https://www.mdpi.com/2076-2615/11/7/2108>
- 699 32. Fernandes HR, Deliza R, Neto OC, Silva CM, Albuquerque NI de, Martins TR, et al.
700 Effect of high hydrostatic pressure on the meat of collared peccaries (*Tayassu tajacu*) with
701 different ages. *Afr J Food Sci.* 30 de setembro de 2022;16(9):215–25.
- 702 33. Gálvez C. H, Arbaiza T, Carcelén C. F, Lucas A. O. *Revista de Investigaciones*
703 *Veterinarias del Perú.* 1999 [citado 5 de setembro de 2024]. VALOR NUTRITIVO DE
704 LAS CARNES DE SAJINO (*Tayassu tajacu*), VENADO COLORADO (*Mazama*
705 *americana*), MAJAZ (*Agouti paca*) Y MOTELO (*Geochelone denticulata*). Disponível
706 em: <https://revistasinvestigacion.unmsm.edu.pe/index.php/veterinaria/article/view/6707>
- 707 34. Jarzyńska G, Falandysz J. Selenium and 17 other largely essential and toxic metals in
708 muscle and organ meats of Red Deer (*Cervus elaphus*) — Consequences to human health.
709 *Environ Int.* 1º de julho de 2011;37(5):882–8.
- 710 35. Johnson HE, Bleich VC, Krausman PR. Mineral deficiencies in tule elk, Owens Valley,
711 California. *J Wildl Dis.* janeiro de 2007;43(1):61–74.
- 712 36. Lima AT. Caracterização Físico Química da Tartaruda da Amazônia de água proveniente
713 de cativeiro e de habitat natural do estado do Amazonas. Universidade Federal do
714 Amazonas; 2009.
- 715 37. Lorenzo JM, Maggolino A, Gallego L, Pateiro M, Serrano MP, Domínguez R, et al.
716 Effect of age on nutritional properties of Iberian wild red deer meat - *Journal of the*
717 *Science of Food and Agriculture* [Internet]. 2019 [citado 5 de setembro de 2024].
718 Disponível em: <https://scijournals.onlinelibrary.wiley.com/doi/abs/10.1002/jsfa.9334>
- 719 38. Rudman M, Leslie AJ, van der Rijst M, Hoffman LC. Quality characteristics of Warthog
720 (*Phacochoerus africanus*) meat. *Meat Sci.* 1º de novembro de 2018;145:266–72.
- 721 39. Neto JV, Bressan MC, Faria PB, Vieira JO e, Santana MTA, Kloster M. Composição
722 centesimal e colesterol da carne de jacaré-do-pantanal (*Caiman yacare* Daudin 1802)
723 oriundo de zoológico e habitat natural. *Ciênc E Agrotecnologia.* agosto de
724 2006;30:701–6.
- 725 40. Landi N, Ragucci S, Di Giuseppe AM, Russo R, Poerio E, Severino V, et al. Nutritional
726 profiling of Eurasian woodcock meat: chemical composition and myoglobin
727 characterization. *J Sci Food Agric.* 2018;98(13):5120–8.
- 728 41. Pérez-Peña PE, Riveros-Montalván MS, Vargas-Arana G, Soria FD, Chumbe JV, Baca
729 YB. Consumo, microbiología y bromatología de la carne silvestre durante la COVID-19
730 en Iquitos, Perú. *Cienc Amaz Iquitos.* 2021;9(2):51–68.
- 731 42. Spiegelhaar N, Martin ID, Tsuji LJS. Indigenous Subarctic Food Systems in Transition:
732 Amino Acid Composition (Including Tryptophan) in Wild-Harvested and Processed

- 733 Meats. *Int J Food Sci.* 27 de junho de 2019;2019:e7096416.
- 734 43. Strazdiņa V, Jemeljanovs A, Šterna V. Nutrition Value of Wild Animal Meat. *Proc Latv*
735 *Acad Sci Sect B Nat Exact Appl Sci.* 31 de outubro de 2013;67(4–5):373–7.
- 736 44. Webb EC, Ryssen JBJ van, Erasmus MEA, McCrindle CME. Copper, manganese, cobalt
737 and selenium concentrations in liver samples from African buffalo (*Syncerus caffer*) in
738 the Kruger National Park. *J Environ Monit.* 29 de novembro de 2001;3(6):583–5.
- 739 45. Lemke S, Delormier T. Indigenous Peoples’ food systems, nutrition, and gender:
740 Conceptual and methodological considerations. *Matern Child Nutr.* novembro de 2017;13
741 *Suppl 3(Suppl 3):e12499.*
- 742 46. Malafaia G, Martins RF, Silva ME. 4 - AVALIAÇÃO DOS EFEITOS DA
743 DEFICIÊNCIA PROTEICA. 2009;4.
- 744 47. Pedraza DF, Sales MC. Estudos realizados no Brasil sobre a deficiência e a
745 suplementação de zinco: ênfase em crianças. *Rev Bras Saúde Materno Infant.* junho de
746 2017;17:217–32.
- 747 48. Cominetti C, Cozzolino SMF. Funções Plenamente Reconhecidas de Nutrientes Selênio.
748 2009; Disponível em: [https://nutritotal.com.br/pro/wp-](https://nutritotal.com.br/pro/wp-content/uploads/2019/03/Fun%C3%A7%C3%B5es_Sele%C3%A7%C3%A3o.pdf)
749 [content/uploads/2019/03/Fun%C3%A7%C3%B5es_Sele%C3%A7%C3%A3o.pdf](https://nutritotal.com.br/pro/wp-content/uploads/2019/03/Fun%C3%A7%C3%B5es_Sele%C3%A7%C3%A3o.pdf)
- 750 49. Jacob MCM, Feitosa IS, Albuquerque UP. Animal-based food systems are unsafe: severe
751 acute respiratory syndrome coronavirus 2 (SARS-CoV-2) fosters the debate on meat
752 consumption. *Public Health Nutr.* dezembro de 2020;23(17):3250–5.
- 753 50. DeClerck FAJ, Fanzo J, Palm C, Remans R. Ecological approaches to human nutrition.
754 *Food Nutr Bull.* março de 2011;32(1 Suppl):S41-50.
- 755 51. Danieli PP, Serrani F, Primi R, Ponzetta MP, Ronchi B, Amici A. Cadmium, lead, and
756 chromium in large game: a local-scale exposure assessment for hunters consuming meat
757 and liver of wild boar. *Arch Environ Contam Toxicol.* novembro de 2012;63(4):612–27.
- 758 52. Marco KCD. Avaliação da exposição ao metilmercúrio e dieta rica em selênio sobre os
759 níveis de óxido nítrico na população da região amazônica [Internet] [Mestrado em
760 Toxicologia]. [Ribeirão Preto]: Universidade de São Paulo; 2007 [citado 5 de março de
761 2024]. Disponível em: [http://www.teses.usp.br/teses/disponiveis/60/60134/tde-25072007-](http://www.teses.usp.br/teses/disponiveis/60/60134/tde-25072007-105904/)
762 [105904/](http://www.teses.usp.br/teses/disponiveis/60/60134/tde-25072007-105904/)
- 763 53. Whiting P, Rutjes AW, Reitsma JB, Bossuyt PM, Kleijnen J. The development of
764 QUADAS: a tool for the quality assessment of studies of diagnostic accuracy included in
765 systematic reviews. *BMC Med Res Methodol.* 10 de novembro de 2003;3(1):25.
- 766 54. Abreu ARC. NÍVEIS DE PROTEÍNA E AMINOÁCIDOS EM DIETAS PARA
767 FRANGOS DE CORTE FÊMEAS ABATIDOS EM DIFERENTES IDADES. 2019;
768 Disponível em:
769 <https://repositorio.ufmg.br/bitstream/1843/31571/1/TESE%20COMPLETA%20.pdf>
- 770 55. Feijó MB da S, Jacob S do C, Mano SB, Fernandes ML, Moraes ML de. Composição
771 Centesimal e Perfil de Minerais da Carne de Avestruz (*Struthio Camellus*). *Rev Assoc*
772 *Bras Nutr - RASBRAN.* 2009;(1):28–34.
- 773 56. Cao T, Jin JP. Evolution of Flight Muscle Contractility and Energetic Efficiency. *Front*
774 *Physiol* [Internet]. 9 de outubro de 2020 [citado 22 de maio de 2024];11. Disponível em:
775 <https://www.frontiersin.org/journals/physiology/articles/10.3389/fphys.2020.01038/full>
- 776 57. Price ER, Bauchinger U, Zajac DM, Cerasale DJ, McFarlan JT, Gerson AR, et al.
777 Migration- and exercise-induced changes to flight muscle size in migratory birds and
778 association with IGF1 and myostatin mRNA expression. *J Exp Biol.* 1º de setembro de
779 2011;214(17):2823–31.
- 780 58. Rodrigues R, Pinto LCG, Crestana S, Pires AK. Ministério da Agricultura, Pecuária e
781 Abastecimento. 2006;
- 782 59. Oliveira IP, Kluthcouski J, Fancelli AL, Neto DD. CONCENTRAÇÕES RESIDUAIS DE

- 783 COBRE, FERRO, MANGANÊS E ZINCO EM LATOSSOLO ROXO EUTRÓFICO
784 SOB DIFERENTES TIPOS DE MANEJO. 2001;
- 785 60. Gaspar A, Silva TJP. Composição nutricional da carne da tartaruga-da-Amazônia
786 (Podocnemis expansa) criada em cativeiro e em idade de abate. São Paulo. 2009;
- 787 61. Ferreira SJF, Luizão FJ, Miranda SÁF, Silva M do SR da, Vital ART. Nutrientes na
788 solução do solo em floresta de terra firme na Amazônia Central submetida à extração
789 seletiva de madeira; Acta Amaz. março de
790 2006;36:59–67.
- 791 62. Silva Jr EC da. SELÊNIO NA CASTANHA-DO-BRASIL (Bertholletia excelsa) E EM
792 SOLOS DA REGIÃO AMAZÔNICA BRASILEIRA [Internet]. Universidade Federal de
793 Lavras; 2016. Disponível em:
794 http://repositorio.ufla.br/jspui/bitstream/1/11520/1/DISSERTA%20c3%87%20c3%83O_Sel%20c3%aanio%20na%20castanha-do-brasil%20%28Bertholletia%20excelsa%29%20e%20em%20solos%20da%20regi%20a3o%20Amaz%20b4nica%20Brasileira.pdf
795
796
797
- 798 63. Wei R, Wang J, Su M, Jia E, Chen S, Chen T, et al. Missing Value Imputation Approach
799 for Mass Spectrometry-based Metabolomics Data. Sci Rep. 12 de janeiro de
800 2018;8(1):663.

801

Variation in Nutritional Composition of Anatomical Parts and Taxonomic Classes of Wild Animals: A Systematic Review Using Data Imputation with Artificial Intelligence.

Context

- The meat of wild animals is crucial for the food security of indigenous peoples and traditional communities.
- Currently, information on the nutritional profile of these foods is scarce.

Methodology

- This study is a systematic review.
- Evaluated the influence of the anatomical part and taxonomic class on the nutritional composition of meat.
- We employed artificial intelligence techniques (SMAPE).




Animal Icons: Mammals (🐷),
Birds (🐦), **Reptiles** (🐍)

**Anatomical Parts: Viscera vs.
Muscles**

Results

It analyzed 26 different species of animals (13 mammal species, 6 bird species, and 7 reptile species) from 11 different countries.



Viscera vs. Muscles:


- 🐦 : the viscera have higher levels of **lipid** and **Fe**.
- 🐷 : the viscera have higher levels of **omega-6**, **Se**, **Fe**, **Mn**, and **Zn**.

Differences between species:

- 🐦 : higher **lipid** levels.
- 🐷 : higher **Zn** levels.
- 🐍 : higher **Se** levels.

Conclusion

Wild meat is an important source of nutrients due to its rich nutritional value. This study is innovative and fills a gap by providing a comprehensive nutritional profile of bushmeat, introducing a methodology to deal with missing data, offering an analytical solution that addresses data scarcity while considering ethical and legal dilemmas in the analysis of wildlife.



*Consider, if feasible to do so, reporting the number of records identified from each database or register searched (rather than the total number across all databases/registers).

**If automation tools were used, indicate how many records were excluded by a human and how many were excluded by automation tools.

From: Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71. doi: 10.1136/bmj.n71. For more information, visit: <http://www.prisma-statement.org/>

Figure 1. Flowchart of the study selection process.