

**Transcriptome-driven Health-status Transversal-predictor Analysis (THTA) using the PBMC
transcriptome for health, food, microbiome and disease markers for understanding the
background and development of lifestyle diseases**

*Tilman Todt (1), Inge van Bussel (2), Lydia Afmann (2), Lorraine Brennan (3), Diana G.
Ivanova (4), Yoana Kiselova-Kaneva (4), E. Louise Thomas (5) and Ralph Rühl (6)*

(1) University of Applied Sciences, Nijmegen, the Netherlands,

(2) Division of Human Nutrition and Health, Wageningen University, Wageningen, the Netherlands,

*(3) UCD School of Agriculture and Food Science, Institute of Food and Health, University College
Dublin, Belfield, Dublin, Ireland,*

*(4) Department of Biochemistry, Molecular Medicine and Nutrigenomics, Medical University of Varna,
Bulgaria,*

(5) Research Centre for Optimal Health, School of Life Sciences, University of Westminster, London, UK,

(6) CISCAREX UG, Berlin, Germany.

Corresponding author:

Dr. Ralph Rühl

CISCAREX UG

Transvaalstr. 27c

D-13351 Berlin

Tel: +49 1522 642 0089

E-mail: ralphruehl@web.de

Abstract

We developed a novel machine-learning artificial intelligence (AI) approach to predict general health and food-intake parameters named Transcriptome-driven Health-status Transversal-predictor Analysis (THTA) with relevance for diabetes markers based on a mathematics-driven and non-transcriptomic biomarker driven approach. The prediction was based on values from food consumption, dietary lipids and their bioactive metabolites, peripheral blood mononuclear cells (PBMC) mRNA-based transcriptome signatures, magnetic resonance imaging (MRI), energy metabolism measurements, microbiome analyses, and baseline clinical parameters in a cohort of 72 subjects. Our novel machine learning approach included transcriptome data from PBMCs as a “one-method” approach to predict 77 general health-status markers for broad stratification of the diabetes phenotype, which are usually necessitating measurements using 16 different methods. The PBMC transcriptome was used to determine these selected 77 basic and background health-status markers in a transversal-predictor establishment group with very high accuracy (Pearson correlations are $r = 0,94$ ranging from 0,88 to 0,98). These collected variables offer a valuable indication to identify which individual factor(s) are mainly targeting diabetes. Based on the “establishment group” prediction approach a further “confirmation group” prediction approach was performed with a predictive potential for these 77 variables of $r = 0,62$ (ranging from 0,30 to 0,99). This “one-method” approach allows monitoring of a large number of health-status variables with relevance for diabetes simultaneously and may enable monitoring of therapeutic and preventive strategies. In summary, this novel technique based on PBMC transcriptomics from human blood offers prediction of a large range of health-related markers, which independently would be obtained in different clinical / research centres at a much higher price.

Key words: diabetes, transcriptomic, lipidomics, metabolomics, predictors, machine learning, regression models, artificial intelligence.

Introduction

Diabetes is a multifaceted disease often developing from obesity towards a diabetic-obesity phenotype (Farag and Gaballa 2011; M. I. Schmidt and Duncan 2003). Many simultaneous lifestyle factors may cause the transition to diabetes like a general unhealthy lifestyle including tobacco smoking, poor sleep habits, high alcohol consumption, low physical activity, and an unhealthy diet (Altobelli et al. 2020; Mozaffarian et al. 2009), with further contributions from a hereditary component which predisposes to diabetes (Goyal et al. 2023). Many dysfunctions in this condition are related to signalling in the glucose, insulin and fatty acids pathways as well as in the inflammation / immune system (Aronoff et al. 2004; M. I. Schmidt and Duncan 2003).

Machine learning artificial intelligence (AI) has been developed to monitor, classify, and predict diabetes (Oikonomou and Khera 2023). These predictive models can be used to study risk factors of type 2 diabetes (Schwarz et al. 2009; Yu et al. 2010; Naz and Ahuja 2020; Heikes et al. 2008) or predict diabetes (Razavian et al. 2015; Zou et al. 2018; Nguyen et al. 2019). However, small sample sizes and biased selection of risk factors may lead to predictive models which perform poorly (Lai et al. 2019). Some models use the PBMC transcriptome, in addition to other analytical methods, though most research focuses on a 'yes/no' answer for diabetes based on a set threshold of clinical diabetes markers (Tonyan et al. 2022). Other approaches for diabetes prediction employ polymorphisms of selected diabetes-relevant genes to indicate increased diabetes prevalence (Witka et al. 2019).

The main recommendation to reduce the burden of diabetes is lifestyle change, with various strategies (Knowler et al. 2002) and methods to both monitor and optimize lifestyle choices. However, long-term lifestyle alterations can be challenging and compliance can be poor (Aladhab and Alabood 2019), while unfortunately quick and easy drug treatment is often preferred (Garedow et al. 2023). Lifestyle interventions often include an "all-at-once-interventions-strategy", which are perceived as quite drastic, uncomfortable and non-preferred by patients and high-risk individuals (S. K. Schmidt et al. 2020). A more detailed evaluation of which selective lifestyle intervention should be altered while having an acceptable modified lifestyle of high-risk individuals will probably result in higher compliance and better outcomes. Selected lifestyle interventions should be chosen when many variables are evaluated at once and can be offered as a "one-method" predictor approach.

A potential approach to identifying clinical and health-status markers is based on the PBMC transcriptome (Costa et al. 2021; Elliott et al. 2014). Previous metabolic phenotyping have shown that the PBMC transcriptome can reflect environmental stress (Reynés et al. 2015) or the effect of specific diets (Bouwens, Afman, and Müller 2007; Caimari et al. 2010; Oliver et al. 2013; de Mello et al. 2012). For clinical diagnostics the PBMC transcriptome can reflect various pathologies (Baine et al. 2011; Chang et al. 2013), and identify biomarkers of obesity (Caimari et al. 2010; Oliver et al. 2013; Manoel-Caetano et al. 2012). It was previously shown that transcriptomic data from PBMCs (Rodney A. Rhoades PhD 2017) reflect the current metabolic and physiological state (Burczynski and Dorner 2006; Liew et al. 2006).

The purpose of our study was to develop models as a “one-method” machine-learning approach for determining basic and background variables like food intake and general health-status markers relevant for diabetes. Our strategy does not start by using known transcriptomic biomarkers / target genes, which are known to be involved in diabetes, but instead focusing on the categorization of the human organism as a “simple” holistic mathematical matrix. We performed a full PBMC transcriptome analysis in 72 volunteers from the EU FP7 NUTRITECH cohort, providing the largest known number of variables per person of any known performed clinical intervention study with additional 10,522 analysed gene transcripts detected in PBMCs being over quantification limit. In total 780 variables were measured by 16 different techniques, including food consumption, nutritional lipids, and their bioactive metabolites, magnetic resonance imaging (MRI), energy metabolism measures, microbiome analyses, and clinical biochemistry (Rundle et al. 2023) in the Nutrittech study, while from these 780 measurements 77 variables as the most diabetes-relevant markers were selected for further analysis.

Materials and methods

The Nutrittech project encompasses comprehensive phenotyping, which includes measurements related to diet, physical activity, body composition, blood biomarkers, and metabolomics, yielding 780 measurements per individual.

Selection of diabetes-related health-status variables for further predictor analysis

To facilitate initial testing of our predictive capabilities, we had to pare this down to a more manageable number of variables, a step taken prior to any machine learning. Our goal was to select a variety of variables linked to the development of a diabetes phenotype. We also attempted to predict variables representative of the wide range of phenotyping methods employed in the Nutritech protocol, including routinely collected physical and clinical biochemical measures, to more sophisticated and expensive analyses, including MRI, microbiome analyses, and multiple lipidomic analyses for a larger range of lipids. In total 28 self-reported dietary intake measures were included relating to plant or animal-sources, sweets/snacks and alcohol (details for categorization added in supplementary material). Moreover, we included routine health assessment measures like blood pressure and physical body measurements based on MRI (fat mass and liver fat), calorimetry, and gut microbiome analyses. Further out of total 780 variables, 77 selected variables per person were chosen with major diabetes-relevance for further targeted analysis, measured by 16 different techniques, including simple methods (eg. body weight and blood pressure; methods 9 and 15). Some variables required advanced and expensive techniques, and are available only at specialized health care facilities or a handful of service laboratories worldwide (methods 2–6; 11-14).

Dataset

Data from the EU FP7 NUTRITECH study (Rundle et al. 2023) were used to develop regression models. The EU FP7 NUTRITECH dataset includes a human cohort of 72 subjects for whom 77 different variables out of 780 available variables and additional PBMC transcriptome data of 10,522 individually expressed genes were determined as described previously (Rundle et al. 2023; Fiamoncini et al. 2018). The cohort was divided in two groups: 32 subjects in the control group and 40 subjects in the intervention group. Data was collected before intervention as well as 13 weeks later after intervention in all participants. The data from the “before intervention group” were used for the “establishment group” model development and evaluation, while the data from the “after intervention group” were used for the “confirmation group” approach.

Modelling and evaluation

A) “Establishment group” (EST) for a transversal prediction analysis

A transversal prediction analysis usually describes the analysis of a larger range of variables which can be predicted based on using one broader, more complex set of variables originating from one method, which is usually a transcriptome analysis. This prediction analysis towards obtaining transversal predictors from one variable is usually done in a group of individuals at the same time in an “establishment group”.

We developed different regression models like the Random Forest (Liaw and Wiener 2002), Elastic Net (Friedman, Hastie, and Tibshirani 2010), and multiple Linear Regression (Hebbali 2024) to evaluate the best prediction model for a broader range of variables using the PBMC transcriptomic data. For each regression analysis, an individual model was developed for each of the 77 variables using data from the “establishment group”.

The following procedure was applied for each regression method: Thirty separate models were developed for each variable. For each model, the data set was randomly split into training and validation sets. The training data set included data from two-thirds of the subjects, and one-third was used for validation. Spearman correlation coefficients were used to select predictors from the training data set. Of the 10,522 transcriptomic variables, a maximum of 50 transcripts were allowed to select predictors. mRNAs with the highest absolute correlation coefficient were selected. Regression was used to create the model. Training of the model with the Elastic Net was performed with cross-validation (with ten convolutions) and optimization of the hyperparameters. No cross-validation and optimization were performed for Random Forest and multiple Linear Regression. Multiple Linear Regression models were built by the stepwise regression function in the R package ‘olsrr’ (Hebbali 2024). The model's root mean square error and the Pearson correlation coefficient (R) were used as evaluation metrics. After training, models were evaluated using the data from the “establishment group”. Performance of the models on the training data and the whole dataset were determined using the Pearson correlation coefficient. Data analyses and model development were performed in R (version 4.2.2) and the R package ‘caret’ (version 6.0-94) (Kuhn 2008). Random forest models in the R package ‘randomForest’ (version 4.7.1.1) (Liaw and Wiener 2002), Elastic net models applied the R package ‘glmnet’ (version 4.1.8) (Friedman, Hastie, and Tibshirani 2010) and multiple linear regression in the R package ‘olsrr’ (version 0.6.0)

(Hebbali 2024). Visualisation of the results was done with the R package 'ggplot2' (version 4.1.8) (Wickham 2016).

B) Applying predictors from the “establishment group” (EST) for a “confirmation group” prediction (CON) analysis

After developing the regression models in the “establishment group” (EST), the models for predicting the outcome were applied to data of a group analysed not at the same time point, named here the “confirmation group”. Predictive performance was measured using the root mean square error and the Pearson correlation coefficient. Of the 30 models established for each variable, the model with the best performance on the “confirmation group” was selected as the final model. Predictive performance of the final models was visualized by comparing the observed values with predicted values and by comparing the overall predictive performance between the three regression methods of Random Forest, Elastic Net, and multiple Linear Regression models.

Results

Predictive models based on machine-learning offer an alternative to existing methods for determining variables with relevance for general health, food intake and targeted in our case for diabetes. One application of this transversal-prediction model is for monitoring individual health-status variables with a novel methodology using one blood sample, followed by a simple highly standardized transcriptomic analysis and a mathematical evaluation instead of using several different complex methodologies.

The first step was the transfer predictor analysis in the “establishment group” starting from the PBMC-transcriptome with 10,522 analysed mRNA transcripts detected, which were analysed being over the quantification limit (van Bussel et al. 2019), using 77 out of 780 variables including food intake and health-related variables for 72 volunteers.

Three different mathematical methods for prediction of 77 variables in the “establishment group” approach were used starting with the Radom Forest-based model (RF) resulting in an average of $r = 0,94$ ranging from 0,88 to 0,98. For prediction of the food diary data (Table 1, variables 1-28) in the transversal prediction approach predictor correlations were in the range of $r = 0,92 - 0,97$ (EST-RF) offering the first indirect calculation of food diary data. Especially summarising data like animal-based food ($r = 0,94$), plant-based food ($r = 0,94$), snacks / sweets ($r = 0,96$), general healthy food ($r = 95$) and general non-healthy food ($r = 0,97$), offer a precise predictor calculation (Figure 1). These predictor correlations are highly relevant to evaluate newly but also already existing large scale databases with PBMC transcriptomic data included but without additional food diary data collection.

The second sets of parameters to be determined were blood lipids (Table 1, variables 29 – 42). The models performed well for variables like the sum of carotenoids (sCARO-sum, variable 33) $r = 0,96$, sum of saturated fatty acids (s-sum SAFAs) $r = 0,94$, sum of monounsaturated fatty acids (s-sum MAFAs) $r = 0,93$, sum of n3-polyunsaturated fatty acids (s-sum n3-PUFAs) $r = 0,93$ and sum of n6-polyunsaturated fatty acids (s-sum n6-PUFAs) $r = 0,94$, sum of pro-inflammatory lipid-mediators (s-sum pro-infl.-lipid mediators) $r = 0,93$ and sum of pro-resolving lipid-mediators (s-sum pro-res.-lipid mediators) $r = 0,88$ are precise predictor calculations of an indirect quantification of the individual lipid levels. These predictor calculations are highly relevant to evaluate newly but also already existing large scale databases with PBMC transcriptomic included to further predict these specific lipids in individuals which are usually are just available using expensive and highly laborious blood lipid analyses.

The third set of new variables (Table 1, variables 43 – 47), which indicates transcriptome-based nuclear hormone selective signalling pathways including RXR- ($r = 0,98$), RAR- ($r = 0,97$), PPAR- ($r = 0,97$), VDR- ($r = 0,97$) and NURR1 / NR4A2-signalling pathways ($r = 0,96$), indicating which nuclear hormone signalling pathway is currently relatively up- or down-regulated.

All health variables including basic clinical chemistry-, basic health-, body composition, and microbiome parameters (Table 1, variables 48 – 77) commonly used to determine individual health-status relevant for

diabesity, were analysed. Of special interest were body composition parameters from MRI analyses including subcutaneous adipose tissue (SAT, $r = 0,93$), intra-abdominal adipose tissue (IAAT, $r = 0,93$), and intrahepatic cellular lipid (indicating liver fat, IHCL, $r = 0,96$). The model was also able to predict microbiome analysis including Firmicutes ($r = 0,95$) and Bifidobacterium ($r = 0,93$).

Other transfer predictor regression models like the Elastic Net (EN) and the Linear Regression model (LR) were tested. Predictor correlations suggested that the Random Forest model performed best with an average of $r = 0,94$, followed by the Elastic Net model $r = 0,80$ and the Linear Regression model $r = 0,77$ (table 2, supplementary tables 2 and 3 and figure 2). Applying all the methodologies and calculating a “best of” predictor analysis offer correlation coefficients of $r = 0,94$. Thus, the Random Forest model always offered the best values for the “establishment group”.

Further, we evaluated the robustness of this methodology by the “establishment group” predictor calculations for variables all determined “at the same day” with variables obtained weeks later in the same person; half of the group maintained their initial lifestyle, and the second group underwent a lifestyle intervention with increased physical activity, an energy-restricted and more balanced diet (Rundle et al. 2023; Fiamoncini et al. 2018). In this second group, the “confirmation group” prediction (CON) approach, we did not dissect these two groups for further calculations but this will be done under future evaluation of the 77 selected health-status parameters for a detailed evaluation of success of a lifestyle change on diabesity.

In these confirmation predictions, the Elastic Net model ($r = 0,63$) provided the best predictive potential, followed by the Random Forest model ($r = 0,62$) and by the Linear Regression model $r = 0,57$ (Table 2 and supplementary tables 2 and 3). Applying all actual methodologies and calculating a “best of” predictor analysis offer values of $r = 0,65$. Consequently, the “best of” predictor calculation offered the best values for the confirmation group mainly based on the Random Forest and Elastic Net models (Figure 3).

Discussion

Machine-learning approaches for health-status monitoring relevant for diabetes (Agliata et al. 2023; Wee et al. 2024) have been developed recently, and many of the existing machine-learning models use common biomarkers / risk factors and variables mainly associated with diabetes (Okada et al. 2022). These prediction models are mainly targeting single variables and are based on preselected known biomarkers (Jangili et al. 2023). Until now, no broader health-status marker array analyses have been performed with a focus on diabetes.

We used data from the EU FP7-NUTRITECH consortium including 72 individuals with 780 analysed individual clinical and lifestyle parameters and additional 10,522 parameters from PBMC transcriptome analyses. For our transfer-predictor analysis we used these 10,522 transcriptomic variables from mRNA transcripts only based on mathematical-relevance and not based on known biomarker functions, which is a new approach. This novel design simply applies data from the PBMC transcriptomics as a “simple” holistic and complex mathematical matrix (Figure 4).

These data from PBMC can easily be obtained from blood samples. A previous study has shown that individual PBMC transcriptomes can reflect obesity-related inflammation (Bories et al. 2012) and lipid metabolism in obesity-associated organs (Konieczna et al. 2015). Our approach determined that our “single method” predictor analysis could predict many individual lifestyle parameters such as food intake, basal and advanced health-status markers that were measured in the NUTRITECH cohort. In addition, as it is a “clean” mathematical methodology where we just focussed on the 77 most relevant health-status variables from a total of 780 parameters measured in the NUTRITECH cohort at one time point. We also selectively tested the possibility that alternative parameters of these 780 parameters can be calculated with similar accuracy using our transversal-predictor analysis (data not shown).

Diabetes mellitus is a multifactorial chronic disease, which can be triggered by many genetic and/or environmental factors (Sun, Yu, and Hu 2014; Kaul and Ali 2016). The number of adults living with diabetes globally has almost quadrupled since 1980 to 422 million and is projected to increase to 693 million by 2045

(Cho et al. 2018). Lifestyle interventions can be very effective in preventing and treating diabetes (Knowler et al. 2002), whereas they often are not well tolerated by participants (S. K. Schmidt et al. 2020). The inclusion of multiple diabetes-related variables may allow the development of targeted, more selective, and potentially more acceptable lifestyle interventions (Zhang et al. 2020). A cost-effective method would allow monitoring to be performed more frequently. In our novel machine-learning approach, we have developed regression models that can be used as low-cost models with good predictive power for continuous monitoring and evaluation of lifestyle interventions.

Furthermore, our approach provides a novel methodology for analysing and monitoring personal health status. Most current approaches use predictive models to distinguish between groups of individuals (e. g. healthy versus diabetic) through classification and use of common risk factors or health-related variables for model development. Our approach does not require the separation of individuals beforehand and uses PBMC transcriptome and a mathematical approach to evaluate many health-status markers concerning individual metabolic and physiological status.

The ability to robustly predict dietary intake, without necessitating the collection and analysis of food intake diaries is an exciting advancement. In the NUTRITECH studies these food dairy data can now easily be used for further correlations with clinical chemistry and this transversal-prediction approach. Especially summarised food groups like healthy or unhealthy dietary patterns, snacks / sweets, animal-based foods, or plant-based foods can be precisely predicted with high accuracy ($r = 0,94 - 0,97$) (Table 2).

Our methodology offers a straightforward and cost-effective approach to predicting and monitoring 77 most relevant health-status markers. In our analysis, the predictive models demonstrated excellent accuracy within the “establishment group” but showed moderate accuracy in the “confirmation group”. The predictive power in the “confirmation group” is influenced mainly by the availability of parameters with a larger distribution pattern and reduces accuracy.

Larger cohort studies are needed to further validate and improve the robustness of our approach. A potential limitation of these studies is their typically more shallow approach to phenotyping, which may lack the comprehensive range of parameters available in studies such as NUTRITECH (Rundle et al. 2023; Fiamoncini et al. 2018). This difference could impact the depth and accuracy of this transversal-predictor analysis when applied to broader datasets. However, these studies would enable the inclusion of additional parameters, such as sex, hormone levels, medication use, disease status, age, and ethnicity, which could enhance the models' predictive power. They would also support the development of specialized models for specific subgroups, including pregnant women, children, athletes, individuals with unique genetic profiles, older adults, and those with specific diseases. These expansions would optimize the methodology, increase predictive accuracy, and broaden its applicability across diverse populations, allowing for more personalized health monitoring and prediction.

In summary, we have developed a novel, “one-method” machine-learning approach for transversal-prediction in an “establishment group”, termed “Transcriptome-driven Health-status Transversal-predictor Analysis” (THAP). This method was further validated in a “confirmation group” consisting of 72 individuals, leveraging PBMC transcriptome data and mathematical processing to predict 77 of the most relevant health-status variables from an original set of 780 variables. This method introduces a novel predictive model that offers a cost-effective approach for monitoring health status markers, aimed at optimizing lifestyle choices and understanding their impact on the development and progression of health conditions, particularly diabetes. Our approach demonstrates that leveraging the individual PBMC transcriptome as a holistic mathematical matrix serves as a robust source for personalized health predictions. This model may also represent a simple yet innovative approach to clinical diagnosis and the management of dietary interventions and pharmacotherapy.

Table 1

	Variable	category	method	subjects	EST-RF R	subjects	CON-RF R
1	Alcohol	food intake measures	1	65	0,95	60	0,75
2	red-meat	"	1	62	0,93	60	0,65
3	meat products	"	1	68	0,93	63	0,52
4	red meat dishes	"	1	62	0,94	59	0,45
5	poultry	"	1	64	0,93	62	0,56
6	poultry dishes	"	1	66	0,94	60	0,62
7	Cheeses	"	1	64	0,92	63	0,45
8	Butter	"	1	66	0,89	61	0,52
9	whole milk	"	1	63	0,92	61	0,44
10	milk beverages	"	1	66	0,94	57	0,64
11	Egg	"	1	63	0,94	61	0,58
12	low fat milk	"	1	65	0,95	62	0,66
13	yoghurt	"	1	68	0,96	63	0,58
14	fish	"	1	68	0,94	62	0,52
15	animal-based food	"	1	67	0,94	63	0,63
16	fruit	"	1	64	0,92	60	0,56
17	fruit juices	"	1	63	0,92	62	0,63
18	soups	"	1	63	0,92	60	0,46
19	vegetables	"	1	63	0,93	60	0,66
20	plant-based food	"	1	62	0,94	58	0,63
21	biscuits	"	1	60	0,96	63	0,53
22	confect	"	1	65	0,88	60	0,63
23	high-energy drinks	"	1	62	0,94	57	0,55
24	ice creams	"	1	69	0,92	64	0,55
25	salty snacks	"	1	66	0,94	61	0,57
26	snacks / sweets	"	1	66	0,96	61	0,60
27	general healthy food	"	1	62	0,95	58	0,50
28	general non-healthy food	"	1	65	0,97	60	0,53
29	sLUT	blood lipids analysis	2	66	0,95	65	0,73
30	sATBC	"	2	64	0,96	63	0,67
31	s9CBC	"	2	65	0,93	64	0,62
32	sAT-LYC	"	2	63	0,92	65	0,65
33	sCARO-sum	"	2	66	0,96	64	0,65
34	s-sum-SAFAs	"	3	66	0,94	66	0,60
35	s-sum-MUFAs	"	3	66	0,93	67	0,40
36	s-s um-n3-PUFAs	"	3	60	0,93	62	0,58
37	s-sum-n6-PUFAs	"	3	69	0,94	69	0,63
38	sROL	"	4	68	0,94	65	0,51
39	sATRA	"	4	64	0,93	62	0,57
40	s9CDHRA	"	4	65	0,95	59	0,65
41	s-s um pro-infl. lipid-mediators	"	5	67	0,93	61	0,38
42	s-s um pro-res. lipid-mediators	"	5	62	0,88	57	0,62
43	RXR-signalling pathway	transcriptomic signatures	6	69	0,98	66	0,94
44	RAR-signalling pathway	"	6	71	0,97	69	0,87
45	PPAR-signalling pathway	"	6	71	0,97	71	0,86
46	VDR-signalling pathway	"	6	70	0,97	72	0,89
47	NURR1-signalling pathways	"	6	69	0,96	68	0,83
48	total-Chol.	basic clinical-chemistry	7	65	0,96	64	0,55
49	HDL-Chol.	"	7	67	0,95	67	0,63
50	LDL-Chol.	"	7	66	0,95	66	0,69
51	sGLUC	"	7	66	0,95	65	0,73
52	sINS	"	7	68	0,95	65	0,71
53	sNEFAs	"	7	69	0,91	68	0,61
54	sTRI	"	7	66	0,95	65	0,55
55	sVitD	"	7	64	0,96	64	0,65
56	sVITK2	"	7	67	0,95	66	0,52
57	eADIQ	ELISA	8	68	0,91	67	0,63
58	eL18	"	8	66	0,95	62	0,64
59	BP-dia	basic health-parameters	9	68	0,96	67	0,68
60	BP-sys	"	9	69	0,94	64	0,63
61	eosinophils	basic clinical chemistry	10	65	0,95	63	0,64
62	monocytes	"	10	67	0,91	64	0,66
63	HOMA2-IR	diabetesity marker	11	68	0,95	67	0,72
64	SAT	MRI	12	61	0,93	62	0,68
65	IAAT	"	12	65	0,93	65	0,63
66	IHCL	"	12	61	0,96	63	0,30
67	indir. calorimetry	energy-metabolism	13	67	0,94	65	0,76
68	RQ	"	14	64	0,88	62	0,73
69	sex	basic health-parameters	15	68	0,98	69	0,99
70	BW	"	15	68	0,92	66	0,70
71	BMI	"	15	65	0,92	66	0,72
72	HC	"	15	66	0,92	63	0,72
73	WC	"	15	64	0,91	66	0,68
74	WHIPR	"	15	69	0,94	68	0,67
75	HR	"	15	64	0,95	62	0,58
76	Firmi cutes	microbiome-analysis	16	55	0,95	59	0,52
77	Bifidobacterium	"	16	57	0,93	60	0,43
	Mean				0,94		0,62

Prediction of variables using Random Forest-based model. Variables were predicted in the “establishment group” (EST) for a transversal prediction starting from the PBMC transcriptome towards the 77 listed variables using the Random Forest-based (RF) regression models and in addition a “confirmation group” prediction (CON) was performed using the same methodology in the same group of volunteers after intervention using the 72 PBMC transcriptomic data of these volunteers. For each variable the following is reported: the number of individual transcriptomics included in the datasets, the number of mRNAs the model is based on, the Pearson correlation coefficient (R) between observed and predicted values, and the experimental method used to determine the variables.

Abbreviations: For food intake measures please check the materials and methods; s – serum; e – ELISA; adj. - adjusted; sLUT – serum lutein; sATBC – serum all-trans- β -carotene, s9CBC – serum 9-cis- β -carotene; sATLYC – serum all-trans-lycopene; CAROsum – sum of total carotenoids in serum; SAFA – saturated fatty acids; MUFA – mono-unsaturated fatty acids; PUFAs – polyunsaturated fatty acids; sROL – retinol; sATRA – serum all-trans-retinoic acid; s9CDHRA – serum 9-cis 13,14-dihydroretinoic acid; pro-inf. – proinflammatory; pro-res. pro-resolving; RXR – retinoid X receptor; RAR – retinoic acid receptor; PPAR – peroxisome proliferator-activated receptor; LXR – liver X receptor; VDR – vitamin D receptor; NURR1 - nuclear receptor related 1 protein/ NR4A2-receptor; Chol – cholesterol; HDL – high density lipoprotein; LDL – low density lipoprotein; GLUC – glucose; INS – insulin; NEFAS – non-esterified fatty acids; TRI – triglycerides; VitD – 25-Hydroxyvitamin D3, VitK2 – vitamin K2, ADIQ – adiponectin, IL18 – interleukin 18, BP-dia – diastolic blood pressure; BP-sys – systolic blood pressure; HOMA2-IR – homeostasis model assessment index; SAT – subcutaneous adipose tissue; IAAT – intraabdominal adipose tissue; IHCL – intrahepatic cellular lipid index; indir calorimetry – indirect calorimetry; RQ – respiratory coefficient; BW – body weight; BMI – body weight index; HC – hip circumference; WC – waist circumference; WHIPR – waist hip ratio; HR – heart rate.

Table 2

	variable	EST-RF R	EST-EN R	EST-LR R	EST-BO R	CON-RF R	CON-EN R	CON-LR R	CON-BO R
1	alcohol	0,95	0,75	0,78	0,95	0,75	0,70	0,57	0,75
2	red-meat	0,93	0,79	0,77	0,93	0,65	0,64	0,62	0,65
3	meat products	0,93	0,84	0,79	0,93	0,52	0,45	0,50	0,52
4	red meat dishes	0,94	0,75	0,73	0,94	0,45	0,40	0,32	0,45
5	poultry	0,93	0,82	0,84	0,93	0,56	0,58	0,44	0,58
6	poultry dishes	0,94	0,89	0,81	0,94	0,62	0,66	0,59	0,66
7	cheeses	0,92	0,67	0,63	0,92	0,45	0,34	0,29	0,45
8	butter	0,89	0,73	0,51	0,89	0,52	0,36	0,41	0,52
9	whole milk	0,92	0,68	0,72	0,92	0,44	0,47	0,33	0,47
10	milk beverages	0,94	0,90	0,83	0,94	0,64	0,56	0,55	0,64
11	egg	0,94	0,72	0,75	0,94	0,58	0,62	0,56	0,62
12	low fat milk	0,95	0,75	0,72	0,95	0,66	0,74	0,66	0,74
13	yoghurt	0,96	0,82	0,77	0,96	0,58	0,68	0,60	0,68
14	fish	0,94	0,83	0,77	0,94	0,52	0,61	0,45	0,61
15	animal-based food	0,94	0,81	0,74	0,94	0,63	0,60	0,50	0,63
16	fruit	0,92	0,68	0,68	0,92	0,56	0,59	0,39	0,59
17	fruit juices	0,92	0,75	0,72	0,92	0,63	0,48	0,41	0,63
18	soups	0,92	0,76	0,69	0,92	0,46	0,39	0,41	0,46
19	vegetables	0,93	0,80	0,86	0,93	0,66	0,66	0,54	0,66
20	plant-based food	0,94	0,80	0,80	0,94	0,63	0,62	0,54	0,63
21	biscuits	0,96	0,81	0,79	0,96	0,53	0,51	0,49	0,53
22	confect	0,88	0,70	0,73	0,88	0,63	0,59	0,58	0,63
23	high-energy drinks	0,94	0,74	0,74	0,94	0,55	0,64	0,55	0,64
24	ice creams	0,92	0,72	0,69	0,92	0,55	0,51	0,51	0,55
25	salty snacks	0,94	0,76	0,72	0,94	0,57	0,59	0,51	0,59
26	snacks / sweets	0,96	0,86	0,81	0,96	0,60	0,69	0,61	0,69
27	general healthy food	0,95	0,85	0,86	0,95	0,50	0,51	0,31	0,51
28	general non-healthy food	0,97	0,88	0,83	0,97	0,53	0,63	0,59	0,63
29	sLUT	0,95	0,80	0,74	0,95	0,73	0,78	0,65	0,78
30	sATBC	0,96	0,73	0,74	0,96	0,67	0,74	0,69	0,74
31	s9CBC	0,93	0,88	0,77	0,93	0,62	0,62	0,63	0,63
32	sAT-LYC	0,92	0,85	0,87	0,92	0,65	0,67	0,56	0,67
33	sCARO-sum	0,96	0,88	0,81	0,96	0,65	0,66	0,58	0,66
34	s-sum-SAFAs	0,94	0,81	0,68	0,94	0,60	0,62	0,63	0,63
35	s-sum-MUFAs	0,93	0,73	0,62	0,93	0,40	0,50	0,43	0,50
36	s-sum-n3-PUFAs	0,93	0,89	0,83	0,93	0,58	0,72	0,60	0,72
37	s-sum-n6-PUFAs	0,94	0,84	0,83	0,94	0,63	0,70	0,63	0,70
38	sROL	0,94	0,78	0,71	0,94	0,51	0,48	0,44	0,51
39	sATRA	0,93	0,78	0,72	0,93	0,57	0,38	0,45	0,57
40	s9CDHRA	0,95	0,87	0,83	0,95	0,65	0,72	0,71	0,72
41	s-sum pro-infl. lipid-mediators	0,93	0,75	0,75	0,93	0,38	0,41	0,20	0,41
42	s-sum pro-res. lipid-mediators	0,88	0,75	0,71	0,88	0,62	0,64	0,49	0,64
43	RXR-signalling pathway	0,98	0,96	0,96	0,98	0,94	0,96	0,96	0,96
44	RAR-signalling pathway	0,97	0,96	0,96	0,97	0,87	0,95	0,95	0,95
45	PPAR-signalling pathway	0,97	0,91	0,91	0,97	0,86	0,87	0,85	0,87
46	VDR-signalling pathway	0,97	0,94	0,92	0,97	0,89	0,91	0,88	0,91
47	NURR1-signalling pathways	0,96	0,94	0,94	0,96	0,83	0,90	0,89	0,90
48	total-Chol.	0,96	0,76	0,74	0,96	0,55	0,65	0,59	0,65
49	HDL-Chol.	0,95	0,80	0,76	0,95	0,63	0,68	0,63	0,68
50	LDL-Chol.	0,95	0,86	0,82	0,95	0,69	0,71	0,63	0,71
51	sGLUC	0,95	0,83	0,76	0,95	0,73	0,76	0,59	0,76
52	sINS	0,95	0,82	0,75	0,95	0,71	0,69	0,63	0,71
53	sNEFAs	0,91	0,67	0,59	0,91	0,61	0,61	0,57	0,61
54	sTRI	0,95	0,78	0,62	0,95	0,55	0,58	0,62	0,62
55	sVITD	0,96	0,82	0,74	0,96	0,65	0,70	0,69	0,70
56	sVITK2	0,95	0,82	0,82	0,95	0,52	0,60	0,53	0,60
57	eADIQ	0,91	0,85	0,77	0,91	0,63	0,70	0,68	0,70
58	eIL18	0,95	0,85	0,72	0,95	0,64	0,51	0,56	0,64
59	BP-dia	0,96	0,83	0,76	0,96	0,68	0,70	0,67	0,70
60	BP-sys	0,94	0,76	0,76	0,94	0,63	0,63	0,53	0,63
61	eosinophils	0,95	0,85	0,82	0,95	0,64	0,61	0,45	0,64
62	monocytes	0,91	0,70	0,68	0,91	0,66	0,59	0,47	0,66
63	HOMA2-IR	0,95	0,79	0,77	0,95	0,72	0,72	0,64	0,72
64	SAT	0,93	0,81	0,78	0,93	0,68	0,73	0,69	0,73
65	IAAT	0,93	0,77	0,76	0,93	0,63	0,69	0,70	0,70
66	IHCL	0,96	0,82	0,83	0,96	0,30	0,28	0,15	0,30
67	indir. calorimetry	0,94	0,87	0,91	0,94	0,76	0,69	0,64	0,76
68	RQ	0,88	0,69	0,67	0,88	0,73	0,67	0,44	0,73
69	sex	0,98	0,94	0,93	0,98	0,99	0,97	0,98	0,99
70	BW	0,92	0,77	0,74	0,92	0,70	0,75	0,73	0,75
71	BMI	0,92	0,69	0,66	0,92	0,72	0,74	0,61	0,74
72	HC	0,92	0,80	0,76	0,92	0,72	0,69	0,53	0,72
73	WC	0,91	0,81	0,78	0,91	0,68	0,74	0,71	0,74
74	WHIPR	0,94	0,85	0,81	0,94	0,67	0,65	0,58	0,67
75	HR	0,95	0,78	0,77	0,95	0,58	0,54	0,52	0,58
76	Firmicutes	0,95	0,92	0,89	0,95	0,52	0,55	0,49	0,55
77	Bifidobacterium	0,93	0,68	0,51	0,93	0,43	0,37	0,39	0,43
	Mean	0,94	0,80	0,77	0,94	0,62	0,63	0,57	0,65

Prediction of variables using Random Forest-based (RF) regression model, the Elastic Net (ER) regression model or the Liner Regression (LR) model and calculating the “best of” (BO) three methods. Summary of the transversal prediction analysis in the “establishment group” (EST) using 77 variables starting from the PBMC transcriptome using the Random Forest-based (RF) regression model, the Elastic Net (ER) regression model or the Liner Regression (LR) model. A further prediction in a “confirmation group” prediction (CON) was performed for the same 77 variables using the same methodology in the same group of volunteers after intervention using the 72 PBMC transcriptomic data of these volunteers. For each variable the following is reported: the number of individual transcriptomics included in the datasets, the number of mRNAs the model is based on, the Pearson correlation coefficient (R) between observed and predicted values, the R-squared (R²) of the model, and the experimental method used to determine the variables. *For abbreviations check table 1.*

Figure legends

Figure 1. Comparison of observed and predicted values of dependent variables. For eight regression models the observed values of the dependent variable is compared to the model predicted values. Models were developed from 72 PBMC original transcriptomics using Random Forest for regression. The regression line is calculated from the observed and predicted values from the training data. The 95% confidence interval of the prediction model is indicated by the dashed lines. The dots represent the observed and predicted values of the measurements before intervention (black) and after intervention (grey). The correlation coefficient R and the percentage of measurements with the 95% confidence interval (CI) is provided. The following dependent variables are reported: (A) general healthy food; (B) general non-healthy food; (C) serum level of 9-*cis*-13,14-dihydroretinoic acid (9CDHRA); (D) serum level of lutein; (E) SAT levels; (F) Firmicutes numbers.

Figure 2. Comparison of the performance of regression methods. For the variable ‘general healthy food’ and ‘general non-healthy food’ regression models were generated using Random Forest (A and B), Elastic Net (C and D), and a standard linear regression method (E and F). Models were developed from 72 PBMC transcriptomic data. The regression line is calculated from the observed and predicted values from the training data. The 95% confidence interval (CI) of the prediction model is indicated by the dashed lines. The dots represent the observed and predicted values of the measurements before intervention (black) and after intervention (GREY). The correlation coefficient R and the percentage of measurements with the 95% CI is provided.

Figure 3. Summarizing figure of the establishment transversal-prediction analysis (on the left side) ranging from the Random Forest, Elastic Net and Linear Regression model prediction analyses as well as “best of” (best) summarising prediction models in addition to confirmation predictions (at the right side) based on the Random Forest, Elastic Net and Linear Regression model prediction analysis as well as “best of” (best) summarising prediction model.

Figure 4. Schematic model of our transversal-predictor analysis

Figure 1.




Figure 2.




Figure 3.





Figure 4.



References

- Agliata, Antonio, Deborah Giordano, Francesco Bardozzo, Salvatore Bottiglieri, Angelo Facchiano, and Roberto Tagliaferri. 2023. "Machine Learning as a Support for the Diagnosis of Type 2 Diabetes." *International Journal of Molecular Sciences* 24 (7): 6775. <https://doi.org/10.3390/ijms24076775>.
- Aladhab, Rawnaq Adil, and Majid Hameed Alabood. 2019. "Adherence of Patients with Diabetes to a Lifestyle Advice and Management Plan in Basra, Southern Iraq." *International Journal of Diabetes and Metabolism* 25 (3–4): 100–105. <https://doi.org/10.1159/000500915>.
- Altobelli, Emma, Paolo Matteo Angeletti, Valerio F. Profeta, and Reimondo Petrocelli. 2020. "Lifestyle Risk Factors for Type 2 Diabetes Mellitus and National Diabetes Care Systems in European Countries." *Nutrients* 12 (9): 2806. <https://doi.org/10.3390/nu12092806>.
- Aronoff, Stephen L., Kathy Berkowitz, Barb Shreiner, and Laura Want. 2004. "Glucose Metabolism and Regulation: Beyond Insulin and Glucagon." *Diabetes Spectrum* 17 (3): 183–90. <https://doi.org/10.2337/diaspect.17.3.183>.
- Baine, Michael J., Subhankar Chakraborty, Lynette M. Smith, Kavita Mallya, Aaron R. Sasson, Randall E. Brand, and Surinder K. Batra. 2011. "Transcriptional Profiling of Peripheral Blood Mononuclear Cells in Pancreatic Cancer Patients Identifies Novel Genes with Potential Diagnostic Utility." Edited by Ludovic Tailleux. *PLoS ONE* 6 (2): e17014. <https://doi.org/10.1371/journal.pone.0017014>.
- Bories, Gael, Robert Caiazzo, Bruno Derudas, Corinne Copin, Violeta Raverdy, Marie Pigeyre, Francois Pattou, Bart Staels, and Giulia Chinetti-Gbaguidi. 2012. "Impaired Alternative Macrophage Differentiation of Peripheral Blood Mononuclear Cells from Obese Subjects." *Diabetes & Vascular Disease Research* 9 (3): 189–95. <https://doi.org/10.1177/1479164111430242>.
- Bouwens, Mark, Lydia A. Afman, and Michael Müller. 2007. "Fasting Induces Changes in Peripheral Blood Mononuclear Cell Gene Expression Profiles Related to Increases in Fatty Acid Beta-Oxidation: Functional Role of Peroxisome Proliferator Activated Receptor Alpha in Human Peripheral Blood Mononuclear Cells." *The American Journal of Clinical Nutrition* 86 (5): 1515–23. <https://doi.org/10.1093/ajcn/86.5.1515>.
- Burczynski, Michael E., and Andrew J. Dorner. 2006. "Transcriptional Profiling of Peripheral Blood Cells in Clinical Pharmacogenomic Studies." *Pharmacogenomics* 7 (2): 187–202. <https://doi.org/10.2217/14622416.7.2.187>.
- Bussel, Inge P. G. van, Parastoo Fazelzadeh, Gary S. Frost, Milena Rundle, and Lydia A. Afman. 2019. "Measuring Phenotypic Flexibility by Transcriptome Time-Course Analyses during Challenge Tests before and after Energy Restriction." *FASEB Journal: Official Publication of the Federation of American Societies for Experimental Biology* 33 (9): 10280–90. <https://doi.org/10.1096/fj.201900148R>.
- Caimari, Antoni, Paula Oliver, Jaap Keijer, and Andreu Palou. 2010. "Peripheral Blood Mononuclear Cells as a Model to Study the Response of Energy Homeostasis-Related Genes to Acute Changes in Feeding Conditions." *Omics: A Journal of Integrative Biology* 14 (2): 129–41. <https://doi.org/10.1089/omi.2009.0092>.
- Chang, David H., John R. Rutledge, Ankur A. Patel, Barbara G. Heerdt, Leonard H. Augenlicht, and Robert J. Korst. 2013. "The Effect of Lung Cancer on Cytokine Expression in Peripheral Blood Mononuclear Cells." *PloS One* 8 (6): e64456. <https://doi.org/10.1371/journal.pone.0064456>.
- Cho, N. H., J. E. Shaw, S. Karuranga, Y. Huang, J. D. da Rocha Fernandes, A. W. Ohlrogge, and B. Malanda. 2018. "IDF Diabetes Atlas: Global Estimates of Diabetes Prevalence for 2017 and Projections for 2045." *Diabetes Research and Clinical Practice* 138 (April): 271–81. <https://doi.org/10.1016/j.diabres.2018.02.023>.
- Costa, Andrea, Bàrbara Reynés, Jadwiga Konieczna, Marian Martín, Miquel Fiol, Andreu Palou, Dora Romaguera, and Paula Oliver. 2021. "Use of Human PBMC to Analyse the Impact of Obesity on Lipid Metabolism and Metabolic Status: A Proof-of-Concept Pilot Study." *Scientific Reports* 11 (1): 18329. <https://doi.org/10.1038/s41598-021-96981-6>.
- Elliott, R. M., B. de Roos, S. J. Duthie, F. G. Bouwman, I. Rubio-Aliaga, L. K. Crosley, C. Mayer, et al. 2014. "Transcriptome Analysis of Peripheral Blood Mononuclear Cells in Human Subjects Following a 36 h

- Fast Provides Evidence of Effects on Genes Regulating Inflammation, Apoptosis and Energy Metabolism." *Genes & Nutrition* 9 (6): 432. <https://doi.org/10.1007/s12263-014-0432-4>.
- Farag, Youssef M.K., and Mahmoud R. Gaballa. 2011. "Diabetes: An Overview of a Rising Epidemic." *Nephrology Dialysis Transplantation* 26 (1): 28–35. <https://doi.org/10.1093/ndt/gfq576>.
- Fiamoncini, Jarlei, Milena Rundle, Helena Gibbons, E. Louise Thomas, Kerstin Geillinger-Kästle, Diana Bunzel, Jean-Pierre Trezzi, et al. 2018. "Plasma Metabolome Analysis Identifies Distinct Human Metabotypes in the Postprandial State with Different Susceptibility to Weight Loss-Mediated Metabolic Improvements." *FASEB Journal: Official Publication of the Federation of American Societies for Experimental Biology* 32 (10): 5447–58. <https://doi.org/10.1096/fj.201800330R>.
- Friedman, Jerome H., Trevor Hastie, and Rob Tibshirani. 2010. "Regularization Paths for Generalized Linear Models via Coordinate Descent." *Journal of Statistical Software* 33 (February):1–22. <https://doi.org/10.18637/jss.v033.i01>.
- Garedow, Aster Wakjira, Tsiyon Mekoya Jemaneh, Addisalem Gebresilase Hailemariam, and Gorfineh Teshome Tesfaye. 2023. "Lifestyle Modification and Medication Use among Diabetes Mellitus Patients Attending Jimma University Medical Center, Jimma Zone, South West Ethiopia." *Scientific Reports* 13 (1): 4956. <https://doi.org/10.1038/s41598-023-32145-y>.
- Goyal, Shiwali, Jyoti Rani, Mohd Akbar Bhat, and Vanita Vanita. 2023. "Genetics of Diabetes." *World Journal of Diabetes* 14 (6): 656–79. <https://doi.org/10.4239/wjd.v14.i6.656>.
- Hebbali, Aravind. 2024. "Olsrr: Tools for Building OLS Regression Models." <https://cran.r-project.org/web/packages/olsrr/index.html>.
- Heikes, Kenneth E., David M. Eddy, Bhakti Arondekar, and Leonard Schlessinger. 2008. "Diabetes Risk Calculator: A Simple Tool for Detecting Undiagnosed Diabetes and Pre-Diabetes." *Diabetes Care* 31 (5): 1040–45. <https://doi.org/10.2337/dc07-1150>.
- Jangili, Shraddha, Hariprasad Vavilala, Gopi Sumanth Bhaskar Boddeda, Suryanaryana Murty Upadhyayula, Ramu Adela, and Srinivasa Rao Mutheneni. 2023. "Machine Learning-Driven Early Biomarker Prediction for Type 2 Diabetes Mellitus Associated Coronary Artery Diseases." *Clinical Epidemiology and Global Health* 24 (November). <https://doi.org/10.1016/j.cegh.2023.101433>.
- Kaul, Nabodita, and Sher Ali. 2016. "Genes, Genetics, and Environment in Type 2 Diabetes: Implication in Personalized Medicine." *DNA and Cell Biology* 35 (1): 1–12. <https://doi.org/10.1089/dna.2015.2883>.
- Knowler, William C., Elizabeth Barrett-Connor, Sarah E. Fowler, Richard F. Hamman, John M. Lachin, Elizabeth A. Walker, David M. Nathan, and Diabetes Prevention Program Research Group. 2002. "Reduction in the Incidence of Type 2 Diabetes with Lifestyle Intervention or Metformin." *The New England Journal of Medicine* 346 (6): 393–403. <https://doi.org/10.1056/NEJMoa012512>.
- Konieczna, Jadwiga, Juana Sánchez, Mariona Palou, Catalina Picó, and Andreu Palou. 2015. "Blood Cell Transcriptomic-Based Early Biomarkers of Adverse Programming Effects of Gestational Calorie Restriction and Their Reversibility by Leptin Supplementation." *Scientific Reports* 5 (1): 9088. <https://doi.org/10.1038/srep09088>.
- Kuhn, Max. 2008. "Building Predictive Models in R Using the Caret Package." *Journal of Statistical Software* 28 (November):1–26. <https://doi.org/10.18637/jss.v028.i05>.
- Lai, Hang, Huaxiong Huang, Karim Keshavjee, Aziz Guergachi, and Xin Gao. 2019. "Predictive Models for Diabetes Mellitus Using Machine Learning Techniques." *BMC Endocrine Disorders* 19 (1): 101. <https://doi.org/10.1186/s12902-019-0436-6>.
- Liaw, Andy, and Matthew Wiener. 2002. "Classification and Regression by randomForest." *R News* 2 (3): 18–22.
- Liew, Choong-Chin, Jun Ma, Hong-Chang Tang, Run Zheng, and Adam A. Dempsey. 2006. "The Peripheral Blood Transcriptome Dynamically Reflects System Wide Biology: A Potential Diagnostic Tool." *The Journal of Laboratory and Clinical Medicine* 147 (3): 126–32. <https://doi.org/10.1016/j.lab.2005.10.005>.
- Manoel-Caetano, Fernanda S., Danilo J. Xavier, Adriane F. Evangelista, Paula Takahashi, Cristhianna V. Collares, Denis Puthier, Maria C. Foss-Freitas, et al. 2012. "Gene Expression Profiles Displayed by Peripheral Blood Mononuclear Cells from Patients with Type 2 Diabetes Mellitus Focusing on Biological Processes Implicated on the Pathogenesis of the Disease." *Gene* 511 (2): 151–60. <https://doi.org/10.1016/j.gene.2012.09.090>.

- Mello, Vanessa Derenji Ferreira de, Marjukka Kolehmanien, Ursula Schwab, Leena Pulkkinen, and Matti Uusitupa. 2012. "Gene Expression of Peripheral Blood Mononuclear Cells as a Tool in Dietary Intervention Studies: What Do We Know so Far?" *Molecular Nutrition & Food Research* 56 (7): 1160–72. <https://doi.org/10.1002/mnfr.201100685>.
- Mozaffarian, Dariush, Aruna Kamineni, Mercedes Carnethon, Luc Djoussé, Kenneth J. Mukamal, and David Siscovick. 2009. "Lifestyle Risk Factors and New-Onset Diabetes Mellitus in Older Adults: The Cardiovascular Health Study." *Archives of Internal Medicine* 169 (8): 798–807. <https://doi.org/10.1001/archinternmed.2009.21>.
- Naz, Huma, and Sachin Ahuja. 2020. "Deep Learning Approach for Diabetes Prediction Using PIMA Indian Dataset." *Journal of Diabetes and Metabolic Disorders* 19 (1): 391–403. <https://doi.org/10.1007/s40200-020-00520-5>.
- Nguyen, Binh P., Hung N. Pham, Hop Tran, Nhung Nghiem, Quang H. Nguyen, Trang T. T. Do, Cao Truong Tran, and Colin R. Simpson. 2019. "Predicting the Onset of Type 2 Diabetes Using Wide and Deep Learning with Electronic Health Records." *Computer Methods and Programs in Biomedicine* 182 (December):105055. <https://doi.org/10.1016/j.cmpb.2019.105055>.
- Oikonomou, Evangelos K., and Rohan Khara. 2023. "Machine Learning in Precision Diabetes Care and Cardiovascular Risk Prediction." *Cardiovascular Diabetology* 22 (1): 259. <https://doi.org/10.1186/s12933-023-01985-3>.
- Okada, Akira, Yohei Hashimoto, Tadahiro Goto, Satoko Yamaguchi, Sachiko Ono, Kayo Ikeda Kurakawa, Masaomi Nangaku, Toshimasa Yamauchi, Hideo Yasunaga, and Takashi Kadowaki. 2022. "A Machine Learning-Based Predictive Model to Identify Patients Who Failed to Attend a Follow-up Visit for Diabetes Care After Recommendations From a National Screening Program." *Diabetes Care* 45 (6): 1346–54. <https://doi.org/10.2337/dc21-1841>.
- Oliver, Paula, Bàrbara Reynés, Antoni Caimari, and Andreu Palou. 2013. "Peripheral Blood Mononuclear Cells: A Potential Source of Homeostatic Imbalance Markers Associated with Obesity Development." *Pflugers Archiv: European Journal of Physiology* 465 (4): 459–68. <https://doi.org/10.1007/s00424-013-1246-8>.
- Razavian, Narges, Saul Blecker, Ann Marie Schmidt, Aaron Smith-McLallen, Somesh Nigam, and David Sontag. 2015. "Population-Level Prediction of Type 2 Diabetes From Claims Data and Analysis of Risk Factors." *Big Data* 3 (4): 277–87. <https://doi.org/10.1089/big.2015.0020>.
- Reynés, Bàrbara, Rubén Díaz-Rúa, Margalida Cifre, Paula Oliver, and Andreu Palou. 2015. "Peripheral Blood Mononuclear Cells as a Potential Source of Biomarkers to Test the Efficacy of Weight-Loss Strategies." *Obesity (Silver Spring, Md.)* 23 (1): 28–31. <https://doi.org/10.1002/oby.20918>.
- Rodney A. Rhoades PhD. 2017. *Medical Physiology: Principles for Clinical Medicine*. 5th edition. Philadelphia: LWW.
- Rundle, Milena, Jarlei Fiamoncini, E Louise Thomas, Suzan Wopereis, Lydia A. Afman, Lorraine Brennan, Christian A. Drevon, et al. 2023. "Diet-Induced Weight Loss and Phenotypic Flexibility Among Healthy Overweight Adults: A Randomized Trial." *The American Journal of Clinical Nutrition* 118 (3): 591–604. <https://doi.org/10.1016/j.ajcnut.2023.07.002>.
- Schmidt, Maria Inês, and Bruce Bartholow Duncan. 2003. "Diabesity: An Inflammatory Metabolic Condition" 41 (9): 1120–30. <https://doi.org/10.1515/CCLM.2003.174>.
- Schmidt, Sabrina K., Liv Hemmestad, Christopher S. MacDonald, Henning Langberg, and Laura S. Valentiner. 2020. "Motivation and Barriers to Maintaining Lifestyle Changes in Patients with Type 2 Diabetes after an Intensive Lifestyle Intervention (The U-TURN Trial): A Longitudinal Qualitative Study." *International Journal of Environmental Research and Public Health* 17 (20): 7454. <https://doi.org/10.3390/ijerph17207454>.
- Schwarz, P. E. H., J. Li, J. Lindstrom, and J. Tuomilehto. 2009. "Tools for Predicting the Risk of Type 2 Diabetes in Daily Practice." *Hormone and Metabolic Research = Hormon- Und Stoffwechselforschung = Hormones Et Metabolisme* 41 (2): 86–97. <https://doi.org/10.1055/s-0028-1087203>.
- Sun, Xue, Weihui Yu, and Cheng Hu. 2014. "Genetics of Type 2 Diabetes: Insights into the Pathogenesis and Its Clinical Application." *BioMed Research International* 2014:926713. <https://doi.org/10.1155/2014/926713>.

- Tonyan, Ziravard N., Yulia A. Nasykhova, Maria M. Danilova, Yury A. Barbitoff, Anton I. Chagalidi, Anastasiia A. Mikhailova, and Andrey S. Glotov. 2022. "Overview of Transcriptomic Research on Type 2 Diabetes: Challenges and Perspectives." *Genes* 13 (7): 1176. <https://doi.org/10.3390/genes13071176>.
- Wee, Boon Feng, Saaveethya Sivakumar, King Hann Lim, W. K. Wong, and Filbert H. Juwono. 2024. "Diabetes Detection Based on Machine Learning and Deep Learning Approaches." *Multimedia Tools and Applications* 83 (8): 24153–85. <https://doi.org/10.1007/s11042-023-16407-5>.
- Wickham, Hadley. 2016. *Ggplot2*. Use R! Cham: Springer International Publishing. <https://doi.org/10.1007/978-3-319-24277-4>.
- Witka, Beska Z., Dede J. Oktaviani, Marcellino Marcellino, Melisa I. Barliana, and Rizky Abdulah. 2019. "Type 2 Diabetes-Associated Genetic Polymorphisms as Potential Disease Predictors." *Diabetes, Metabolic Syndrome and Obesity* 12 (December):2689–2706. <https://doi.org/10.2147/DMSO.S230061>.
- Yu, Wei, Tiebin Liu, Rodolfo Valdez, Marta Gwinn, and Muin J. Khoury. 2010. "Application of Support Vector Machine Modeling for Prediction of Common Diseases: The Case of Diabetes and Pre-Diabetes." *BMC Medical Informatics and Decision Making* 10 (March):16. <https://doi.org/10.1186/1472-6947-10-16>.
- Zhang, Yanbo, Xiong-Fei Pan, Junxiang Chen, Lu Xia, Anlan Cao, Yuge Zhang, Jing Wang, et al. 2020. "Combined Lifestyle Factors and Risk of Incident Type 2 Diabetes and Prognosis among Individuals with Type 2 Diabetes: A Systematic Review and Meta-Analysis of Prospective Cohort Studies." *Diabetologia* 63 (1): 21–33. <https://doi.org/10.1007/s00125-019-04985-9>.
- Zou, Quan, Kaiyang Qu, Yamei Luo, Dehui Yin, Ying Ju, and Hua Tang. 2018. "Predicting Diabetes Mellitus With Machine Learning Techniques." *Frontiers in Genetics* 9:515. <https://doi.org/10.3389/fgene.2018.00515>.